

University of Groningen

WNT5A/5B signaling represses functional responses in lung epithelial progenitors

Wu, Xinhui; Bos, Isabella; Kistemaker, Loes; Gosens, Reinoud

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

2018

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Wu, X., Bos, I., Kistemaker, L., & Gosens, R. (2018). *WNT5A/5B signaling represses functional responses in lung epithelial progenitors*. Abstract from ATS 2018, San Diego, California, United States.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

WNT5a/5b Signaling Represses Functional Responses in Lung Epithelial Progenitors

X. Wu¹, E. M. Dijk², J. Ng-Blichfeldt¹, S. Bos³, L. E. M. Kistemaker¹, R. Gosens⁴; ¹Molecular Pharmacology, University of Groningen, Groningen, Netherlands, ²Molecular Pharmacology, Rijksuniversiteit Groningen, Groningen, Netherlands, ³University of Groningen, Groningen, Netherlands, ⁴Univ of Groningen, Groningen, Netherlands.

Corresponding author's email: x.wu@rug.nl

Background: Chronic obstructive pulmonary disease (COPD) represents a worldwide concern with high morbidity and mortality, and is believed to be driven in part by accelerated ageing of the lung. Although several studies indicate a role for non-canonical WNT signaling, in particular of WNT-5A and WNT-5B, in ageing and COPD, the precise impact of non-canonical WNT signaling on lung repair remains poorly understood. We hypothesized that WNT-5A/5B might impact on alveolar epithelial repair. **Methods:** Lungs were harvested for precision-cut lung slices (PCLS) or the generation of lung organoids. PCLS were treated with recombinant WNT-5A or WNT-5B (500 ng/mL) or vehicle control. Lung organoids recapitulate various features of the lung providing an in vitro model system for studying regenerative mechanisms of adult lung epithelium. Lung organoids were established by co-culturing epithelial cells (EpCAM⁺/CD45⁻/CD31⁻) with CCL206 fibroblasts in Matrigel. **Results:** The gene expression levels of the non-canonical ligands WNT-5A and WNT-5B were increased in whole lung tissue from aged WT mice and their expression correlated with the senescence marker p16. In PCLS derived from young WT mice, treatment with WNT-5A had no effect on gene expression of the type I and type II epithelial cell markers Aqp5 and Sftpc; interestingly, WNT-5B treatment significantly decreased gene expression of Aqp5 (mean difference = 36.35% ± 10.93%, compared to vehicle control, p < 0.05,) and Sftpc (mean difference = 46.25% ± 5.76%, compared to vehicle control, p < 0.01). In addition, WNT-5B significantly decreased expression of the canonical WNT target gene, Axin2 (mean difference = 83.13% ± 9.65%, compared to vehicle control, p < 0.05). The number of organoids visible 7 days after culture was significantly decreased by WNT-5A and WNT-5B (p < 0.05). WNT-5A significantly decreased the number of airway type organoids (p < 0.01), whereas WNT-5B selectively repressed alveolar type organoid formation (p < 0.05). Immunofluorescence studies confirmed that numbers of acetylated- α tubulin⁺ organoids were significantly decreased by both WNT-5A and WNT-5B, however, numbers of SPC⁺ organoids were significantly decreased by WNT-5B (p < 0.05). Neither WNT-5A nor WNT-5B stimulation affected the size of lung organoids measured on day -14. **Conclusions:** Non-canonical WNT signaling is correlated with ageing. The non-canonical WNT ligands WNT-5A and WNT-5B repress gene expression of alveolar epithelial cell markers in lung slices, and functionally inhibit lung organoid formation. We speculate that such a mechanism may contribute to defective alveolar repair in ageing and COPD.

This abstract is funded by: CSC

Am J Respir Crit Care Med 2018;197:A7624
Internet address: www.atsjournals.org

Online Abstracts Issue