

## University of Groningen

### Is the choice of walking speed in people with a lower limb amputation dependent on the relative aerobic load?

Wezenberg, Daphne; Slootman, Hans J.; Faber, Willemein; van der Woude, L. H. V.; de Haan, Arnold; Houdijk, Han

*Published in:*  
Revalidata

**IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.**

*Document Version*  
Publisher's PDF, also known as Version of record

*Publication date:*  
2010

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Wezenberg, D., Slootman, H. J., Faber, W., van der Woude, L. H. V., de Haan, A., & Houdijk, H. (2010). Is the choice of walking speed in people with a lower limb amputation dependent on the relative aerobic load? *Revalidata*, 154, 23.

#### Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

#### Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

# Is the choice of walking speed in people with a lower limb amputation dependent on the relative aerobic load?

D. Wezenberg, J.R. Sloopman, W.X.M. Faber, L.H.V. van der Woude, A. de Haan, H. Houdijk

Although increasing the aerobic capacity is a widely accepted goal in the rehabilitation of people after lower limb amputation, as yet it is unclear to which extent aerobic capacity plays an important role in regaining walking ability. This study investigated the relative aerobic load of people after lower limb amputation during walking at different speeds. Eleven trans-tibial and five trans-femoral traumatic amputees and twenty-one weight and aged matched controls participated in this research (age  $61 \pm 6.3$  years). Peak oxygen consumption was measured while

walking at preferred walking speed (PWS) and at speeds 15 and 30% higher and while performing a discontinuous graded one-legged peak cycling test. The fraction of the peak oxygen consumption needed to walk is referred to as 'relative aerobic load'. Relative aerobic load at PWS did not differ between groups ( $p=0.814$ ). The PWS was, however, significantly lower in both patient groups compared to controls ( $p<0.001$ ). When the speed of the patient groups was matched with the PWS of the control group the relative aerobic load was significantly

higher in the patients compared to controls ( $p=0.047$ ). It was concluded that patients after lower limb amputation choose a lower PWS corresponding to a relative aerobic load similar to able-bodied controls. Research with healthy subjects has postulated that the PWS is the speed requiring the least oxygen per distance walked. However, preliminary data will be presented suggesting that this might not be the case, considering this speed might entail a relative aerobic load above the acceptable limit in patients.

---

*Drs. D. Wezenberg, Heliomare R&D, Wijk aan Zee*

*Drs. J.R. Sloopman, Heliomare rehabilitation, Wijk aan Zee*

*Drs. W.X.M. Faber, Heliomare rehabilitation, Wijk aan Zee*

*Prof. Dr. L.H.V. van der Woude, Center for Human Movement Sciences, University Medical Center Groningen*

*Prof. Dr. A. de Haan, Research Institute MOVE, Faculty of Human Movement Sciences, VU University, Amsterdam*

*Dr. H. Houdijk, Research Institute MOVE, Faculty of Human*

*Movement Sciences, VU University, Amsterdam & Heliomare R&D*