

University of Groningen

## Advances of treatment in atypical cartilaginous tumours

Dierselhuis, Edwin Frank

**IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.**

*Document Version*

Publisher's PDF, also known as Version of record

*Publication date:*

2019

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Dierselhuis, E. F. (2019). *Advances of treatment in atypical cartilaginous tumours*. [Thesis fully internal (DIV), University of Groningen]. Rijksuniversiteit Groningen.

### Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.


### Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

# CHAPTER I


## General introduction



## INTRODUCTION

### CHONDROSARCOMA


Sarcomas are malignant tumours of mesenchymal origin that arise in soft tissue and bone. They differ from common types of cancer, such as breast cancer, which derive from epithelial cells and are called carcinomas. Sarcomas are relatively uncommon, representing about 1% of all new cancer diagnoses in the United States.<sup>1</sup> In the Netherlands, about 150 cases of primary bone malignancies are diagnosed each year.<sup>2</sup> Chondrosarcomas (CS) are a very heterogeneous group of cartilage matrix-producing tumours. They are most commonly seen in patients aged 40-70 and are the third most common primary bone malignancy.<sup>3</sup> The majority of cases (85%) are *de novo* central tumours, inside of the bone,<sup>3</sup> yet can also arise secondarily from the surface of the bone from an osteochondroma – peripheral CS – or in the presence of Ollier disease or Maffucci syndrome. All these tumours form a wide range of malignant potential. The benign counterpart – enchondroma – is often an asymptomatic, locally non-aggressive tumour. On the other side of the spectrum, dedifferentiated CS is a very high-grade tumour with very poor survival<sup>4</sup> (Figure 1). In general, chondroid tumours are poorly vascularised and have a low percentage of dividing cells. This makes them relatively insensitive to radiation and/or chemotherapy, and the mainstay of treatment for malignant cartilaginous tumours is surgery.<sup>3,4</sup>



**FIGURE 1.** The spectrum of cartilaginous tumours, ranging from benign enchondroma to dedifferentiated chondrosarcoma

## ATYPICAL CARTILAGINOUS TUMOUR

An atypical cartilaginous tumour (ACT) is a cartilage-producing tumour located in the bone with intermediate malignant potential – also known as low-grade or grade I chondrosarcoma<sup>5</sup> (Figure 2). Most cases are incidental findings, when patients come for evaluation of other joint- or bone-related complaints. With increased usage of MRI and CT-scanning, incidence of the disease has risen in recent decades.<sup>6-8</sup> This is why it is important for orthopaedic surgeons to be aware of this entity and its possible treatment options.



**FIGURE 2.** Typical MRI image of an atypical cartilaginous tumour (ACT) in the proximal humerus, showing a large lesion and wall-to-wall filling but no signs of higher-grade aggressiveness



**FIGURE 3.** Tumour in the diaphysis of the femur, treated by intercalary resection, and reconstructed by allograft with plate and nail fixation.

ACTs tend to be only locally aggressive, although incidental metastasising has been reported in literature.<sup>9,10</sup> Historically a correct diagnosis has been notoriously difficult to make, since histology and imaging alone are not always conclusive and have shown high inter-observer variability.<sup>11</sup> In some cases the tumour evolved to a high grade after local recurrence.<sup>12</sup> This clearly has a negative impact on patient survival, and wide resection has long been advocated in order to achieve adequate surgical margins.<sup>13</sup> However, nowadays

a paradigm shift has taken place in literature towards more local (intralesional) surgery.<sup>14</sup> It may be that more aggressive tumour biology *after* local recurrence is the result of an undertreated high-grade tumour, rather than a direct consequence of the local recurrence in itself. With improved imaging modalities and a better understanding of the natural behaviour of these tumours, case series have been published that show excellent survival after intralesional surgery by curettage of the tumour.<sup>15-23</sup> Given the seemingly relatively mild nature of ACT but the potential morbidity of the current surgical strategies, one can wonder whether the cure is not worse than the disease. Minimally invasive treatment might thus be a step towards an ideal treatment regime: local control leading to excellent oncological outcome, no compromise on functional results, and ideally performed in day care. This concept was developed by our group after an apparently benign lesion treated by RFA turned out to be a cartilaginous malignancy.<sup>24</sup>

## SURGICAL TECHNIQUES AND THEIR CONSEQUENCES

As described above, different surgical techniques have been applied for treatment of ACT, every single one with its specific characteristics and advantages/disadvantages. More details are provided below for each particular technique.

### WIDE RESECTION

In this technique whole segments of bone (including joints) are removed in order to attain extensive surgical margins. This will sometimes lead to amputation if neurovascular bundles cannot be saved, reconstruction is not feasible, or soft-tissue coverage cannot be achieved. If limbs can be salvaged but joints are lost, endoprosthesis such as total knee arthroplasty (TKA) or total hip arthroplasty (THA) is needed. This often requires specially designed tumour prostheses rather than conventional arthroplasties. If segments of bone between joints are removed (intercalary resection), reconstruction is done with autologous bone (e.g. vascularised fibula) and/or allograft (Figure 3).

Wide resection, regardless of the reconstructive possibilities, often leads to functional deficiencies. Amputees will not be the only ones suffering from functional loss, as in reconstructive surgery too muscle function is often (temporarily) lost and weight bearing is prohibited for several months.<sup>25,26</sup> Finally, due to the scope of the surgery, operating time and large wound bed there is a considerable risk of postoperative infection, nerve damage, fracture and thromboembolic events.<sup>26-28</sup>


### INTRALESIONAL SURGERY (CURETTAGE)

In curettage the tumour is removed while leaving the surrounding bone virtually intact. Only a small cortical window is created to have access to the tumour. The lesion is removed using a curette, traditionally under fluoroscopy guidance (Figure 4). To improve surgical margins, several local adjuvants are available. Most common are the application of phenol ( $C_6H_5OH$ ) with ethanol washout, polymethylmethacrylate (PMMA) and the use of liquid nitrogen ( $LN_2$ ). In an *in vitro* model, cytotoxic effects were found for concentrations of 1.5% phenol and 42.5% ethanol.<sup>29</sup> Furthermore, 96% ethanol is capable of reducing phenol levels for safe washout of the cavity. PMMA is often used in orthopaedic surgery, primarily to fixate endoprostheses. It is also used to fill defects, as it enhances early weight bearing. Another possible advantage is its necrotising effect due to the exothermic chemical reaction during hardening, where temperatures over 80°C are reached. It is estimated that the surgical margins are hereby enlarged by 2 to 5 mm in cancellous bone.<sup>30</sup> Cryosurgery using liquid nitrogen is a more potent adjuvant, as 7 to 12 mm of extra bone tissue are necrotised.<sup>31</sup> However, it is not widely used as it has drawbacks like temporary nerve damage and increased risk of fracturing.<sup>32</sup>

Curettage preserves the integrity of the bone and joint dramatically compared to *en bloc* resection or amputation. As a result, functional outcome is significantly better after curettage than after resection in retrospective comparisons.<sup>25-26</sup> Nevertheless, this technique also has its drawbacks, as complications such as fracturing or infection do occur.<sup>26-28</sup> In addition, hospital admission is needed and extremities are often protected from weight bearing for several weeks to months.




**FIGURE 4.** Postoperative image of an ACT treated by curettage and PMMA filling of the defect.



**FIGURE 5.** Working field of an RFA needle<sup>35</sup>.

### RADIOFREQUENCY ABLATION (RFA)

In radiofrequency ablation (RFA) a high-frequency alternating current heats tissue to approximately 80°C.<sup>33</sup> An electromagnetic field is created which results in vibration of molecules, leading to heat due to friction in about a 3-cm bony zone<sup>34</sup> (Figure 5). As temperatures rise above 46°C coagulation necrosis takes place, with almost instantaneous cell death at 60°C and beyond.<sup>36</sup> RFA can be applied percutaneously under computed tomography (CT) guidance under general or spinal anaesthesia (Figure 6). The technique was originally developed successfully for solid organ tumours such as hepatocellular carcinoma.<sup>37-38</sup> Over two decades ago it was also introduced in orthopaedics and has become the gold standard for osteoid osteoma, with primary and secondary success rates of 79-96% and 97-100% respectively.<sup>39-42</sup> It has proven to be a precise, safe and relatively inexpensive treatment tool for other bony lesions as well, such as chondroblastoma and metastases.<sup>43-46</sup> Major advantages are that it allows early weight bearing and can be performed in day care. This technique also has potential complications, mainly burning of the skin or fracturing.<sup>24,47</sup>



**FIGURE 6.** Per-operative image of CT guided RFA procedure of an ACT in the distal femur

## AIM AND OVERVIEW OF THE THESIS

This thesis aims to analyse in two parts the results of current practice of ACT treatment and to investigate new treatment modalities.

### PART I

To date, no prospective studies have been published that could help design an adequate treatment algorithm for ACT in the long bones. There are only low-evidence retrospective studies available, with widespread publication dates, surgical indications and applied techniques. Hence in the absence of high-level evidence we first aim to review all available literature and meta-analysis data in a Cochrane Review (**Chapter II**). Next, we will analyse our own experience of treating ACT by intralesional surgery (**Chapter III**). As computer-assisted surgery (CAS) has also been introduced in the field of oncologic orthopaedics, we will also evaluate its value compared to fluoroscopy in the treatment of ACT (**Chapter IV**). CAS offers the surgeon real-time feedback on its whereabouts during surgery, potentially decreasing residual tumour rates and enhancing disease-free survival. Moreover, patients as well as surgeons are protected from X-rays during surgery.



## PART II

Minimally invasive treatment using RFA is studied in part II of the thesis. In **Chapters V and VI** we provide insight into the efficacy of RFA in the treatment of ACT, together with evaluation of imaging modalities for follow-up. In **Chapter V** we also investigate functional outcome by means of musculoskeletal tumour society (MSTS) scores after RFA compared to intralesional surgery. In **Chapter VI** we analyse the learning curve of applying this new technique. A general discussion and the implications of our studies are provided in **Chapter VII**, and the thesis is summarised in **Chapter VIII**.

---

## REFERENCES

---

1. Borden EC, Baker LH, Bell RS, et al. Soft tissue sarcomas of adults: state of the translational science. *Clin. Cancer Res.* 9 (6): 1941–56 <https://www.oncoline.nl/beentumoren>
2. Gelderblom H, Hogendoorn PC, Dijkstra SD, van Rijswijk CS, Krol AD, Taminiau AH, et al. The clinical approach towards chondrosarcoma. *Oncologist* 2008 Mar;13(3):320-9.
3. Nota SP, Braun Y, Schwab JH, van Dijk CN, Bramer JA. The Identification of Prognostic Factors and Survival Statistics of Conventional Central Chondrosarcoma. *Sarcoma* 2015:623746
4. Hogendoorn P. B., Bovee J. M., Nielsen G. P. Chondrosarcoma (grades I-III), including primary and secondary variants and periosteal chondrosarcoma. In: Fletcher C. D. M., Bridge J. A., Hogendoorn P. C. W., Mertens F., editors. *World Health Organization Classification of Tumours of Soft Tissue and Bone*. Vol. 5. Lyon, France: IARC; 2013. p. p. 264.
5. Hong ED, Carrino JA, Weber KL, Fayad LM. Prevalence of shoulder enchondromas on routine MR imaging. *Clin Imaging.* 2011 Sep-Oct;35(5):378
6. Kransdorf MJ, Peterson JJ, Bancroft LW. MR imaging of the knee: incidental osseous lesions. *Radiol Clin North Am.* 2007 Nov;45(6):943-5
7. van Praag VM, Rueten-Budde AJ, Dijkstra PDS, Study group, van de Sande MAJ. Bone and Soft tissue tumours (WeBot) Incidence, outcomes and prognostic factors during 25 years of treatment of chondrosarcomas. *Surgical Oncology* 27 (2018) 402-8
8. Leerapun T, Hugate RR, Inwards CY, Scully SP, Sim FH. Surgical management of conventional grade I chondrosarcoma of long bones. *Clin Orthop Relat Res* 2007 Oct;463:166-72
9. Gunay C, Atalar H, Hapa O, Basarir K, Yildiz Y, Saglik Y. Surgical management of grade I chondrosarcoma of the long bones. *Acta Orthop Belg.* 2013 Jun;79(3):331-7
10. Skeletal Lesions Interobserver Correlation among Expert Diagnosticians (SLICED) Study Group. Reliability of histopathologic and radiologic grading of cartilaginous neoplasms in long bones. *J Bone Joint Surg Am.* 2007 Oct;89(10):2113-23.
11. Schwab JH, Wenger D, Unni K, Sim FH. Does local recurrence impact survival in low-grade chondrosarcoma of the long bones? *Clin Orthop Relat Res.* 2007 Sep;462:175-80.
12. Fiorenza F, Abudu A, Grimer RJ, Carter SR, Tillman RM, Ayoub K, Mangham DC, Davies AM. Risk factors for survival and local control in chondrosarcoma of bone. *J Bone Joint Surg Br.* 2002 Jan;84(1):93-9.

13. Hickey M, Farrokhyar F, Deheshi B, Turcotte R, Ghert M. A systematic review and meta-analysis of intralesional versus wide resection for intramedullary grade I chondrosarcoma of the extremities. *Ann Surg Oncol*. 2011 Jun;18(6):1705-9.
14. van der Geest IC, de Valk MH, de Rooy JW, Pruszczynski M, Veth RP, Schreuder HW. Oncological and functional results of cryosurgical therapy of enchondromas and chondrosarcomas grade 1. *J Surg Oncol* 2008 Nov 1;98(6):421-6.
15. Hanna SA, Whittingham-Jones P, Sewell MD, Pollock RC, Skinner JA, Saifuddin A, et al. Outcome of intralesional curettage for low-grade chondrosarcoma of long bones. *Eur J Surg Oncol* 2009 Dec;35(12):1343-7.
16. Kim W, Han I, Kim EJ, Kang S, Kim H. Outcomes of curettage and anhydrous alcohol adjuvant for low-grade chondrosarcoma of long bone. *Surgical Oncology* 2015;24:89-94.
17. Kim W, Lee JS, Chung HW. Outcomes after extensive manual curettage and limited burring for atypical cartilaginous tumour of long bone. *Bone Joint J* 2018 Feb;100-B(2):256-261.
18. Meftah M, Schult P, Henshaw RM. Long-term results of intralesional curettage and cryosurgery for treatment of low-grade chondrosarcoma. *J Bone Joint Surg Am*. 2013 Aug 7;95(15):1358-64.
19. Mermerkaya MU, Bekmez S, Karaaslan F, Danisman M, Kosemehmetoglu K, Gedikoglu G, Ayvaz M, Tokgozoglu AM. Intralesional curettage and cementation for low-grade chondrosarcoma of long bones: retrospective study and literature review. *World Journal of Surgical Oncology* 2014;12:336-41.
20. Mohler DG, Chiu R, McCall DA, Avedian RS. Curettage and Cryosurgery for Low-grade Cartilage Tumors Is Associated with Low Recurrence and High Function Is Associated with Low Recurrence and High Function. *Clin Orthop Relat Res* 2010;(468):2765-73.
21. Souna BS, Belot N, Duval H, Langlais F, Thomazeau H. No recurrences in selected patients after curettage with cryotherapy for grade I chondrosarcomas. *Clin Orthop Relat Res*. 2010 Jul;468(7):1956-62.
22. Verdegaal SH, Brouwers HF, van Zwet EW, Hogendoorn PC, Taminiau AH. Low-grade chondrosarcoma of long bones treated with intralesional curettage followed by application of phenol, ethanol, and bone-grafting. *J Bone Joint Surg Am* 2012 Jul 3;94(13):1201-1207.
23. Dierselhuis EF, Jutte PC, van der Eerden PJ, Suurmeijer AJ, Bulstra SK. Hip fracture after radiofrequency ablation therapy for bone tumors: two case reports. *Skeletal Radiol* 2010;39:1139-1143.

24. Donati D, Colangeli S, Colangeli M, Di Bella C, Bertoni F. Surgical treatment of grade I central chondrosarcoma. *Clin Orthop Relat Res.* 2010 Feb;468(2):581-9
25. Aarons C, Potter BK, Adams SC, Pitcher JD Jr, Temple HT. Extended intralesional treatment versus resection of low-grade chondrosarcomas. *Clin Orthop Relat Res.* 2009 Aug;467(8):2105-11.
26. Campanacci DA, Scoccianti G, Franchi A et al. Surgical treatment of central grade 1 chondrosarcoma of the appendicular skeleton. *J Orthopaed Traumatol* 2013;14:101-107.
27. Etchebehere M, de Camargo OP, Croci AT, et al. Relationship between surgical procedure and outcome for patients with grade I chondrosarcomas. *Clinics (Sao Paulo)* 2005;60:121-6.
28. Verdegaal SH, Corver WE, Hogendoorn PC, Taminiau AH. The cytotoxic effect of phenol and ethanol on the chondrosarcoma-derived cell line OUMS-27: an in vitro experiment. *J Bone Joint Surg Br.* 2008 Nov;90(11):1528-32.
29. Malawer MM, Marks MR, McChesney D, et al. The effect of cryosurgery and polymethylmethacrylate in dogs with experimental bone defects comparable to tumor defects. *Clin Orthop* 1988;226:299-310
30. Marcove RC, Stovell PB, Huvos AG, Bullough PG. The use of cryosurgery in the treatment of low and medium grade chondrosarcoma: a preliminary report. *Clin Orthop* 1977;122:147-56
31. Schreuder HWB, Keijser LC, Veth RPH. Beneficial effects of cryosurgical treatment: benign and low-grade malignant bone tumors in 120 patients. *Ned Tijdschr Geneeskd* 1999;143:2275-81
32. Patterson EJ, Scudamore CH, Owen DA, Nagy AG, Buczkowski AK. Radiofrequency ablation of porcine liver in vivo: effects of blood flow and treatment time on lesion size. *Ann Surg* 1998;227:559-565.
33. Rachbauer F, Mangat J, Bodner G, Eichberger P, Krismer M. Heat distribution and heat transport in bone during radiofrequency catheter ablation. *Arch Orthop Trauma Surg* 2003 Apr;123(2-3):86-90.
34. Cool-tip<sup>™</sup> RF Ablation System E Series; <http://www.medtronic.com/covidien/products/ablation-systems/cool-tip-rf-ablation-system-e-series>. Accessed 15.05., 2017.
35. Goldberg SN. Radiofrequency tumor ablation: principles and techniques. *European Journal of Ultrasound* 2001;13(2):129-147.

36. Chen MS, Li JQ, Zheng Y, Guo RP, Liang HH, Zhang YQ, Xiao JL, et al. A prospective randomized trial comparing percutaneous local ablative therapy and partial hepatectomy for small hepatocellular carcinoma. *Annals of Surgery* 2006; 243: 321-8.
37. Livraghi T, Meloni F, Stasi M Di, Rolle E, Solbiati L, Tinelli C, Rossi S. Sustained complete response and complications rates after radiofrequency ablation of very early hepatocellular carcinoma in cirrhosis: Is resection still the treatment of choice? *Hepatology* 2008; 47: 82-9.
38. Lindner NJ, Ozaki T, Roedl R, Gosheger G, Winkelmann W, Wortler K. Percutaneous radiofrequency ablation in osteoid osteoma. *J Bone Joint Surg Br.* 2001;83(3):391-6.
39. Akhlaghpour S, Tomasian A, Arjmand Shabestari A, Ebrahimi M, Alinaghizadeh MR. Percutaneous osteoid osteoma treatment with combination of radiofrequency and alcohol ablation. *Clin Radiol.* 2007;62(3):268-73.
40. Cioni R, Armillotta N, Bargellini I, Zampa V, Cappelli C, Vagli P et al. CT-guided radiofrequency ablation of osteoid osteoma: longterm results. *Eur Radiol.* 2004;14(7):1203-8.
41. Woertler K, Vestring T, Boettner F, Winkelmann W, Heindel W, Lindner N. Osteoid osteoma: CT-guided percutaneous radiofrequency ablation and follow-up in 47 patients. *J Vasc Interv Radiol.* 2001;12(6):717-22.
42. Rybak LD, Rosenthal DI, Wittig JC. Chondroblastoma: radiofrequency ablation: alternative to surgical resection in selected cases. *Radiology* 2009;251:599-604.
43. Goetz MP, Callstrom MR, Charboneau JW, Farrell MA, Maus TP, Welch TJ, et al. Percutaneous image-guided radiofrequency ablation of painful metastases involving bone: a multicenter study. *J Clin Oncol.* 2004;22(2):300-6.
44. Belfiore G, Tedeschi E, Ronza FM, Belfiore MP, Della VT, Zeppetella G, et al. Radiofrequency ablation of bone metastases induces long-lasting palliation in patients with untreatable cancer. *Singapore Med J.* 2008;49(7):565-70.
45. Kashima M, Yamakado K, Takaki H, Kaminou T, Tanigawa N, Nakatsuka A, et al. Radiofrequency ablation for the treatment of bone metastases from hepatocellular carcinoma. *AJR Am J Roentgenol.* 2010;194(2):536-41.
46. Finstein JL, Hosalkar HS, Ogilvie CM, Lackman RD. Case reports: an unusual complication of radiofrequency ablation treatment of osteoid osteoma. *Clin Orthop Relat Res* 2006;448:248-51.



