

University of Groningen

## RNA-seq-based profiling of extracellular vesicles in plasma reveals a potential role of miR-122-5p in asthma

Bahmer, Thomas; Krauss-Etschmann, Susanne; Buschmann, Dominik; Behrends, Jochen; Watz, Henrik; Kirsten, Anne-Marie; Pedersen, Frauke; Waschki, Benjamin; Fuchs, Oliver; Pfaffl, Michael W.

*Published in:*  
Allergy

*DOI:*  
[10.1111/all.14486](https://doi.org/10.1111/all.14486)

**IMPORTANT NOTE:** You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

*Document Version*  
Publisher's PDF, also known as Version of record

*Publication date:*  
2021

[Link to publication in University of Groningen/UMCG research database](#)

### *Citation for published version (APA):*

Bahmer, T., Krauss-Etschmann, S., Buschmann, D., Behrends, J., Watz, H., Kirsten, A-M., Pedersen, F., Waschki, B., Fuchs, O., Pfaffl, M. W., von Mutius, E., Rabe, K. F., Hansen, G., Kopp, M. V., Koenig, I. R., & Bartel, S. (2021). RNA-seq-based profiling of extracellular vesicles in plasma reveals a potential role of miR-122-5p in asthma. *Allergy*, 76(1), 366-371. <https://doi.org/10.1111/all.14486>

### **Copyright**

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

### **Take-down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



# RNA-seq-based profiling of extracellular vesicles in plasma reveals a potential role of miR-122-5p in asthma

To the Editor,

Asthma is a heterogeneous disease encompassing several distinct sub-phenotypes with different etiologies and treatment responses,<sup>1</sup> but we are lacking markers to differentiate patient subgroups.

MicroRNAs (miRNAs) can regulate gene expression post-transcriptionally. Due to their high stability in body fluids, their easy detection, and their functional relevance in asthma, we and others have proposed extracellular miRNAs in noninvasive clinical samples as biomarkers for asthma.<sup>2,3</sup> Yet, extracellular miRNA profiles can be confounded by unspecific release of miRNAs from dying cells. Transfer of miRNAs by extracellular vesicles (EVs) is, however, supposedly a selective communication mechanism,<sup>4</sup> and EV-miRNA levels have been shown to be altered in bronchoalveolar lavage (BAL) fluid of patients with asthma.<sup>5</sup> In this pilot study, we profiled the EV-miRNA signature in plasma of patients with mild-to-moderate (AM) or severe eosinophilic asthma (AS) (as defined by ERS/ATS guidelines<sup>6</sup>) and healthy control (HC) subjects (Tables S1 and S2).

Therefore, we isolated small EVs (EVs) (<200 nm) by size exclusion chromatography (SEC) (qEV, Izon Bioscience) from 1 mL plasma of 45 adult subjects with AM (n = 15) or AS (n = 14), and HC (n = 16) (Table S1). All study participants were enrolled in the all age asthma (ALLIANCE) cohort, a multi-center longitudinal asthma patient cohort of the German Center for Lung Research (DZL).<sup>7</sup>

SEC-isolated particles were confirmed to be <200 nm by nanoparticle tracking analysis (NTA) (ZetaView PMX 110, Particle Metrix) and were equal in concentration and size distribution across groups (Figure 1A). The median particle concentration isolated from 1 mL of plasma was  $8.6 \times 10^9$  particles/mL (interquartile range [IQR]:  $4.5 \times 10^9$ - $1.4 \times 10^{10}$ ) for HC,  $1.2 \times 10^{10}$  particles/mL (IQR:  $6.6 \times 10^9$ - $2.8 \times 10^{10}$ ) for AM, and  $1.0 \times 10^{10}$  particles/mL (IQR:  $3.9 \times 10^9$ - $3.6 \times 10^{10}$ ) for AS. Furthermore, SEC fractions 7-9 used for sequencing were without protein contamination (Figure 1B) and expressed different amounts of typical small EV markers CD63, CD81, and/or CD9 (Figure S1).

EV-RNA was isolated (miRNeasy Micro Kit; Qiagen) and subjected to RNA sequencing (RNA-seq) (HiSeq 2500, Illumina) of small RNAs (below 35 nt) according to Ref.<sup>8</sup> While all samples achieved high Phred scores (>30) indicating excellent sequencing quality (Figure

S2A), 46.42% of reads were short or unmapped (39.11%) (Figure 1C). From all mapped reads, 17.15% were classified as miRNAs, while 81.08% were ribosomal RNAs. In total, we detected 139 distinct EV miRNAs with  $\geq 10$  reads (AM: 115; AS: 114; HC: 128) (Figure S2B, Tables S3-S5), and 35 of which had read counts  $\geq 50$ . Unsupervised clustering and principal component analysis did not separate the different groups (Figure S2C, D), indicating that there is no difference between asthma patients and healthy controls based on total EV-miRNA expression. However, miR-122-5p was significantly increased in all patients with asthma (log<sub>2</sub> fold change (log<sub>2</sub>FC) = 1.74, false discovery rate (FDR) = 0.03) and in AS (log<sub>2</sub>FC = 1.77, FDR = 0.02) compared with healthy controls, while miR-3168 was decreased (log<sub>2</sub>FC = -1.28; FDR = 0.05) (Figure 1D). In AS, we found similar trend for miR-191-5p (log<sub>2</sub>FC = 0.39, FDR = 0.12) (Figure 1E).



In a bivariate analysis, miR-191-5p normalized read counts correlated negatively with FEV<sub>1%pred</sub> (Spearman's  $R = -.38$ ;  $P = .013$ ) and lymphocyte percentage in blood ( $R = -.5$ ;  $P = .0006$ ) (Figure 1F). We found a positive correlation with blood neutrophil ( $R = .48$ ;  $P = .0012$ ) counts and a similar trend for blood eosinophil counts ( $R = .29$ ;  $P = .058$ ). miR-3168 read counts correlated significantly with blood eosinophils ( $R = -.28$ ,  $P = .012$ ), neutrophils ( $R = -.3$ ,  $P = .05$ ), and lymphocytes ( $R = .38$ ,  $P = .012$ ). Read counts of miR-122-5p positively correlated with both eosinophil ( $R = .32$ ;  $P = .034$ ) and neutrophil ( $R = .39$ ;  $P = .0094$ ) counts in blood, and trendwise with blood lymphocytes ( $R = -.28$ ;  $P = .065$ ) (Figure 1F). We do acknowledge that some correlation coefficients are quite low ( $R < .3$ ) which could be due to the small sample size and needs to be confirmed in larger future studies.

Next, we isolated EVs from additional plasma samples of the same subjects and 10 new subjects per group (Figure 2A), and found a strong trend for an increase of miR-122-5p (FC = 1.84;  $P = .07$ ) in severe asthma, and in all asthma subjects (FC = 1.96;  $P = .09$ ) (Figure 2A) by RT-qPCR, confirming the sequencing results. miR-191-5p did not differ significantly among the groups, and miR-3168 was not detectable by RT-qPCR. Of note, in a pilot approach both miRNAs could also be detected in EVs isolated from sputum supernatant samples of four healthy control and ten asthma subjects, and we have a first hint that they are increased in asthma compared

Thomas Bahmer and Susanne Krauss-Etschmann contributed equally to this study.

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

© 2020 The Authors. Allergy published by European Academy of Allergy and Clinical Immunology and John Wiley & Sons Ltd



**(D) all Asthma vs HC**


miRNA	baseMean	log2FoldChange	pvalue	FDR
hsa-miR-122-5p	1441.75	1.74	4.60E-05	0.03
hsa-miR-206	0.93	1.87	1.97E-02	0.99
hsa-miR-148a-3p	450.36	0.63	5.52E-03	0.99
hsa-miR-941	14.50	0.61	6.44E-03	0.99
hsa-miR-223-5p	8.59	0.53	2.23E-02	0.99
hsa-miR-191-5p	69.32	0.29	7.23E-03	0.99
hsa-miR-486-5p	1184.81	-0.39	1.89E-02	0.99
hsa-miR-3168	1327.95	-1.28	1.48E-04	0.05
hsa-miR-6842-3p	1.27	-1.38	2.36E-02	0.99

**(E) AM vs HC**


miRNA	baseMean	log2FoldChange	pvalue	FDR
hsa-miR-206	0.79	2.05	2.86E-02	1.00
hsa-miR-576-3p	0.80	1.51	6.64E-02	1.00
hsa-miR-3652	1.67	1.26	4.61E-02	1.00
hsa-miR-184	83.13	1.19	5.78E-02	1.00
hsa-miR-28-5p	2.52	1.04	1.77E-02	1.00
hsa-miR-941	14.04	0.82	3.30E-04	0.20
hsa-miR-122-5p	685.37	0.66	1.56E-02	1.00
hsa-miR-2110	2.66	-1.12	2.00E-02	1.00
hsa-miR-3168	1389.62	-1.37	6.69E-04	0.21
hsa-miR-6842-3p	1.28	-1.87	1.88E-02	1.00

**(E) AS vs HC**

miRNA	baseMean	log2FoldChange	pvalue	FDR
hsa-miR-122-5p	1230.18	1.77	3.21E-05	0.02
hsa-miR-191-5p	71.15	0.39	3.81E-04	0.12
hsa-miR-223-5p	8.72	0.69	1.06E-02	0.73
hsa-miR-148a-3p	398.11	0.47	6.56E-03	0.73
hsa-miR-143-3p	60.94	0.45	1.04E-02	0.73
hsa-miR-let-7e-5p	40.70	-0.37	7.32E-03	0.73
hsa-miR-409-3p	17.28	-0.88	1.03E-02	0.73
hsa-miR-3168	1626.92	-1.16	9.62E-03	0.73
hsa-miR-4433b-3p	4.63	-1.25	1.21E-02	0.75
hsa-miR-134-5p	1.50	-2.28	6.64E-03	0.73



**FIGURE 1** small RNA sequencing from plasma EVs. A, EV concentration (left panel), size distribution, and median size (right panels) in plasma EV isolations from healthy controls (HC,  $n = 16$ ), mild-to-moderate (AM,  $n = 15$ ), or severe asthma (AS,  $n = 14$ ) assessed by nanoparticle tracking analysis (lines represent median). B, Protein content of qEV fractions in mg/mL, assessed by microBCA ( $n = 45$  per fraction, mean  $\pm$  SD). C, Sequencing distribution of small RNA classes in %, with relative distribution within mapped reads (right) (D and E) 10 top-regulated miRNAs (sorted by log<sub>2</sub> fold change) in asthma patients vs HC (D) and AM vs HC (left) and AS vs HC (right) (E); FDR = false discovery rate. F, Spearman's correlations of miR-122-5p (upper panels), miR-3168 (middle panels), and miR-191-5p (lower panels) read counts with clinical characteristics of all included subjects ( $n = 45$ ), and each graph depicts correlation co-efficient  $R$  and  $P$ -value



**FIGURE 2** EV-miRNA from plasma correlates with clinical characteristics of asthma. A and B, RT-qPCR for miR-122-5p and miR-191-5p normalized to miR-21 levels (reference) in (A) plasma EVs from healthy controls (HC,  $n = 26$ ), all asthma ( $n = 51$ ), mild-to-moderate (AM,  $n = 25$ ) or severe asthma (AS,  $n = 26$ ), or (B) sputum supernatant EVs from healthy controls (HC,  $n = 4$ ) and patients with asthma ( $n = 10$ ). All Mann-Whitney U to HC or AM. \* $P \leq .05$ , \*\* $P \leq .01$ , \*\*\* $P \leq .001$ . Graphs depict fold changes to healthy controls with lines indicating the median. (C) Ingenuity pathway analysis showing association of miR-122-5p and miR-191-5p with quantity of leukocytes, function of antigen-presenting cells, activation of antigen-presenting cells, function of Th2 cells, quantity of regulatory T lymphocytes, and differentiation of Th2 cells



with healthy controls (miR-122-5p: FC = 2.90;  $P = .014$ ; miR-191-5p: FC = 2.69,  $P = .014$ ) (Figure 2B).

Ingenuity Pathway Analysis (IPA) of predicted targets of miR-122-5p and miR-191-5p revealed a concise network, containing the biological functions *quantity of leukocytes*, *function of antigen-presenting cells*, *activation of antigen-presenting cells*, *function of Th2 cells*, *quantity of regulatory T lymphocytes*, and *differentiation of Th2 cells* (Figure 2C). This is a first hint that miR-122-5p (and potentially miR-191-5p) could influence immune cell function upon uptake. This could systemically perpetuate the asthmatic phenotype and should be confirmed in larger studies, where also the specificity for asthma vs a general immune reaction should be assessed.

The relatively low number of significantly altered miRNAs here might be due to the isolation of highly pure EVs by SEC. This is in line with previous reports,<sup>8</sup> but SEC isolation is superior to other methods in separating EVs from contaminating proteins.<sup>9</sup> Further, we have here focused on eosinophilic asthma of different severities, hampering comparisons of different asthma sub-phenotypes. This will require considerably larger patient numbers and was thus beyond the scope of this study.

In summary, in relatively well-controlled asthma the total miRNA expression in plasma EVs is not different to controls. miR-122-5p is increased in plasma and sputum supernatant EVs derived from patients with (severe) asthma, and this miRNA correlated with immune cell types in the blood. Combined with the IPA-predicted role in lymphocyte differentiation and function, it is intriguing to speculate that this miRNA can sub-differentiate different forms of asthma, such as neutrophilic from eosinophilic asthma. This should be investigated in larger asthma cohort studies with a broad spectrum of clinically well-defined phenotypes and different treatment regimen also including steroid-naïve patients.

## ACKNOWLEDGMENTS

The authors thank all included patients for their contribution. Furthermore, we thank Regine Wieland, Petra Hundack-Winter, Susann Prange, Vera Veith, and Lukas Hundack for their excellent support in patient recruitment and study logistics.

## CONFLICT OF INTEREST

This study was supported by an unrestricted grant (Next Generation Award) from Bencard Allergie GmbH to SB, which did not influence the content of the manuscript or the conclusions drawn. The authors declare no further conflict of interest relating to this manuscript.

## AUTHOR CONTRIBUTIONS

TB, HW, A-MK, FP, BW, OF E.vM., KFR, GH, and MVK designed and conducted the clinical study including patient recruitment and processing of plasma samples. DB, MWP, J.B, S.K-E., and SB performed experiments and critically analyzed the data. IRK performed the bivariate analysis with clinical data. SB had primary responsibility for the experimental study design and writing of the manuscript. All authors have contributed to discussion of the data and writing of the manuscript, and approved the final version.

## FUNDING INFORMATION

This study was supported by an unrestricted grant (Next Generation Award; Bencard Allergie GmbH) to SB. The general costs of patient recruitment and study procedures are covered by unrestricted grants of the German Federal Ministry of Education and Research (BMBF) as part of the funding of the German Center for Lung Research (DZL) (for further details please see Ref. 7).


## ETHICAL APPROVAL

The study was approved by the local ethics committee of the Medical School Luebeck, Schleswig-Holstein (Germany, Az. 12-215), and is registered at clinicaltrials.gov (Identifier: NCT02419274). All participants gave their written informed consent.

## REFERENCES

1. Wenzel SE. Asthma phenotypes: the evolution from clinical to molecular approaches. *Nat Med*. 2012;18:716-725.
2. Heffler E, Allegra A, Pioggia G, Picardi G, Musolino C, Gangemi S. MicroRNA profiling in asthma: potential biomarkers and therapeutic targets. *Am J Respir Cell Mol Biol*. 2017;57:642-650.
3. Milger K, Götschke J, Krause L, et al. Identification of a plasma miRNA biomarker signature for allergic asthma: a translational approach. *Allergy*. 2017;72(12):1962-1971.
4. Valadi H, Ekström K, Bossios A, Sjöstrand M, Lee JJ, Lötvald JO. Exosome-mediated transfer of mRNAs and microRNAs is a novel mechanism of genetic exchange between cells. *Nat Cell Biol*. 2007;9:654-659.
5. Levänen B, Bhakta NR, Torregrosa Paredes P, et al. Altered microRNA profiles in bronchoalveolar lavage fluid exosomes in asthmatic patients. *J Allergy Clin Immunol*. 2013;131(3):894-903.
6. Chung KF, Wenzel SE, Brozek JL, et al. International ERS/ATS guidelines on definition, evaluation and treatment of severe asthma. *Eur Respir J*. 2014;43:343-373.
7. Fuchs O, Bahmer T, Weckmann M, et al. The all age asthma cohort (ALLIANCE) - from early beginnings to chronic disease: a longitudinal cohort study. *BMC Pulm Med*. 2018;18:140.
8. Buschmann D, Kirchner B, Hermann S, et al. Evaluation of serum extracellular vesicle isolation methods for profiling miRNAs by next-generation sequencing. *J Extracell Vesicles* 2018;7:1481321.
9. Stranska R, Gysbrechts L, Wouters J, et al. Comparison of membrane affinity-based method with size-exclusion chromatography for isolation of exosome-like vesicles from human plasma. *J Transl Med*. 2018;16:1.

Thomas Bahmer<sup>1,2</sup>  
Susanne Krauss-Etschmann<sup>2,3</sup>   
Dominik Buschmann<sup>4</sup>   
Jochen Behrends<sup>5</sup>  
Henrik Watz<sup>6</sup>  
Anne-Marie Kirsten<sup>6</sup>  
Frauke Pedersen<sup>1,6</sup>  
Benjamin Waschki<sup>1,7</sup>  
Oliver Fuchs<sup>8,9</sup>  
Michael W. Pfaffl<sup>4</sup>   
Erika von Mutius<sup>10</sup>  
Klaus F. Rabe<sup>1,2</sup>  
Gesine Hansen<sup>11</sup>  
Matthias V. Kopp<sup>8,9</sup>

Inke R. König<sup>12</sup>  
Sabine Bartel<sup>3,13</sup> 

<sup>1</sup>Pneumology, LungenClinic Grosshansdorf, Member of the German Center for Lung Research (DZL), Airway Research Center North (ARCN), Grosshansdorf, Germany

<sup>2</sup>University Hospital Schleswig-Holstein, Campus Kiel, Member of the German Center for Lung Research (DZL), Airway Research Center North (ARCN), Kiel, Germany

<sup>3</sup>Leibniz Lung Center, Research Center Borstel, Member of the German Center for Lung Research (DZL), Airway Research Center North (ARCN), Borstel, Germany

<sup>4</sup>Division of Animal Physiology and Immunology, School of Life Sciences Weihenstephan, Technical University of Munich, Munich, Germany

<sup>5</sup>Flow Cytometry Core Unit, Leibniz Lung Center Borstel, Member of the German Center for Lung Research (DZL), Airway Research Center North (ARCN), Borstel, Germany

<sup>6</sup>Pulmonary Research Institute, LungenClinic Grosshansdorf, Member of the German Center for Lung Research (DZL), Airway Research Center North (ARCN), Grosshansdorf, Germany

<sup>7</sup>Department of General and Interventional Cardiology, University Heart Center Hamburg, Hamburg, Germany

<sup>8</sup>Inselspital Bern, University Children's Hospital, Bern, Switzerland

<sup>9</sup>Department of Pediatric Pulmonology and Allergology, Children's Hospital at the University of Luebeck, Member of the German Center for Lung Research (DZL), Airway Research Center North (ARCN), Luebeck, Germany

<sup>10</sup>Dr. von Hauner Children's Hospital, Helmholtz Center Munich, Comprehensive Pneumology Center – Munich (CPC-M), Member of the German Center for Lung Research (DZL), Munich, Germany

<sup>11</sup>Department of Pediatric Pulmonology, Allergology and Neonatology, Hannover Medical School, Member of the German Center for Lung Research (DZL), Biomedical Research in Endstage and Obstructive Lung Disease (BREATH), Hannover, Germany

<sup>12</sup>Institute of Medical Biometry and Statistics, University of Luebeck, Member of the German Center for Lung Research (DZL), Airway Research Center North (ARCN), Luebeck, Germany

<sup>13</sup>Department of Pathology and Medical Biology, University of Groningen, University Medical Center Groningen, GRIAC Research Institute, Groningen, The Netherlands

#### Correspondence

Sabine Bartel, University Medical Center Groningen, Hanzeplein 1, 9700 RB Groningen, The Netherlands.

Email: s.r.bartel@umcg.nl

#### ORCID

Susanne Krauss-Etschmann  <https://orcid.org/0000-0001-5945-5702>

Dominik Buschmann  <https://orcid.org/0000-0003-0460-6459>

Michael W. Pfaffl  <https://orcid.org/0000-0002-3192-1019>

Sabine Bartel  <https://orcid.org/0000-0002-9163-795X>

#### SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section.