

University of Groningen

The Unruptured Intracranial Aneurysm Treatment Score as a Predictor of Aneurysm Growth or Rupture

Molenberg, Rob; Aalbers, Marlien W; Mazuri, Aryan; Luijckx, Gert Jan; Metzemaekers, Jan D M; Groen, Rob J M; Uyttenboogaart, Maarten; van Dijk, J Marc C

Published in:
European Journal of Neurology

DOI:
[10.1111/ene.14636](https://doi.org/10.1111/ene.14636)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2021

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Molenberg, R., Aalbers, M. W., Mazuri, A., Luijckx, G. J., Metzemaekers, J. D. M., Groen, R. J. M., Uyttenboogaart, M., & van Dijk, J. M. C. (2021). The Unruptured Intracranial Aneurysm Treatment Score as a Predictor of Aneurysm Growth or Rupture. *European Journal of Neurology*, 28(3), 837-843. <https://doi.org/10.1111/ene.14636>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.



Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

ORIGINAL ARTICLE

The Unruptured Intracranial Aneurysm Treatment Score as a predictor of aneurysm growth or rupture

Rob Molenberg¹  | Marlien W. Aalbers¹ | Aryan Mazuri² | Gert Jan Luijckx³ |
Jan D. M. Metzemaekers¹ | Rob J. M. Groen¹ | Maarten Uyttenboogaart^{2,3} |
J. Marc C. van Dijk¹ 

¹Departments of Neurosurgery, University of Groningen, University Medical Center Groningen, Groningen, the Netherlands

²Departments of Radiology, University of Groningen, University Medical Center Groningen, Groningen, the Netherlands

³Departments of Neurology, University of Groningen, University Medical Center Groningen, Groningen, the Netherlands

Correspondence

Rob Molenberg, Department of Neurosurgery AB71, University Medical Center Groningen, PO Box 33.001, 9700 RB Groningen, the Netherlands.
Email: r.molenberg@umcg.nl

Abstract

Background and purpose: The Unruptured Intracranial Aneurysm Treatment Score (UIATS) was built to harmonize the treatment decision making on unruptured intracranial aneurysms. Therefore, it may also function as a predictor of aneurysm progression. In this study, we aimed to assess the validity of the UIATS model to identify aneurysms at risk of growth or rupture during follow-up.

Methods: We calculated the UIATS for a consecutive series of conservatively treated unruptured intracranial aneurysms, included in our prospectively kept neurovascular database. Computed tomography angiography and/or magnetic resonance angiography imaging at baseline and during follow-up was analyzed to detect aneurysm growth. We defined rupture as a cerebrospinal fluid or computed tomography-proven subarachnoid hemorrhage. We calculated the area under the receiver operator curve, sensitivity, and specificity, to determine the performance of the UIATS model.

Results: We included 214 consecutive patients with 277 unruptured intracranial aneurysms. Aneurysms were followed for a median period of 1.3 years (range 0.3–11.7 years). During follow-up, 17 aneurysms enlarged (6.1%), and two aneurysms ruptured (0.7%). The UIATS model showed a sensitivity of 80% and a specificity of 44%. The area under the receiver operator curve was 0.62 (95% confidence interval 0.46–0.79).

Conclusions: Our observational study involving consecutive patients with an unruptured intracranial aneurysm showed poor performance of the UIATS model to predict aneurysm growth or rupture during follow-up.

KEYWORDS

growth, intracranial aneurysm, predictors, subarachnoid hemorrhage, UIATS

INTRODUCTION

Unruptured intracranial aneurysms constitute a relatively frequently encountered neurovascular disease, with a prevalence of around 3%

in the general population [1,2]. Aneurysm rupture leading to a subarachnoid hemorrhage (SAH) is associated with high morbidity and mortality rates [3]. In this respect, proactive treatment via aneurysm clipping or coiling is seen as the cornerstone of rupture prevention.

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

© 2020 The Authors. *European Journal of Neurology* published by John Wiley & Sons Ltd on behalf of European Academy of Neurology

Because elective treatment also bears procedural risks, it should be balanced against the estimated risk of future aneurysm rupture. Therefore, conservative treatment may be the preferred treatment strategy for aneurysms deemed low risk. Typically, short-term follow-up of such aneurysms is indicated, enabling the detection of potential aneurysm growth and facilitating timely reevaluation of elective treatment [4].

Various risk-assessment tools have been introduced to assist in the treatment decision making of unruptured aneurysms. The Unruptured Intracranial Aneurysm Treatment Score (UIATS) is relatively new and represents the most extensive risk-assessment tool [5]. The UIATS is a Delphi consensus-based model, consisting of 29 variables compromising patient-, aneurysm-, as well as treatment-specific variables. Points are given for each variable in the model, either in favor of elective treatment or conservative management. The result is a recommendation to treat or not to treat an aneurysm (Table 1). A point difference smaller than three results in an inconclusive recommendation. Despite its extensiveness, its clinical predictive performance and external validity remain largely unknown.

The primary objective of this study was to determine the ability of the UIATS model to identify unruptured intracranial aneurysms at risk of growth or rupture during follow-up.

METHODS

Study population

All adult patients with a conservatively treated, unruptured intracranial aneurysm, assessed between 1997 and 2017, were consecutively included in our prospectively kept neurovascular database. The decision for conservative management of the aneurysm was made by the multidisciplinary neurovascular group at our tertiary referral center. We excluded extradural aneurysms and patients without follow-up imaging. Patients signed an informed consent as required by our University Medical Center Groningen Research Ethical Board.

Data collection

We used questionnaires and chart reviews to acquire data regarding the 29 variables described in the UIATS model (Table 1). The questionnaires were mainly used to ask if patients experienced a reduced quality of life due to fear of rupture, as well as double checking UIATS-related patient-specific risk factors potentially not mentioned in the patients' referral information. We omitted the variables aneurysm growth and aneurysm de novo formation on serial imaging, because we calculated UIATS recommendations at baseline. Aneurysm measurements were based on three-dimensional (3D) time-of-flight magnetic resonance angiography with 0.6-mm slice thickness or contrast-enhanced computed

tomography angiography with 0.75-mm slice thickness. We measured aneurysm height, width, and neck diameter at baseline and during follow-up to detect aneurysm growth. Aneurysm shape was evaluated using 3D reconstructions, with an irregular shape defined as the presence of blebs, wall protrusions, or multiple lobes [6]. Growth was defined as either an increase in size of at least 1 mm in at least one direction, or a change in aneurysm shape [7]. We defined rupture as a computed tomography- and/or cerebrospinal fluid-proven SAH. For all unstable aneurysms (i.e., those with growth or rupture), we also calculated the PHASES score. PHASES stands for the variables population, hypertension, age, size of the aneurysm, earlier SAH from another aneurysm, and site of the aneurysm, which can be used to estimate the 5-year risk of rupture [8]. We used the PHASES score to evaluate the estimated risk distribution of the unstable aneurysms at baseline.

Statistical analysis

UIATS recommendations were determined by calculating the sum of points for each strategy (treatment vs. conservative). Aneurysms with an inconclusive recommendation were excluded from statistical analysis. Aneurysms with a treatment recommendation showing progression during follow-up were considered as true positives, whereas those receiving a conservative recommendation showing progression during follow-up were considered as false negatives. We used the Fisher exact test for group comparisons of categorical data, and the Student *t* test or Mann-Whitney *U* test for continuous data. The level of statistical significance was set at 0.05. We calculated the sensitivity, specificity, positive and negative predictive value, and the area under the receiver operator curve (AUC) to assess the performance of the UIATS model to discriminate between aneurysms with and without progression. Statistical analyses were performed using SPSS (version 23.0; IBM Corp.).

RESULTS

A total of 214 patients (277 unruptured aneurysms) with a conservative treatment recommendation were assessed. The majority of patients (74%) were female. Aneurysms were followed for a median period of 1.3 years (range 0.3–11.7 years). During follow-up, 17 (6.1%) aneurysms enlarged, and two (0.7%) aneurysms ruptured. Median duration to the occurrence of growth or rupture was 2.45 years (range 0.8–9.8 years). Based on the UIATS model, a total of 88 (32%) aneurysms received a treatment recommendation, 65 (23%) received a conservative recommendation, and for 124 (45%) the UIATS recommendation was inconclusive (Table 2). The majority of unstable aneurysms had a relatively low estimated 5-year rupture risk based on the PHASES score at baseline, generally favoring conservative treatment. One large aneurysm of the anterior communicating artery with a high PHASES score of 16 was treated conservatively due to the patients' older age, intra-aneurysmal thrombosis,

TABLE 1 The Unruptured Intracranial Aneurysm Treatment Score variables

Variables	Categories	Points in favor of treatment	Points in favor of conservative management
Age	<40 years	4	
	40–60 years	3	
	61–70 years	2	
	71–80 years	1	
	>80 years	0	
Risk factor incidence	Previous SAH from a different aneurysm or SAH	4	
	Familial intracranial aneurysms or SAH	3	
	Japanese, Finnish, Inuit ethnicity	2	
	Current cigarette smoking	3	
	Hypertension	2	
	Autosomal polycystic kidney disease	2	
	Current drug abuse	2	
	Current alcohol abuse	1	
Clinical symptoms related to UIA	Cranial nerve deficit	4	
	Clinical or radiological mass effect	4	
	Thromboembolic events from the aneurysm	3	
	Epilepsy	1	
Other	Reduced quality of life due to fear of rupture	2	
	Aneurysm multiplicity	1	
Life expectancy due to chronic and/or malignant diseases	<5 years		4
	5–10 years		3
	>10 years		1
Comorbid disease	Neurocognitive disorder		3
	Coagulopathies, thrombophilic diseases		2
	Psychiatric disorder		2
Maximum diameter	≤3.9 mm	0	
	4.0–6.9 mm	1	
	7.0–12.9 mm	2	
	13.0–24.0 mm	3	
	≥25 mm	4	
Morphology	Irregularity or lobulation	3	
	Size ratio >3 or aspect ratio >1.6	1	
Aneurysm location	Basilar bifurcation	5	
	Vertebral/basilar artery	4	
	ACoA or PCoA	2	
Other	Aneurysm growth on serial imaging	4	
	Aneurysm de novo formation on serial imaging	3	
	Contralateral steno-occlusive vessel disease	1	

(Continues)

TABLE 1 (Continued)

Variables	Categories	Points in favor of treatment	Points in favor of conservative management
Age-related risk	<40 years		0
	40–60 years		1
	61–70 years		3
	71–80 years		4
	>80 years		5
Aneurysm size-related risk	<6.0 mm		0
	6.0–10 mm		1
	10.1–20.0 mm		3
	>20 mm		5
Aneurysm complexity-related risk	High		3
	Low		0
Intervention-related risk	Constant		5

Note: Adapted from Etminan et al. [5]. A point difference ≥ 3 results in a recommendation favoring the treatment strategy with the highest number of points. If the difference is ≤ 2 , an inconclusive recommendation is given.

Abbreviations: ACoA, anterior communicating artery; ADPKD, autosomal dominant polycystic kidney disease; PCoA, posterior communicating artery; SAH, subarachnoid hemorrhage; UIA, unruptured intracranial aneurysm.

TABLE 2 Distribution of UIATS recommendations for stable and unstable aneurysms

UIATS recommendation	Stable, n = 258	Unstable, n = 19
Repair	80 (31)	8 (42)
Conservative management	63 (24)	2 (11)
Inconclusive	115 (45)	9 (47)

Note: Data are displayed as n (%).

Abbreviation: UIATS, Unruptured Intracranial Aneurysm Treatment Score.

and significant comorbidities. No clear differences between PHASES scores and UIATS recommendations could be identified (Figure 1).

Besides older age, high aneurysm complexity risk (mainly due to small aneurysm size) was the only variable associated with favoring a conservative recommendation (Table 3). Variables associated with favoring a treatment recommendation were: (i) a previous SAH from a different aneurysm, (ii) familial intracranial aneurysms or SAH, (iii) current smoking, (iv) current alcohol abuse, (v) reduced quality of life due to fear of rupture, and (vi) aneurysm multiplicity. Subsequent analysis of the UIATS model showed a sensitivity of 80%, a specificity of 44%, a positive predictive value of 9%, and a negative predictive value of 97% (Table 2). The AUC of the receiver operator curve was 0.62 (95% confidence interval [CI] 0.46–0.79), implicating poor performance of the UIATS model to discriminate between stable and unstable aneurysms.

DISCUSSION

We evaluated whether the UIATS model is valid to predict growth or rupture of unruptured intracranial aneurysms during follow-up. The

results of our study show that aneurysm progression during follow-up could not be reliably predicted by the UIATS model. In addition, almost one-third of aneurysms remaining stable during follow-up received a treatment recommendation. On the other hand, nearly half of the aneurysms in this study received an inconclusive recommendation, whereas the number of aneurysms with growth or rupture was highest in this category.

To our knowledge, two other studies investigated the predictive performance of the UIATS model. The first study performed a retrospective analysis of 212 consecutive SAH patients at their institution [9]. They calculated UIATS recommendations for all aneurysms to determine which aneurysms would have received a treatment recommendation (true positives) or a conservative treatment recommendation (false negatives). Analysis demonstrated a relatively low sensitivity of 44% in detecting aneurysms at risk of rupture. The second study described 142 patients with 182 unruptured intracranial aneurysms diagnosed between 1956 and 1978 [10]. Patients were followed for a median period of 21 years (range 0.8–52.3 years). The UIATS demonstrated poor performance to predict rupture, with an AUC of 0.62 (95% CI 0.50–0.73). It should be noted that these aneurysms were almost all incidental findings after an SAH from a different aneurysm, and UIATS points given due to their Finnish nationality also increased the likelihood of a UIATS recommendation favoring treatment. We therefore believe our study adds to this by providing a more contemporary cohort of patients. In our large group of aneurysms, a significant proportion of aneurysms was discovered incidentally on brain imaging in patients without a previous SAH, and follow-up angiography was performed to identify potential aneurysm growth.

In contrast to other models used in the assessment of unruptured aneurysms (e.g., the PHASES score), the UIATS model is not based on longitudinal collected data regarding risk factors




FIGURE 1 PHASES scores and UIATS recommendations of unstable aneurysms (dots). The legend shows the estimated 5-year risk of rupture of the associated PHASES score [8]. PHASES, population, hypertension, age, size of the aneurysm, earlier subarachnoid hemorrhage from another aneurysm, and site of the aneurysm; UIATS, Unruptured Intracranial Aneurysm Treatment Score.

of aneurysm progression [8]. Instead, a Delphi consensus-based approach was used in constructing the model, and the model was not validated. The developers of the UIATS model state that the model enables clinicians to appreciate what highly informed individuals in the cerebrovascular field would advise in a particular patient based on current data and uncertainties. They also state that its applicability and clinical accuracy remains to be investigated. In this respect, multiple cerebrovascular institutions compared the UIATS treatment recommendation with decisions made by their own multidisciplinary teams [11–13]. Despite all studies using different statistical methods of comparing the UIATS recommendation to their own treatment decisions, they all show limited agreement between the two.

Of interest, in accordance with the results presented in this article, most studies investigating the UIATS show only a minor role of aneurysm size in the determination of the UIATS treatment recommendation [11,12]. This is remarkable, because aneurysm size is a well-studied variable and one of the strongest predictors of rupture [14]. It seems to result from the fact that aneurysm size is both represented as a variable favoring aneurysm treatment

as well as favoring conservative management, whereas different size categories are also used for the two. Consequently, the effect of aneurysm size on the UIATS stays between one point in favor of aneurysm repair and two points in favor of conservative management, for the full range of aneurysm sizes [15]. The result is an almost nonexistent effect of aneurysm size on UIATS recommendation.

Our study does have some limitations. First, treatment decisions for individual aneurysms by our multidisciplinary neurovascular working group may be different than other neurovascular centers, potentially limiting the generalizability of our results. Second, our focus on conservatively treated aneurysms might underestimate the performance of the UIATS model to predict progression due to the inclusion of mostly aneurysms deemed low risk by our working group. Hence, included aneurysms are relatively small, and the majority of aneurysms are located in the anterior circulation, because posterior circulation aneurysms are frequently deemed more high risk and therefore more frequently treated. Finally, the mean duration of follow-up was relatively short, preventing assessment of the UIATS performance over a longer period of follow-up.

TABLE 3 The UIATS patient and aneurysm characteristics

Variable	Total group, N = 277	UIATS treatment recommendation			p value*
		Repair, n = 88	Conservative, n = 65	Inconclusive, n = 124	
Mean age ± SD, years	57 ± 10	51 ± 8	64 ± 9	57 ± 10	0.33
Risk factor incidence					
Previous SAH from a different aneurysm	109 (39)	53 (60)	16 (25)	40 (32)	<0.01
Familial intracranial aneurysms or SAH	48 (17)	24 (27)	6 (9)	18 (15)	0.01
Current cigarette smoking	122 (44)	65 (74)	9 (14)	48 (39)	<0.01
Hypertension	133 (48)	49 (56)	30 (46)	54 (44)	0.26
ADPKD	10 (4)	5 (6)	1 (2)	4 (3)	0.24
Current alcohol abuse	12 (4)	9 (10)	1 (2)	2 (2)	0.045
Clinical symptoms related to UIA					
Epilepsy	1 (0.4)	1 (1)	0	0	1.0
Other					
Reduced quality of life due to fear of rupture	68 (25)	38 (43)	10 (15)	20 (16)	<0.01
Aneurysm multiplicity	180 (65)	70 (80)	32 (49)	78 (63)	<0.01
Reduced life expectancy	2 (0.7)	0	1 (2)	1 (1)	0.43
Comorbid disease					
Neurocognitive disorder	2 (0.7)	0	2 (3)	0	0.18
Psychiatric disorder	4 (1.4)	0	3 (5)	1 (1)	0.08
Morphology					
Mean maximum diameter ± SD, mm	4.6 ± 2.7	4.1 ± 3.4	5.3 ± 2.8	4.7 ± 2.8	0.1
Irregularity or lobulation	23 (8.3)	9 (10)	2 (3)	12 (10)	0.12
Size ratio >3 or aspect ratio >1.6	60 (22)	19 (22)	11 (17)	30 (24)	0.54
Location					
BA bifurcation	13 (4.7)	7 (8)	1 (2)	5 (4)	0.14
ACoA or PCoA	57 (21)	16 (18)	12 (18)	29 (23)	1.0
Other					
Contralateral steno-occlusive vessel disease	1 (0.4)	0	0	1 (1)	N/A
High aneurysm complexity-related risk	148 (53)	33 (38)	52 (80)	63 (51)	<0.01

Note: Data are displayed as n (%) unless indicated otherwise. The following variables were omitted from this table because no patients or aneurysms were included in these categories: Japanese, Finnish, or Inuit ethnicity; current drug abuse; cranial nerve deficit; clinical or radiological mass effect; thromboembolic events from the aneurysm; coagulopathies; thrombophilic diseases; and aneurysms located at the vertebral/basilar artery.

Abbreviations: ACoA, anterior communicating artery; ADPKD, autosomal dominant polycystic kidney disease; BA, basilar artery; N/A, not applicable; PCoA, posterior communicating artery; SAH, subarachnoid hemorrhage; UIA, unruptured intracranial aneurysm; UIATS, Unruptured Intracranial Aneurysm Treatment Score.

*Statistical significance of the difference between repair and conservative.

CONCLUSIONS

This observational study involving patients with unruptured intracranial aneurysms who underwent follow-up imaging at our institution demonstrated poor performance of the UIATS model to predict aneurysm progression during follow-up.

CONFLICT OF INTEREST

None.

DATA AVAILABILITY STATEMENT

The data that support the findings of this study are available from the corresponding author upon reasonable request.

AUTHOR CONTRIBUTIONS

Rob Molenberg: conceptualization (equal); dataCuration (equal); formalAnalysis (equal); investigation (equal); methodology (equal); projectAdministration (equal); writingOriginalDraft (equal). **Marlien**

W. Aalbers: conceptualization (equal); dataCuration (equal); investigation (equal); methodology (equal); writingReviewEditing (equal). **Aryan Mazuri:** dataCuration (equal); writingReviewEditing (equal). **Gert Jan Luijckx:** dataCuration (equal); writingReviewEditing (equal). **Jan D. M. Metzemaekers:** dataCuration (equal); writingReviewEditing (equal). **Rob J. M. Groen:** dataCuration (equal); writingReviewEditing (equal). **Maarten Uyttenboogaart:** conceptualization (equal); dataCuration (equal); methodology (equal); writingReviewEditing (equal). **J. Marc C. van Dijk:** conceptualization (equal); dataCuration (equal); methodology (equal); supervision (equal); writingReviewEditing (equal).

ORCID

Rob Molenberg  <https://orcid.org/0000-0002-6621-2028>

J. Marc C. van Dijk  <https://orcid.org/0000-0002-0814-5680>

REFERENCES

1. Bos D, Poels MM, Adams HH, et al. Prevalence, clinical management, and natural course of incidental findings on brain MR images: the population-based Rotterdam scan study. *Radiology*. 2016;281(2):507-515.
2. Vlak MH, Algra A, Brandenburg R, Rinkel GJ. Prevalence of unruptured intracranial aneurysms, with emphasis on sex, age, comorbidity, country, and time period: a systematic review and meta-analysis. *Lancet Neurol*. 2011;10(7):626-636.
3. Nieuwkamp DJ, Setz LE, Algra A, Linn FH, de Rooij NK, Rinkel GJ. Changes in case fatality of aneurysmal subarachnoid haemorrhage over time, according to age, sex, and region: a meta-analysis. *Lancet Neurol*. 2009;8(7):635-642.
4. Molenberg R, Aalbers MW, Metzemaekers JDM, et al. Clinical relevance of short-term follow-up of unruptured intracranial aneurysms. *Neurosurg Focus*. 2019;47(1):E7.
5. Etminan N, Brown RD, Beseoglu K, et al. The unruptured intracranial aneurysm treatment score: a multidisciplinary consensus. *Neurology*. 2015;85(10):881-889.
6. Backes D, Vergouwen MDI, Velthuis BK, et al. Difference in aneurysm characteristics between ruptured and unruptured aneurysms in patients with multiple intracranial aneurysms. *Stroke*. 2014;45(5):1299-1303.
7. Backes D, Rinkel GJE, Greving JP, et al. ELAPSS score for prediction of risk of growth of unruptured intracranial aneurysms. *Neurology*. 2017;88(17):1600-1606.
8. Greving JP, Wermer MJ, Brown RD Jr, et al. Development of the PHASES score for prediction of risk of rupture of intracranial aneurysms: a pooled analysis of six prospective cohort studies. *Lancet Neurol*. 2014;13(1):59-66.
9. Hernández-Durán S, Mielke D, Rohde V, Malinova V. Is the unruptured intracranial aneurysm treatment score (UIATS) sensitive enough to detect aneurysms at risk of rupture? *Neurosurg Rev*. 2020. <http://dx.doi.org/10.1007/s10143-020-01246-x>
10. Juvela S. Treatment scoring of unruptured intracranial aneurysms. *Stroke*. 2019;50(9):2344-2350.
11. Ravindra VM, de Havenon A, Gooldy TC, et al. Validation of the unruptured intracranial aneurysm treatment score: comparison with real-world cerebrovascular practice. *J Neurosurg*. 2018;129(1):100-106.
12. Smedley A, Yusupov N, Almousa A, Solbach T, Toma AK, Grieve JP. Management of incidental aneurysms: comparison of single Centre multi-disciplinary team decision making with the unruptured incidental aneurysm treatment score. *Br J Neurosurg*. 2018;32(5):536-540.
13. Hernández-Durán S, Mielke D, Rohde V, Malinova V. The application of the unruptured intracranial aneurysm treatment score: a retrospective, single-center study. *Neurosurg Rev*. 2018;41(4):1021-1028.
14. Wiebers DO. Unruptured intracranial aneurysms: natural history, clinical outcome, and risks of surgical and endovascular treatment. *Lancet*. 2003;362(9378):103-110.
15. Fahed R, Darsaut TE. The Delphi Oracle and the management of aneurysms. *J NeuroIntervent Surg*. 2017;9(e1):e1-e2.

How to cite this article: Molenberg R, Aalbers MW, Mazuri A, et al. The Unruptured Intracranial Aneurysm Treatment Score as a predictor of aneurysm growth or rupture. *Eur J Neurol*. 2021;28:837-843. <https://doi.org/10.1111/ene.14636>