



University of Groningen

Unraveling the knots of gaseous Cosmic Web filaments at z 3 through H-alpha emission observations

Cantalupo, Sebastiano; Fossati, Matteo; Fumagalli, Michele; Gallego, Sofia; Lilly, Simon J.; Mackenzie, Ruari; Marino, Raffaella Anna; Maseda, Michael; Matthee, Jorryt; Nanayakkara, Themiya

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date: 2021

Link to publication in University of Groningen/UMCG research database

Citation for published version (APA):

Cantalupo, S., Fossati, M., Fumagalli, M., Gallego, S., Lilly, S. J., Mackenzie, R., Marino, R. A., Maseda, M., Matthee, J., Nanayakkara, T., Pezzulli, G., Steidel, C. C., & de Beer, S. D. C. (2021, Mar 1). Unraveling the knots of gaseous Cosmic Web filaments at z 3 through H-alpha emission observations. https://ui.adsabs.harvard.edu/abs/2021jwst.prop.1835C

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): http://www.rug.nl/research/portal. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Download date: 07-06-2022



1835 - Unraveling the knots of gaseous Cosmic Web filaments at $z\sim3$ through Halpha emission observations

Cycle: 1, Proposal Category: GO

INVESTIGATORS

Name	Institution	E-Mail
Prof. Sebastiano Cantalupo (PI) (ESA Member)	Universita di Milano-Bicocca	sebastiano.cantalupo@unimib.it
Prof. Charles C. Steidel (CoI) (US Admin CoI)	California Institute of Technology	ccs@astro.caltech.edu
Dr. Ruari Mackenzie (CoI) (ESA Member)	ETH Zurich	mruari@phys.ethz.ch
Prof. Simon J. Lilly (CoI) (ESA Member)	Eidgenossiche Technische Hochschule (ETH)	simon.lilly@phys.ethz.ch
Ms. Sofia Gallego (CoI) (ESA Member)	ETH Zurich	sofiag.gallego@gmail.com
Dr. Raffaella Anna Marino (CoI) (ESA Member)	Eidgenossiche Technische Hochschule (ETH)	marinor@phys.ethz.ch
Dr. Jorryt Matthee (CoI) (ESA Member)	ETH Zurich	matthee@strw.leidenuniv.nl
Dr. Gabriele Pezzulli (CoI) (ESA Member)	Kapteyn Astronomical Institute	pezzulli@astro.rug.nl
Ms. Stephanie Dorothy Catherine de Beer (CoI) (ES A Member)	Universita di Milano-Bicocca	s.debeer@campus.unimib.it
Dr. Michael Maseda (CoI) (ESA Member)	Universiteit Leiden	maseda@strw.leidenuniv.nl
Dr. Themiya Nanayakkara (CoI)	Swinburne University of Technology	themiyananayakkara@gmail.com
Prof. Michele Fumagalli (CoI) (ESA Member)	Universita di Milano-Bicocca	michele.fumagalli@unimib.it
Dr. Matteo Fossati (CoI) (ESA Member)	Universita di Milano-Bicocca	matteo.fossati@unimib.it

OBSERVATIONS

Folder Observati	ion Label	Observing Template	Science Target	
Observation Folder	er			
1	MQN03_Plan1	NIRSpec MultiObject Spectroscopy	(1) MQN03-PA28	
2	MQN01 Plan1	NIRSpec MultiObject Spectroscopy	(2) MQN01-PA45	

ABSTRACT

Our cosmological model predicts that most of the matter in the universe is distributed in a network of filaments - the Cosmic Web - in which galaxies form and evolve. Because most of this material is very diffuse, its direct imaging has for long remained elusive, leaving many questions still open, e.g.: what are the morphological and kinematical properties of the Cosmic Web on both small (kpc) and large (Mpc) scales? How do galaxies get their gas from the Cosmic Web? Here, we tackle these questions with an innovative method to detect in emission the gaseous Cosmic Web using bright quasars as "cosmic flashlights". In particular, we propose to observe in H-alpha emission two fields at z~3 which contain the largest Cosmic Web filaments – over 4 cMpc in length - discovered so far in deep MUSE Ly-alpha emission searches around bright quasars. Because Ly-alpha is affected by radiative transfer which change both its spatial and spectral distribution, non-resonant H-alpha observations are fundamental in order to directly constrain both the filament densities and kinematics. The filament projected angular sizes are perfectly suited for NIRSpec-MOS which can trace the filaments over their full length capturing, at the same time, several embedded galaxies. Our H-alpha observations will probe structures within the filaments on scales smaller than a few physical kpc directly constraining both their density and kinematics. By relating these quantities to the kinematics and distance from associated galaxies, our result will be fundamental to informing a new generation of theoretical and numerical models in order to reveal the physics of intergalactic gas accretion and galactic outflows.

OBSERVING DESCRIPTION

The observations for this program consist of NIRSpec Multi-Object-Spectroscopy using the G235H grating and F170LP filter within two different fields, labeled MQN01 and MQN03, centered on two bright quasars at z~3.2. These two fields contain the largest contiguous Cosmic Web filaments discovered so far in Ly-alpha emission around bright quasars.

The primary observational goal of this proposal is to detect the H-alpha emission produced by both gas and galaxies within these filaments. The emission is expected to be at a wavelength of about 2.7 microns. The region of interest in both fields extend to about 90" and they can therefore fit in a single MSA quadrant in each field. In particular, the MSA configurations are designed to follow the most interesting filamentary structures using a set of "slitlets" with a length not smaller than 3 MSA. The ideal PA which orients the MSA vertical direction parallel to the direction of the main filaments is within the schedulable range for both fields. Given the extended nature of our target, a range of PAs can be used as well in order to cover the filamentary structures and an optimal configuration can be provided once the PA is assigned. For similar reasons, continuum pre-imaging and Target Acquisition are not needed for our observations. A small nod and sub-pixel dither pattern will be used to mitigate the effect of open shutters, light-leakages and detector defects, as well as to improve PSF sampling. Open shutters in other quadrants and the fix slits will be used to obtain a "master background".

JWST Proposal 1835 (Created: Tuesday, March 30, 2021 at 3:02:47 PM Eastern Standard Time) - Overview

Observations for each field can be executed in single visits of duration 12.2 hours including overheads, estimated with the JWST ETC and APT in order to reach the required SN level. The observations, which are mostly read-out noise limited, are split in 21 exposures of about 27 minutes each per field in order to avoid excessive cosmic ray hits. In order to reduce data volume and flow, we will use the "NRSIRS2" read-out pattern splitting each integrations in 22 groups. Given the faint and diffuse nature of our targets, saturation and persistence are not an issue for our observations.

Proposal 1835 - Targets - Unraveling the knots of gaseous Cosmic Web filaments at z~3 through H-alpha emission observations

<u></u>	posai	1000 Targets	Childrening the kiloto of gaseous o	Comine Web mamorite at 2 C times	agir i i dipila difilodidi obodive	2110110
	#	Name	Target Coordinates	Targ. Coord. Corrections	Miscellaneous	
l	(1)	MQN03-PA28	RA: 00 44 33.7558 (11.1406492d)			
ပ္ပ			Dec: -26 11 45.50 (-26.19597d)			
gets			Equinox: J2000			
Tar	Comment Descripti					
xed	(2)	MQN01-PA45	RA: 00 41 32.5500 (10.3856250d)			
ΙĚ			Dec: -49 36 20.95 (-49.60582d)			
I —			Equinox: J2000			
l	Commen					
	Descripti	ion=[]				

Proposal 1835 - Observation 1 - Unraveling the knots of gaseous Cosmic Web filaments at z~3 through H-alpha emission observations

	<u> </u>	ervation i - Omav	ching the knots of	i gaseous Cosinic Wei	maments at 2~5	tillough n-aipha en	nission observations				
ation	Proposal 1835, Observation	n 1: MQN03_Plan1					Tue Mar 30 20:02:47 GMT 2021				
ı į	Diagnostic Status: Warning										
Į≅	Observing Template: NIRSpec MultiObject Spectroscopy										
Se											
Observ											
H	(MON03 Plan1 (Obs 1)) Wa	arning (Form): Config c1 (#1)	has 6 primary slit traces af	fected by failed open shutters.							
س	· -			fected by failed open shutters.							
<u>.</u> 2				fected by failed open shutters.							
ostic				-							
guc		(MQN03_Plan1 (Obs 1)) Warning (Form): Config c1 (#4) has 6 primary slit traces affected by failed open shutters.									
ag	(MQN03_Plan1 (Obs 1)) Warning (Form): Config c1 (#5) has 6 primary slit traces affected by failed open shutters. (MQN03_Plan1 (Obs 1)) Warning (Form): Config c1 (#6) has 6 primary slit traces affected by failed open shutters.										
Dia											
		-		fected by failed open shutters.							
<u></u>		Overheads are provisional un			α	3.61 13					
l ä	# Name	Target Co		Targ. Coord.	Corrections	Miscellaneous					
ρ	(1) MQN03-PA2		33.7558 (11.1406492d)								
Targets		Dec: -26 1	1 45.50 (-26.19597d)								
		Equinox: J	2000								
xed											
Ϊ́Ξ	Description=[]										
9	TA Method	Obtain Confirmation Images	Science Aperture	Primary Candidate List	Filler Candidate List	Spectral Overlap Map	Spectral Overlap Threshold				
<u> </u>											
Template	NONE	No	MSA Center	MQN03-PA28 (23 sources)		jwst-nirspec-g235h	1.2				

Proposal 1835 - Observation 1 - Unraveling the knots of gaseous Cosmic Web filaments at z~3 through H-alpha emission observations

	#	Exposure Specification	MSA Configuration	Nod Pattern	Pointing	Dispersion Offset (Shutters)	Cross-Dispersion Offset (Shutters)	Total Dithers	Total Integrations	Total Exposure Time
	1	1 (G235H/F170LP)	c1	1 Shutter Slitlet	11.161054 Degrees			1	3	4858.1
					26.1865930555553 4 Degrees					
	2	1 (G235H/F170LP)	c1	1 Shutter Slitlet	11.161054 Degrees		0.05	1	3	4858.1
					26.1865930555553 4 Degrees					
nts	3	1 (G235H/F170LP)	c1	1 Shutter Slitlet	11.161054 Degrees		-0.05	1	3	4858.1
Spectral Elements					26.18659305555553 4 Degrees					
	4	1 (G235H/F170LP)	c1	1 Shutter Slitlet	11.161054 Degrees		0.1	1	3	4858.1
ectra					26.1865930555553 4 Degrees					
S	5	1 (G235H/F170LP)	c1	1 Shutter Slitlet	11.161054 Degrees		-0.1	1	3	4858.1
					26.18659305555553 4 Degrees					
	6	1 (G235H/F170LP)	c1	1 Shutter Slitlet	11.161054 Degrees		0.15	1	3	4858.1
					26.1865930555553 4 Degrees					
	7	1 (G235H/F170LP)	c1	1 Shutter Slitlet	11.161054 Degrees		-0.15	1	3	4858.1
					- 26.1865930555553 4 Degrees					

Aperture PA Range 13 to 43 Degrees (V3 234.50765991 to 264.50765991000003)

MSA Planned Aperture PA 28.0 to 28.0 Degrees (V3 249.50765991 to 249.50765991)

Proposal 1835 - Observation 2 - Unraveling the knots of gaseous Cosmic Web filaments at z~3 through H-alpha emission observations

	Proposal 1835, Observation		elling the knots of	gaseous Cosmic Web	maments at 2~3	tillough n-aipha en	Tue Mar 30 20:02:47 GMT 2021					
흥	Diagnostic Status: Warnin	~ =					Tue Mai 30 20.02.47 GWH 2021					
Observation	U	pec MultiObject Spectroscopy										
	(MQN01_Plan1 (Obs 2)) Warning (Form): Config c1 (#1) has 4 primary slit traces affected by failed open shutters.											
	(MQN01_Plan1 (Obs 2)) W	Varning (Form): Config c1 (#2)	has 4 primary slit traces affe	ected by failed open shutters.								
တ	(MQN01_Plan1 (Obs 2)) W	'arning (Form): Config c1 (#3)	has 4 primary slit traces affe	ected by failed open shutters.								
ostic	(MQN01_Plan1 (Obs 2)) W	'arning (Form): Config c1 (#4)	has 4 primary slit traces affe	ected by failed open shutters.								
l so	(MQN01_Plan1 (Obs 2)) W	Varning (Form): Config c1 (#5)	has 2 master background sh	utters affected by failed open or clos	ed shutters.							
g	(MQN01_Plan1 (Obs 2)) Warning (Form): Config c1 (#5) has 2 primary slit traces affected by failed open shutters.											
Dia	(MQN01_Plan1 (Obs 2)) W	Varning (Form): Config c1 (#6)	has 4 primary slit traces affe	ected by failed open shutters.								
1	(MQN01_Plan1 (Obs 2)) W	'arning (Form): Config c1 (#7)	has 2 master background sh	utters affected by failed open or clos	ed shutters.							
	(MQN01_Plan1 (Obs 2)) W	Varning (Form): Config c1 (#7)	has 2 primary slit traces affe	ected by failed open shutters.								
<u> </u>	(Visit 2:1) Warning (Form):	Overheads are provisional un	til the Visit Planner has been	run.								
Targets	# Name	Target Co	ordinates	Targ. Coord. (Corrections	Miscellaneous						
Įğ	(2) MQN01-PA	45 RA: 00 41	32.5500 (10.3856250d)									
<u>a</u>		Dec: -49 3	6 20.95 (-49.60582d)									
Fixed	Comments: Description=[]											
late	TA Method	Obtain Confirmation Images	Science Aperture	Primary Candidate List	Filler Candidate List	Spectral Overlap Map	Spectral Overlap Threshold					
	NONE	No	MSA Center	MQN01-PA45 (29 sources)		jwst-nirspec-g235h	1.2					

Proposal 1835 - Observation 2 - Unraveling the knots of gaseous Cosmic Web filaments at z~3 through H-alpha emission observations

	#	Exposure Specification	MSA Configuration	Nod Pattern	Pointing	Dispersion Offset (Shutters)	Cross-Dispersion Offset (Shutters)	Total Dithers	Total Integrations	Total Exposure Time
	1	1 (G235H/F170LP)	c1	1 Shutter Slitlet	10.38745320833333 4 Degrees - 49.63508083333335 Degrees			1	3	4858.1
	2	1 (G235H/F170LP)	c1	1 Shutter Slitlet	10.38745320833333 4 Degrees - 49.63508083333335 Degrees		0.05	1	3	4858.1
Spectral Elements	3	1 (G235H/F170LP)	c1	1 Shutter Slitlet	10.38745320833333 4 Degrees - 49.63508083333335 Degrees		-0.05	1	3	4858.1
ectral El	4	1 (G235H/F170LP)	c1	1 Shutter Slitlet	10.38745320833333 4 Degrees - 49.63508083333335 Degrees		0.1	1	3	4858.1
Sp	5	1 (G235H/F170LP)	c1	1 Shutter Slitlet	10.38745320833333 4 Degrees - 49.63508083333335 Degrees		-0.1	1	3	4858.1
	6	1 (G235H/F170LP)	c1	1 Shutter Slitlet	10.38745320833333 4 Degrees - 49.63508083333335 Degrees		0.15	1	3	4858.1
	7	1 (G235H/F170LP)	c1	1 Shutter Slitlet	10.38745320833333 4 Degrees - 49.63508083333335 Degrees		-0.15	1	3	4858.1
Special Requirements			251.50765991 to 281.: egrees (V3 266.50765		765991000003)					