









# BMJ Open Factors associated with accessing and utilisation of healthcare and provision of health services for residents of slums in low and middle-income countries: a scoping review of recent literature

Ji-Eun Park <sup>1,2</sup>, Peter Kibe <sup>3</sup>, Godwin Yeboah <sup>4</sup>, Oyinlola Oyeboode,<sup>1</sup> Bronwyn Harris <sup>1</sup>, Motunrayo M Ajisola,<sup>5</sup> Frances Griffiths <sup>1,6</sup>, Navneet Aujla,<sup>1,7</sup> Paramjit Gill <sup>1</sup>, Richard J Lilford <sup>8</sup>, Yen-Fu Chen <sup>1</sup> The Improving Health in Slums Collaborative

**To cite:** Park J-E, Kibe P, Yeboah G, *et al*. Factors associated with accessing and utilisation of healthcare and provision of health services for residents of slums in low and middle-income countries: a scoping review of recent literature. *BMJ Open* 2022;**12**:e055415. doi:10.1136/bmjopen-2021-055415

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2021-055415>).

Received 12 July 2021

Accepted 25 March 2022



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY. Published by BMJ.

For numbered affiliations see end of article.

## Correspondence to

Dr Yen-Fu Chen;  
y-f.chen@warwick.ac.uk

## ABSTRACT

**Objective** To identify factors associated with accessing and utilisation of healthcare and provision of health services in slums.

**Design** A scoping review incorporating a conceptual framework for configuring reported factors.

**Data sources** MEDLINE, Embase, CINAHL, Web of Science and the Cochrane Library were searched from their inception to December 2021 using slum-related terms.

**Eligibility criteria** Empirical studies of all designs reporting relevant factors in slums in low and middle-income countries.

**Data extraction and synthesis** Studies were categorised and data were charted according to a preliminary conceptual framework refined by emerging findings. Results were tabulated and narratively summarised.

**Results** Of the 15 469 records retrieved from all years, 4368 records dated between 2016 and 2021 were screened by two independent reviewers and 111 studies were included. The majority (63 studies, 57%) were conducted in Asia, predominantly in India. In total, 104 studies examined healthcare access and utilisation from slum residents' perspective while only 10 studies explored provision of health services from providers/planners' perspective (three studies included both). A multitude of factors are associated with accessing, using and providing healthcare in slums, including recent migration to slums; knowledge, perception and past experience of illness, healthcare needs and health services; financial constraint and competing priorities between health and making a living; lacking social support; unfavourable physical environment and locality; sociocultural expectations and stigma; lack of official recognition; and existing problems in the health system.

**Conclusion** The scoping review identified a significant body of recent literature reporting factors associated with accessing, utilisation and provision of healthcare services in slums. We classified the diverse factors under seven broad categories. The findings can inform a holistic approach to improving health services in slums by tackling barriers at different levels, taking

## Strengths and limitations of this study

- ⇒ We conducted literature search in multiple databases using generic terms related to slums to ensure that a wide range of relevant studies was captured.
- ⇒ A conceptual framework explaining factors associated with accessing and utilisation of healthcare by slum residents as well as provision of healthcare in slums was developed and used to categorise identified studies and factors.
- ⇒ We examined barriers and facilitators of accessing healthcare and service provision from the perspectives of both demand side (slum residents) and supply side (healthcare providers and service planners).
- ⇒ Only studies published in academic journals between 2016 and 2021 in English language were included, and methodological quality of each included study was not examined because of time constraint.
- ⇒ We did not explore the complex relationships and interactions between various factors in different contexts at different slum locations, but our mapping of these factors to the conceptual framework should facilitate further in-depth analyses.

into account local context and geospatial features of individual slums.

**Systematic review registration number** <https://osf.io/694t2>.

## INTRODUCTION

Rapid urbanisation has resulted in a growing number of residents in slums<sup>1</sup> who face ongoing problems such as unemployment, poor sanitation, lack of transport, high level of crime and haphazard development.<sup>2</sup> In 2018, over 1 billion people were living in slum-like conditions, and Central, South and South-East Asia and sub-Saharan Africa

accounted for 80% of them.<sup>1</sup> Even though various definitions of slums exist, there is no universally agreed definition of what constitutes ‘a slum’, and the term itself is widely debated and contested.<sup>3,4</sup> For the purpose of this scoping review, we refer to slums as densely populated areas characterised by lack of basic services, substandard housing, overcrowding, unhealthy living condition, insecure tenure and poverty,<sup>4,5</sup> taking into account the crucial concepts of place and space that are important in shaping health outcomes and community access to health services in these urban settings.<sup>4</sup>

Previous studies have reported various risk factors affecting health of slum residents such as physical environment,<sup>6</sup> sanitation,<sup>7</sup> social capital<sup>8,9</sup> and water governance,<sup>10</sup> and have observed in some cases that slum residents have worse health status compared with non-slum urban and/or rural residents. For example, Ezeh *et al* found that children living in slums had higher mortality than rural and non-slum urban populations.<sup>3</sup> Poorer height for age for children<sup>11</sup> and higher prevalence of childhood illnesses and malnutrition<sup>12</sup> have also been observed in slums compared with non-slum urban and rural settings. In addition, slum residents are susceptible to unhealthy behaviours.<sup>13,14</sup> Living in slums has been found to be associated with low physical activity,<sup>13</sup> poor diet<sup>14</sup> and poor knowledge about the cause and preventability of diseases.<sup>15</sup>

Despite the unfavourable health status and environment, and consequently the potential high level of healthcare needs, previous studies showed that slum residents were less likely to seek and use healthcare services than their non-slum counterparts in the cities.<sup>16,17</sup> Slum residents have been found to have lower rates of healthcare utilisation in antenatal services<sup>16</sup> and services for non-communicable diseases<sup>17</sup> compared with residents of urban ‘formal’ settings. One study in Iran showed that only about half of slum households that required outpatient services could use them.<sup>18</sup> Another study in Haiti also reported that one-third of slum households were not able to access medical care for their children when it was needed in the past year.<sup>19</sup>

While the health status and needs of slum residents have been described in previous reviews,<sup>3,20</sup> factors associated with healthcare-seeking behaviour and healthcare utilisation of slum residents and factors related to the provision of health services in slums have not been systematically examined (with the exception of immunisation services).<sup>21</sup> This scoping review aims to fill in these evidence gaps and inform efforts to improve healthcare delivery to people in slums.

## METHODS

This scoping review was performed according to current best practice guidance.<sup>22</sup> The broad question of interest was: ‘*What factors are associated with slum residents’ accessing and utilisation of health care and/or the provision of health services in slum settings in low and middle income countries*

(LMICs)?’. The protocol for this review was registered in Open Science Framework.<sup>23</sup>

## Literature search and study selection

A broad search of five databases (MEDLINE, Embase, CINAHL, Web of Science and the Cochrane Library) was conducted in April 2020 and updated in December 2021. Searches were limited to English language. Key terms related to slums were used: slum or slums or ghetto or ghettos or informal settlement\$ or shantytown\$ or shanty town\$ or favela\$ (online supplemental appendix 1). We did not include terms related to other concepts in order to maximise the sensitivity of our searches. In addition, we searched the organisational websites of Slum Dwellers International, UN HABITAT, United Nations and WHO but did not identify relevant studies.<sup>24–27</sup>


Records retrieved from databases (after duplicates were removed) were initially screened by one reviewer (J-EP) and those which did not meet the inclusion criteria were disregarded. After that, a second reviewer (PK, GY, OO) examined the remaining records independently based on titles and abstracts. When the decisions of two reviewers differed, the discrepancy was resolved based on full texts and/or by discussion with a third reviewer (Y-FC) or the broader review team. This study-screening process started from records of the most recent years (ie, in the past 3 years) and then proceeded to prior years. Due to the larger than expected volume of the literature, we eventually screened records between 2016 and 2021 and did not cover earlier records in order to synthesise and present the findings from latest evidence in a timely fashion to inform the wider project hosting this review.<sup>28,29</sup>

## Inclusion and exclusion criteria

A study was included when it: (1) described factors related to slum residents’ accessing or utilisation of healthcare or the provision of health services in slums; and (2) was conducted in relation to slums in LMICs. Only articles written in English were included. A study was excluded when it was a commentary, opinion or narrative review; described slum residents’ utilisation of health services or the provision of health services without exploring the associated factors; investigated informal care at home; or included mixed slum and non-slum populations without separately reporting data for slum residents or investigating residency in slums as a factor for healthcare access.

During our updated search in December 2021, we found several studies reporting healthcare utilisation<sup>28</sup> and provision related to COVID-19 in slums.<sup>30</sup> These studies were not included in this scoping review, since the factors associated with healthcare utilisation and health service provision under the pandemic situation are dramatically different and warrant a separate synthesis.

We included both primary studies and systematic reviews that examine data collected empirically and that derive their findings based on the data. Both quantitative and qualitative studies (and by extension, mixed methods studies) were considered. Even though slums have existed



**Figure 1** Preliminary framework for factors influencing slum residents' healthcare-seeking behaviour and utilisation of health services and the provision of services in slum settings.


in both high-income countries and LMICs, the context may be quite different between these countries. For example, while all slums are vulnerable to natural disasters such as tropical cyclones, the impact of these could be far more severe in slums of LMICs due to the different socioeconomic contexts.<sup>31</sup> In this review, we focused on settings in LMICs and excluded studies conducted in high-income countries.

#### Study coding and data extraction/charting

Eligible studies were coded and data extracted/charted according to a prespecified preliminary framework shown in figure 1. The preliminary framework was developed by the review authors based on existing conceptual models

related to healthcare access and service delivery<sup>32–35</sup> and was modified during the scoping review process to accommodate new factors/themes identified from the literature. The refined conceptual framework is shown in figure 2.

Based on the refined conceptual framework, each eligible study was coded as being associated with one or more of the three phenomena of interest, namely slum residents' healthcare accessing (which covered perception of needs/desire for care, healthcare seeking and healthcare reaching as defined by Levesque *et al*<sup>34</sup>), healthcare utilisation and provision of health services (which covered various arrangements related to service delivery) in slum settings (figure 2).



**Figure 2** Updated framework of factors influencing healthcare-seeking behaviour/healthcare utilisation/provision of healthcare services in slums.

In addition, to facilitate the organisation of complex evidence in this review, diverse factors reported in the included studies were initially classified into six different categories according to the preliminary framework shown in [figure 1](#). However, during the data-charting process, we realised that many factors such as perception of symptoms and experience from past use of healthcare services did not fit into one of these six original categories. A new category of ‘cognitive and experiential factors’ was therefore added to the refined conceptual framework ([figure 2](#)) to reflect the emerging themes, which include seven categories:

1. Personal and biological factors: these relate to personal characteristics of slum residents, including age, sex, ethnicity and the nature and severity of health conditions.
2. Cognitive and experiential factors: these relate to personal awareness, knowledge, perception, attitude, belief and experience, etc, formed through cognitive process based on upbringing and past events.
3. Socioeconomic factors: these include income and wealth, economic hardship/poverty and economic opportunities, marital status, education, crime, social capital (such as bonding, trust and reciprocity between close relatives, neighbours and community members),<sup>36</sup> use of technologies for social and economic purposes, commercial and charitable organisations and activities.
4. Physical environment: this covers natural environment such as proximity to a health facility, built environment and infrastructure such as water supply, transport and mobile/internet networks, as well as weather conditions and environmental pollutions.
5. Cultural and religious factors: these include cultural and religious beliefs and activities, and local and national customs.
6. Legal, political and policy factors: these include government policies and issues related to legal, justice and political systems.
7. Health system factors: these relate to historical and current organisation and provision of healthcare that may impact on provision and delivery of health services in individual slum communities and the services experienced by slum residents.

In addition to the ‘cognitive and experiential factors’ category, another major difference between the preliminary ([figure 1](#)) and refined ([figure 2](#)) conceptual framework relates to the definition of healthcare access. Our preliminary framework adopted the definition by Levesque and colleagues, who defined healthcare access as ‘the possibility to identify healthcare needs, to seek healthcare services, to reach the healthcare resources, to obtain or use health care services, and to actually be offered services appropriate to the needs for care’.<sup>34</sup> However, during our study-screening and data-charting process, we found that it would be helpful to make a distinction between the process of ‘accessing’ healthcare (which covers gaining awareness of needs, forming

an intention to seek healthcare and taking an action to reach healthcare) and the actual receipt and utilisation of healthcare (‘accessed care’) when examining empirical evidence, as healthcare needs could only be met when the latter occurs and this depends on both factors related to service users (demand side) and factors related to service providers/planners (supply side). Therefore, we separated out utilisation of healthcare from ‘accessing health care’ to highlight that it requires a match between demand and supply side factors.

Data on study population, study design, country in which the study was conducted, methodology and associated factors were extracted using a data-charting spreadsheet which was developed and continuously updated as the review progressed by two of the reviewers (J-EP and Y-FC). Whether a study was conducted exclusively within slums and whether a comparison was made between slum and non-slum urban or rural residents were also noted. Coding of phenomena and factors and data charting were conducted by one reviewer (J-EP) and checked by a second reviewer (PK, GY, OO, Y-FC). Disagreements were discussed between reviewers until consensus was reached.


### Patient and public involvement

Given the focus of this scoping review on published literature, we did not directly involve residents and service providers/planners from slum settings. Nevertheless, our wider project has a work package that specifically engages with slum residents and service providers and planners,<sup>28</sup> and early plans and findings of this review were shared with the wider project team who provided comments based on their experiences of community engagement.

## RESULTS

The reporting of this review follows the Preferred Reporting Items for Systematic Reviews and Meta-Analyses extension for Scoping Reviews.<sup>37</sup> Using the search strategy described earlier, a total of 15 469 records were retrieved from the initial and updated searches (MEDLINE 4688, Embase 5090, Web of Science 3553, Cochrane 381, CINAHL 1757), with 9916 records remaining after excluding duplicates. Two additional articles<sup>18 38</sup> were identified from references of the included studies. As described earlier, screening was limited to the 4368 records published from 2016 onwards.

A total of 111 articles were included in this scoping review ([figure 3](#)). Thirty-two studies reported factors associated with healthcare accessing of slum residents, 73 studies reported factors related to healthcare service utilisation and 10 articles reported the factors related to provision of healthcare services in slums (four studies reported factors related to more than one phenomenon of interest). Seventy-four of the 111 studies were quantitative studies, 21 studies were qualitative studies and 14 studies were undertaken using mixed methods. The remaining two studies were systematic reviews. A total of



**Figure 3** Flow chart.

42 (38%) studies were conducted in India, followed by Kenya (14 studies, 13%) (table 1).

Participants, country, study design, methodology, observed phenomena and outcomes, and factors of interests for each study are described in online supplemental tables 1–3. Online supplemental table 1 shows 32 studies reporting factors associated with general healthcare-seeking behaviours; healthcare seeking for children or women; slum residents’ preference for healthcare providers; and healthcare seeking related to HIV testing. Online supplemental table 2 presents various factors reported in 73 studies related to general healthcare utilisation as well as use of specific services such as childhood immunisation, maternal healthcare and possession of health insurance. In online supplemental table 3, ten studies reporting factors related to the provision of health services in slums are summarised. Key findings are described below.

**Demand side: factors associated with healthcare accessing and healthcare utilisation of slum residents**

We found 104 articles which identified many different factors affecting healthcare accessing and utilisation. These factors are often inter-related and exert their influence at different levels (eg, from personal, family to community level) in different circumstances. We classified various factors into seven categories (figure 2). Factors particularly relevant to slum settings and other commonly identified factors within each category are highlighted below.

**Personal and biological factors:** The common factors associated with healthcare accessing and utilisation included intrinsic factors such as age,<sup>21 39–55</sup> sex<sup>18 21 41 45 51 53 55–58</sup>

**Table 1** Characteristics of included studies

Category	Subcategory	Studies, n (%)
Publication year	2016	22 (20)
	2017	17 (15)
	2018	23 (21)
	2019	22 (20)
	2020	18 (16)
	2021	9 (8)
Analysis method	Quantitative	74 (67)
	Qualitative	21 (19)
	Mixed methods	14 (13)
	Narrative synthesis	2 (2)
Study location	Asia	
	India	42 (38)
	Bangladesh	9 (8)
	Nepal	4 (4)
	Pakistan	3 (3)
	Myanmar	2 (2)
	Iran	2 (2)
	Sri Lanka	1 (1)
	South America	
	Brazil	7 (6)
	Peru	2 (2)
	Africa	
	Kenya	14 (13)
	Ethiopia	7 (6)
	Malawi	4 (4)
	Uganda	3 (3)
South Africa	2 (2)	
Sierra Leone	1 (1)	
Nigeria	1 (1)	
Egypt	1 (1)	
Zambia	1 (1)	
Namibia	1 (1)	
Ghana	1 (1)	
North America		
Haiti	1 (1)	
Multiple nations	1 (1)	
Healthcare services in slums*	Healthcare accessing	32
	Healthcare service utilisation	73
	Provision of healthcare services	10
Total		111 (100)

\*One study reported factors related to both healthcare accessing and healthcare utilisation and three studies reported factors related to both healthcare utilisation and provision of healthcare services.

and ethnicity,<sup>21</sup> familial factors such as birth order of the sick child,<sup>21 46 59–61</sup> as well as personal health and type of illness,<sup>45 58</sup> disability<sup>47</sup> and morbidity,<sup>21 51 62 63</sup> and the specific features of the health condition.<sup>52 55 64</sup> Slum residents are more likely to seek healthcare services when sick



children are younger,<sup>48 49 52 55</sup> but evidence on the association between mother's age and child's vaccination was inconsistent.<sup>21 39</sup> Healthcare seeking and utilisation were different by sex, but the association was context dependent. Several studies reported higher healthcare utilisation among female slum dwellers,<sup>18 51 55 57</sup> while other studies showed male children had higher vaccination coverage<sup>56</sup> and incurred more medical expenditure.<sup>58</sup> Major life events such as recent migration<sup>21 49 65-67</sup> and relocation<sup>48</sup> into slums tend to be associated with lower healthcare seeking and utilisation. Recency of migration to slums was also related to lower uptake of Rashtriya Swasthya Bima Yojana (RSBY), a national health insurance programme run by the Indian government for poor families.<sup>68</sup> People with specific symptoms (such as fever, tachypnoea, persistent vomiting),<sup>52 55 64</sup> disability<sup>47</sup> and illnesses including chronic disease<sup>21 51 53 62 63</sup> tend to use healthcare services more. Although people with tobacco habit were less likely to participate in breast cancer screening, they were more likely to take part when they had family history of cancer or history of cancer screening.<sup>69</sup> Lower birth order of the child was associated with increased utilisation of hospitals for childbirth,<sup>21 46 59-61</sup> while the use of family planning service<sup>46</sup> and out-of-pocket expenditure was higher in multigravida than primigravida.<sup>62</sup>

**Cognitive and experiential factors:** These factors were not included in our initial conceptual framework but rather emerged inductively from our data. Consequently, their identification led us to revise the conceptual framework for this scoping review. A wide range of factors formed through cognitive processes and influenced by individual's upbringing, past experience and surrounding environment were reported to be associated with both healthcare seeking and healthcare utilisation of slum residents. Perception,<sup>39 50 52 67 70-76</sup> knowledge<sup>52 77-82</sup> and experience of symptoms and illnesses<sup>50</sup> were commonly found to influence healthcare seeking and utilisation. Mothers who experienced child death and subsequently planned pregnancy showed higher use of antenatal healthcare services.<sup>50</sup> When people perceived the symptom or disease to be serious they tend to seek healthcare services.<sup>52 70 72 76</sup> Although lack of knowledge could be a barrier to accessing healthcare services,<sup>77 78</sup> one study showed caregivers with good knowledge of child danger signs were less likely to seek healthcare services timely.<sup>52</sup> People perceiving their health status as good showed lower odds of having insurance,<sup>83</sup> but awareness and knowledge of health problems led people to use healthcare services.<sup>39 73 75 79 80 82</sup> Home remedy and home management delayed healthcare-seeking behaviour.<sup>52 70 81 84</sup> In addition, perception,<sup>21 42 63 75 83 85-89</sup> knowledge<sup>21 42 48 60 67 74 81 85 90-95</sup> and experience of healthcare services<sup>39 49 59 61 66 69 74 79 86 96 97</sup> including fear and distrust of healthcare services,<sup>21 38 67 71 74 75 78 88 98-100</sup> and preference related to care provider's gender<sup>87 101</sup> were frequently cited factors. Provider shopping associated with distrust of healthcare providers and denial of diagnosis delayed first care seeking and treatment initiation of

patients with pulmonary tuberculosis in India.<sup>70</sup> Perception or experience of healthcare services also affected uptake or renewal of health insurance.<sup>83 94</sup>

**Socioeconomic factors:** Socioeconomic status was associated with utilisation of healthcare services,<sup>21 39 40 46 56 61 81 102</sup> and even though one study showed that slum residents of lower socioeconomic class were more likely to enrol in health insurance than slum residents of higher socioeconomic class,<sup>40</sup> the latter were more likely to use healthcare services.<sup>39 46 56 61</sup> One study reported higher public hospital visits (compared with private hospital visits) among lower socioeconomic status.<sup>103</sup> Income and wealth<sup>21 41 48-50 60 65 67 80 104-108</sup> including financial constraint<sup>19213845737678848689100102109-113</sup> featured prominently. Higher education level<sup>39 48 55-58 60 61 65-67 69 80-82 106-108 114-116</sup> and higher income<sup>21 48-50 55 57 60 65 67 80 104 106-108 117</sup> were associated with more seeking and utilisation of healthcare services. With some exceptions,<sup>66 104</sup> previous studies reported that employed slum residents tend to seek and use healthcare services more frequently than unemployed slum residents and housewives.<sup>48 50 65 66 69 80 82 83 118</sup> Even though married people tend to seek and use more healthcare services,<sup>18 69</sup> the reported influence of family type was inconsistent.<sup>39 50 115</sup> Female slum residents in nuclear family used more antenatal services than those in joint family type,<sup>50</sup> but female slum residents in joint family type used more postnatal service<sup>39</sup> and immunisation service for their children.<sup>115</sup> Smaller family size used more maternal healthcare services,<sup>66</sup> and bigger households had higher odds of having health insurance.<sup>68</sup> The socioeconomic challenges faced by slum residents also manifested as competing priorities<sup>73 93 119</sup> and lack of time<sup>21 100 120</sup> for healthcare seeking and utilisation, because they did not want to or could not afford to miss work and lose income,<sup>21 70</sup> which can be exacerbated by lack of social support.<sup>74 76 87 93 99 121</sup>

**Physical environment.** Slum residents considered proximity of healthcare facilities,<sup>21 38 46 76 80 84 85 95 108 122-126</sup> transport such as travel assistance,<sup>76</sup> lack of transportation,<sup>38 82 102</sup> traffic congestion<sup>127</sup> and environment of residence area when they sought and used healthcare services. Long distance from health facility,<sup>38 54 76 84 101 122</sup> no transportation or travel assistance<sup>38 82 102</sup> and unsafe environment of residential area such as darkness at night were reported as barriers to reaching healthcare facilities.<sup>99</sup>

**Cultural and religious factors:** These included religion<sup>41 56 59 69 75 114 128</sup>; sociocultural influence<sup>93 101</sup> such as exposure to media<sup>79 97</sup>; stigma associated with unplanned/extramarital pregnancy,<sup>78 93</sup> postpartum depression<sup>129</sup> and other illnesses such as contagious skin disease, barrenness and female sexually related problems<sup>101</sup>; and use of traditional/home medicine.<sup>75 86 99</sup> Women in slums could not go to hospital because they had difficulties in disclosing the symptoms, postponed their health issues because of their responsibilities at home and engaged in self-treatment practices such as home remedies recommended by grandmother and friends because

of sociocultural influences towards healthcare-seeking behaviour.<sup>77</sup> Women in Ethiopia reported not returning to postnatal care due to religious and cultural expectation for the mother and the baby to stay home for 80 days after birth.<sup>93</sup> One Indian survey showed that some women could not seek healthcare services during labour since their husband or family did not allow that.<sup>38</sup>

**Legal, political and policy factors:** Type of slums (in terms of official recognition and availability of basic facilities) and possession of a ration card were found to be associated with uptake of the Indian RSBY national health insurance programme.<sup>68</sup> One study reported that slum residents could not seek healthcare facilities for abortion because of the perceived illegality of abortion.<sup>78</sup>

**Health system factors:** Slum residents were also influenced by many factors related to health systems when they sought healthcare. These included accessibility associated with the location<sup>21 101</sup> and timing of services<sup>21 84 86</sup>; quality of healthcare services<sup>38 66 85 86 101 112 124</sup> such as delay in advising patients to go for related tests or referral,<sup>70</sup> likelihood of receiving appropriate examination<sup>84 109</sup> and adverse events.<sup>75</sup> Slum residents considered service organisation including medical turnover,<sup>130</sup> availability of supplies/healthcare workers,<sup>46 84 109 112</sup> attitude of healthcare providers,<sup>86</sup> type of healthcare facilities<sup>38 85 116 131</sup> and waiting time.<sup>72 84-86 109 111 132</sup> Slum residents tend to seek government and non-governmental organisation facility<sup>117</sup> and avoid private hospitals<sup>38</sup> for healthcare services. Healthcare utilisation was higher among slum residents with healthcare insurance than those without it,<sup>18 82</sup> and households with higher quarterly out-of-pocket healthcare expenditure had lower scores for an index of access to primary healthcare.<sup>133</sup>

In an Ethiopian study, some participants reported unavailability of female birth attendants as a reason for not delivering at healthcare facilities<sup>112</sup> (table 2).

### Supply side: provision of healthcare services

Ten articles described factors associated with provision of healthcare services in slums from the service providers' perspective. None of the studies reported personal and biological factors. Factors related to other categories are summarised below.

**Cognitive and experiential factors:** Odhiambo *et al* reported slum residents' fear of side effects, size of tablet and misconceptions regarding treatment as the factors hindering drug administration activities by healthcare workers for a deworming programme in Kenya.<sup>134</sup> On the other hand, this study also reported a high demand for drugs from slum residents in the final year of this programme because people realised that free treatment was to be ended.<sup>134</sup>

**Socioeconomic factors:** Effective community mobilisation was a facilitator<sup>134</sup> whereas poor community support<sup>135</sup> and insufficient time allocated for providers to implement healthcare programmes<sup>134</sup> were barriers for provision of healthcare services in slums. In the deworming programme mentioned above, community

health workers reported that direct observation of slum residents taking deworming drugs after meals was sometimes not feasible because slum residents skipped meals or ate late at night due to food shortage.<sup>134</sup> Some slum residents demanded money to take the deworming drugs, either to facilitate purchase of food or to have their own share of the money that they perceived the community health workers would be paid by the programme if they complied with taking the drugs.<sup>134</sup>

**Physical environment:** Poor sanitation,<sup>134 136</sup> presence of rodents and no pavement<sup>136</sup> and bushy and unprotected environment<sup>134</sup> were reported as factors making the provision of healthcare services difficult in slums.

**Cultural and religious factors:** Religious beliefs and mistrust of interventions,<sup>134</sup> lack of a shared understanding of the needs, purposes and consequences of family planning and pregnancy-related services among slum residents and healthcare providers<sup>73</sup> were the barriers for healthcare service provision. In the previous deworming programme, portrayal of unrelated death being linked to the programme and related negative publicity affected participants' compliance.<sup>134</sup>

**Legal and political factors and policy:** Devolution of service delivery through downward transfer of funds and responsibilities from central/national government level to elected local bodies; management by professional managerial and technical cadres; tight organisation of public health services; and professional support from the state directorate of public health were found to strengthen public health service provision in Chennai slums compared with Delhi.<sup>137</sup> One study reported that policies affected healthcare provision negatively because of staff shortage arising from change and suspension of the appointment of health promoters, which led to overwork and lack of time to provide required care by other healthcare staff.<sup>132</sup> In Brazil, home visits for the provision of healthcare services were hampered because slum residents could not present documents required to register for healthcare.<sup>136</sup> On the other hand, giving priority to socially less developed areas for strengthening the Family Health System in Brazil might have been associated with better service coverage for slum residents with tuberculosis compared with their urban non-slum counterparts.<sup>138</sup>

**Health system factors:** Pay scale of frontline healthcare workers,<sup>135</sup> knowledge of intervention area by community health workers,<sup>134</sup> issues related to rigid task assignment by service managers,<sup>136</sup> requirement to follow standardised protocol,<sup>136</sup> demands from the management,<sup>136</sup> work burden<sup>132 136</sup> and no incentive,<sup>135</sup> insufficient time,<sup>134</sup> attitude<sup>73</sup> and support of healthcare providers,<sup>135</sup> ill-defined geographical boundary of service with unserved areas and left-out urban slum pockets<sup>139</sup> were associated with healthcare service provision in slums.

Lack of community-based care (such as school-based education for reproductive health and community support networks for women),<sup>132</sup> unreliable immunisation and household data<sup>128</sup>; and inefficient utilisation of funds,<sup>128</sup> affordability (price) and availability of

**Table 2** Factors associated with healthcare accessing and healthcare utilisation in slums from service user's (demand side) perspective

Factors	Healthcare accessing	Healthcare utilisation
Personal and biological factors		
Age	(-) Age <sup>52-55</sup>	(±) Age <sup>21 39-51</sup> ; (+) age of household head <sup>18</sup>
Gender	(±) Sex <sup>53 55 57 58</sup>	(±) Sex <sup>18 21 41 45 51 56 58</sup> ; (male) sex of household head <sup>133</sup>
Ethnicity		Ethnicity <sup>21</sup>
Migration		(-) Recent migration <sup>21 49 65-68</sup> ; (-) relocation <sup>48</sup> ; (-) return to home village <sup>21</sup>
Biological	(+) Symptoms such as fever, tachypnoea, chest in drawing, persistent vomiting <sup>52 55 64</sup> ; having disease <sup>53</sup>	Type of illness <sup>45 58</sup> ; (+) having a disability <sup>47</sup> ; (+) morbidity <sup>21 51 62 63</sup>
Other personal	(-) Tobacco habits <sup>69</sup> ; (+) family history of cancer and history of cancer screening <sup>69</sup>	(-) Birth order of sick child <sup>21 46 59-61</sup> ; (-) parity <sup>42 46 62 148</sup>
Cognitive and experiential factors		
Knowledge/experience of symptoms and illnesses	(+) Perception of symptoms <sup>70</sup> or illness <sup>52 72 76</sup> ; (±) knowledge of symptom/disease <sup>52 77 78</sup> ; (-) denial and complacency <sup>71</sup>	(+) Experience of child death <sup>50</sup> ; (+) planned pregnancy <sup>50</sup> ; (+) perceived health status <sup>83</sup> and health problem <sup>39 73 75</sup> ; (+) knowledge of symptom <sup>82</sup> ; disease <sup>79 80</sup>
Ability/experience in handling health-related conditions and perceived needs for accessing health services	(+) Awareness of the need for healthcare services <sup>38 53 55</sup> ; (-) home remedies <sup>70</sup> or management of childhood illness <sup>52 84</sup>	(+) Perceived needs for healthcare services <sup>21 67 74 75 90 93 112 128</sup> ; (-) home delivery <sup>81</sup>
Perception/knowledge/experience/preference of health services	(-) Fear of mistreatment <sup>71 100</sup> and (-) doubts about medical care <sup>38 78 99</sup> ; gender-induced affordability <sup>101</sup> ; (-) provider shopping <sup>70</sup>	(positive) Perception of healthcare services <sup>21 42 63 75 83 85-89</sup> and providers <sup>21 84 93 111 124</sup> ; (+) knowledge of health services <sup>21 42 48 60 67 74 81 85 90-93</sup> or facilities <sup>21 94 95</sup> ; (+) previous use of related healthcare services <sup>39 42 49 59 61 66 79 96 97</sup> ; (-) bad experiences of friends and relatives at healthcare facilities <sup>94</sup> ; (-) misunderstanding or fear <sup>21 67 74 75 88 98 128</sup> ; gender healthcare worker preference <sup>87</sup> ; (-) side effect <sup>81</sup> ; lack of trust <sup>45</sup>
Socioeconomic factors		
Socioeconomic status	(-) Social class <sup>103</sup> ; social group (caste) of caregiver <sup>53</sup>	(+) Socioeconomic status <sup>21 39 40 46 56 61 81 102</sup> ; caste <sup>108 114</sup> ; (-) insecure or poor residential background <sup>21 46 68 81</sup> ; (+) possession of ration card <sup>68</sup>
Marital status	(married) Marital status <sup>69</sup>	(married) Marital status <sup>18 41</sup> ; duration of marriage <sup>42</sup>
Family composition and living arrangement	(-) Family size <sup>53</sup>	(±) Family type <sup>39 50 115 126</sup> ; (±) family size <sup>66 68</sup> ; (-) number of children in household <sup>21 44 48</sup> ; (+) number of male children <sup>149</sup> ; (+) housing condition <sup>21</sup>
Education	(+) Education <sup>53 55 57 69</sup>	(+) Education <sup>39 41-43 48 56 58 60 61 65-67 80-82 106-108 114-116 126 148</sup> ; (±) husband education <sup>44 50</sup> ; (+) mother's education and literacy <sup>21 43 46 56 59</sup>

Continued



Table 2 Continued

Factors	Healthcare accessing	Healthcare utilisation
Income and wealth	(+) Income <sup>55 117</sup> ; (+) wealth <sup>54 57</sup> ; (-) inability to afford care <sup>19 38 76 78 84 89 100 109 110</sup>	(+) Income <sup>41 49 50 67 80 104</sup> ; (+) wealth <sup>21 48 60 65 105-108</sup> ; (-) financial constraint <sup>21 45 73 86 102 111-113</sup>
Occupation	(+) Occupation <sup>53 57 69</sup>	(+) Employment <sup>21 65 83 118 126</sup> ; (±) occupation <sup>48 66 80 82 104 148</sup> ; (±) occupation of spouse <sup>50 60</sup> or household <sup>68</sup>
Social support	(-) Difficulty in reaching services (security risk at night) <sup>99</sup> ; (+) accompanying person <sup>76</sup> ; decision-making person for seeking healthcare <sup>53</sup>	(+) Family support <sup>74 87</sup> ; (+) social connectedness <sup>93</sup> ; (+) socioeconomic support <sup>121</sup> ; permission for immunisation by decision-maker <sup>128</sup>
Competing priorities/lack of time	(-) Competing priorities (ability to work and income) <sup>119</sup> ; (-) not want to miss work <sup>70</sup> ; (-) lack of time <sup>100 120</sup>	(-) Competing priorities <sup>73 81 93</sup> ; (-) risk of lost income <sup>21</sup> ; (-) parents being too busy <sup>21</sup>
Physical environment		
Distance from health facility	Proximity of healthcare facilities <sup>38 54 76 84 122</sup> ; geographical distance of formal healthcare <sup>101</sup>	(-) Distance from health facility <sup>21 46 80 85 95 108 123-126</sup>
Transport	(+) Travel assistance <sup>76</sup> ; (-) no transportation <sup>38</sup>	(-) Lack of transportation <sup>82 95 102</sup> ; (-) variability in traffic congestion <sup>127</sup>
Environment of residence area	(-) Difficulty in reaching services (darkness at night) <sup>99</sup>	Residential background <sup>21 68 105</sup>
Cultural and religious factors		
Religion	Religion <sup>69</sup>	Religion <sup>41 56 59 75 114 128</sup>
Sociocultural influence	(-) Stigma <sup>78 101 129</sup> ; mother tongue <sup>69</sup> ; (-) difficulties in disclosing the symptoms, (-) neglecting behaviours and sociocultural influences <sup>77</sup> ; (+) cultural competency of care <sup>101</sup> ; (+) easy communication <sup>101</sup> ; living with the burden of cultural expectations <sup>101</sup> ; (-) no permission to seek care from family <sup>38</sup>	(-) Exposure to media <sup>79 97</sup> ; stigma <sup>128</sup> ; (-) cultural expectation for women after birth and fear of stigma for pregnancy out of wedlock <sup>93</sup>
Tradition	(-) Traditional medicine <sup>99</sup>	(-) Traditional remedies <sup>75</sup> ; (-) home remedies <sup>86</sup>
Legal, political and policy factors		
Legal issues	(-) Perceived illegality of abortion <sup>78</sup>	Type of slums and possession of a ration card <sup>79</sup>
Health system factors		
Accessibility	(+) Ease of access <sup>101</sup> ; (-) late facility opening times <sup>84</sup>	(-) Limited access to the services due to location <sup>90 93</sup> ; (-) timing of services <sup>21 86</sup> ; household visit by health workers <sup>21</sup>
Quality and safety of services	Quality of treatment and expected outcome of therapies <sup>38 101</sup> ; (-) delay in advising related tests <sup>70</sup> ; referral <sup>70</sup> ; optimal examination <sup>84 109</sup> ; (-) provider shopping <sup>70</sup>	Quality of service <sup>66 85 86 91 112 124</sup> ; (-) adverse events <sup>75</sup>
Charges for health services	(+) Insurance coverage of both public and private providers and of extended family members <sup>150</sup>	(-) Average out-of-pocket healthcare expenditure <sup>133</sup> ; healthcare insurance <sup>18 63 82</sup>

Continued



Table 2 Continued

Factors	Healthcare accessing	Healthcare utilisation
Service organisation and delivery arrangement	(-) Medical turnover and overload or healthcare providers <sup>130</sup> ; (+) government/NGO facility <sup>117</sup> ; (-) private hospital <sup>38</sup> ; early engagement by healthcare workers <sup>54</sup>	Attitude of healthcare providers <sup>86</sup> <sup>95</sup> ; mode of delivery <sup>39 56 59 62 81 151</sup> ; (-) hospitals refused to accept health insurance cards <sup>94</sup>
Facility and resources	Availability of medicines and supplies <sup>84 109</sup> ; (-) lack of healthcare facilities <sup>144</sup>	Type of healthcare facility <sup>40 85 95 116 131</sup> <sup>133</sup> ; inadequate resources <sup>90</sup> ; (+) number of available healthcare workers <sup>46</sup> ; (-) unavailability of female birth attendants <sup>112</sup>
Waiting time	(-) Waiting time <sup>72 84 109</sup>	(-) Waiting time <sup>85 86 111 132</sup>

(-) Negative association; (±) inconsistent/conflicting evidence or context dependent; (+) positive association. NGO, non-governmental organisation.

medicine,<sup>140</sup> limited medical supplies<sup>73 135</sup> and infrastructural facilities,<sup>135</sup> inadequate space and equipment,<sup>136 139</sup> suboptimal training of staff,<sup>139</sup> insufficient availability of logistics and health manpower<sup>139</sup> also affected service provision (table 3).

### Comparison between slums and other settings

Seven studies which met our inclusion criteria also included data from non-slum urban and/or rural areas and potentially allowed exploration of factors associated with healthcare access across different settings. Key findings from these studies are summarised in table 4.

These recent studies showed a mixed and dynamic picture of healthcare access across slum and other settings and reported various factors associated with this. For example, the proportion of young children fully immunised was found to be lower in slums compared with non-slum urban setting but was higher than rural settings in Nigeria. Nevertheless, the coverage improved over time across all settings.<sup>59</sup> While many common factors associated with full immunisation of young children were identified, giving birth in health facilities (as opposed to home) had a larger positive effect on subsequent immunisation coverage in slums compared with non-slum urban and rural settings.<sup>59</sup> A narrowing of gaps in delivery by skilled birth attendants between slum and non-slum urban settings over time and a reverse of the trend from having lower usage to higher usage of modern contraceptive methods by married women in slums versus urban non-slums were reported in Bangladesh.<sup>46</sup> Slum residents reported financial issues being the main reason for not taking prescribed drugs whereas getting better was the cited main reason for urban non-slum residents in Iran.<sup>113</sup> Better coverage of services and higher rates of treatment completion were reported for patients with tuberculosis in slums compared with non-slum urban setting in two studies in Brazil,<sup>51 138</sup> where a higher priority given to enhancing the Family Health System in socially less developed areas in recent years was suggested to be

a likely factor associated with better service provision in slums<sup>138</sup> (table 4).

## DISCUSSION

### Statement of principle findings

This scoping review of recent literature examined the demand side factors associated with slum residents' healthcare accessing and utilisation, as well as supply side factors associated with provision of health services in slums. We found over 104 studies related to the former but only 10 studies related to the latter. We identified different factors associated with accessing, utilisation and provision of health services in slums, and mapped them to a conceptual framework developed and refined for this review into seven broad categories (figure 2).

### Findings in the context of existing literature

Even though previous reviews have investigated factors associated with healthcare access in various settings,<sup>141 142</sup> to our best knowledge this scoping review is the first that has examined wide-ranging factors across different service areas of healthcare in slums. Our findings are consistent with previous studies which highlighted common factors associated with healthcare seeking and utilisation such as age, income and education.<sup>141 143</sup> We identified several factors that are particularly pertinent in slum settings, such as costs of healthcare,<sup>19 21 73 76 78 84 89 100 102 109–111</sup> lack of time due to slum residents' competing priorities<sup>21 100 120</sup> and issues arising from adverse physical environment,<sup>82 102 134 136</sup> security,<sup>99 136</sup> fear of formal registration due to distrust of the authorities<sup>136</sup> and proximity of healthcare facilities.<sup>21 76 80 84 85 108 122–125</sup> In addition, included studies showed that the effects of a given factor may differ between slum, urban non-slum and rural settings.<sup>59</sup>

Healthcare cost is a major barrier between the intention to seek care and actual utilisation of services.<sup>109 144</sup> Healthcare provision supported by tax-based financing and/or various forms of social and private insurance

**Table 3** Factors associated with provision of healthcare services in slums from service provider's (supply side) perspective

Cognitive and experiential factors	
Perception/knowledge/experience/preference of health services	Fear of side effects, size of tablet and misconceptions regarding treatment, high demand for drugs in the final year of treatment <sup>134</sup>
Socioeconomic factors	
Income and wealth	Difficulty in directly observing deworming treatment at mealtime due to food shortage <sup>134</sup>
Social support	Effective community mobilisation <sup>134</sup> ; poor community support <sup>135</sup> ; non-involvement of community members and urban local bodies <sup>139</sup> ; absence of community members during the drug administration exercise <sup>134</sup> ; demand for incentives by community members to take deworming drugs <sup>134</sup>
Physical environment	
Environment of residence area	Environment (sanitation, territory) <sup>136</sup> ; unsanitary environmental conditions <sup>134</sup> ; inaccessibility (filthy and bush environment) <sup>134</sup>
Cultural and religious factors	
Religion	Religious beliefs and mistrust of interventions <sup>134</sup>
Sociocultural influence	Lack of shared understanding of the problems in community <sup>73</sup> ; unrelated death and the associated negative publicity (of a deworming programme) by the media <sup>134</sup>
Legal, political and policy factors	
Policy issues	Devolution of service delivery transferring funds and responsibilities to elected local bodies <sup>137</sup> ; management by professional managerial and technical cadres <sup>137</sup> ; tight organisation of public health services <sup>137</sup> ; professional support from the state directorate of public health <sup>137</sup> ; healthcare policies <sup>132</sup> ; policy prioritising low social development areas <sup>138</sup>
Legal issues	Fear of requirement for formal registration <sup>136</sup>
Health system factors	
Cost	Pay scale of frontline healthcare workers <sup>135</sup> ; medicine price <sup>140</sup>
Quality and safety of services	Knowledge of intervention area by community health workers <sup>134</sup>
Service organisation and delivery arrangement	Issues related to assignment of tasks <sup>136</sup> ; requirement to follow standardised protocol <sup>136</sup> ; demands from the management <sup>136</sup> ; work overload <sup>132 136</sup> ; underperformance of staff <sup>128</sup> ; documentation work/work burden/no incentive for work <sup>135</sup> ; insufficient time <sup>134</sup> ; attitude of healthcare providers <sup>73</sup> ; lack of supportive staff <sup>135</sup> ; community health worker familiarity with households led to warm reception <sup>134</sup> ; opportunity to integrate mass drug administration with other health interventions <sup>134</sup> ; presence of community health workers and their supervisory structure, and points of referral for serious side effects <sup>134</sup> ; restriction of range of services <sup>139</sup> ; unserved areas and left-out urban slum pockets <sup>139</sup> ; poor monitoring and supervision <sup>139</sup> ; unreliable immunisation and household data <sup>128</sup>
Facility and resources	Community-based care <sup>132</sup> ; inefficient utilisation of funds <sup>128</sup> ; affordability and availability of medicine <sup>140</sup> ; limited medical supplies <sup>73 135</sup> ; infrastructural facilities <sup>135</sup> ; inadequate space and equipment <sup>136</sup> ; suboptimal training of staff <sup>139</sup> ; insufficient availability of space, logistics and health manpower <sup>139</sup>

that reduce out-of-pocket expenditure at point of care could be potential measures to overcome this barrier and help achieve universal coverage goals. Limited evidence showed that initiatives prioritising primary healthcare coverage in slums could improve access,<sup>138</sup> but there is insufficient evidence from studies included in this review to determine the best model of healthcare financing for improving healthcare access and coverage in slum settings.

Although possession of/coverage by health insurance was associated with higher levels of utilisation of health services among slum residents,<sup>18 82</sup> studies showed that uptake of government-run public insurance among slum residents was low.<sup>68 83</sup> This may be attributed to lack of awareness, difficulties in navigating through the health system and in obtaining official proof of identity required for enrolment<sup>68</sup> and poor quality of care and range of services offered.<sup>68 83</sup> Even among slum residents covered

**Table 4** Studies that examined factors associated with healthcare seeking and utilisation in both urban slum and non-slum urban and rural settings

Study and location	Differences in healthcare access	Associated factors
Kalyango <i>et al</i> <sup>150</sup> Kampala City, Uganda	<i>Preferences and willingness to pay for health insurance</i> Households in non-slum communities had a high preference for health insurance plans covering chronic illnesses and major surgeries to other plans.	Coverage of extended family (vs restricted enrolment of children); coverage of both private and public providers (vs private only).
Obanewa and Newell <sup>59</sup> Nationwide, Nigeria	<i>Fully immunised child coverage (FIC)</i> Proportion in slum lower than urban non-slum but higher than rural; proportions increased between 2003 and 2013 across all three settings.	From multivariable regression*: year, birth order, antenatal attendance, maternal education level, religion, maternal age at child's birth, media exposure, region of the country, interaction between place of residence and place of delivery.
Angeles <i>et al</i> <sup>46</sup> Multiple cities, Bangladesh	<i>Use of modern contraceptive methods</i> Proportion changed from being lower in slums in 2006 to being higher in slums in 2013 compared with urban non-slums.  <i>Delivery by skilled birth attendant</i> Proportion substantially lower in slums compared with urban non-slums but the gaps narrowed over time.	From multivariable regression*: parity, mother's age, mother's educational attainment, socioeconomic status, interaction (slum×time period).  From multivariable regression*: residing in slums, parity, mother's age, mother's educational attainment, length of stay in current city of residence, socioeconomic status, number of available community health workers, distance from health facility, interaction (slum×time period).
Islam <sup>106</sup> Multiple cities, Bangladesh	<i>Antenatal care visits</i> 'there was a large inequality' between slum and urban non-slum (detail not reported).  <i>Using contraceptive methods</i> 'Prevalence rate higher among slum women' than urban non-slum women.	Level of educational attainment, wealth index of the household.  Not reported.
Tabrizi <i>et al</i> <sup>113</sup> Tabriz, Iran	<i>Utilisation of health services in the past 30 days</i> Similar utilisation overall, but with lower proportion received needed health services and used private clinics, higher use of vaccination and maternal health services, and lower use of services for heart failure and hypertension for slum residents compared with urban non-slum.  <i>Home care services</i> Very little use both in slum and urban non-slum areas.  <i>Prescribed drug during last visit to health facilities</i> Lower proportion for slum versus urban non-slum.  <i>Not taking drugs prescribed</i> Higher proportion for slum versus urban non-slum.	High cost of services.  High cost of services.  Not reported.  Main reason: financial problems for slum versus getting better/feeling well for non-slum urban.
Snyder <i>et al</i> <sup>51</sup> Rio de Janeiro, Brazil	<i>Directly observed treatment coverage for tuberculosis (TB)</i> Higher for slum versus urban non-slum patients with TB.  <i>Abandonment of TB treatment</i> Lower for slum versus urban non-slum patients with TB.	Not examined.  From multivariable regression*: residency in a slum, sex, age, extrapulmonary clinical disease, HIV/AIDS, interaction (directly observed treatment×residency in a slum).
Prado Junior <i>et al</i> <sup>138</sup> Rio de Janeiro, Brazil	<i>Coverage under Family Health System for patients with TB</i> Higher for slum versus urban non-slum.	Giving the Family Health Strategy priority to coverage of areas with lower social development.

\*From the model with most comprehensive adjustment including residency in slum as one of the variables; only factors that were statistically significant (at 5% level) are shown.

by health insurance, access to care was often refused and additional charges were frequently requested.<sup>94</sup> Policies that aim to improve access to healthcare services among slum residents through public health insurance will need to address these challenges.

Several studies reported lack of time and competing priorities as a factor affecting healthcare-seeking behaviour<sup>100 119 120</sup> and health service utilisation.<sup>21 73 93</sup>

This suggests a delicate balance between factors that individual slum residents have to strike when making decisions on healthcare seeking and utilisation. van der Heijden *et al* showed that health was considered as an asset for working ability in slums,<sup>119</sup> but paradoxically the ability to work often seems to impede healthcare seeking for health issues. This highlights the importance of considering slum residents' interest and priorities when

providing healthcare services and promoting healthcare utilisation in slums.

### Strengths and weaknesses of the review

This scoping review has several strengths. We conducted a comprehensive literature search using generic terms related to slums with few other restrictions. The search was therefore likely to be sensitive for identifying relevant literature. Contemporary methodological guidelines for undertaking scoping reviews were followed,<sup>22</sup> and a conceptual framework which was adapted based on emerging findings was used to facilitate the organisation of evidence.

The review has enabled theory building and refinement of a conceptual framework. Our preliminary framework included six categories (figure 1). During data coding and extraction, it emerged that many studies reported perception, knowledge and experience of slum residents being associated with their healthcare seeking and utilisation. We subsequently classified these factors as cognitive and experiential factors, which primarily consist of three subcategories: knowledge/experience of illness, perceived needs for accessing healthcare services and perception/experience of healthcare services. These factors were influenced by other factors included in our original conceptual framework, but highlighted the crucial links between those factors and the ultimate actions by individual slum residents to access health services. Future interventions to promote health service utilisation for slum residents<sup>145</sup> could make use of our framework to develop programme theories and map out causal pathways.

This review also has some limitations. Given time constraint, we were only able to examine the most recent literature published in English in academic journals, and have not examined the methodological quality of individual studies (which we noted to be quite varied) in detail. We attempted some preliminary synthesis to configure the identified evidence but have not explored the complex relationship between the factors identified and their interplay with the context of individual slums in depth. Nevertheless, the findings from this scoping review will provide a good foundation for further syntheses.

### Methodological considerations

A number of challenges in the process of classifying and coding data are worth mentioning. First, access to healthcare has been conceptualised and defined in various ways in previous studies. The WHO suggested six building blocks of a health system including service delivery, health workforce, health information systems, access to essential medicines, financing and leadership/governance to strengthen health systems<sup>146</sup> and, in its report, defined access to healthcare as public responsibility for ensuring all citizens' entitlements to the protection of their health beyond simply a proportion of a target population that benefits from an intervention, towards universal coverage.<sup>146</sup> They also pointed

out system constraints such as financial access difficulty, physical access difficulty, low knowledge and skills, poorly motivated staff, weak leadership and management, ineffective intersectoral action and partnership as barriers to access.<sup>146</sup> The WHO's definition and conceptual framework focus on health system-level factors and would be particularly useful when examining supply side factors, which seem to be understudied based on our findings. As described in the Methods section, we primarily adopted the conceptual model of healthcare access developed by Levesque and colleagues given our shared focus on service users. However, in our conceptual model, we separated the dynamic stages of 'accessing' healthcare from the actual 'accessed' healthcare (utilisation) to highlight the crucial match required between the demand side and supply side factors to facilitate utilisation of healthcare when there is a need.

Several factors associated with healthcare accessing and utilisation can be viewed from different perspectives and therefore potentially be coded under different categories. For example, barriers for healthcare seeking and utilisation related to costs can be considered as socioeconomic issues from the slum dwellers' perspective but can also be viewed as health system issues for not offering the services in an affordable way. Indeed, previous access frameworks suggested that access is created and negotiated in a dynamic interchange between households/communities and healthcare workers/systems (ie, demand and supply) on each access dimension.<sup>34 147</sup> In such situations, we tried to code a factor under the category that most directly reflects the original data through discussions within the review team (in the example of healthcare cost, the factor was coded primarily under socioeconomic factors rather than health system factors when the factor was reported by slum residents as a barrier); otherwise, more than one category was coded (eg, bad experience from previous utilisation of health services was coded both as a cognitive and experiential factor and a health system factor).

### Implication for research and practice

The multitude of factors identified in this review are often inter-related and interacting, and span across personal, family, community and society levels. For example, the association between occupation and healthcare utilisation was reported in several studies.<sup>48 60 66 80 82 104</sup> The effect of predominantly casual work undertaken by slum residents on their healthcare access could be mediated through working hours, income level, knowledge of health and available services, etc. There is also possibility that occupation was associated with health status and hence needs for healthcare services, instead of/in addition to behaviour of using healthcare services. Teasing out the complicated relationships between various determinants and their interaction with the diverse contexts of slums will require in-depth analysis and a more holistic approach to synthesising the evidence. Given the unique features of individual slums, service planners and policy makers will need to examine these relationships with due

consideration to the context specific to each locality and geospatial features and neighbourhood effects that characterise slum settings.<sup>4</sup>

We found far fewer studies that have examined health service providers' perspective than studies that have investigated factors associated with accessing healthcare from slum residents' perspective. There may be scope for greater research and policy attention to supply side factors, including experiences and practices of local frontline healthcare providers, availability of healthcare facilities and infrastructure and policy to support them in order to overcome the many barriers highlighted from both supply and demand sides.

Although only six of the included studies explored factors associated with healthcare access or health service provision across slum and non-slum settings, they showed a generally encouraging picture that access to and provision of healthcare are continuously evolving (and often improving) in slums and other settings, and equality between different settings is not beyond reach.

## CONCLUSION

This scoping review summarises a large body of recent literature evaluating factors associated with seeking and utilisation of healthcare by slum residents, but found substantially fewer studies examining factors associated with provision of health services from providers' perspective. Recent migration into slums; knowledge, perception (including misconception and distrust) and past experience of illness, healthcare needs and health services; financial constraint, competing priorities and inadequacy of social support; adverse physical environment and unfavourable locality; sociocultural expectations and stigma; lack of official recognition; and various problems in existing health system all contribute towards the challenges faced by slum residents. Future research and policy aimed at improving healthcare services in slums should pay more attention to supply side issues ranging from individual healthcare providers and practices to structural and policy-level factors to tackle different barriers faced by slum residents, which in turn need to be evaluated holistically and take into account local context and geospatial features of slums.

### Author affiliations

<sup>1</sup>Warwick Medical School, University of Warwick, Coventry, UK

<sup>2</sup>KM Data Division, Korea Institute of Oriental Medicine, Daejeon, Republic of Korea

<sup>3</sup>Health and Systems for Health, African Population and Health Research Center, Nairobi, Kenya

<sup>4</sup>Information and Digital Group, University of Warwick, Coventry, UK

<sup>5</sup>Sociology, University of Ibadan, Ibadan, Nigeria

<sup>6</sup>Centre for Health Policy, University of the Witwatersrand, Johannesburg-Braamfontein, South Africa

<sup>7</sup>Population Health Sciences Institute, Newcastle University, Newcastle upon Tyne, UK

<sup>8</sup>Institute of Applied Health Research, University of Birmingham, Birmingham, UK

**Twitter** Godwin Yeboah @GodwinYeboah and Yen-Fu Chen @yfchen12

**Collaborators** The Improving Health in Slums Collaborative: African Population and Health Research Centre (APHRC), Nairobi, Kenya: Pauline Bakibinga, Caroline Kabaria, Ziraba Kasiira, Peter Kibe, Lyagamula Kisia, Catherine Kyobutungi, Nelson Mbaya, Blessing Mberu, Shukri Mohammed, Anne Njeri. Aga Khan University, Karachi, Pakistan: Iqbal Azam, Romaina Iqbal, Ahsana Nazish, Narjis Rizvi. Independent University, Dhaka, Bangladesh: Syed Shifat Ahmed, Nazratun Choudhury, Omar Rahman, Rita Yusuf. Nigerian Academy of Sciences, Lagos, Nigeria: Doyin Odubanjo. University of Ibadan, Ibadan, Nigeria: Motunrayo Ayobola, Olufunke Fayeahun, Akinyinka Omigbodun, Mary Osuh, Eme Owoaje, Olalekan Taiwo. University of Birmingham, Birmingham, UK: Richard Lilford, Jo Sartori, Samuel Watson. University of Lancaster, Lancaster, UK: Peter Diggie. University of Warwick, Coventry, UK: Navneet Aujla, João Porto de Albuquerque, Yen-Fu Chen, Paramjit Gill, Frances Griffiths, Bronwyn Harris, Jason Madan, Oyinlola Oyeboade, Ji-Eun Park, Simon Smith, Grant Tregonning, Olalekan Uthman, Ria Wilson, Godwin Yeboah.

**Contributors** J-EP, BH, MMA, FG and Y-FC conceptualised the scoping review. J-EP carried out literature searches. J-EP, PK, GY, OO and Y-FC participated in study screening and coding. J-EP and Y-FC performed data charting and drafted the initial manuscript. NA, PG and RJL provided critical input during the drafting of the manuscript. All authors commented on and contributed to the revision of subsequent versions and approved the final version for submission. Y-FC is the guarantor for this article.

**Funding** This research is funded by the NIHR Global Health Research Unit on Improving Health in Slums using UK aid from the UK government to support global health research. MMA gratefully acknowledges the support provided by the Warwick Institute of Advanced Study Global Challenges Research Fund Fellowship (IAS/32013/1914). FG receives funding as South Africa Research Chair in Health Policy and Systems from the National Research Foundation, South Africa. RJL is supported by the NIHR Applied Research Collaboration (ARC) West Midlands, UK. Y-FC is supported by Warwick Evidence, which is a Technology Assessment Review team funded by the NIHR Evidence Synthesis Programme. PG is NIHR Senior Investigator and supported by the NIHR ARC West Midlands. Upon submission, NA had joined the Population Health Sciences Institute, Newcastle University (UK).

**Disclaimer** The views expressed in this publication are those of the author(s) and not necessarily those of the NIHR or the UK Department of Health and Social Care.

**Competing interests** None declared.

**Patient and public involvement** Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

**Patient consent for publication** Not required.

**Ethics approval** This realist synthesis included literature that is available in the public domain and did not involve the collection of personal data.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Data availability statement** All data relevant to the study are included in the article or uploaded as supplementary information.

**Supplemental material** This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy, redistribute, remix, transform and build upon this work for any purpose, provided the original work is properly cited, a link to the licence is given, and indication of whether changes were made. See: <https://creativecommons.org/licenses/by/4.0/>.

### ORCID iDs

Ji-Eun Park <http://orcid.org/0000-0002-2932-5373>

Peter Kibe <http://orcid.org/0000-0002-9027-9054>

Godwin Yeboah <http://orcid.org/0000-0003-4618-3175>

Bronwyn Harris <http://orcid.org/0000-0003-4695-008X>

Frances Griffiths <http://orcid.org/0000-0002-4173-1438>

Paramjit Gill <http://orcid.org/0000-0001-8756-6813>

Richard J Lilford <http://orcid.org/0000-0002-0634-984X>

## REFERENCES

- United Nations. *The sustainable development goals report*, 2019.
- Kosamkar A. Problems in urban society. *EPRA Int J Multidiscip Res* 2020;6.
- Ezeh A, Oyebo O, Satterthwaite D, et al. The history, geography, and sociology of slums and the health problems of people who live in slums. *Lancet* 2017;389:547–58.
- Lilford R, Kyobutungi C, Ndugwa R, et al. Because space matters: conceptual framework to help distinguish slum from non-slum urban areas. *BMJ Glob Health* 2019;4:e001267.
- Zulu EM, Beguy D, Ezeh AC, et al. Overview of migration, poverty and health dynamics in Nairobi City's slum settlements. *J Urban Health* 2011;88:S185–99.
- Michiani MV, Asano J. Physical upgrading plan for slum riverside settlement in traditional area: a case study in Kuin Utara, Banjarmasin, Indonesia. *Front Archit Res* 2019;8:378–95.
- Winter SC, Dreifelbis R, Dzombo MN, et al. A mixed-methods study of women's sanitation utilization in informal settlements in Kenya. *PLoS One* 2019;14:e0214114.
- Salinas DA, Fouts HN, Neitzel CL, et al. Young children's social networks in an informal urban settlement in Kenya: examining network characteristics among Kamba, Kikuyu, Luo, and Maasai children. *J Cross Cult Psychol* 2019;50:639–58.
- Prayitno G, Sari N, Putri IK. Social capital in poverty alleviation through PRO-POOR tourism concept in slum area (case study: Kelurahan jodipan, Malang City). *GEOMATE* 2019;16:131–7.
- Khalil D. The flexible governance of water in Cairo's informal areas. *Water* 2019;11:1644.
- Pörtner CC, Su Y-H. Differences in child health across rural, urban, and slum areas: evidence from India. *Demography* 2018;55:223–47.
- Mberu BU, Haregu TN, Kyobutungi C, et al. Health and health-related indicators in slum, rural, and urban communities: a comparative analysis. *Glob Health Action* 2016;9:33163.
- Mlangeni L, Makola L, Naidoo I, et al. Factors associated with physical activity in South Africa: evidence from a national population based survey. *Open Public Health J* 2018;11:516–25.
- Nunn RL, Kehoe SH, Chopra H, et al. Dietary micronutrient intakes among women of reproductive age in Mumbai slums. *Eur J Clin Nutr* 2019;73:1536–45.
- Samal J. Perception and knowledge of tuberculosis and its services among slum dwellers in Chhattisgarh. *Indian J Respir Care* 2017;6:828–31.
- Khan MZ, Shujaa MD, Iftikhar H. Utilization of ante-natal services among reproductive age women of Bahawalpur. *Indo Am J Pharm Sci* 2018;5:11355–65.
- Duy Kien V, Van Minh H, Bao Giang K, et al. Horizontal inequity in public health care service utilization for non-communicable diseases in urban Vietnam. *Glob Health Action* 2014;7:24919.
- Amiresmaili M, Yazdi-Feyzabadi V, Heidarijamebozorgi M. Health services utilization among slum dwellers: an experience from Iran. *J Educ Health Promot* 2019;8:210.
- McNairy ML, Tymejczyk O, Rivera V, et al. High burden of non-communicable diseases among a young slum population in Haiti. *J Urban Health* 2019;96:797–812.
- Sverdluk A. Ill-Health and poverty: a literature review on health in informal settlements. *Environ Urban* 2011;23:123–55.
- Crocker-Buque T, Mindra G, Duncan R, et al. Immunization, urbanization and slums - a systematic review of factors and interventions. *BMC Public Health* 2017;17:556.
- Peters MDJ, Godfrey C, Mclnerney P. Chapter 11: Scoping reviews. In: Aromataris E, Munn Z, eds. *Joanna Briggs Institute reviewer's manual*, 2017.
- Center for Open Science. Slum health healthcare access and provision of services scoping review protocol 05082020.pdf (version: 1), 2020. Available: <https://osf.io/mj6kp/> [Accessed 9 Jun 2021].
- United Nations. United nations 2021. Available: <https://www.un.org/en> [Accessed 25 Dec 2021].
- World Health Organization. World Health organization 2021. Available: <https://www.who.int> [Accessed 25 Dec 2021].
- SDI. SDI 2021. Available: <https://sdinet.org> [Accessed 25 Dec 2021].
- UN HABITAT. UN-Habitat for a better urban future 2021. Available: <https://unhabitat.org/> [Accessed 25 Dec 2021].
- Ahmed SAKS, Ajisola M, Azeem K, et al. Impact of the societal response to COVID-19 on access to healthcare for non-COVID-19 health issues in slum communities of Bangladesh, Kenya, Nigeria and Pakistan: results of pre-COVID and COVID-19 lockdown stakeholder engagements. *BMJ Glob Health* 2020;5:e003042.
- Improving Health in Slums Collaborative. A protocol for a multi-site, spatially-referenced household survey in slum settings: methods for access, sampling frame construction, sampling, and field data collection. *BMC Med Res Methodol* 2019;19:109.
- George CE, Inbaraj LR, Rajukutty S, et al. Challenges, experience and coping of health professionals in delivering healthcare in an urban slum in India during the first 40 days of COVID-19 crisis: a mixed method study. *BMJ Open* 2020;10:e042171.
- Peduzzi P, Chatenoux B, Dao H, et al. Global trends in tropical cyclone risk. *Nat Clim Chang* 2012;2:289–94.
- Effective practice and organisation of care (EPOC). EPOC taxonomy, 2015. Available: <https://epoc.cochrane.org/epoc-taxonomy> [Accessed 27 Jul 2020].
- Lavis JN, Wilson MG, Moat KA, et al. Developing and refining the methods for a 'one-stop shop' for research evidence about health systems. *Health Res Policy Syst* 2015;13:10.
- Levesque J-F, Harris MF, Russell G. Patient-centred access to health care: conceptualising access at the interface of health systems and populations. *Int J Equity Health* 2013;12:18.
- National Cancer Institute. *Theory at a glance: a guide for health promotion practice*, 2005.
- Mpanje D, Gibbons P, McDermott R. Social capital in vulnerable urban settings: an analytical framework. *Int J Humanitarian Action* 2018;3:4.
- Tricco AC, Lillie E, Zarin W, et al. PRISMA extension for scoping reviews (PRISMA-ScR): checklist and explanation. *Ann Intern Med* 2018;169:467–73.
- Pune Municipal Corporation. City health plan 2016 - 2020, 2016. Available: <https://www.healthynewbornnetwork.org/hnn-content/uploads/Pune-City-Health-Plan.pdf> [Accessed 10 Jun 2021].
- Upadhyai N, Gupta SK. Utilization of postnatal care services and factors affecting it among women of urban slums in Dehradun, Uttarakhand. *Indian J Community Health* 2019;31:470–6.
- Mendhe HG, David R, Singh D, et al. Universal health insurance coverage and utilization among women in urban slum of Rajnandgaon, Chhattisgarh. *J Family Med Prim Care* 2021;10:1313–9.
- Iyalomhe FO, Adekola PO, Cirella GT. Community-based health financing: empirical evaluation of the socio-demographic factors determining its uptake in Awka, Anambra state, Nigeria. *Int J Equity Health* 2021;20:235.
- Yadav K, Agarwal M, Shukla M, et al. Unmet need for family planning services among young married women (15-24 years) living in urban slums of India. *BMC Womens Health* 2020;20:187.
- de Araújo Veras AAC, da Fonseca Lima EJ, Caminha MdeFC, et al. Vaccine uptake and associated factors in an irregular urban settlement in northeastern Brazil: a cross-sectional study. *BMC Public Health* 2020;20:1152.
- Roja VR, Narayanan P, Sekaran VC, et al. Living environment and health of under-five children in urban slums of a coastal region in South India. *Ghana Med J* 2020;54:238–44.
- Vora K, Saiyed S, Shah AR, et al. Surgical unmet need in a low-income area of a metropolitan City in India: a cross-sectional study. *World J Surg* 2020;44:2511–7.
- Angeles G, Ahsan KZ, Streatfield PK, et al. Reducing inequity in urban health: have the Intra-urban differentials in reproductive health service utilization and child nutritional outcome narrowed in Bangladesh? *J Urban Health* 2019;96:193–207.
- Renzaho AMN, Kamara JK, Georgeou N, et al. Sexual, reproductive health needs, and rights of young people in slum areas of Kampala, Uganda: a cross sectional study. *PLoS One* 2017;12:e0169721.
- Hong L, Kakoly NS, Abedin J, et al. Effect of household relocation on child vaccination and health service utilisation in Dhaka, Bangladesh: a cross-sectional community survey. *BMJ Open* 2019;9:e026176.
- Lae WL, Jayasvasti I, Mongkolchati A. Utilization of immunization service and predictors among under 3-year-old children in urban slums of Chanmyathazi township, Mandalay, Myanmar. *J Med Assoc Thai* 2018;101:1085–92.
- Sharma D, Pokharel HP, Budhathoki SS, et al. Antenatal health care service utilization in slum areas of Pokhara Sub-Metropolitan City, Nepal. *J Nepal Health Res Council* 2016;14:39–46.
- Snyder RE, Marlow MA, Phuphanich ME, et al. Risk factors for differential outcome following directly observed treatment (DOT) of slum and non-slum tuberculosis patients: a retrospective cohort study. *BMC Infect Dis* 2016;16:494.
- Lungu EA, Darker C, Biesma R. Determinants of healthcare seeking for childhood illnesses among caregivers of under-five children in

- urban slums in Malawi: a population-based cross-sectional study. *BMC Pediatr* 2020;20:20.
- 53 Mohanty P, Satpathy SK, Patnaik S, *et al*. Do the caregivers of under-fives have proper health care seeking behaviour for their children? A study from urban slums of a City in eastern India. *J Nepal Paediatr Soc* 2021;41:169–76.
- 54 Pakhare A, Joshi A, Anwar R, Kumar S, *et al*. Linkage to primary-care public health facilities for cardiovascular disease prevention: a community-based cohort study from urban slums in India. *BMJ Open* 2021;11:e045997.
- 55 Kerai S, Nisar I, Muhammad I, *et al*. A community-based survey on health-care utilization for pneumonia in children in Peri-Urban slums of Karachi, Pakistan. *Am J Trop Med Hyg* 2019;101:1034–41.
- 56 Devasenapathy N, Ghosh Jerath S, Sharma S, *et al*. Determinants of childhood immunisation coverage in urban poor settlements of Delhi, India: a cross-sectional study. *BMJ Open* 2016;6:e013015.
- 57 Kar SS, Kalaiselvi S, Archana R, *et al*. Is rule of halves still an occurrence in South India: findings from community-based survey in a selected urban area of Puducherry. *J Postgrad Med* 2017;63:232–6.
- 58 Mishra S, Kusuma YS, Babu BV. Treatment-seeking and out-of-pocket expenditure on childhood illness in a migrant tribal community in Bhubaneswar, Odisha state, India. *Paediatr Int Child Health* 2017;37:181–7.
- 59 Obanewa OA, Newell ML. The role of place of residency in childhood immunisation coverage in Nigeria: analysis of data from three DHS rounds 2003–2013. *BMC Public Health* 2020;20:123.
- 60 Shrestha S, Shrestha DK. Utilization of maternal health care services among mothers residing at slum area. *J Nepal Health Res Counc* 2019;17:193–9.
- 61 Nayaz A, Ahmed MS, Sahu PC. Preference and practices regarding place of childbirth in the slums of a City in northern India. *J Pioneer Med Sci* 2016;6:33–6.
- 62 Sahu KS, Bharati B. Out-Of-Pocket health expenditure and sources of financing for delivery, postpartum, and neonatal health in urban slums of Bhubaneswar, Odisha, India. *Indian J Public Health* 2017;61:67–73.
- 63 Wambiya EOA, Otieno PO, Mutua MK, *et al*. Patterns and predictors of private and public health care utilization among residents of an informal settlement in Nairobi, Kenya: a cross-sectional study. *BMC Public Health* 2021;21:850.
- 64 Mishra S, Kusuma YS, Babu BV. Mother's recognition of and treatment triggers for common childhood illnesses among migrant Santal tribe living in Bhubaneswar, Odisha, India. *J Trop Pediatr* 2017;63:301–6.
- 65 Razzaque A, Clair K, Chin B, *et al*. Association of time since migration from rural to urban slums and maternal and child outcomes: Dhaka (North and South) and Gazipur City Corporations. *J Urban Health* 2020;97:158–70.
- 66 Geddam JB, Ponna SN, Kommu PR. Utilization of maternal health services by the migrant population living in the non-notified slums of Hyderabad City, India. *Indian J Community Health* 2017;29:29–38.
- 67 Singh S, Sahu D, Agrawal A, *et al*. Ensuring childhood vaccination among slums dwellers under the National Immunization Program in India - Challenges and opportunities. *Prev Med* 2018;112:54–60.
- 68 Kusuma YS, Pal M, Babu BV. Health insurance: awareness, utilization, and its determinants among the urban poor in Delhi, India. *J Epidemiol Glob Health* 2018;8:69–76.
- 69 Kulkarni S, Mishra G, Dussane R. Determinants of compliance to breast cancer screening and referral among women from urban slums in India. *Eur J Cancer* 2016;54:S51.
- 70 Mistry N, Rangan S, Dholakia Y, *et al*. Durations and delays in care seeking, diagnosis and treatment initiation in uncomplicated pulmonary tuberculosis patients in Mumbai, India. *PLoS One* 2016;11:e0152287.
- 71 Thomson KA, Telfer B, Opondo Awiti P, *et al*. Navigating the risks of prevention of mother to child transmission (PMTCT) of HIV services in Kibera, Kenya: barriers to engaging and remaining in care. *PLoS One* 2018;13:e0191463.
- 72 Kamati M, Godman B, Kibuule D. Prevalence of self-medication for acute respiratory infections in young children in Namibia: findings and implications. *J Res Pharm Pract* 2019;8:220–4.
- 73 Kaba M, Taye G, Getachew S. Perceived barriers to health care for residents in vulnerable urban centers of Ethiopia. *Ethiop J Health Dev* 2020;34:4–11.
- 74 Verma H, Sagili KD, Zachariah R, *et al*. Do incentivised community workers in informal settlements influence maternal and infant health in urban India? *Public Health Action* 2017;7:61–6.
- 75 Pugliese-Garcia M, Heyerdahl LW, Mwamba C, *et al*. Factors influencing vaccine acceptance and hesitancy in three informal settlements in Lusaka, Zambia. *Vaccine* 2018;36:5617–24.
- 76 Ramagiri R, Kannuri NK, Lewis MG, *et al*. Evaluation of whether health education using video technology increases the uptake of screening for diabetic retinopathy among individuals with diabetes in a slum population in Hyderabad. *Indian J Ophthalmol* 2020;68:S37–41.
- 77 Evangeline Goonewardena C, Fernandopulle R. Women's understanding and cultural practices related to vaginal discharge: a qualitative study. *Nurs Midwifery Stud* 2018;7:74–80.
- 78 Jayaweera RT, Ngui FM, Hall KS, *et al*. Women's experiences with unplanned pregnancy and abortion in Kenya: a qualitative study. *PLoS One* 2018;13:e0191412.
- 79 Kusuma YS, Kaushal S, Garg R, *et al*. Birth preparedness and determinants of birth place among migrants living in slums and slum-like pockets in Delhi, India. *Sex Reprod Healthc* 2018;16:160–6.
- 80 Adane M, Mengistie B, Mulat W, *et al*. Utilization of health facilities and predictors of health-seeking behavior for under-five children with acute diarrhea in slums of Addis Ababa, Ethiopia: a community-based cross-sectional study. *J Health Popul Nutr* 2017;36:9.
- 81 Shrestha S, Shrestha M, Wagle RR, *et al*. Predictors of incompleteness of immunization among children residing in the slums of Kathmandu Valley, Nepal: a case-control study. *BMC Public Health* 2016;16:970.
- 82 Owusu-Ansah FE, Tagbor H, Togbe MA. Access to health in City slum dwellers: the case of Sodom and Gomorrah in Accra, Ghana. *Afr J Prim Health Care Fam Med* 2016;8:e1–7.
- 83 Otieno PO, Wambiya EOA, Mohamed SF, *et al*. Prevalence and factors associated with health insurance coverage in resource-poor urban settings in Nairobi, Kenya: a cross-sectional study. *BMJ Open* 2019;9:e031543.
- 84 Lungu EA, Biesma R, Chirwa M, *et al*. Healthcare seeking practices and barriers to accessing under-five child health services in urban slums in Malawi: a qualitative study. *BMC Health Serv Res* 2016;16:410.
- 85 Owiti A, Oyugi J, Essink D. Utilization of Kenya's free maternal health services among women living in Kibera slums: a cross-sectional study. *Pan Afr Med J* 2018;30:86.
- 86 Madan NV. Ethnographic perspectives on slum-dwelling women's access to primary care: the case of Pune, India. *Urbanities* 2019;9:114–30.
- 87 Wairiuko JM, Cheboi SK, Ochieng GO. Access to healthcare services in informal settlement: perspective of the elderly in Kibera slum Nairobi-Kenya. *Ann Med Health Sci Res* 2017;7:5–9.
- 88 Martínez Pérez G, Cox V, Ellman T, *et al*. 'I Know that I Do Have HIV but Nobody Saw Me': Oral HIV Self-Testing in an Informal Settlement in South Africa. *PLoS One* 2016;11:e0152653.
- 89 Sudhinaraset M, Beyeler N, Barge S, *et al*. Decision-making for delivery location and quality of care among slum-dwellers: a qualitative study in Uttar Pradesh, India. *BMC Pregnancy Childbirth* 2016;16:148.
- 90 Sendo EG, Chauke ME, Ganga-Limando M. Why some women who attend focused antenatal care fail to deliver in health facilities: a qualitative study of women's perspectives from slums of Addis Ababa, Ethiopia. *BMJ Open* 2020;10:e039189.
- 91 Sendo EG, Chauke ME, Ganga-Limando M. Women's perspectives on the measures that need to be taken to increase the use of health-care facility delivery service among slums women, Addis Ababa, Ethiopia: a qualitative study. *Reprod Health* 2021;18:174.
- 92 Manandhar K, Bajcharya K, Prajapati R, *et al*. Prevalence and predictors of incomplete immunization among children residing in the slums of Kathmandu Valley: a community based door-to-door survey. *Kathmandu Univ Med J* 2018;16:8–13.
- 93 Kaba M, Taye G, Gizaw M. Maternal health service utilization in urban slums of selected towns in Ethiopia: qualitative study. *Ethiop J Health Dev* 2017;31:96–102.
- 94 Gupta S. Awareness and utilization of Rashtriya Swasthaya bimA Yojana and its implications for access to health care by the poor in slum areas of Delhi. *Health Syst* 2017;6:242–59.
- 95 Chauhan M, Saxena S. Barriers in utilization of public health services by elderly slum dwellers in Jaipur City. *Indian J Public Health Res Develop* 2020;11:730–6.
- 96 Oluoch P, Orwa J, Lugalia F, *et al*. Application of psychosocial models to home-based testing and counseling (HBTC) for increased uptake and household coverage in a large informal urban settlement in Kenya. *Pan Afr Med J* 2017;27:285.



- 97 Atusiimire LB, Waiswa P, Atuyambe L, *et al.* Determinants of facility based-deliveries among urban slum dwellers of Kampala, Uganda. *PLoS One* 2019;14:e0214995.
- 98 Iberico MM, Montoya R, Valiente B, *et al.* Uptake and utilization of shantytown tuberculosis preventive therapy in a Peruvian Peri-urban shantytown. *Ann Glob Health* 2016;82:366.
- 99 Hutain J, Perry HB, Koffi AK, *et al.* Engaging communities in collecting and using results from verbal autopsies for child deaths: an example from urban slums in Freetown, Sierra Leone. *J Glob Health* 2019;9:010419.
- 100 Misra V, Vashist P, Singh SS, *et al.* Awareness and eye health-seeking practices for cataract among urban slum population of Delhi: the North India eye disease awareness study. *Indian J Ophthalmol* 2017;65:1483–8.
- 101 Das M, Angeli F, Krumeich AJSM, *et al.* The gendered experience with respect to health-seeking behaviour in an urban slum of Kolkata, India. *Int J Equity Health* 2018;17:24.
- 102 Sheehy G, Aung Y, Foster AM. "She learned it from her mother and grandmother": women's experiences with delivery and post-partum practices in peri-urban Yangon, Myanmar. *Matern Child Health J* 2016;20:854–61.
- 103 Angeli F, Ishwardat ST, Jaiswal AK, *et al.* Socio-cultural sustainability of private healthcare providers in an Indian slum setting: a bottom-of-the-pyramid perspective. *Sustainability* 2018;10:4702.
- 104 Rahman S. A comparative study of intrauterine contraceptive device utilization among currently married women in a rural area of Rani block and urban slums of Guwahati City. *Int J Sci Study* 2016;4:55–9.
- 105 Mutua MK, Mohamed SF, Iddi S, *et al.* Do inequalities exist in the disadvantaged populations? Levels and trends of full and on-time vaccination coverage in two Nairobi urban informal settlements. *Glob Epidemiol* 2020;2:100044.
- 106 Islam M. Use of reproductive health care services among urban slum women in Bangladesh. *Eur J Public Health* 2018;28:67.
- 107 Jolly SP, Rahman M, Afsana K, *et al.* Evaluation of maternal health service indicators in urban slum of Bangladesh. *PLoS One* 2016;11:e0162825.
- 108 Sadhna S, Kajal J, Debabratta R, *et al.* Utilisation of maternal health services and its predictors in slum population. *Acta Medica International* 2016;3:56–62.
- 109 Lungu EA, Guda Obse A, Darker C, *et al.* What influences where they seek care? Caregivers' preferences for under-five child healthcare services in urban slums of Malawi: a discrete choice experiment. *PLoS One* 2018;13:e0189940.
- 110 Nasrin M, Sarker MNI, Huda N. Determinants of health care seeking behavior of pregnant slums dwellers in Bangladesh. *Med Sci* 2019;23:35–41.
- 111 Castiglione D, Lovasi GS, Carvalho MS. Perceptions and uses of public and private health care in a Brazilian Favela. *Qual Health Res* 2018;28:159–72.
- 112 Getachew S, Kaba M, Gizaw M. Health service access, utilization and prevailing health problems in the urban vulnerable sections of Ethiopia. *Ethiop J Health Dev* 2020;34:12–23.
- 113 Tabrizi JS, Farahbakhsh M, Bazargani HS. Health services utilization and responsiveness: a comparison of slum and Non-slum regions in Tabriz, Iran. *Med Sci* 2018;22:577–82.
- 114 Agrawal D, Tyagi N, Dhakar JS. Awareness and utilization of geriatric welfare schemes among urban elderly population of district Gautam Budh Nagar. *Indian J Community Health* 2019;31:315–21.
- 115 Dasgupta P, Bhattacharjee S, Mukherjee A, *et al.* Vaccine hesitancy for childhood vaccinations in slum areas of Siliguri, India. *Indian J Public Health* 2018;62:253–8.
- 116 Bayou YT, Mashalla YS, Thupayagale-Tshweneagae G. The adequacy of antenatal care services among slum residents in Addis Ababa, Ethiopia. *BMC Preg Child* 2016;16:527–8.
- 117 Aleemi AR, Khaliq H, Faisal A. Challenges and patterns of seeking primary health care in slums of Karachi: a disaster lurking in urban shadows. *Asia Pac J Public Health* 2018;30:479–90.
- 118 Viramgami AP, Verma PB, Vala MC, *et al.* A cross-sectional study to assess reproductive and child health profile of working women residing in urban slums of Rajkot City. *Indian J Community Med* 2019;44:313–6.
- 119 van der Heijden J, Gray N, Stringer B, *et al.* 'Working to stay healthy', health-seeking behaviour in Bangladesh's urban slums: a qualitative study. *BMC Public Health* 2019;19:600.
- 120 Gaiha SM, Gillander Gadin K. 'No time for health': exploring couples' health promotion in Indian slums. *Health Promot Int* 2020;35:70–81.
- 121 Wingfield T, Tovar MA, Huff D, *et al.* Socioeconomic support to improve initiation of tuberculosis preventive therapy and increase tuberculosis treatment success in Peru: a household-randomised, controlled evaluation. *Lancet* 2017;389:S16.
- 122 Muralidharan A. Constrained choices? menstrual health and hygiene needs among adolescents in Mumbai slums. *Indian J Gend Stud* 2019;26:12–39.
- 123 MacPherson P, Khundi M, Nliwasa M, *et al.* Disparities in access to diagnosis and care in Blantyre, Malawi, identified through enhanced tuberculosis surveillance and spatial analysis. *BMC Med* 2019;17:21.
- 124 Černauskas V, Angeli F, Jaiswal AK, *et al.* Underlying determinants of health provider choice in urban slums: results from a discrete choice experiment in Ahmedabad, India. *BMC Health Serv Res* 2018;18:473.
- 125 Schultz JS, Muema S, Ouma A. Timeliness of vaccination of vaccination in an urban slum in Nairobi, Kenya. *Am J Trop Med Hyg* 2017;95:567.
- 126 Sharma S, Sarathi Mohanty P, Omar R, *et al.* Determinants and utilization of maternal health care services in urban slums of an Industrialized City, in Western India. *J Family Reprod Health* 2020;14:95–101.
- 127 Ahmed S, Adams AM, Islam R, *et al.* Impact of traffic variability on geographic accessibility to 24/7 emergency healthcare for the urban poor: a GIS study in Dhaka, Bangladesh. *PLoS One* 2019;14:e0222488.
- 128 Yazdani AT, Muhammad A, Nisar MI, *et al.* Unveiling and addressing implementation barriers to routine immunization in the peri-urban slums of Karachi, Pakistan: a mixed-methods study. *Health Res Policy Syst* 2021;19:55.
- 129 Williams A, Sarker M, Ferdous ST. Cultural attitudes toward postpartum depression in Dhaka, Bangladesh. *Med Anthropol* 2018;37:194–205.
- 130 Athié K, Dowrick C, Menezes ALdoA, *et al.* Anxious and depressed women's experiences of emotional suffering and help seeking in a Rio de Janeiro favela. *Cien Saude Colet* 2017;22:75–86.
- 131 Kuria Ng'endo, Reid A, Owiti P, *et al.* Compliance with follow-up and adherence to medication in hypertensive patients in an urban informal settlement in Kenya: comparison of three models of care. *Trop Med Int Health* 2018;23:785–94.
- 132 Mataboge MLS, Beukes S, Nolte AGW. The experiences of clients and healthcare providers regarding the provision of reproductive health services including the prevention of HIV and AIDS in an informal settlement in Tshwane. *Health SA Gesondheid* 2016;21:67–76.
- 133 Otieno PO, Wambiya EOA, Mohamed SM, *et al.* Access to primary healthcare services and associated factors in urban slums in Nairobi-Kenya. *BMC Public Health* 2020;20:981.
- 134 Odhiambo GO, Musuva RM, Odieri MR, *et al.* Experiences and perspectives of community health workers from implementing treatment for schistosomiasis using the community directed intervention strategy in an informal settlement in Kisumu City, Western Kenya. *BMC Public Health* 2016;16:986.
- 135 Patil SK, Ahmed MM. Assessment of integrated child development services in urban slums of Belagavi City, Karnataka. *Indian J Public Health Res Dev* 2016;7:208–11.
- 136 Agonig RC, Carvalho SM, Freire MAM, *et al.* The production of care in the routine of family health teams. *Rev Bras Enferm* 2018;71:2659–65.
- 137 Das Gupta M, Dasgupta R, Kugananthan P, *et al.* Flies without borders: lessons from Chennai on improving India's municipal public health services. *J Dev Stud* 2020;56:907–28.
- 138 Prado Junior JC, Virgilio TC, Medronho RdeA. Cure rates for tuberculosis in the municipality of Rio de Janeiro, Brazil, in 2012 compared with coverage by, and time of establishment of, family health units, and socio-economic and demographic factors. *Cien Saude Colet* 2016;21:1491–8.
- 139 Banerjee S, Selvaraj K, Bandyopadhyay K, *et al.* Urban health and nutrition day or only immunisation day? Barriers and bottlenecks in implementing urban health and nutrition day in an urban primary health centre of Nagpur, central India. *J Mother Child* 2021;25:51–60.
- 140 Ongarora D, Karumbi J, Minnaard W, *et al.* Medicine prices, availability, and affordability in private health facilities in low-income settlements in Nairobi County, Kenya. *Pharmacy* 2019;7:40.
- 141 Liu Y, Kong Q, Yuan S, *et al.* Factors influencing choice of health system access level in China: a systematic review. *PLoS One* 2018;13:e0201887.
- 142 Banke-Thomas OE, Banke-Thomas AO, Ameh CA. Factors influencing utilisation of maternal health services by adolescent mothers in Low-and middle-income countries: a systematic review. *BMC Pregnancy Childbirth* 2017;17:65.



- 143 Mendoza-Sassi R, Béria J. Health services utilization: a systematic review of related factors. *Cad Saude Publica* 2001;17:819–32.
- 144 Wekesah FM, Kyobutungi C, Grobbee DE, *et al.* Understanding of and perceptions towards cardiovascular diseases and their risk factors: a qualitative study among residents of urban informal settings in Nairobi. *BMJ Open* 2019;9:e026852.
- 145 Lilford RJ, Oyebode O, Satterthwaite D, *et al.* Improving the health and welfare of people who live in slums. *The Lancet* 2017;389:559–70.
- 146 World Health Organization. Strengthening health systems to improve health outcomes, 2007. Available: <https://apps.who.int/iris/handle/10665/43918> [Accessed 25 Dec 2021].
- 147 McIntyre DI, Thiede M, Birch S. Access as a policy-relevant concept in low- and middle-income countries. *Health Econ Policy Law* 2009;4:179–93.
- 148 Kardalkar S, Sherkhane MS. Utilization of antenatal care package among women of urban slums. *Indian J Public Health Res Develop* 2020;11:517–23.
- 149 Abd El Fatah SAM, El Habashy EM, Ismail HAH. Role of receipt of antenatal care in subsequent contraceptive use at primary health care centres serving slum areas of Cairo, Egypt. *Eur J Contracept Reprod Health Care* 2019;24:356–61.
- 150 Kalyango E, Kananura RM, Kiracho EE. Household preferences and willingness to pay for health insurance in Kampala City: a discrete choice experiment. *Cost Eff Resour Alloc* 2021;19:21.
- 151 Sharma S, Verma PB, Virangami AP, *et al.* Analysis of out-of-pocket expenditure in utilization of maternity care services in urban slums of Rajkot City, Gujarat. *Indian J Community Med* 2018;43:215–9.