

Coral reef resilience differs among islands within the Gulf of Mannar, southeast India following successive coral bleaching events

Diraviya Raj, K. ; Aeby, Greta; Mathews, G. ; Williams, Gareth J.; Caldwell, Jamie; Laju, R. L. ; Selva Bharath, M. ; Dinesh Kumar, P. ; Arasamuthu, A. ; Gladwin Gnana Asir, N. ; Wedding, Lisa; Davies, Andrew; Moritsch, Monica ; Patterson Edward, J. K.

Coral Reefs

DOI:
[10.1007/s00338-021-02102-0](https://doi.org/10.1007/s00338-021-02102-0)

Published: 01/08/2021

Peer reviewed version

[Cyswllt i'r cyhoeddiad / Link to publication](#)

Dyfyniad o'r fersiwn a gyhoeddwyd / Citation for published version (APA):

Diraviya Raj, K., Aeby, G., Mathews, G., Williams, G. J., Caldwell, J., Laju, R. L., Selva Bharath, M., Dinesh Kumar, P., Arasamuthu, A., Gladwin Gnana Asir, N., Wedding, L., Davies, A., Moritsch, M., & Patterson Edward, J. K. (2021). Coral reef resilience differs among islands within the Gulf of Mannar, southeast India following successive coral bleaching events. *Coral Reefs*, 40(4), 1029-1044. <https://doi.org/10.1007/s00338-021-02102-0>

Hawliau Cyffredinol / General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

1 **Title:** Coral reef resilience differs among islands within the Gulf of Mannar, southeast
2 **India following successive coral bleaching events**

3

4 **Authors:** K. Diraviya Raj¹, Greta S. Aeby², G. Mathews¹, Gareth J. Williams³, Jamie
5 Caldwell^{4,5}, R. L. Laju¹, M. Selva Bharath¹, P. Dinesh Kumar¹, A. Arasamuthu¹, N. Gladwin
6 Gnana Asir¹, Lisa Wedding⁶, Andrew J. Davies⁷, Monica Moritsch⁸ and J. K. Patterson Edward¹

7 ¹ Suganthi Devadason Marine Research Institute, Tuticorin, Tamil Nadu, India

8 ² Department of Biology and Environmental Science, Qatar University, Doha, Qatar

9 ³ School of Ocean Sciences, Bangor University, Anglesey, UK

10 ⁴ Hawaii Institute of Marine Biology, University of Hawaii at Manoa, Honolulu, HI, USA

11 ⁵ ARC Centre of Excellence for Coral Reef Studies, James Cook University, Townsville,
12 Australia

13

14 ⁶ School of Geography and the Environment, University of Oxford, Oxford, United Kingdom

15 ⁷ Department of Biological Sciences, University of Rhode Island, Kingston, RI, USA

16 ⁸ United States Geological Survey, Western Geographic Science Center, Moffett Field,
17 California, USA

18

19 **Corresponding author:** diraviyam_raj@yahoo.co.in

20

21 **Keywords:** successive bleaching events, coral resilience, Gulf of Mannar, southeast India, coral
22 morphotypes, island-specific response, chlorophyll-a, reef fish

23

24 **Abstract**

25 We used a 12-year data set of benthic cover (2005-2017), spanning two bleaching events,
26 to assess changes in benthic cover and coral community composition along 21 islands within
27 Gulf of Mannar (GoM), southeast India. Overall, between 2005 and 2017 reefs had a
28 simultaneous decrease in relative coral cover (avg. = -36%) and increase in algal cover (avg. =
29 +45%). Changes in benthic cover were not consistent among islands, ranging from -34% to +5%
30 for coral cover and from -0.3% to +50% for algae. There was a spatial gradient in coral
31 mortality, which increased among islands from west to east. However, there was a disconnect
32 between coral loss and subsequent increases in algae. Algal cover increased more on islands in
33 west GoM where coral loss was minimal. Environmental co-factors (coral cover, percent
34 bleaching, degree heating weeks, fish densities, Chl-a, pollution) explained >50% of the benthic
35 cover responses to successive bleaching. Coral survival was favored on islands with higher fish
36 densities and chlorophyll-a levels and increases in algal cover were associated with higher
37 measures of pollution from terrestrial runoff. Coral morphotypes differed in their response
38 following successive bleaching resulting in changes in the relative abundance of different coral
39 morphotypes. Existing climate projections (RCP8.5) indicate a 22-year gap in the onset of annual
40 severe bleaching (ASB) for reefs in the east versus west GoM and ASB was ameliorated for all
41 reefs under the RCP4.5 projections. There is limited knowledge of the resilience of GoM reefs
42 and this study identifies coral morphotypes and reefs that are most likely to, recover or decline,
43 from successive bleaching, in the context of forecasts of the frequency of future bleaching events
44 in GoM.

45

46

47 **Introduction**

48 Coral reefs are one of the most sensitive ecosystems to climate change and repeated mass
49 coral bleaching events caused by ocean warming (Heron et al. 2016; Hughes et al. 2018) are
50 fundamentally altering coral reefs as we know them (Williams and Graham 2019). Based on
51 global averages, ocean temperatures in 2015-2017 were the highest temperatures recorded since
52 the 1800s (Blunden and Arndt 2019) and resulted in a 3-year global coral bleaching event
53 (Hughes et al. 2018; Eakin et al. 2019). Bleaching occurs when there is a breakdown between
54 corals and their symbiotic microalgae (zooxanthellae) (Vidal-Dupiol et al. 2009) and can result
55 in extreme nutritional stress for corals (Muscatine 1990). Thermally stressed corals have reduced
56 growth, reproductive output, higher disease susceptibility and increased risk of mortality,
57 depending on the duration of the heat event (Baker et al. 2008). Coral loss subsequently effects
58 other organisms that depend on coral reefs for food and shelter (Glynn 1985; Sano 2004;
59 Bellwood et al. 2006; Baker et al. 2008). Coral bleaching also changes the balance between reef
60 accretion and erosion (Cantin and Lough 2014), resulting in a loss of reef topographic
61 complexity and rugosity (Perry and Alvarez-Filip 2019).

62 As ocean waters continue to warm under climate change, bleaching events are expected
63 to become more frequent and severe, giving coral reefs little time to recover between
64 disturbances (van Hooidonk et al. 2016). Coral reefs show spatial heterogeneity in the severity of
65 coral bleaching and degree of recovery (Graham et al. 2015; Hughes et al. 2018; Safaie et al.
66 2018), which is influenced by factors such as bleaching severity, coral community structure,
67 abundance of herbivores, maintenance of biodiversity, exposure to secondary stressors and
68 gradients in oceanography and climate (Baker et al. 2008; Graham et al. 2015; Safaie et al. 2018;
69 McClanahan et al. 2019; Head et al. 2019). For example, reefs in the Seychelles were more

70 likely to recover to coral dominance following mass coral bleaching if they were in deeper water
71 and had more abundant herbivore populations (Graham et al. 2015). There is still much
72 uncertainty surrounding coral reef responses to successive bleaching events and gathering data
73 on the effects of recurrent bleaching on coral reefs is important to understand which coral
74 species, reefs and regions are most likely to display resistance or resilience to climate change.

75 Using a 12-year data set of benthic cover (2005-2017), spanning two bleaching events,
76 the long-term benthic cover and coral community composition of reef sites were assessed along
77 21 islands within Gulf of Mannar, southeast India. Gulf of Mannar (GoM) reefs were first
78 impacted by bleaching in 1998, where 89% of the coral bleached and 23% subsequently died
79 (Arthur 2000). More recently in 2010, thermal stress caused 10% bleaching and 9.7% mortality
80 (Edward et al. 2012), and in 2016 resulted in 24% bleaching and 16% mortality (Edward et al.
81 2018). Our study examined the resilience of these reefs in response to successive bleaching
82 events. Changes in benthic cover and coral community composition was examined following the
83 two recent bleaching events, in terms of which coral morphotypes drove changes in coral
84 community composition, and what environmental conditions were associated with changes in
85 coral and algal cover following bleaching. Finally, global climate model predictions were used to
86 assess future annual severe bleaching conditions for reefs in Gulf Mannar associated with global
87 climate change.

88

89 **Methods**

90 **Survey methods**

91 Four monitoring sites were established at 21 islands in the Gulf of Mannar (GoM), India
92 in 2005 (Fig. 1 and Supplemental Table 1), which have been resurveyed annually through 2017

93 (Edward et al. 2008a, 2008b, 2012, 2018). At each site, three 20 m transects were laid parallel to
94 shore with a minimum of a 20-meter gap between transects (3 transects/site * 4 sites/island =252
95 transect/year). Along each transect, substrate characteristics were recorded using line-intercept
96 method with corals recorded by growth forms (morphotypes) and further categorized by corals
97 within Acroporidae versus corals in other families following English et al. (1997) (Table 2).
98 Other substrate categories included soft corals, algae (macroalgae and algal turf), crustose
99 coralline algae, abiotic (sand, rock and old dead corals) and others (sponges, sea anemones,
100 ascidia, zoanthids, crinoids, oysters, hydroids, and bryozoans). Annual surveys were conducted
101 between October and December and additional surveys were conducted at the same sites
102 between April and June during bleaching events in 2010 and 2016. Timing of surveys during the
103 bleaching events were based on sea surface temperatures (SST) indicating water temperatures
104 were passing the bleaching threshold for corals in GoM (30°C) (Edward et al. 2018) and rapid
105 surveys conducted at representative sites during the elevated SST time periods. In this manner,
106 we were able to resurvey the reefs as bleaching was approximately at its peak. At each survey
107 date, sites were relocated via GPS coordinates allowing the same area of the reef to be surveyed
108 but transect placement was random rather than along permanent markers.

109

110 **Environmental variables**

111 Environmental variables that could affect bleaching, mortality or recovery of benthic
112 populations were measured *in situ* during annual surveys or derived from remotely sensed data
113 (Table 2). Water clarity was measured at each site with a 20 cm Secchi disc and divided by
114 maximum bottom depth to standardize across sites. Sedimentation was assessed annually in
115 2005-2008 and 2013-2017 using four replicate PVC sediment traps (10 cm height x 8 cm

116 diameter) per island. Traps were secured adjacent to the reef, 20 cm above the bottom, and
117 collected after 10 to 15 days. Samples were dried at 70°C and weighed to calculate milligrams of
118 sediment deposited per cm² per day. At the island-level, sedimentation varied little through time,
119 therefore, mean sedimentation values per island were used in statistical analyses. Reef fish
120 densities were recorded using visual census along six belt transects (50 x 5 m) per island between
121 April 2014 and March 2015. Annual maximum degree heating week (DHW) values at 5-km
122 spatial resolution were obtained for each island for the two bleaching years, 2010 and 2016, from
123 NOAA's Coral Reef Watch (Liu et al. 2005, NOAA Coral Reef Watch 2019). Maximum
124 monthly chlorophyll-a values (as a proxy for phytoplankton biomass) at 4-km spatial resolution
125 for each island and bleaching year were obtained from monthly chlorophyll-a observations from
126 NASA MODIS-Aqua (NASA 2014, 2017). Missing data at a given sampling site and date were
127 excluded from calculation of the maximum. To account for human impacts on reefs, the 2015
128 human population for India (WorldPop 2017) was measured at 100-m spatial resolution. For
129 each coast-adjacent grid cell, the total number of people living within 10km of each survey point
130 was found using the Zonal Statistics 2 Toolbox in ArcMap 10.1 (Environmental Systems
131 Research, Inc.). To assess the relative impact of coastal populations on each island, the Inverse
132 Distance Weighted method was employed, where grid cells farther away from an island received
133 less weight than closer cells. The population in each cell was divided by the square of the
134 distance of that cell from the survey point to produce a weighted population measure for each
135 survey point.

136

137

138

139 **Data analysis**

140 *Analyses of benthic changes following successive bleaching events*

141 To determine whether the mean proportional change in coral and algae cover differed
142 across islands, ANOVAs were used. In the analysis of variance (ANOVAs) the response variable
143 was the difference in coral or algae cover between 2005 and 2017 for each transect and the
144 predictor variable (group) was island. Numerous environmental, ecological and human factors
145 are hypothesized to affect coral bleaching resistance and resilience (West and Salm 2002; Obura
146 2005). We assessed the role of potential environmental drivers in explaining variability in
147 benthic cover responses to the 2010 and 2016 bleaching events using beta regression models.
148 The response variables in the beta regression models were calculated as the change in coral or
149 algae cover on a transect from before and after each bleaching event (i.e., substrate change
150 between 2009 and 2010 and 2015 and 2016), which varied from -100 to 100% and were scaled
151 between 0 and 1. The initial predictor variables considered in the beta regression models
152 included a suite of environmental variables hypothesized to affect coral's ability to resist
153 bleaching or recover following a bleaching event (Table 1), however, variables with greater than
154 50% correlation (based on Pearson's correlation coefficient) were removed from the analysis.
155 Random effects of transect, nested within site, nested within island, were also included in the
156 beta regression models. Stepwise forward selection was used to select the optimal model,
157 sequentially adding variables to the nested random effects that reduced the AIC value by more
158 than two. Model fit was further assessed by calculating R^2 values for model predicted outcomes
159 with observations of substrate change. The beta regression models were conducted in R
160 statistical software (hereafter referred to as R) using the glmmTMB function and package
161 (Magnusson et al. 2017). The magnitude of change in coral or algae cover following the 2010

162 bleaching event was compared with the changes following the 2015 event using paired t-tests,
163 with data paired by island.

164

165 ***Coral community changes following successive bleaching***

166 Permutational multivariate analysis of variance (PERMANOVA) were used to determine
167 whether the coral community composition (based on proportion of different morphotypes)
168 shifted following each bleaching event and across islands. For each island, a PERMANOVA was
169 conducted where the response variable was the Bray-Curtis dissimilarity matrix from the raw
170 percentage morphotype cover values for each transect and the predictor variable was time period
171 with nested random effects of transect within site. There were three time periods assessed,
172 including the period: i) before the 2010 bleaching event (2005-2009), ii) following the 2010
173 bleaching event and before the 2016 bleaching event (2010-2015), and iii) following the 2016
174 bleaching event (2016-2017). PERMANOVAs were conducted using the vegan package (Dixon
175 2003) in R.

176 An indicator species analysis was used to identify morphotypes driving differences in the
177 coral community composition across time periods and islands. An indicator species analysis
178 identifies species assemblages that are characteristic of specific groups (i.e., in this study group
179 refers to time period). This is done by combining species relative abundance and relative
180 frequency of occurrence across all combinations of groups, where the index is maximized when
181 a species is only found in a single group and is present in all samples associated with that group.
182 The indicator species analysis was conducted using the multipatt function in the indicpecies
183 package (De Caceres 2013) in R. Code for the ANOVAs, beta regression models,

184 PERMANOVAs, and indicator species analysis are available in the following github repository:
185 https://github.com/jms5151/Coral_times_series_Gulf_of_Mannar.

186

187 **Climate projections for Gulf of Mannar**

188 Future bleaching frequency for the Gulf of Mannar (GoM) was analyzed using global
189 coral bleaching predictions from van Hooidonk et al. (2016). Down-scaled (4-km resolution)
190 climate model projections of predicted ocean surface warming over the coming decades were
191 used to assess the 21 islands in the GoM. Ocean warming was predicted from an ensemble of
192 Coupled Model Intercomparison Project phase 5 models using emissions pathways RCP8.5 (high
193 CO₂ emissions) and RCP4.5. Emissions scenario RCP4.5 represents lower emissions mid-
194 century than will eventuate if pledges made following the 2015 Paris Climate Change
195 Conference (COP21) become reality (van Hooidonk et al. 2016). From the van Hooidonk et al.
196 (2016) data layers (available at:
197 https://coralreefwatch.noaa.gov/climate/projections/downscaled_bleaching_4km/index.php), we
198 calculated the decade in which reefs across the GoM are predicted to start bleaching twice per
199 decade and 10 times per decade, referred to as Annual Severe Bleaching (ASB) were determined
200 (van Hooidonk et al. 2016). ASB translates to an exceedance of 8 Degree Heating Weeks
201 (DHWs) projected to occur in each of the 10 years per decade; 8 DHWs is higher than the mean
202 optimum world-wide bleaching predictor of 6.1 DHWs (i.e. at 8 DHWs thermal stress will be
203 sufficiently great for bleaching to occur) (van Hooidonk and Huber 2009).

204

205

206

207 **Results**

208 **Island-specific responses following successive bleaching**

209 Between 2005 and 2017 average coral cover (all islands combined) declined by 36% and
210 average algae cover increased by 45% with changes in percent cover occurring after each
211 bleaching event and a greater magnitude of benthic response after the severe 2016 event (Fig. 2).
212 Following the 2016 bleaching event, there was an average of 6.5% more coral cover loss ($t(20)=-$
213 -3.67 , $p<0.01$) and an average increase of 9.2% more algal cover ($t(20)=3.16$, $p<0.01$) than
214 following the 2010 event. However, changes were not consistent among islands, ranging from -
215 34% to +5% for coral cover and from -0.3% to +50% for algae cover (Fig. 3). There was a
216 spatial gradient in coral mortality with islands in the eastern part of GoM losing more coral than
217 islands in the western part but there was a disconnect between coral loss and subsequent
218 increases in algae cover. The four western-most islands, Vaan, Koswari, Kariyachalli, and
219 Villanguchalli had minimal to no reductions in coral cover (+5%, +2%, -10%, -5%), respectively
220 but the largest increases in algae cover (+31%, +53% +39%, +51%), respectively (Fig. 3).

221 Changes in benthic cover following each bleaching event were significantly different
222 among islands for coral (ANOVA, $F(20, 231) = 26.21.854$, $p < 0.001$) and algae cover
223 (ANOVA, $F(20, 231) = 31.84$, $p < 0.001$) and by examining the effect of each environmental
224 covariate on benthic cover change, while accounting for all other variables in the model, we
225 found these changes were associated with specific environmental drivers. Coral mortality was
226 lower on islands with higher fish densities and higher chlorophyll-a levels whereas mortality was
227 greater on islands with more bleaching and higher than average coral cover prior to bleaching
228 (Fig. 4). The optimal model explained 53% of the spatial variability in changes in coral cover
229 among islands (Table 2). It must be noted that the total loss of coral cover is probably due to a

230 combination of direct mortality caused by bleaching, as well as subsequent mortality suffered by
231 corals, which have been stressed or suffered partial mortality, leaving them more vulnerable to
232 algal overgrowth and diseases (West and Salm 2003). Algae cover following bleaching events
233 increased on islands with more terrestrial runoff pollution, and higher thermal stress, with lower
234 increases in algae found on islands with higher initial algae cover and higher chlorophyll-a levels
235 (Fig. 5). The optimal model explained 51% of the spatial variability in algal cover change (Table
236 2).

237 **Taxon-specific resilience to successive bleaching**

238 Coral morphotypes differed in their response following successive bleaching through
239 time. All coral morphotypes showed coral loss following the 2010 bleaching with cover
240 stabilizing or even slightly increasing up until the 2016 bleaching. However, after the severe
241 2016 bleaching, most coral morphotypes had an even greater loss in coral cover, but there were
242 two types, encrusting Acroporidae (ACE) and submassive corals (CS), which increased in
243 absolute cover (Fig. 6). Hence, the proportional contribution of each morphotype to the overall
244 coral community changed between 2005 and 2017 by either having a greater increase or decrease
245 in cover relative to other morphotypes or by keeping cover constant while other morphotypes
246 changed in abundance (Fig. 7). The largest reduction among morphotypes was for digitate
247 Acroporidae (ACD) which represented 17.1% of the community in 2005 and 5.1% of the
248 community in 2017. In contrast, mounding corals (CM) had the largest increase from 31.1% of
249 the community in 2005 to 42.8% in 2017. Ultimately, coral communities in GoM shifted
250 following successive bleaching with some coral morphotypes relative “winners” and others as
251 relative “losers” (Fig. 7).

252

253 **Coral community shifts differ among islands**

254 For all islands, the coral community shifted significantly following the severe 2016
255 bleaching event (Supplemental Table 2) but the degree of change differed among islands (Fig. 8).
256 Some islands showed a more distinct shift in community structure after the 2016 bleaching event
257 (orange polygons relative to the grey and blue polygons in Fig. 8) (e.g., Mulli) whereas others
258 were less extreme (e.g., Manoliputti). The coral types exerting the strongest influence on spatial
259 variations in community structure also differed among islands especially following the 2016
260 bleaching (Fig. 9). However, some consistencies were evident such as encrusting Acroporidae
261 (ACE) which increased in abundance at 16 of the 21 islands and foliose corals (CF) which
262 decreased in abundance at all but two islands (Fig. 9).

263 **Climate projections for Gulf of Mannar**

264 The downscaled climate projections showed that all islands across the GoM are predicted
265 to experience annual severe bleaching (ASB) under a high emissions scenario (RCP8.5) prior to
266 2070 and bleaching twice per decade prior to 2060 (Fig. 10). However, the projections also
267 highlighted local-scale (10s km) spatial variability in the expected frequency of severe bleaching
268 events (Fig. 10). For a high emissions scenario (RCP8.5), the onset of ASB showed a clear east
269 to west gradient, with reefs towards the eastern end of the GoM (the islands of Shingle,
270 Krusadai, Pullivasal and Poomarichan) all predicted to experience ASB before 2045. Moving
271 west, the onset of ASB generally occurs later and by Nallathanni Island the onset is pushed to
272 2061. Three islands towards the far western end (Koswari, Vilanguchalli, Kariyachalli) are not
273 expected to experience ASB until 2067. The patterns for severe bleaching twice per decade
274 generally show the same east to west gradient (Fig. 10). The reduced emissions scenario RCP4.5,

275 has clear ameliorating effects, and means the majority of islands would not experience ASB (or
276 even severe bleaching twice per decade) between now and 2070. Exceptions to this pattern are
277 the four islands towards the far eastern end of the GoM, which are still predicted to experience
278 ASB prior to 2089 and severe bleaching twice per decade prior to 2070 under RCP4.5 (Fig. 10).

279

280 **Discussion**

281 Following successive bleaching events in the GoM in 2010 and 2016, reefs generally
282 exhibited the classic paradigm of a simultaneous decrease in coral cover and increase in algae
283 cover. However, there were contrasting responses among the 21 islands to the multiple bleaching
284 events: some islands lost significant coral cover over time while others were able to maintain
285 their cover. Thermal stress, expressed in degree heating weeks (DHW), experienced by GoM
286 reefs was lower in 2010 (range 2.4-4.8) compared to 2016 (range 6.4-9.3), but varied among
287 islands in both years, even though all islands had reefs at similar depths (<5m). As expected,
288 maximum DHW and percent bleaching were significant factors explaining change in benthic
289 cover among islands. However, other environmental variables also impacted coral bleaching,
290 mortality or recovery. Lower coral mortality following bleaching events was found on islands
291 with higher fish densities and chlorophyll a levels. Reef fish play a critical role in maintaining
292 ecosystem function and resilience of coral reef habitats (Graham et al. 2011). Grazing by
293 herbivores generates reductions in algal cover that promotes recovery of corals (Mumby et al.
294 2006; [Burkepile and Hay 2008](#)) and maintaining fish diversity can mitigate threats from coral
295 disease (Raymundo et al. 2009). Chlorophyll-a is a proxy for phytoplankton biomass and thus
296 ocean surface primary productivity (Gove et al. 2016; Coelho et al. 2017). Historically,
297 chlorophyll-a has been used as a proxy for water quality and eutrophication, and excess coastal

298 nutrients can reduce coral cover and promote macroalgal cover, particularly in human populated
299 regions (Fabricius 2005; Wooldridge 2009). Reefs exposed to higher nutrients can also
300 experience more severe bleaching (Woodridge 2009; Woolridge and Done 2009; Vega-Thurber
301 et al. 2014). However, the relationship between a reef's response to thermal stress and
302 "nutrients" is more nuanced than this (D'Angelo and Wiedenmann 2014; Williams et al. 2019).
303 Chlorophyll-a is also strongly correlated with the abundance of zooplankton which represents
304 key food sources for reef-building corals (Fox et al. 2018) that can promote their spatial
305 dominance (Williams et al. 2015; Aston et al. 2019), their resilience to coral bleaching (Grottoli
306 et al. 2006) and overall ecosystem function (Graham et al. 2018).

307 Higher concentrations of phytoplankton in the water might also offer some degree of
308 protection to corals by limiting the amount of ultraviolet radiation (UVR) reaching colonies.
309 UVR can directly damage corals (Lesser 1996; Anderson et al., 2001; Baruch et al. 2005) and is
310 a synergistic factor increasing bleaching severity during thermal stress events (Torregiani and
311 Lesser 2007; Ferrier-Pages et al. 2007). Factors that ameliorate the amount of UVR reaching
312 coral colonies could reduce bleaching or other harmful effects of UVR exposure. For example,
313 Iluz et al. (2008) found that bleaching-related colony mortality within warmer lagoon waters was
314 lower than colonies on surrounding slopes in cooler water. Lagoon waters had high turbidity due
315 to seagrass leachate which attenuated UVR and protected corals from further bleaching.

316 Given the increasing gradient in coral mortality from west to east, it was surprising that
317 increases in algae cover did not follow the same spatial pattern as coral mortality as has been
318 found in other studies (e.g., coral loss is followed by subsequent increases in algal cover; Diaz-
319 Pulido et al. 2009). Instead, the highest increases in algal cover occurred on the four islands in
320 the west of GoM that had the lowest levels of coral loss (Vaan, Koswari, Kariyachalli,

321 Vilanguchalli). These four islands maintained or lost little coral suggesting that increases in algae
322 among islands were not necessarily linked to reductions in coral cover. Coral reef ecosystem
323 recovery patterns occur against the background of local stressors. The four islands with the
324 highest increases in algal cover were closest to the main population center of Tuticorin and a
325 major sewer outfall for the region (Meiaraj and Jeyapriya, 2019), and we found terrestrial runoff
326 pollution as an important factor explaining spatial differences in increased algae cover among
327 islands. Increasing levels of algae are already a problem for reefs in GoM (Jeevamani et al.
328 2013, Bharath et al. 2017) and may prove problematic for reef resilience, as algae can directly
329 overgrow corals, trap sediment, prevent coral settlement and potentially harbor coral pathogens
330 (Smith et al. 2006; Mumby et al. 2007; McClanahan et al. 2012; Vega-Thurber et al. 2012).
331 Climate change-related coral mortality is unavoidable here, but local management actions can
332 improve conditions allowing reefs to better recover. For example, algal growth could be
333 minimized by reducing pollution or enhancing herbivore populations, which in turn will
334 maximize the potential for coral regrowth and for the establishment of juvenile corals. This
335 study identifies islands prone to coral mortality and/or algal overgrowth following bleaching
336 events, providing direction for potential mitigation.

337 Variability in bleaching and mortality may have also arisen from local acclimation of
338 corals to heat stress. Reefs within GoM undergo mild bleaching almost every summer that also
339 varies among islands (Edward et al. 2012, 2018). This may have resulted in coral populations at
340 some islands with a higher heat tolerance. Other studies have suggested that historical
341 temperature variability affects corals' physiological tolerance under thermal stress (McClanahan
342 et al. 2004; Oliver and Palumbi 2011) with surviving populations better adapted to withstanding
343 further thermal stress events (Carilli et al. 2012; Palumbi et al. 2014). Conversely, Hughes et al.

344 (2018) found no evidence for a protective effect of past bleaching (e.g. from acclimation or
345 adaptation) along the Great Barrier Reef, Australia. They found that reefs with higher bleaching
346 scores in 1998 or 2002 did not experience less severe bleaching in 2016.

347 Bleaching and subsequent mortality patterns can differ among coral genera (Edmunds
348 1994; Hoegh-Guldberg 1999; Marshall and Baird 2000) and growth forms (Loya et al. 2001; Iluz
349 et al. 2008), and have been attributed to numerous coral host factors (Loya et al. 2001; Brown et
350 al. 2002; Grottelli et al. 2006; Visram and Douglas 2007; Baird et al. 2009) and/or the density or
351 types of *Symbiodinium* residing within the coral host (Bhagooli and Yakovleva 2004; Sampayo
352 et al. 2008; Howells et al. 2012; Cunning and Baker 2013). Similarly, within GoM, coral
353 morphotypes varied in bleaching severity and mortality following each bleaching event, resulting
354 in a change in community structure. As some coral species died after the bleaching events, the
355 more bleaching tolerant species increased in relative abundance in the community. Massive
356 corals had the largest relative increase in the community likely because this morphotype has
357 coral taxa known to be stress tolerant such as *Porites*, *Dipsastraea* and *Favites*, (Stafford-Smith
358 1993; Riegl 1999; Burt et al. 2013) among others. Digitate Acroporidae had the largest losses as
359 these coral morphotypes contained the more thermally sensitive coral taxa, *Acropora* spp. and
360 *Montipora* spp. (Marshall and Baird 2000; Kayanne et al. 2002; McClanahan et al. 2004). It is
361 important to note that for the current study, most coral morphotypes included multiple coral
362 genera, which can differ in bleaching susceptibility regardless of their growth form (Baird and
363 Marshall 2002; McClanahan et al. 2004). As the coral communities in GoM shift through time,
364 so may the risk to reefs from different threats. As an example, *Montipora* spp., are becoming a
365 larger component of the GoM coral community and *Montipora* spp. are known to be susceptible
366 to outbreaks of tissue loss disease in GoM (Raj et al. 2016). In contrast, foliose corals (CF)

367 which are important in providing habitat for fish and other marine species decreased in
368 abundance. As coral communities continue to change through time, it would be advantageous
369 for managers to re-evaluate local threats to GoM reefs.

370 A key issue for the potential resilience of all reefs is the frequency of disturbance events
371 and whether sufficient time for recovery of mature coral assemblages can occur. When reefs
372 bleach annually, reef recovery becomes highly unlikely (van Hooidonk et al. 2016). Islands
373 across GoM are not predicted to experience annual severe bleaching (ASB) under a high
374 emissions scenario (RCP8.5) until after 2040. Reefs are then expected to have a distinct east to
375 west gradient in timing of ASB with a 22-year gap predicted between the onset of ASB on reefs
376 in east versus west GoM. This predicted spatial pattern of thermal stress among islands is
377 consistent with what was found during the 2010 and 2016 bleaching events, and provides some
378 indication of how GoM reefs might respond to repetitive bleaching events in the future. The
379 western reefs will become increasingly important, constituting spatial refugia (van Hooidonk et
380 al. 2013) for corals which is critical for reef recovery via larval transport (Hock et al. 2017). The
381 predictions of ASB for GoM are greatly improved under the reduced emissions scenario
382 (RCP4.5) with only the four most vulnerable northern islands predicted to experience ASBs and
383 then not until after 2070. This is potentially good news for GoM and provides strong motivation
384 for global policy makers to take steps to limit carbon emissions to mitigate global climate
385 change. There is limited knowledge on the resilience of GoM reefs and this study identifies the
386 coral morphotypes and reefs that are most likely to, recover or decline, from successive
387 bleaching events. We forecast the frequency of future bleaching events for individual islands
388 providing guidance for future management.

389

390 **Acknowledgements**

391 Climate projections for GoM were done under the Third NATCOM (National
392 Communication) to the United Nations Framework Convention on Climate Change (UNFCCC)
393 project of Ministry of Environment, Forest and Climate Change (MoEF & CC), Government of
394 India. The authors are thankful to the Ministry of Environment, Forests and Climate Change,
395 Government of India (GOI); Tamil Nadu Forest Department (TNFD), Government of Tamil
396 Nadu (GOTN); Department of Environment, GOTN; Gulf of Mannar Biosphere Reserve Trust,
397 GOTN; Coral Reef Degradation in Indian Ocean, Sweden; and V.O.Chidambaranar Port Trust
398 (VOCPT), GOI for funding support. Thanks are also due to Chief Wildlife Warden, TNFD;
399 Wildlife Warden, Gulf of Mannar Marine National Park (GOMMNP), GOTN; and Chairman,
400 VOCPT, GOI for research permissions to carry out coral reef surveys and monitoring within the
401 Marine National Park and harbour area and to Suganthi Devadason Marine Research Institute for
402 logistical support.

403 **Conflict of interest statement**

404 On behalf of all authors, the corresponding author states that there is no conflict of interest.

405 **References**

- 406 Anderson S, Zepp R, Machula J, Santavy D, Hansen L, Mueller E (2001) Indicators of UV
407 exposure in corals and their relevance to global climate change and coral bleaching. *Hum*
408 *Ecol Risk Assess* 7:1271-1282
409
- 410 Arthur R (2000) Coral bleaching and mortality in three Indian reef regions during an El Niño
411 southern oscillation event. *Curr Sci* 79:1723-1729
412
- 413 Aston EA, Williams GJ, Green JM, Davies AJ, Wedding LM, Gove JM, Jouffray JB, Jones TT,
414 Clark J (2019) Scale-dependent spatial patterns in benthic communities around a tropical
415 island seascape. *Ecography* 42:578-590

416
417 Baird A, Marshall P (2002) Mortality, growth and reproduction in scleractinian corals following
418 bleaching on the Great Barrier Reef. *Mar Ecol Prog Ser* 237:133–141.
419
420 Baird AH, Bhagooli R, Ralph PJ, Takahashi S (2009) Coral bleaching: the role of the host.
421 *Trends Ecol Evol* 24:16–20
422
423 Baker AC, Glynn PW, Riegl B (2008) Climate change and coral reef bleaching: An ecological
424 assessment of long-term impacts, recovery trends and future outlook. *Estuar Coast Shelf*
425 *Sci* 80:435–471
426
427 Baruch R, Avishai N, Ranopwitz C (2005) UV incites diverse levels of DNA breaks in different
428 cellular compartments of a branching coral species. *Journal of Experimental Biology*
429 208:843–848
430
431 Bellwood DR, Hoey AS, Ackerman JL, Depczynski M (2006) Coral bleaching, reef fish
432 community phase shifts and the resilience of coral reefs. *Glob Chang Biol* 12:1587–1594
433
434 Bhagoolil R, Yakovleva I (2004) Differential bleaching susceptibility and mortality patterns
435 among four corals in response to thermal stress. *Symbiosis* 37:121-136
436
437 Bharath MS, Raj KD, Mathews G, Edward JKP (2017) Increasing macroalgae pose a threat to
438 corals in Koswari Island of Gulf of Mannar, southeast India. *Journal of Aquatic Biology*
439 *& Fisheries* 5:177-183
440
441 Blunden J, Arndt DS (2019) State of the Climate in 2018. *Bull Am Meteorol Soc* 100: Si–S305
442 [doi:10.1175/2019BAMSstateoftheClimate.2]
443
444 Brown BE, Downs CA, Dunne RP, Gibb SW (2002) Exploring the basis of thermotolerance in
445 the reef coral *Goniastrea aspera*. *Mar Ecol Prog Ser* 242:119-129
446
447 Burkepile DE, Hay ME (2008) Herbivore species richness and feeding complementarity affect
448 community structure and function on a coral reef. *Proc Natl Acad Sci U S A* 105:16201-
449 16206.
450
451 Burt JA, Feary DA, Van Lavieren H (2013) Persian Gulf reefs: an important asset for climate
452 science in urgent need of protection. *Ocean Challenge* 20:49-56
453
454 Cantin NE, Lough JM (2014) Surviving coral bleaching events: *Porites* growth anomalies on the
455 Great Barrier Reef. *PLoS One* 9:e88720 [doi.org/10.1371/journal.pone.0088720]
456
457 Carilli J, Donner SD, Hartmann AC (2012) Historical temperature variability affects coral
458 response to heat stress. *PLoS One* 7:e34418 [doi.org/10.1371/journal.pone.0088720.t001]
459

460 Coelho C, Heim B, Foerster S, Brosinsky A, de Araújo J (2017) In situ and satellite observation
461 of CDOM and chlorophyll-a dynamics in small water surface reservoirs in the Brazilian
462 semiarid region. *Water* 9:913 [doi: 10.3390/w9120913]
463

464 Cunning R, Baker AC (2013) Excess algal symbionts increase the susceptibility of reef corals to
465 bleaching. *Nat Clim Chang* 3: 259-262
466

467 D'Angelo C, Wiedenmann J (2014) Impacts of nutrient enrichment on coral reefs: new
468 perspectives and implications for coastal management and reef survival. *Curr Opin*
469 *Environ Sustain* 7: 82-93
470

471 De Cáceres M (2013) How to use the indicpecies package (ver.1.6.7). [http://www2.uaem.mx/r-](http://www2.uaem.mx/r-mirror/web/packages/indicpecies/vignettes/indicpeciesTutorial.pdf)
472 [mirror/web/packages/indicpecies/vignettes/indicpeciesTutorial.pdf](http://www2.uaem.mx/r-mirror/web/packages/indicpecies/vignettes/indicpeciesTutorial.pdf)
473

474 Diaz-Pulido G, McCook LJ, Dove S, Berkelmans R, Roff G, Kline DI, Weeks S, Evans RD,
475 Williamson DH, Hoegh-Guldberg O (2009) Doom and boom on a resilient reef: climate
476 change, algal overgrowth and coral recovery. *PloS one* 4: p.e5239
477 [doi.org/10.1371/journal.pone.0005239]
478

479 Dixon P (2003) VEGAN, a package of R functions for community ecology. *Appl Veg Sci* 14:
480 927-930

481 Vidal-Dupiol J, Adjeroud M, Roger E, Foure L, Duval D, Mone Y, Ferrier-Pages C, Tambutte,
482 E, Tambutte S, Zoccola D, Allemand D, Mitta G (2009) Coral bleaching under thermal
483 stress: putative involvement of host/symbiont recognition mechanisms. *BMC Physiol* 9
484 [doi: 10.1186/1472-6793-9-14]

485 Eakin CM, Sweatman HPA, Brainard RE (2019) The 2014–2017 global-scale coral bleaching
486 event: insights and impacts. *Coral Reefs* 38:539–545
487

488 Edmunds PJ (1994) Evidence that reef-wide patterns of coral bleaching may be the result of the
489 distribution of bleaching-susceptible clones. *Marine Biology* 121:137-142
490

491 Edward JKP, Mathews G, Raj KD, Tamelander J (2008a) Coral reefs of the Gulf of Mannar,
492 Southeastern India - observations on the effect of elevated SST during 2005-2008, in:
493 Riegl, B., Dodge, R.E., (Eds.), *Proc 11th Int Coral Reef Symp* 2:1286-1288
494

495 Edward JKP, Mathews G, Patterson J, Kumar RR, Wilhelmsson D, Tamelander J, Linden O
496 (2008b) Status of coral reefs of the Gulf of Mannar, Southeastern India, Obura, D.,
497 Tamelander, J., Linden, O. (Eds.). *Ten years after bleaching – facing the consequences of*
498 *climate change in the Indian Ocean, CORDIO Status Report 2008, Mombasa, pp 45-60*
499

500 Edward JKP, Mathews G, Raj KD, Thinesh T, Patterson J, Tamelander J (2012) Coral reefs of
501 Gulf of Mannar, India - signs of resilience, Yellowlees, D., Hughes, T.P. (Eds.), *Proc*
502 *11th Int Coral Reef Symp* 18F
503

504 Edward JKP, Mathews G, Raj KD, Laju RL, Bharath MS, Arasamuthu A, Kumar PD, Bilgi, DS,
505 Malleshappa H (2018) Coral mortality in the Gulf of Mannar, Southeastern India, due to
506 bleaching caused by elevated Sea temperature in 2016. *Curr Sci* 114:1967-1972
507

508 English S, Wilkinson C, Baker V (1997) *Survey Manual for Tropical Marine Resources*, 2nd
509 edition. Australian Institute of Marine Science, Townsville
510

511 Fabricius KE (2005) Effects of terrestrial runoff on the ecology of corals and coral reefs: review
512 and synthesis. *Mar Pollut Bull* 50:125–146
513

514 Ferrier-Pagès C, Richard C, Forcioli D, Allemand D, Pichon M, Shick JM (2007) Effects of
515 temperature and UV radiation increases on the photosynthetic efficiency in four
516 Scleractinian coral species. *Biol Bull* 213:76–87
517

518 Fox MD, Williams GJ, Johnson MD, Radice VZ, Zgliczynski BJ, Kelly EL, Rohwer FL, Sandin,
519 SA, Smith JE (2018) Gradients in primary production predict trophic strategies of
520 mixotrophic corals across spatial scales. *Curr Biol* 28:3355-3363
521

522 Glynn P (1985) El Nino-associated disturbance to coral reefs and post disturbance mortality by
523 *Acanthaster planci*. *Mar Ecol Prog Ser* 26:295–300
524

525 Gove JM, McManus MA, Neuheimer AB, Polovina JJ, Drazen JC, Smith CR, Merrifield MA,
526 Friedlander AM, Ehses JS, Young CW, Dillon AK (2016) Near-island biological hotspots
527 in barren ocean basins. *Nat Commun* 7:1-8
528

529 Graham NA, Chabanet P, Evans RD, Jennings S, Letourneur Y, Aaron MacNeil M, McClanahan
530 TR, Öhman MC, Polunin NV, Wilson SK (2011) Extinction vulnerability of coral reef
531 fishes. *Ecol Lett* 14:341-348
532

533 Graham NA, Jennings S, MacNeil MA, Mouillot D, Wilson SK (2015) Predicting climate-driven
534 regime shifts versus rebound potential in coral reefs. *Nature* 518:94-97
535

536 Graham NA, Wilson SK, Carr P, Hoey AS, Jennings S, MacNeil MA (2018) Seabirds enhance
537 coral reef productivity and functioning in the absence of invasive rats. *Nature* 559:250-
538 253
539

540 Grottoli AG, Rodrigues LJ, Palardy JE (2006) Heterotrophic plasticity and resilience in bleached
541 corals. *Nature* 440:1186–1189
542

543 Head CE, Bayley DT, Rowlands G, Roche RC, Tickler DM, Rogers AD, Koldewey H, Turner
544 JR, Andradi-Brown DA (2019) Coral bleaching impacts from back-to-back 2015–2016
545 thermal anomalies in the remote central Indian Ocean. *Coral Reefs* 38:605-618
546

547 Heron SF, Maynard JA, van Hooidonk R, Eakin CM (2016) Warming trends and bleaching
548 stress of the world’s coral reefs 1985–2012. *Sci Rep* 6:38402
549 [doi.org/10.1038/srep38402]

550
551 Hock K, Wolff NH, Ortiz JC, Condie SA, Anthony KR, Blackwell PG, Mumby PJ (2017)
552 Connectivity and systemic resilience of the Great Barrier Reef. *PLoS Biol* 15:e2003355
553 [doi.org/10.1371/journal.pbio.2003355]
554
555 Hoegh-Guldberg O (1999) Climate change, coral bleaching and the future of the world's coral
556 reefs. *Mar Freshw Res* 50:839-866
557
558 Howells EJ, Beltran VH, Larsen NW, Bay LK, Willis BL, van Oppen MJH (2012) Coral thermal
559 tolerance shaped by local adaptation of photosymbionts. *Nat Clim Chang* 2:116–120
560
561 Hughes TP, Anderson KD, Connolly SR, Heron SF, Kerry JT, Lough JM, Baird AH, Baum JK,
562 Berumen ML, Bridge TC, Claar DC, Eakin CM, Gilmour JP, Graham NAJ, Harrison H,
563 Hobbs JPA, Hoey AS, Hoogenboom M, Lowe RJ, McCulloch MT, Pandolfi JM,
564 Pratchett M, Schoepf V, Torda G, Wilson SK (2018) Spatial and temporal patterns of
565 mass bleaching of corals in the Anthropocene. *Science* 359:80–83
566
567 Iluz D, Vago R, Chadwick NE, Hoffman R, Dubinsky Z (2008) Seychelles lagoon provides
568 corals with a refuge from bleaching. *Research Letters in Ecology* 2008:1–4
569
570 Jeevamani JJJ, Kamalakannan B, Nagendran NA, Chandrasekaran S (2013) Climate change
571 induced coral bleaching and algal phase shift in reefs of the Gulf of Mannar, India, in:
572 Nautiyal S., Rao K.S., Kaechele H., Raju K.V., Schaldach R. (Eds.), *Knowledge systems*
573 *of societies for adaptation and mitigation of impacts of climate change*. Book series:
574 *Environmental Science and Engineering*. Springer, Heidelberg, New York, Dordrecht,
575 London, pp 87–94
576
577 Kayanne H, Harii S, Ide Y, Akimoto F (2002) Recovery of coral populations after the 1998
578 bleaching on Shiraho Reef, in the southern Ryukyus, NW Pacific. *Mar Ecol Prog Ser*
579 239:93–103
580
581 Lesser MP (1996) Elevated temperatures and ultraviolet radiation cause oxidative stress and
582 inhibit photosynthesis in symbiotic dinoflagellates. *Limnol Oceanogr* 41:271–283
583
584 Liu GA, Strong E, Skirving W, Arzayus LF (2005) Overview of NOAA coral reef watch
585 program's near-real time satellite global coral bleaching monitoring activities. *Proc 10th*
586 *Int Coral Reef Symp* 1:1783-1793.
587
588 Loya Y, Sakai K, Yamazato K, Nakano Y, Sambali H, van Woesik R (2001) Coral bleaching:
589 the winners and the losers. *Ecol Lett* 4:122–131
590
591 Magnusson A, Skaug H, Nielsen A, Berg C, Kristensen K, Maechler M, van Benthem K, Bolker
592 B, Brooks M, (2017) Package 'glimm TMB'. R Package Version 0.2. 0.
593 <http://cran.nexr.com/web/packages/glimmTMB/glimmTMB.pdf>
594

595 Marshall PA, Baird AH (2000) Bleaching of corals on the Great Barrier Reef: differential
596 susceptibilities among taxa. *Coral Reefs* 19:155–163
597

598 McClanahan TR, Baird AH, Marshall PA, Toscano MA (2004) Comparing bleaching and
599 mortality responses of hard corals between southern Kenya and the Great Barrier Reef,
600 Australia. *Mar Pollut Bull* 48:327–335
601

602 McClanahan TR, Donner SD, Maynard JA, MacNeil MA, Graham NAJ, Maina J, Baker AC,
603 Alemu IJB, Beger M, Campbell SJ, Darling ES, Eakin CM, Heron SF, Jupiter SD,
604 Lundquist CJ, McLeod E, Mumby PJ, Paddock MJ, Selig ER, van Woessik R (2012)
605 Prioritizing key resilience indicators to support coral reef management in a changing
606 climate. *PLoS One* 7: e42884 [doi.org/10.1371/journal.pone.0042884]
607

608 McClanahan T, Schroeder R, Friedlander A, Vigliola L, Wantiez L, Caselle JE, Graham NA,
609 Wilson S, Edgar GJ, Stuart-Smith RD, Oddenyo RM (2019) Global baselines and
610 benchmarks for fish biomass: comparing remote reefs and fisheries closures. *Mar Ecol*
611 *Prog Ser* 612:167– 92
612

613 Meiaraj C Jeyapriya SP (2019) Marine water quality studies at Tuticorin harbour coastal area.
614 *Indian J Geomarine Sci* 48:943-946
615

616 Mumby PJ, Dahlgren CP, Harborne AR, Kappel CV, Micheli F, Brumbaugh DR, Holmes KE,
617 Mendes JM, Broad K, Sanchirico JN, Buch K (2006) Fishing, trophic cascades, and the
618 process of grazing on coral reefs. *Science* 311:98-101
619

620 Mumby PJ, Harborne AR, Williams J, Kappel CV, Brumbaugh DR, Micheli F, Holmes KE,
621 Dahlgren CP, Paris CB, Blackwell PG (2007) Trophic cascade facilitates coral
622 recruitment in a marine reserve. *P Natl Acad Sci U S A* 104:8362–8367
623

624 Muscatine L (1990) The role of symbiotic algae in carbon and energy flux in reef corals. *Coral*
625 *reefs* 25:1-29
626

627 NASA (2014) Moderate-resolution Imaging Spectroradiometer (MODIS) Aqua Chlorophyll
628 Data; 2014 Reprocessing. NASA Goddard Space Flight Center, Ocean Ecology
629 Laboratory, Ocean Biology Processing Group. NASA, USA. [doi:
630 10.5067/AQUA/MODIS/L3B/CHL/2014]
631

632 NASA (2017) Chlorophyll Concentrations (1-month – AQUA/MODIS). NASA Earth
633 Observations. NASA Goddard Space Flight Center, Ocean Ecology Laboratory, Ocean
634 Biology Processing Group.
635 https://neo.sci.gsfc.nasa.gov/view.php?datasetId=MY1DMM_CHLORA&year=2017
636

637 NOAA Coral Reef Watch (2019) NOAA Coral Reef Watch Version 3.1 Daily 5km Satellite
638 Regional Virtual Station Time Series Data for Southeast Florida, Mar. 12, 2013-Mar. 11,
639 2014. <https://coralreefwatch.noaa.gov/product/vs/data.php>
640

641
642 Obura DO (2005) Resilience and climate change: lessons from coral reefs and bleaching in the
643 Western Indian Ocean. *Estuar Coast Shelf Sci* 63: 353-372
644
645 Oliver TA, Palumbi SR (2011) Do fluctuating temperature environments elevate coral thermal
646 tolerance? *Coral Reefs* 30:429–440
647
648 Palumbi SR, Barshis DJ, Traylor-Knowles N, Bay RA (2014). Mechanisms of reef coral
649 resistance to future climate change. *Science* 344:895-898
650
651 Perry CT, Alvarez-Filip L (2019) Changing geo-ecological functions of coral reefs in the
652 Anthropocene. *Funct Ecol* 33:976-988
653
654 Raj K, Aeby GS, Mathews G, Bharath M, Rajesh S, Laju R, Arasamuthu A, Kumar D, Patterson
655 Edward JK (2016) Patterns in the abundance of fish and snail corallivores associated with
656 an outbreak of acute tissue loss disease on the reefs of Vaan Island in the Gulf of Mannar,
657 India. *Proc 13th Int Coral Reef Symp* 85-99
658
659 Raymundo LJ, Halford AR, Maypa AP, Kerr AM (2009) Functionally diverse reef-fish
660 communities ameliorate coral disease. *P Natl Acad Sci U S A* 106:17067-17070
661
662 Riegl B (1999) Corals in a non-reef setting in the southern Arabian Gulf (Dubai, UAE): fauna
663 and community structure in response to recurring mass mortality. *Coral Reefs* 18:63-73
664
665 Safaie A, Silbiger NJ, McClanahan TR, Pawlak G, Barshis DJ, Hench JL, Rogers JS, Williams
666 GJ, Davis KA (2018) High frequency temperature variability reduces the risk of coral
667 bleaching. *Nat commun* 9:1-12
668
669 Sampayo EM, Ridgway T, Bongaerts P, Hoegh-Guldberg O (2008) Bleaching susceptibility and
670 mortality of corals are determined by fine-scale differences in symbiont type. *P Natl*
671 *Acad Sci U S A* 105:10444–10449
672
673 Sano M (2004) Short-term effects of a mass coral bleaching event on a reef fish assemblage at
674 Iriomote Island, Japan. *Fisheries Science* 70:41–46
675
676 Smith JE, Shaw M, Edwards RA, Obura D, Pantos Sala E, Sandin SA, Smriga S, Hatay M,
677 Rohwer FL (2006) Indirect effects of algae on coral: algae-mediated, microbe-induced
678 coral mortality. *Ecol Lett* 9:835–845
679
680 Stafford-Smith MG (1993) Sediment-rejection efficiency of 22 species of Australian
681 scleractinian corals. *Marine Biology* 115:229-243
682
683 Torregiani JH, Lesser MP (2007) The effects of short-term exposures to ultraviolet radiation in
684 the Hawaiian Coral *Montipora verrucosa*. *J Exp Mar Biol Ecol* 340:194–203
685

686 [Van Hooidonk R, Huber M \(2009\) Quantifying the quality of coral bleaching predictions. Coral](#)
687 [Reefs 28: 579-587](#)
688

689 Van Hooidonk R, Maynard JA, Planes S (2013) Temporary refugia for coral reefs in a warming
690 world. *Nat Clim Chang* 3: 508-511
691

692 Van Hooidonk R, Maynard J, Tamelander J, Gove J, Ahmadi G, Raymundo L, Williams G,
693 Heron SF, Planes S (2016) Local-scale projections of coral reef futures and implications
694 of the Paris Agreement. *Sci Rep* 6: 39666 [doi: 10.1038/srep39666]
695

696 Vega Thurber R, Burkepile DE, Correa AMS, Thurber AR, Shantz AA, Welsh R, Pritchard C,
697 Rosales S (2012) Macroalgae decrease growth and alter microbial community structure
698 of the reef-building coral, *Porites astreoides*. *PLoS One* 7:e44246 [doi:
699 10.1371/journal.pone.0044246]
700

701 Vega Thurber RL, Burkepile DE, Fuchs C, Shantz AA, McMinds R, Zaneveld JR (2014)
702 Chronic nutrient enrichment increases prevalence and severity of coral disease and
703 bleaching. *Glob Chang Biol* 20:544-554
704

705 Visram S, Douglas AE (2007) Resilience and acclimation to bleaching stressors in the
706 scleractinian coral *Porites cylindrica*. *J Exp Mar Biol Ecol* 349:35-44
707

708 West JM, Salm RV (2003) Resistance and resilience to coral bleaching: implications for coral
709 reef conservation and management. *Conserv Biol* 17: 956-967
710

711 Williams GJ, Graham NA (2019) Rethinking coral reef functional futures. *Funct Ecol* 33:942-
712 947
713

714 Williams GJ, Gove JM, Eynaud Y, Zgliczynski BJ, Sandin SA (2015) Local human impacts
715 decouple natural biophysical relationships on Pacific coral reefs. *Ecography* 38:751-761
716

717 Williams GJ, Graham NA, Jouffray JB, Norström AV, Nyström M, Gove JM, Heenan A,
718 Wedding LM (2019) Coral reef ecology in the Anthropocene. *Funct Ecol* 33:1014-1022
719

720 Wooldridge SA (2009) Water quality and coral bleaching thresholds: Formalizing the linkage for
721 the inshore reefs of the Great Barrier Reef, Australia. *Mar Pollut Bull* 58:745–751
722

723 Wooldridge SA, Done TJ (2009) Improved water quality can ameliorate effects of climate
724 change on corals. *Ecol Appl* 19:1492-1499
725

726 WorldPop. 2017. India 100m Population, Version 2. University of Southampton. [doi:
727 10.5258/SOTON/WP00532]
728
729

730

731 **Figure legends**

732 **Fig. 1.** Map of Gulf of Mannar, India showing long-term monitoring sites surveyed between
733 2005 and 2017 and re-surveyed during the bleaching events in 2010 and 2016.

734

735 **Fig. 2.** Changes in average coral and algae cover on reefs subsequent to the 2010 and 2016
736 bleaching events (marked with arrows). Bleaching occurred in the Summer of each year and
737 annual surveys were conducted in the following November of each year. The data show the long-
738 term outcome from the bleaching events. Annual surveys were conducted on coral reefs at 21
739 islands within Gulf of Mannar. Data reflect mean and standard error of the mean (SEM) of
740 average island values.

741

742 **Fig. 3.** Change in mean coral and algae cover between 2005 and 2017 for islands surveyed
743 within Gulf of Mannar. This data reflects the island-specific outcome for benthic cover due to
744 successive bleaching events. Islands are ordered from west to east.

745

746 **Fig. 4.** Scatter plots showing modeled relationships (lines with shaded confidence intervals)
747 between changes in coral cover subsequent to bleaching events in relation to different co-factors
748 overlaid with survey data (points). The overlaid data points show the data used to create the
749 model and how well they fit the model. The black line and shaded confidence interval ($\pm 1.96 *$
750 SE) shows the marginal effect of the beta regression model for a) coral bleaching, b) pre-
751 bleaching coral cover, c) reef fish densities, and d) chlorophyll-a concentration. The marginal
752 effects show the predicted change in coral cover associated with a change in an ecological

753 covariate while accounting for all other factors included in the model (e.g., other ecological
754 drivers and nested random effects).

755

756 **Fig. 5.** Scatter plots showing significant modeled relationships (lines with shaded confidence
757 intervals) between percent change in algal cover subsequent to bleaching events in relation to
758 different co-factors overlaid with survey data (points). The overlaid data points show the data
759 used to create the model and how well they fit the model. The black line and shaded confidence
760 interval ($\pm 1.96 * SEM$) shows the marginal effect of the beta regression model for a) annual
761 maximum degree heating weeks, b) pre-bleaching algal cover, c) chlorophyll-a concentration,
762 and d) terrestrial runoff pollution. The marginal effects show the predicted change in algal cover
763 associated with a change in an ecological covariate while accounting for all other factors in the
764 model (e.g., other ecological drivers and nested random effects).

765

766 **Fig. 6.** Differences in cover among coral morphotypes through time. 21 islands were surveyed
767 each year. Note that the y-axis units differ among morphotypes. CS=submassive coral,
768 ACE=encrusting Acroporidae, CB=branching coral, ACT=table Acroporidae, ACD=digitate
769 Acroporidae, CE=encrusting coral, CF=foliose coral, ACB=branching Acroporidae,
770 CM=massive coral. Grey vertical lines delineate the bleaching years.

771

772 **Fig. 7.** Shifts in the proportional contribution of different coral morphotypes to overall coral
773 communities in response to multiple bleaching events through time. Data show the proportion of
774 the mean coral community represented by each morphotype in 2005 when surveys began and
775 2017 at the end of study. CS=submassive coral, ACE=encrusting Acroporidae, CB=branching

776 coral, ACT=table Acroporidae, ACD=digitate Acroporidae, CE=encrusting coral, CF=foliose
777 coral, ACB=branching Acroporidae, CM=massive coral.

778

779 **Fig. 8.** NMDS plot where the points indicate mean coral community composition for every
780 transect, site, and year within an island based on a Bray-Curtis dissimilarity matrix. The convex
781 hulls (polygons with shaded interiors) outline the multidimensional niche space of coral
782 community composition in the three time blocks of interest: before the 2010 bleaching event
783 (2005-2009, grey convex hulls), after the 2010 bleaching event and before the 2016 bleaching
784 event (2010-2015, blue convex hulls), and after the 2016 bleaching event (2016-2017, orange
785 convex hulls). Islands are ordered from west to east.

786

787 **Fig. 9.** Coral morphotypes influencing shifts in coral communities across islands following the
788 2010 and 2016 bleaching events. The left panel indicates morphotypes that significantly differed
789 before and after the 2010 bleaching event (i.e., 2005-2009 compared with 2010-2015). The right
790 panel indicates the morphotypes that significantly differed before and after the 2016 bleaching
791 event (i.e., 2010-2015 compared with 2016-2017). Morphotypes were identified as less or more
792 common based on indicator species analyses performed by island. Islands are ordered from west
793 to east. CS=submassive coral, ACE=encrusting Acroporidae, CB=branching coral, ACT=table
794 Acroporidae, ACD=digitate Acroporidae, CE=encrusting coral, CF=foliose coral,
795 ACB=branching Acroporidae, CM=massive coral.

796

797 **Fig. 10.** The predicted frequency of future bleaching events differs across islands within Gulf of
798 Mannar. Downscaled (4-km resolution) climate projections of predicted ocean surface warming

799 (from van Hooidonk et al. 2016) across the Gulf of Mannar (GoM) in the coming decades and
800 the subsequent year in which the onset of severe bleaching every 5 years (red) and annual severe
801 bleaching (blue) conditions is predicted to occur under RCP8.5 (high emissions) and a reduced
802 emissions scenario RCP4.5. Note the high local-scale (10s km) variation seen in the projections
803 across the GoM, with a clear gradient in the timing of bleaching onset from east to west for both
804 RCP8.5 and RCP4.5. Note that for RCP4.5, several islands do not experience severe bleaching
805 every 5 years (islands 1-11, 13, 15) or annual severe bleaching (islands 1-17) within the 83-year
806 modeled period (2006-2089).

807

808

Table 1. Predictor variables with their description and units used to model potential environmental drivers of coral reef resilience within Gulf of Mannar, India

Variable	Description and units	Min	Max	Data Source
Water clarity	Secchi disc divided by maximum water depth in meters	0.25	1.10	Reef survey
Sedimentation rate	mg of sediment per cm ² /day	33.94	47.39	Reef survey
Fish density	# fish per 250m ²	213.80	1346.20	Reef survey
Surface runoff pollution	Modeled diffusive plumes in 2013 based on impervious surface runoff from watershed	0.41	78.1	Halpern (2013) Ocean Health Index https://knb.ecoinformatics.org/#view/doi:10.5063/F1S180FS
Mean bleaching in 2010	% coral cover bleached during 2010 bleaching event	8	48	Reef survey
Mean bleaching in 2016	% coral cover bleached during 2016 bleaching event	27	99	Reef survey
Chlorophyll-a in 2010	Maximum chlorophyll <i>a</i> concentration (mg/m ³) observed at survey locations March-June of year	1.91	3.45	NASA MODIS-Aqua (https://oceandata.sci.gsfc.nasa.gov/MODIS-Aqua/Mapped/Monthly/4km/chlor_a/)
Chlorophyll-a in 2016	Maximum chlorophyll <i>a</i> concentration (mg/m ³) observed at survey locations March-June of year	1.55	2.82	NASA MODIS-Aqua (https://oceandata.sci.gsfc.nasa.gov/MODIS-Aqua/Mapped/Monthly/4km/chlor_a/)

Degree Heating Weeks (DHW) 2010	Maximum degree heating week values at 4 km spatial resolution for the 2010 bleaching year	2.40	4.78	NOAA Coral Reef Watch (ftp://ftp.star.nesdis.noaa.gov/pub/sod/mecb/crw/data/5km/v3.1/nc/v1.0/annual)
Degree Heating Weeks (DHW) 2016	Maximum degree heating week values at 4 km spatial resolution for the 2016 bleaching year	6.44	9.27	NOAA Coral Reef Watch (ftp://ftp.star.nesdis.noaa.gov/pub/sod/mecb/crw/data/5km/v3.1/nc/v1.0/annual)
Pop10k	Estimated number of people living within a 10km radius of the survey point in 2015 divided by the distance to coast squared	0.00	0.03	WorldPop (http://www.worldpop.org.uk/data/summary/?doi=10.5258/SO_TON/WP00532)

Table 2. Coefficient values for model covariates. Blank values indicate the covariate was not included in the model. Initial cover refers to the pre-bleaching percent of coral and algal cover for the coral and algal models, respectively. Human population refers to the human population size within 10 km of the nearest coastline, divided by the distance to the nearest coast squared to account for the hypothesized decreasing influence of humans with distance. Percent coral bleaching and maximum degree heating weeks were tested separately the models because they are colinear. The R^2 value was calculated based on in-sample model predictions.

Covariate	Percent change in coral cover model	Percent change in algal cover model
Intercept	-0.26	0.19
Initial cover (%)	-0.15	-0.13
Percent coral bleaching	-0.10	
Maximum DHW		0.07
Fish density	0.05	
Human population		
Chlorophyll-a concentration	0.03	-0.09
Pollution (impervious surface runoff)		0.13
R²	0.53	0.51

**Title: Coral reef resilience differs among islands within the Gulf of Mannar, southeast
India following successive coral bleaching events**

Journal Name: Coral Reefs

Authors: K. Diraviya Raj¹, Greta S. Aeby², G. Mathews¹, Gareth J. Williams³, Jamie Caldwell^{4,5}, R. L. Laju¹, M. Selva Bharath¹, P. Dinesh Kumar¹, A. Arasamuthu¹, N. Gladwin Gnana Asir¹, Lisa Wedding⁶, Andrew J. Davies⁷, Monica Moritsch⁸ and J. K. Patterson Edward¹

¹ Suganthi Devadason Marine Research Institute, Tuticorin, Tamil Nadu, India

² Dept of Biology and Environmental Science, Qatar University, Doha, Qatar

³ School of Ocean Sciences, Bangor University, Anglesey, UK

⁴ Hawaii Institute of Marine Biology, University of Hawaii at Manoa, Honolulu, HI, USA

⁵ ARC Centre of Excellence for Coral Reef Studies, James Cook University, Townsville, Australia

⁶ School of Geography and the Environment, University of Oxford, Oxford, United Kingdom

⁷ Department of Biological Sciences, University of Rhode Island, Kingston, RI, USA

⁸ United States Geological Survey, Western Geographic Science Center, Moffett Field, California, USA

Corresponding author: diraviyam_raj@yahoo.co.in

Supplemental Table 1. Site coordinates and average depths of long-term monitoring sites in Gulf of Mannar, India.

Island Name	GPS	Avg. Depth (m)
Shingle	9°14'43" N, 79°14'14" E	2.5
Krusadai	9°14'44" N, 79°13'21" E	2.9
Pullivasal	9°15'1" N, 79°11'49" E	2.1
Poomarichan	9°14'58" N, 79°10'51" E	2.8
Manoliputti	9°13'2" N, 79°9'2" E	2.4
Manoli	9°13'8" N, 79°8'11" E	2.9
Hare	9°12'19" N, 79°5'31" E	3.1
Mulli	9°11'24" N, 78°58'22" E	2.9
Valai	9°11'10" N, 78°56'20" E	2.1
Thalaiyari	9°11'4" N, 78°55'58" E	2.8
Appa	9°9'59" N, 78°49'22" E	3.5
Poovarasampatti	9°9'34" N, 78°45'1" E	3.1
Valimunai	9°9'17" N, 78°43'44" E	2.6
Anaipar	9°9'13" N, 78°41'26" E	3.3
Nallathanni	9°6'25" N, 78°34'21" E	3.1
Puluvnichalli	9°6'5" N, 78°32'26" E	3.8
Upputhanni	9°5'8" N, 78°29'37" E	3.8
Vilanguchalli	8°56'3" N, 78°16'5" E	4.0
Kariyachalli	8°57'9" N, 78°15'0" E	3.1
Koswari	8°51'55" N, 78°13'29" E	3.3
Vaan	8°50'4" N, 78°12'40" E	2.3

Supplemental Table 2. Coral categories used in the surveys. Corals were grouped by growth form and further classified by whether they were within the Family Acroporidae or not

Coral morphotype	Genus
Branching Acroporidae (ACB)	<i>Acropora</i>
Table Acroporidae (ACT)	<i>Acropora</i>
Digitate Acroporidae (ACD)	<i>Acropora, Montipora</i>
Foliose Acroporidae (ACF)	<i>Montipora</i>
Encrusting Acroporidae (ACE)	<i>Montipora, Astreopora</i>
Massive coral (CM)	<i>Pachyseris, Siderastrea, Pseudosiderastrea, Coscinaraea, Goniopora, Porites, Dipsastraea, Favites, Goniastrea, Platygyra, Leptoria, Hydnohora, Leptastrea, Cyphastrea, Galaxea, Acanthastrea, Lobophyllia, Symphyllia</i>
Submassive coral (CS)	<i>Pocillopora, Madracis, Pavona, Pseudosiderastrea, Psammocora, Goniopora, Porites, Favites, Platygyra, Plesiastrea</i>
Branching coral (CB)	<i>Pocillopora, Tubastrea, Dendrophyllia</i>
Foliose coral (CF)	<i>Pavona, Pachyseris, Echinopora, Mycedium, Merulina, Turbinaria</i>
Encrusting coral (CE)	<i>Madracis, Pavona, Culicia, Cyphastrea</i>

Supplemental Table 3. Results from PERMANOVA examining shifts in coral community structure after bleaching events for each of the 21 islands in Gulf of Mannar

Island	factor	degrees of freedom	Sum of squares	R ²	F	Pr(>F)
Vaan	time period	2	1.60574643	0.21697135	22.2921245	0.001
Vaan	site	1	0.63809627	0.08622072	17.7170208	0.001
Vaan	site:transect	1	0.15066316	0.02035788	4.18322811	0.002
Vaan	residual	139	5.00622436	0.67645006		
Vaan	total	143	7.40073021	1		
Koswari	time period	2	5.29320566	0.39689784	48.4134067	0.001
Koswari	site	1	0.36120807	0.02708429	6.6074566	0.001
Koswari	site:transect	1	0.08335401	0.00625009	1.52476656	0.187
Koswari	residual	139	7.59867604	0.56976779		
Koswari	total	143	13.3364438	1		
Kariyachalli	time period	2	1.37953716	0.20499781	24.1340865	0.001
Kariyachalli	site	1	1.33054293	0.19771731	46.5539299	0.001
Kariyachalli	site:transect	1	0.04672711	0.0069436	1.63491937	0.141
Kariyachalli	residual	139	3.97271438	0.59034128		
Kariyachalli	total	143	6.72952157	1		
Vilanguchalli	time period	2	4.64115672	0.32471194	36.7177369	0.001
Vilanguchalli	site	1	0.83268545	0.05825766	13.1753039	0.001
Vilanguchalli	site:transect	1	0.03444268	0.00240973	0.54497509	0.722
Vilanguchalli	residual	139	8.78486583	0.61462067		
Vilanguchalli	total	143	14.2931507	1		
Upputhanni	time period	2	2.1548811	0.32180454	36.0460736	0.001
Upputhanni	site	1	0.34218066	0.0511004	11.4477493	0.001
Upputhanni	site:transect	1	0.04437995	0.00662759	1.48474354	0.186
Upputhanni	residual	139	4.15480028	0.62046746		
Upputhanni	total	143	6.69624199	1		
Puluvnichalli	time period	2	1.05893256	0.16817832	15.0447782	0.001
Puluvnichalli	site	1	0.23271769	0.03695993	6.61267054	0.001
Puluvnichalli	site:transect	1	0.11305213	0.01795479	3.21237483	0.014
Puluvnichalli	residual	139	4.89178448	0.77690696		
Puluvnichalli	total	143	6.29648686	1		
Ilathanni	time period	2	5.4373574	0.50391653	79.8603195	0.001
Ilathanni	site	1	0.53340866	0.04943457	15.6687091	0.001
Ilathanni	site:transect	1	0.08746229	0.00810572	2.56917675	0.056

Ilathanni	residual	139	4.73196628	0.43854318		
Ilathanni	total	143	10.7901946	1		
Aipar	time period	2	1.90702688	0.23616153	25.1330115	0.001
Aipar	site	1	0.82014997	0.10156536	21.6177746	0.001
Aipar	site:transect	1	0.07444108	0.0092186	1.96214185	0.085
Aipar	residual	139	5.27347741	0.65305451		
Aipar	total	143	8.07509535	1		
Valimui	time period	2	2.25865834	0.29289849	31.3834953	0.001
Valimui	site	1	0.30075868	0.03900181	8.35793396	0.001
Valimui	site:transect	1	0.1500974	0.01946434	4.17113188	0.005
Valimui	residual	139	5.00188883	0.64863536		
Valimui	total	143	7.71140324	1		
Poovarasapatti	time period	2	5.00890496	0.39460551	50.4366576	0.001
Poovarasapatti	site	1	0.52527738	0.04138177	10.5784541	0.001
Poovarasapatti	site:transect	1	0.25716599	0.02025974	5.17901347	0.005
Poovarasapatti	residual	139	6.9021008	0.54375298		
Poovarasapatti	total	143	12.6934491	1		
Appa	time period	2	6.37198003	0.54628072	95.6347472	0.001
Appa	site	1	0.4916078	0.04214638	14.7567279	0.001
Appa	site:transect	1	0.17004172	0.01457797	5.10418954	0.009
Appa	residual	139	4.63066641	0.39699494		
Appa	total	143	11.664296	1		
Thalaiyari	time period	2	3.73006146	0.45355055	63.7462317	0.001
Thalaiyari	site	1	0.29847869	0.03629302	10.2019187	0.001
Thalaiyari	site:transect	1	0.12885715	0.01566817	4.40430164	0.006
Thalaiyari	residual	139	4.06673876	0.49448826		
Thalaiyari	total	143	8.22413606	1		
Valai	time period	2	1.65256361	0.21717098	21.684996	0.001
Valai	site	1	0.5193052	0.06824428	13.6286812	0.001
Valai	site:transect	1	0.14120117	0.01855589	3.70569306	0.007
Valai	residual	139	5.29643495	0.69602885		
Valai	total	143	7.60950492	1		
Mulli	time period	2	5.62602738	0.60763258	112.443325	0.001
Mulli	site	1	0.08804552	0.00950926	3.51940396	0.033
Mulli	site:transect	1	0.0674704	0.00728706	2.69696395	0.079
Mulli	residual	139	3.47738651	0.37557111		
Mulli	total	143	9.25892981	1		
Hare	time period	2	3.97559027	0.48140399	68.4163525	0.001
Hare	site	1	0.21788835	0.02638409	7.49932728	0.001

Hare	site:transect	1	0.02628597	0.00318297	0.90471608	0.444
Hare	residual	139	4.0385597	0.48902896		
Hare	total	143	8.25832429	1		
Manoli	time period	2	4.22224657	0.39292522	49.3842283	0.001
Manoli	site	1	0.50996171	0.04745739	11.9292254	0.001
Manoli	site:transect	1	0.07136396	0.00664118	1.66937382	0.146
Manoli	residual	139	5.9421023	0.5529762		
Manoli	total	143	10.7456745	1		
Manoliputti	time period	2	1.51615987	0.22239881	33.5030669	0.001
Manoliputti	site	1	2.10074049	0.30814837	92.8414614	0.001
Manoliputti	site:transect	1	0.0552236	0.00810051	2.44058692	0.098
Manoliputti	residual	139	3.14517807	0.46135231		
Manoliputti	total	143	6.81730203	1		
Poomarichan	time period	2	1.69195673	0.44575721	59.9576948	0.001
Poomarichan	site	1	0.11596432	0.03055157	8.21883096	0.003
Poomarichan	site:transect	1	0.02653748	0.00699147	1.88081168	0.107
Poomarichan	residual	139	1.96123272	0.51669975		
Poomarichan	total	143	3.79569124	1		
Pullivasal	time period	2	3.9829144	0.49760685	74.0370311	0.001
Pullivasal	site	1	0.23744466	0.02966523	8.82755474	0.001
Pullivasal	site:transect	1	0.04494083	0.0056147	1.67077923	0.162
Pullivasal	residual	139	3.73883916	0.46711322		
Pullivasal	total	143	8.00413904	1		
Krusadai	time period	2	4.55764249	0.45376639	63.3874619	0.001
Krusadai	site	1	0.36636973	0.03647637	10.1909035	0.002
Krusadai	site:transect	1	0.12287412	0.01223355	3.41785398	0.025
Krusadai	residual	139	4.99714208	0.49752369		
Krusadai	total	143	10.0440284	1		
Shingle	time period	2	7.03448031	0.54860441	90.4821765	0.001
Shingle	site	1	0.31796479	0.02479741	8.1797503	0.001
Shingle	site:transect	1	0.0668213	0.00521125	1.71900023	0.177
Shingle	residual	139	5.4032341	0.42138693		
Shingle	total	143	12.8225005	1		