

MANKELOW, J., RYAN, C.G., TAYLOR, P.C. et al. [2022]. International, multi-disciplinary, cross-section study of pain knowledge and attitudes in nursing, midwifery and allied health professions students. *BMC medical education* [online], (accepted).

International, multi-disciplinary, cross-section study of pain knowledge and attitudes in nursing, midwifery and allied health professions students.

MANKELOW, J., RYAN, C.G., TAYLOR, P.C. et al.

2022

This is the accepted manuscript for an article that will be published in BMC Medical Education. Once published, the version of record will be available from <https://bmcmmededuc.biomedcentral.com/>

1 International, multi-disciplinary, cross-section study of pain knowledge and
2 attitudes in nursing, midwifery and allied health professions students

- 3
- 4 Jagjit Mankelow^{a, m}
- 5 Cormac G. Ryan^a
- 6 Paul C. Taylor^a
- 7 Maire-Brid Casey^b
- 8 Jenni Naisby^c
- 9 Kate Thompson^d
- 10 Joseph G. McVeigh^e
- 11 Chris Seenan^f
- 12 Kay Cooper^g
- 13 Paul Hendrick^h
- 14 Donna Brownⁱ
- 15 William Gibson^j
- 16 Mervyn Travers^{j,k}
- 17 Norelee Kennedy^l
- 18 Cliona O’Riordan^l
- 19 Denis Martin^a

- 20
- 21 a. Teesside University, England
- 22 b. University College Dublin, Ireland
- 23 c. Northumbria University, England
- 24 d. Leeds Beckett University, England
- 25 e. University College Cork, Ireland
- 26 f. Glasgow Caledonian University, Scotland
- 27 g. Robert Gordon University, Scotland
- 28 h. University of Nottingham, England
- 29 i. University of Ulster, Northern Ireland
- 30 j. The University of Notre Dame Australia, Western Australia
- 31 k. Curtin University, Western Australia
- 32 l. University of Limerick, Ireland
- 33 m. Corresponding author

34

35 **Abstract**

36

37 **Background:** Persistent pain is a highly prevalent, global cause of disability.
38 Research suggests that many healthcare professionals are not well equipped
39 to manage pain, and this may be attributable at least in part to undergraduate
40 education. The primary aim of this study was to quantify and compare first
41 and final year nursing, midwifery and allied health professional (NMAHP)
42 students’ pain related knowledge and attitudes.. The secondary aim was to
43 explore what factors influence students’ pain related knowledge and attitudes.

44 **Methods:** In this cross-sectional study, 1154 first and final year healthcare
45 students, from 12 universities in five different countries completed the Revised

46 Neurophysiology of Pain Quiz (RNPQ) [knowledge] and the Health Care
47 Providers Pain and Impairment Relationship Scale (HC-PAIRS) [attitudes] .

48 **Results:** Physiotherapy was the only student group with statistically and
49 clinically improved pain related knowledge [mean difference, 95% CI] (3.4, 3.0
50 to 3.9, p=0.01) and attitudes (-17.2, -19.2 to 15.2, p=0.01) between first and
51 final year. Pain education teaching varied considerably from course to course
52 (0 to 40 hours), with greater levels of pain related knowledge and attitudes
53 associated with higher volumes of pain specific teaching.

54 **Conclusions:** There was little difference in pain knowledge and attitudes
55 between all first and final year NMAHP students other than physiotherapy.
56 This suggests that for most NMAHP disciplines, undergraduate teaching has
57 little or no impact on students' understanding of pain. There is an urgent need
58 to enhance pain education provision at the undergraduate level in NMAHPs.

59 The study protocol was prospectively registered at ClinicalTrials.Gov
60 (NCT03522857), <https://clinicaltrials.gov/ct2/show/NCT03522857>.

61

62

63

64 **BACKGROUND**

65

66 Pain is amongst the most common reason patients engage with health care^{1,2,3}. Pain, the
67 unpleasant sensory and emotional experience associated with actual or potential tissue
68 damage, can be classified by duration of symptoms as acute, sub-acute or chronic pain^{4,5}.
69 High rates of pain are present globally. For example, chronic pain affects 28 million people in
70 the UK alone⁶ and is often associated with significant disability⁷. Similarly, over three million
71 Australians identify as living with chronic pain. The economic burden amounts to AUD 73.2
72 billion each year including AUD 48.3 billion in lost productivity⁸. But the issue cannot be
73 adequately captured by dollars lost. Chronic pain negatively affects quality of life affecting
74 physical, mental, and social health⁹. The Prevalence Impact and Cost of Chronic Pain
75 (PRIME) study conducted in Ireland reported a chronic pain prevalence rate of 35.5%. Over
76 37% of those with pain reported moderate to severe pain-related disability¹⁰.

77

78 Multiple disciplines are involved in the management of pain, therefore it is vital that all health
79 care professionals (HPC) in every health care discipline are well equipped to manage this
80 problem and have a good knowledge of pain and positive attitudes towards function in those
81 with pain. Furthermore, it is imperative that this management is evidence-based and
82 guideline-compliant to ensure consistent high-quality care which is individualised^{11,12}.

83

84 Existing research suggests that many HCPs across the disciplines are not well equipped to
85 manage pain. Non-evidence based and inconsistent patterns of pain management occur

86 frequently in various health care settings which results in the high use of resources¹³⁻¹⁶.
87 Clinicians often do not feel confident or able to treat patients with persistent pain¹⁷⁻¹⁹.
88 Furthermore, there is evidence to suggest that HCPs' attitudes about the functional ability of
89 people in pain influences their management recommendations, and this in turn influences
90 patients' attitudes about pain and their health outcomes²⁰⁻²⁴. Patients often have a biomedical
91 understanding of their pain and link it to structural damage. These attitudes seem to be
92 influenced by their HCPs' pain knowledge and attitudes which are often also biomedical.^{25,26}
93 It is important that HCPs' pain attitudes and knowledge are evidence-based¹². However, it is
94 widely recognised that this is not always the case. It has been suggested that a part of this
95 problem may be the absence of adequate pain education in pre-registration training²⁷⁻²⁹.
96 Knowledge is accepted as a component of attitudes, which are key indicators of behaviour³⁰.
97 It has been proposed that improved understanding of pain amongst clinicians would improve
98 the delivery of evidence-based care, leading to better patient outcomes³¹.

100 The inadequacy of pain education in health care curricula has been observed throughout
101 Europe, New Zealand and Australia, the USA and Canada³²⁻³⁴. The first step towards
102 addressing the deficiency in pain education among HCPs would be to assess current pain
103 understanding amongst HCP students. A number of studies have explored this issue, however,
104 these studies are generally limited to single institutions, discrete regions or only a small
105 number of health care disciplines, reducing the generalisability of the findings³⁵⁻³⁹. If some
106 disciplines were found to have poorer pain-related understanding than others, this difference
107 could be explored, and pain education resources could be targeted accordingly.

108
109 The primary aim of this study was to quantify and compare nursing, midwifery and allied
110 health professional (NMAHP) students' knowledge and attitudes about pain management in
111 the first and the final year of their studies across a range of disciplines in multiple institutions
112 and countries. The secondary aim was to explore some of the factors that may influence
113 students' pain related knowledge and attitudes towards the functional ability of people with
114 pain.

115 116 **Method**

117 118 *Design*

119
120 In this observational, cross-sectional study the attitudes and knowledge of first and final year
121 NMAHP students were collected using two questionnaires to establish the change during
122 undergraduate health care degree courses. The attitudes and knowledge of students were
123 compared. The questionnaires were administered in the first semester for first years and as
124 close as possible to the completion of the degree course in the case of final year students.
125 Data on participants' age, gender, and year of study and course of study were collected.

126 127 **Ethics**

128
129 Ethical approval for this study was initially granted by Teesside University's (TU) School of
130 Health and Social Care Research Ethics and Governance Committee local ethics project
131 number 114/17. Each of the other eleven collaborating Universities obtained permission from
132 their respective University's research ethics and governance committee. The study protocol
133 was prospectively registered at ClinicalTrials.Gov NCT03522857,
134 <https://clinicaltrials.gov/ct2/show/NCT03522857>.

135 136 **Participants and recruitment**

137
138 First year and final year BSc and MSc pre-registration students were recruited between the
139 period of October 2017 to September 2019, from 12 universities and six disciplines across
140 Australia, England, Northern Ireland, the Republic of Ireland and Scotland. NMAHP

141 disciplines were selected based on those frequently involved in pain management, and
142 included physiotherapy, occupational therapy, paramedics, diagnostic radiography,
143 midwifery, and nursing. To meet the inclusion criteria for participation, individual students
144 needed to be in the first or final year of their studies within one of the aforementioned
145 disciplines.

146
147 Collaborating universities were invited to take part through informal networks, via on-site
148 academics acting as local pain education “champions”. Pain champions disseminated the
149 recruitment invitation to local programme leaders for delivery to students and either
150 disseminated and collected surveys physically or directed students to the online survey. A
151 reminder email was sent two weeks later. Additionally, where possible, the local champions
152 delivered short presentations to student groups to raise awareness of the study. Paper
153 questionnaires were made available at these presentations and a confidential drop box at a
154 different location from the distribution site was provided for questionnaire collection. The site
155 of questionnaire distribution and collection were kept separate in order to ensure that students
156 did not feel obliged to participate in the study. Participants were asked to complete the survey
157 only once when they received a reminder email. The participant information sheet explained
158 to prospective participants that consent was implied by completion of the survey.

159
160 Participating universities were invited to provide information about the extent and format of
161 pain education within the disciplines surveyed. Where possible respondents were asked to
162 quantify the time spent teaching pain education specifically and whether this involved one-off
163 lectures or complete modules with credit values. This data was then compiled and categorised
164 according to hours of pain education delivery. It was agreed that the public would be blind to
165 students University of study, so that institutional variation is quantifiable but specific
166 institutions could not be directly compared.

167 **Outcome measures**

168
169 The survey contained two questionnaires: 1) the 12-item Revised Neurophysiology
170 Questionnaire RNPQ⁴⁰ to measure pain knowledge, and 2) the 13-item Health Care Providers
171 Pain and Impairment Relationship Scale HC-PAIRS⁴¹ to measure attitudes towards chronic
172 pain. These questionnaires together were estimated to take less than 10 minutes to complete.

173 *The Revised Neurophysiology of Pain Questionnaire RNPQ*

174
175
176
177 This 12-item questionnaire was used to assess knowledge of pain neurophysiology.
178 Responses are marked ‘yes’, ‘no’ or ‘undecided’ the latter being important to prevent
179 respondents from guessing the answer. Scores range from 0-12 with high scores indicating a
180 good knowledge of pain neurophysiology. The RNPQ was developed from the original 19-
181 item Neurophysiology of Pain Test⁴². It was found to have reasonable internal consistency
182 person separation index =0.84 and good test-retest reliability with an intra-class correlation
183 coefficient value of ICC =0.97. The RNPQ has now been used consistently in patient, student,
184 clinician and clinical administration staff studies since its inception⁴³⁻⁴⁷. Furthermore, it is a
185 discipline generic rather than a discipline specific questionnaire, therefore fit for a multi-
186 disciplinary group. There is no established minimally clinically important difference MCID
187 for the RNPQ. However, this can be tentatively estimated as half the baseline SD presented in
188 previous studies⁴⁸⁻⁵¹. Based upon data from Catley *et al.* (2013)⁴⁰ the MCID for RNPQ
189 knowledge was set at 0.9 points or 7.3%.

190 *The 13-item modified Health Care Providers Pain and Impairment Relationship Scale HC-PAIRS*

191
192
193
194 The modified HC-PAIRS⁴¹ measures HCPs’ attitudes towards patients with chronic pain and
195 their functional ability. It features a 7-point Likert scale in 13-items with scores ranging from

196 13 to 91, the lower score indicates a more positive attitude towards pain. Psychometric
197 properties of the HC-PAIRS are well established. Excellent internal consistency has been
198 demonstrated Cronbach's $\alpha=0.92^{52}$ as well as good test-retest reliability [ICC=0.84] 95%
199 confidence interval 0.78-0.89. Latimer, Maher and Refshauge (2004)⁵³ also observed its
200 adequate responsiveness to change. Overall, the psychometric properties of the HC-PAIRS
201 are superior to other tools and hence it is consistently widely used^{52, 54-56}. A previous study
202 about student HCPs estimated an MCID of 4.5 for the HC-PAIRS⁵⁷. However, Dworkin *et al.*
203 (2008)⁵¹ advise that MCIDs should be population specific, thus, for this study, the MCID was
204 set at 4.2 points 4.6% based upon half the baseline values for HC-PAIRS data from student
205 HCPs (Colleary *et al.* 2017)⁴³. Originally designed to question attitudes about chronic low
206 back pain Houben *et al.* (2004)⁴¹ suggest that it is a good measure of chronic pain generically.
207

208 **Data analysis**

209
210 Missing data for the HC-PAIRS was managed as follows: data sets were retained if they were
211 full sets or had only one answer missing^{41,57}. Those with more than one unanswered question
212 were discarded from the data set. Missing answers were replaced with a neutral response, 4⁵⁷.
213 There are no recommendations within the literature regarding how missing data from the
214 RNPQ should be handled. Thus, for consistency, a similar approach to that of the HC-PAIRS
215 was taken in that a single missing answer in a questionnaire was replaced with a '0' value
216 indicating an incorrect answer. Questionnaires with more than once missing answer were
217 discarded.
218

219 Data were analysed using SPSS version 26.0. The data were found to have a normal
220 distribution after a visual inspection of histograms and Q-Q plots, and statistical analysis via
221 the Shapiro-Wilk test. Descriptive statistics are presented as the mean and 1SD. Data were
222 analysed using two-way ANOVA with year of study first or final, and discipline of degree
223 Physiotherapy; Occupational therapy; Nursing; Midwifery; Paramedic; Radiographer as
224 independent variables for the HC-PAIRS and RNPQ separately. The interaction effects of the
225 two independent variables year of study*discipline of degree were also investigated. In
226 addition, a series of post-hoc independent samples t-tests were undertaken to identify where
227 differences lay between individual disciplines and the first and final year of study in each
228 discipline. Correlation analyses were also undertaken as part of a secondary analysis to
229 explore the association between hours of pain education teaching, and knowledge and/or
230 attitude scores, adjusting for age, gender, year of study and discipline. A p-value of <0.05 was
231 considered statistically significant.
232

233 **Results**

234 **Response rate**

235
236
237 There were 1156 respondents from the 12 universities out of 4067 invitations to participate,
238 representing a 28% response rate. Eight incomplete paper questionnaires were removed for
239 HC-PAIRS six sets and RNPQ two sets as they were almost entirely incomplete. In addition,
240 162 RPNQ questionnaire data sets were removed as an incorrect version of the questionnaire
241 was accidentally circulated due to human error. This left 1154 respondents who completed
242 and returned surveys adequately, and whose data were analysed. Fifteen of these respondents
243 had left one question unanswered in one of their surveys, nine in the HC-PAIRS
244 questionnaire and eight in RNPQ.
245

246 Participants had a mean (SD) age of 26 (8) years, were predominantly female 82% and
247 studying at BSc level 83%. A breakdown of surveys returned can be seen in Table 1, by
248 University and by discipline. Nursing students were categorised together irrespective of
249 speciality as not all respondents disclosed their area of speciality. Some universities returned
250 more surveys than others, and some disciplines had a higher response rate than others, with

251 physiotherapists and nursing students returning the largest numbers of surveys. The overall
 252 response rate was lower amongst final year students except in nursing which was heavily
 253 dominated by a strong return at one University.

254
 255
 256
 257

Table 1. Number of respondents per University and breakdown of number of respondents in first and final year by discipline

University Code	Number of responses	Disciplines Surveyed	First year respondents	Final year respondents
1	11	Occupational therapists	43	34
2	134			
3	8	Physiotherapists	266	104
4	514			
5	12	Paramedics	68	9
6	126			
7	51	Midwives	32	11
8	47			
9	11	Nurses	235	312
10	120			
11	97	Diagnostic radiographers	31	9
12	23			
Total	1154	Total	675	479

258
 259
 260
 261

HC-PAIRS

262 The two-way ANOVA for HC-PAIRS found a significant independent effect of both year of
 263 study $p=0.001$ and discipline $p=0.001$. Table 2 lists the mean HC-PAIRS attitude scores for
 264 individual professions. First year mean values ranged from 54.4 to 60.0 lower values
 265 indicating more positive attitudes. In final year they ranged from 37.5 to 56.1. Between first
 266 and final year the greatest improvement in attitudes to pain was shown by physiotherapy
 267 students, with a mean difference 95% confidence interval [CI] of -17.2 [-19.2 to -15.2] points.
 268 All of the other professions showed clinically insignificant, less than or equal to the MCID,
 269 and statistically insignificant changes from first to final year. This is with the exception of
 270 nursing which showed a clinically insignificant but statistically significant improvement -2.2
 271 [3.6 to -0.7] $p=0.03$. A two-way ANOVA revealed that there was a statistically significant
 272 interaction ($p<0.01$) between the effects of the two independent variables year of
 273 study*discipline of degree.

274
 275
 276
 277


Table 2 HC-PAIRS, pain attitude scores for first and final year by profession

Profession n=total number	1 st Year Mean (SD)	Final Year Mean (SD)	Mean Difference	95% CI	P-value
OT n=77	56.4 (8.6)	52.8 (7.3)	-3.7	-7.4 to 0.1	0.51
Physiotherapy n=370	54.7 (8.8)	37.5 (9.1)	-17.2	-19.2 to -15.2	0.01*

Paramedics n=77	55.7 (8.2)	52.1 (8.3)	-3.6	-9.4 to 2.3	0.23
Midwifery n=43	60.0 (9.6)	56.1 (8.6)	-3.9	-10.5 to 2.7	0.24
Nursing n=547	57.1 (8.0)	55.0 (8.5)	-2.2	-3.6 to -0.7	0.03*
Diagnostic Radiography n=40	54.4 (9.0)	51.6 (8.9)	-2.9	-9.7 to 4.1	0.40

278 **Legend:** SD, standard deviation; CI, confidence interval; HC-PAIRS, Health Care Providers
 279 Pain and Impairment Relationship Scale. P-values were calculated using independent t-tests.
 280 * Indicates statistical significance at $p < 0.05$.

281
 282 As physiotherapy was the only discipline that showed a clinically and statistically significant
 283 change from first to final year, secondary analysis was carried out within that discipline to
 284 explore if all universities performed equally well as shown in Figure 1. Seven of the eight
 285 universities, which had first and final year respondents, showed a difference between the year
 286 groups, exceeding the MCID of -4.2, ranging from -8 to -23 units. University 6 had a mean
 287 change of less than -4.2. This may have been an artefact of the very small number of
 288 respondents from this sub-group. There were only 17 first year respondents and only two final
 289 year respondents thus it was not representative of the final year. Two universities, codes 5 and
 290 9, had only first year participants and not final years; one University did not have any
 291 physiotherapy respondents code 12.



293
 294 **Figure 1** First and final year mean SD HC-PAIRS scores for physiotherapy cohorts in
 295 Universities 1-11 12 did not include any physiotherapists

296
 297 **RNPQ**

298
 299 Two-way ANOVA for RNPQ found a significant independent effect of year of study $p = 0.044$
 300 and discipline $p = 0.025$. Table 3 lists the mean RNPQ knowledge scores for individual

301 professions with higher scores indicating better knowledge of pain neurophysiology. The
 302 minimum mean (SD) score in the first year was 5.7 (2.0) and the maximum was 7.3 (1.8).
 303 Final year scores ranged from a minimum of 5.7 (2.1) and maximum 9.1 (2.0). The biggest
 304 improvement in pain knowledge between first and final year is shown by physiotherapy
 305 students with a change of 3 points, a difference which was statistically significant $p=0.01$. All
 306 the other professions showed clinically small less than or equal to the MCID and statistically
 307 insignificant differences from first to final year. A two-way ANOVA revealed that there was
 308 a statistically significant interaction ($p<0.01$) between the effects of the two independent
 309 variables year of study*discipline of degree.

310
 311
 312


Table 3 RNPQ pain knowledge scores for first and final year by profession

Profession Total numbers, n=	1 st Year Mean (SD)	Final Year Mean (SD)	Mean Difference	95% CI	P-value
Occupational Therapy n=77	5.9 (1.8)	6.4 (1.6)	0.5	0.3 to 1.3	0.26
Physiotherapy n=370	5.7 (2.0)	9.1 (2.0)	3.4	3.0 to 3.9	0.01*
Paramedics n=77	6.1 (1.5)	5.7 (2.1)	-0.4	-0.9 to 1.8	0.48
Midwifery n=43	6.1 (2.0)	7.00 (1.4)	0.9	0.6 to 2.3	0.24
Nursing n=547	5.9 (2.0)	6.2 (2.0)	0.3	0.1 to 0.7	0.06
Diagnostic Radiography n=40	7.3 (1.8)	6.0 (2.1)	-1.3	-0.4 to 3.0	0.13

313
 314
 315
 316
 317
 318
 319
 320
 321
 322
 323
 324
 325

Legend: RNPQ, revised Neurophysiology Questionnaire; SD, standard deviation; CI, confidence interval; P-values were calculated using independent t-tests. * Indicates statistical significance at $p<0.05$.

Once again, as they were the only discipline to have demonstrated a statistical and clinical difference between first and final year cohorts, secondary analysis of the physiotherapy data were carried out to explore if some universities made greater gains than others. The minimum mean difference was 1.1 95%CI [2.9 to 5.2] and the maximum mean difference was 4.7 [4.0 to 5.3] see Figure 2. Thus, the size of pain knowledge improvement was not consistently high in all physiotherapy cohorts at all of the universities sampled, but always exceeded the MCID of 0.9 points.



326
 327
 328
 329
 330
 331
 332
 333
 334
 335
 336
 337
 338
 339
 340
 341
 342
 343


Figure 2 First and final year mean RNPQ scores for physiotherapy cohorts in Universities 1-11 12 did not include any physiotherapists

Secondary analysis

Multiple linear regression analyses were completed to explore the association between hours of pain education in all of the disciplines studied, and knowledge and attitude scores respectively, adjusting for age, gender, year of study and discipline.

For both dependent variables, pain knowledge and pain attitudes, hours of pain education teaching was found to be an independent predictor though the strength of the relationship was small (RNPQ β value=0.11, $p=0.01$ and HC-PAIRS β value =0.15, $p=0.001$).

The amount of focused pain teaching at the time of data collection varied considerably between universities and disciplines. Figure 3 reflects this difference with physiotherapy departments generally delivering the greatest amounts of pain education teaching.



344
345 **Figure 3** Approximate hours of pain education teaching in each discipline and University
346

347 **Discussion**
348

349 There has been recent suggestion that there is a need to shift understanding about pain on a
350 societal level in order to optimise and contemporise care⁵⁸. HCPs will be a key sector of
351 society to focus upon as they will influence the pain understanding of others. Furthermore,
352 targeting HCP students, whose understanding may be more malleable, may be the optimal
353 point at which to target HCPs. An important step in this process is to survey pain attitudes and
354 knowledge amongst future health care workers to quantify current levels of understanding and
355 identify if training could be enhanced. Accordingly, this study compared the pain knowledge
356 and attitudes in first and final year students, across six disciplines, at 12 institutions, in five
357 countries. To date, this is the largest, international cross-sectional study to quantify the
358 knowledge and attitudes about pain amongst NMAHP students. There were differences in
359 pain knowledge and attitudes between year of study and between disciplines. There was also
360 a year of study*discipline interaction effect. Of the six disciplines, physiotherapy had the
361 greatest mean differences between the first and final year for both the RNPQ and the HC-
362 PAIRS which were clinically and statistically significant. In contrast there was little
363 difference between first and final year values for both knowledge and attitudes scores in the
364 other disciplines.
365

366 The nursing cohorts showed the least improvement in attitudes with a mean difference of -2.2,
367 well below the MCID of 4.2 of all disciplines, yet statistical testing showed the difference to
368 be significant $p=0.03$. It is likely that this was due to the larger sample size for the nursing
369 group and thus greater statistical power. However, the magnitude of the difference is well
370 below the MCID and thus likely to be clinically unimportant.
371

372 Direct comparison with existing literature is difficult as a large portion of the literature uses
373 different outcome measures, and studies using similar outcome measures include
374 physiotherapy students only. The improvement in attitudes for physiotherapy students over
375 the duration of a degree programme, as measured by the HC-PAIRS, in this study are greater
376 than previously reported^{36, 60, 61}, but scores were not quite as high as the changes measured in
377 RCTs following targeted, brief pain science education interventions directly addressing

378 knowledge and attitudes in physiotherapists⁴³ and NMAHPs⁴⁷. This suggests there is scope
379 for greater changes on the observed degree programmes in this study.

380
381 Whilst Carroll *et al.*, (2020)³⁹ found greater improvement amongst their nursing cohorts'
382 attitudes (1.6% - 7% amongst different nursing specialities) than in this study, 2.4%, our
383 findings accord with Amponsah *et al.* (2020)⁶² and Leahy *et al.* (2019)⁶³ that final year nurses
384 have considerable deficits in pain knowledge and attitudes. Mukoka, Olivier and Ravat
385 (2019)⁶² found more positive attitudes in their nursing and occupational therapy students but
386 not as positive among their physiotherapy students. Overall the findings from this study
387 generally concur with the existing body of literature that suggests there is a deficiency in pain
388 knowledge and attitudes towards pain in final year HCP students. Many previous studies
389 noted an improvement in HCP students' knowledge and attitudes from first to final year^{38, 62,}
390 ⁶⁵, and while we found this among physiotherapy students, it was not the case overall.
391 Worryingly, Ryan *et al.*, (2010)⁶¹ noted that non-health care students demonstrated a 3.9 point
392 3.7% mean difference in HC-PAIRS 15-point questionnaire from first to final year. This is
393 similar if not better than the difference seen for the health care students in the current study,
394 apart from physiotherapy students. The comparatively poor difference in pain attitudes
395 demonstrated for most disciplines other than physiotherapy in this study may be attributable
396 in part to a biomedical model-based curricula impeding the natural small biopsychosocial
397 shift with time seen in the non-health care programme sample studied by Ryan *et al.* (2010)⁶¹.

398
399 There were larger volumes of pain specific teaching on the physiotherapy courses relative to
400 the other NMAHP disciplines in the current study (Figure 3). This is perhaps unsurprising as
401 physiotherapists may be perceived to play a larger role in pain management than some of the
402 other disciplines. The larger differences between first and final year in physiotherapy are
403 likely in part due to the higher volumes of pain specific teaching. Within our data, there was a
404 moderate/high correlation between difference in attitudes and knowledge and higher volumes
405 of pain teaching $r=0.5, p=0.16$ and $r=0.7, p=0.03$ respectively. This provides a rationale for
406 larger volumes of pain teaching within NMAHP curricula.

407
408 An additional factor influencing student pain knowledge and attitudes that has not been
409 explored in this study is the effect of clinical placements. This aspect of health care education
410 warrants further investigation as it may positively or negatively⁶⁶ influence pain management
411 behaviours.

412
413 Thompson *et al.* (2018)²⁷ propose an array of reasons that inhibit the implementation of
414 effective pain education into pre-registration health care programs. These authors suggest that
415 all health care disciplines have different curricula pressures placed upon them by internal and
416 external bodies, and pain education may not yet be recognised as a priority topic for these
417 health care disciplines. Furthermore, professional opportunities to manage pain are not always
418 the focus of some disciplines and some disciplines may play a larger role in the care pathway
419 than others and thus arguably may need higher levels of knowledge and attitudes relative to
420 other disciplines. However, each discipline involved in this study may encounter people with
421 pain directly and as such it is important that they all have appropriate knowledge and attitudes
422 to provide patients with clear and consistent high quality basic pain management advice For
423 example, in diagnostic radiography patient interaction may be limited, nevertheless, even if
424 interactions are brief, correct communication is critical^{67,68}. Kyei *et al.* (2014)⁶⁹ observe the
425 need for good radiographer communication skills because there is only a short time frame
426 available to establish a relationship with patients. Furthermore, the reports that an extended
427 scope radiographer may be required to complete are often shown to patients and it is
428 important that these report any anomalies within the context of age-related changes and the
429 possibility that an individual's pain may not always be linked to the findings⁷⁰⁻⁷³. Ultimately,
430 failures from a key team member in a pain management multi-disciplinary team can affect the
431 pain management efforts of the whole team and thus patient outcome.

432

433 **Limitations**

434

435 The observational, cross-sectional nature of this study means that no claim of cause and effect
436 can be made. Measuring students in the first and final year meant it was impossible to identify
437 at what points in training pain knowledge and attitudes changed, and thus understand what
438 aspects of training may influence change. Future studies should employ a longitudinal design,
439 measuring students yearly to identify potential triggers for improving knowledge and attitudes
440 towards pain, taking into account student placements and their impact. In addition, a
441 longitudinal study would help to establish if the cross-sectional differences seen in this study
442 are comparable to changes in the same cohort of students followed over the course of their
443 degree. There is a need for pain management behaviours resulting from education to be
444 investigated specifically, though changes in knowledge and attitudes can be predictors of
445 behaviour Ajzen, (2020)³⁰.

446

447 Some universities and disciplines returned more responses than others, thus there may be a
448 response bias in this snapshot of pain knowledge and attitudes in students.

449

450 There was not an *a priori* sample size calculation. Instead, the researchers attempted to recruit
451 as many participants as possible from the institutions involved. As such it is possible that the
452 study is underpowered for some disciplines and may explain the lack of statistical differences
453 between first and final year students for some disciplines. However, the magnitude of the
454 differences between first and final year, would be less likely to be influenced by sample size
455 and those differences were small and well below the MCID for all except the physiotherapy
456 group.

457

458 In a small minority of cases the number of participants in sub-groups were very small. In such
459 cases the sub-analysis was exploratory and should be interpreted with caution.

460

461 The differing sample sizes may have been due, in part, to final year students being on clinical
462 placements at different times, and thus being less receptive to email invitations to participate
463 in this study. Other factors may have been survey fatigue; the National Students Survey NSS
464 was underway in the England, Scotland and Northern Ireland at a similar time to data
465 collection, as well as individual module feedback surveys at many universities. Despite this,
466 every attempt was made to access final year students at the end of their degree programme,
467 including extending the study for a further year of data collection.

468

469

470 Participant self-selection may have influenced sample size. The pain champion at each of the
471 universities may not have equally reflected all disciplines. The majority of pain champions
472 were physiotherapists. This may have accounted for the larger numbers of physiotherapists
473 relative to other disciplines for example only two universities represented paramedic training
474 whilst 11 universities represented physiotherapy. Arguably medical doctors, such as general
475 practitioners GPs and anaesthesiologists, will have more involvement in pain management
476 than some NMAHPs and it would be illuminating to include this health care discipline in
477 future studies of student knowledge and/or attitudes.

478

479 In one quarter of the physiotherapy courses investigated there was up to 40 hours of pain
480 education teaching and this is reflected in the difference in knowledge and attitudes in first
481 and final year physiotherapy students. This volume of teaching may not be reflective of all
482 physiotherapy courses, and may inflate the overall variance between disciplines. Furthermore,
483 the time spent teaching pain education is of interest, but the content of that education is also
484 important (Mankelov *et al.* 2021). This study did not investigate the content of pain
485 education being delivered and future studies should investigate the impact of educational
486 content on pain related knowledge and attitudes.

486

487 **Conclusions**

488

489 To date, this is the largest investigation of HCP student pain related knowledge and attitudes
490 amongst NMAHPs, including 12 universities and six disciplines in five countries. Only
491 physiotherapy students showed statistically and clinically significant improvements in pain
492 related attitudes and knowledge from first to final year. The differences were correlated with
493 the volume of pain teaching received. Given that clinicians with more positive attitudes
494 towards pain are more likely to make evidence-based recommendations, in turn improving
495 patient outcomes, this study highlights the need to improve NMAHP pain education.

496

497

498 **Declarations**

499

500 **Author Contribution**

501 JM, PT, CR and DM conceived the idea. All authors collected data and JM, CR and DM
502 analysed the data under supervision of. JM, PT, CR and DM drafted the paper. All authors
503 contributed to the interpretation of results and in making an important intellectual
504 contribution to the manuscript. All authors read and approved the final manuscript.

505

506 **Funding Source**

507 The Musculoskeletal Association for Chartered Physiotherapists has permitted funding to be
508 used for dissemination of this study.

509

510 **Competing interests Statement**

511 There are no competing interests for any contributing authors.

512

513 **Availability of Data and Materials Statement**

514

515 The datasets used and/or analysed during the current study are available from the
516 corresponding author on reasonable request.

517

518 **Ethics approval and consent to participate**

519

520 Ethical approval for this study was granted by Teesside University's TU School of Health and
521 Social Care Research Ethics and Governance Committee local ethics project number 114/17.
522 Each of the other eleven collaborating universities obtained permission from their respective
523 University's research ethics and governance committee. Informed consent was obtained from
524 all subjects. All methods were carried out in accordance with relevant guidelines and
525 regulations.

526

527 **Consent for publication**

528

529 Not applicable.

530

531 **Acknowledgements**

532

533 The authors acknowledge the contribution of colleagues at each university who assisted data
534 collection.

535

536 **Abbreviations**

537

538 ANOVA – Analysis of variance

539 AUD – Australian Dollar

540 BSc – Bachelor of Science

541 CI – Confidence Interval

542 GBD – Global Burden of Disability

543 GP – general practitioner
544 HCP – Health Care Professionals
545 HC-PAIRS – Health Care Providers Pain and Impairment Relationship Scale
546 IASP – International Association for the Study of Pain
547 MCID – Minimally clinically important difference
548 N - number
549 NMAHP – Nursing, Midwifery and Allied Health Professionals
550 NSS – National Student Survey
551 PRIME – Prevalence Impact and Cost of Chronic Pain
552 RNPQ – Revised Neuro Physiology Questionnaire
553 SD – Standard deviation
554 SPSS – Statistical Package for the Social Sciences
555 TU – Teesside University
556 UK – United Kingdom
557
558
559
560
561

562 **References**

- 563
564 1. Mäntyselkä P, Kumpusalo E, Ahonen R, Kumpusalo A, Kauhanen J, Viinamäki H, et al.
565 Pain as a reason to visit the doctor: a study in Finnish primary health care. *Pain*. 2001; doi:
566 10.1016/s0304-39590000361-4.
567
568 2. Finley CR, Chan DS, Garrison S, Korownyk C, Kolber MR, Campbell S, et al. What are
569 the most common conditions in primary care? Systematic review. *Canadian Family*
570 *Physician*. 2018;64(11):832–840.
571
572 3. Lim YZ, Chou L, Au RTM, Seneviwickrama KLMD, Cicuttini FM, Briggs AM. People
573 with low back pain want clear, consistent and personalised information on prognosis,
574 treatment options and self-management strategies: a systematic review. *Journal of*
575 *Physiotherapy*. 2019; 65(3):124–135.
576
577 4. IASP Subcommittee on Taxonomy. Pain terms: a list with definitions and notes on usage.
578 Recommended by the IASP Subcommittee on Taxonomy. *Pain*. 1979;6:249–52.
579
580 5. Jess M-A, Hamilton S, Ryan C, Wellburn S, Alexanders J, Spence D, Martin D. Exploring
581 the origin of pain subclassification, with emphasis on low back pain: a systematic scoping
582 review. *JBIS Database of Systematic Reviews and Implementation Reports*. 2020; doi:
583 10.11124/JBISRIR-D-19-00383.
584
585 6. Fayaz A, Croft P, Langford RM, Donaldson LJ, Jones GT. Prevalence of chronic pain in
586 the UK: A systematic review and meta-analysis of population studies, *BMJ Open*. 2016; doi:
587 10.1136/bmjopen-2015-010364.
588
589 7. GBD 2017 Global Burden of Disease Study. Seattle. 2017. [Accessed on 1 Mar 2021].
590 Available from:
591 http://www.healthdata.org/sites/default/files/files/policy_report/2019/GBD_2017_Booklet.pdf
592 [%0Ahttp://ghdx.healthdata.org/sites/default/files/record-attached-](http://ghdx.healthdata.org/sites/default/files/record-attached-files/IHME_GBD_2017_DISABILITY_WEIGHTS_Y2018M11D08.XLSX)
593 [files/IHME_GBD_2017_DISABILITY_WEIGHTS_Y2018M11D08.XLSX](http://ghdx.healthdata.org/sites/default/files/record-attached-files/IHME_GBD_2017_DISABILITY_WEIGHTS_Y2018M11D08.XLSX).
594
595 8. Deloitte Access Economics. The Cost of Pain in Australia: PainAustralia. 2019. [Accessed
596 on 1 Mar 2021]. Available from: [https://www.painaustralia.org.au/static/uploads/files/the-](https://www.painaustralia.org.au/static/uploads/files/the-cost-of-pain-in-australia-final-report-12mar-)
597 [14](https://www.painaustralia.org.au/static/uploads/files/the-cost-of-pain-in-australia-final-report-12mar-</p></div><div data-bbox=)

wfxbrfyboams.pdf?fbclid=IwAR0OaQTiCWV6XNqTWIM3Yg6uO_w2_q7J6Pzj9rj0M2svm
osNtJ-FhuDGqOo.

598
599
600

601 9. Dueñas M, Ojeda B, Salazar A, Mico JA, Failde I. A review of chronic pain impact on
602 patients, their social environment and the health care system. *Journal of Pain Research*. 2016;
603 doi: 10.2147/JPR.S105892.
604

605 10. Raftery MN, Sarma K, Murphy AW, la Harpe DD, Normand C, McGuire BE. Chronic
606 pain in the Republic of Ireland—Community prevalence, psychosocial profile and predictors
607 of pain-related disability: Results from the Prevalence, Impact and Cost of Chronic Pain
608 PRIME study. Part 1. *Pain*. 2011;1525:1096–103.
609

610 11. Gardner T, Refshauge K, McAuley J, Hübsher M, Goodall S, Smith L. Goal setting
611 practice in chronic low back pain. What is current practice and is it affected by beliefs and
612 attitudes? *Physiotherapy Theory and Practice*. 2017;3410:795–805.
613

614 12. Stevans JM, Delitto A, Samannaaz SK, Patterson CG, Smith CN, Schneider MJ, et al.
615 Risk Factors Associated With Transition From Acute to Chronic Low Back Pain in US
616 Patients Seeking Primary Care. *JAMA Neurology*. 2021; doi:
617 10.1001/jamanetworkopen.2020.37371.
618

619 13. Toye F, Seers K, Allcock N, Briggs M, Carr E, Andrews JA, Barker K. Patients
620 experiences of chronic non-malignant musculoskeletal pain: A qualitative systematic review.
621 *British Journal of General Practice*. 2013; doi: 10.3399/bjgp13X675412.
622

623 14. Hart OR, Uden RM, McMullan JE, Ritchie MS, Williams TD, Smith BH. A study of
624 National Health Service management of chronic osteoarthritis and low back pain. *Primary
625 Health Care Research & Development*. 2015;162:157–166.
626

627 15. Arumugam V, MacDermid JC, Walton D, Grewal R. Attitudes, knowledge and behaviors
628 related to evidence-based practice in health professionals involved in pain management.
629 *International Journal of Evidence-Based Healthcare*. 2018;162:107–118.
630

631 16. Cowell I, O'Sullivan P, O'Sullivan K, Poyton R, McGregor A, Murtagh G. Perceptions of
632 physiotherapists towards the management of non-specific chronic low back pain from a
633 biopsychosocial perspective: A qualitative study. *Musculoskeletal Science and Practice*.
634 Elsevier Ltd. 2018;38:113–119.
635

636 17. Upshur C, Luckmann R, Savageau J. Primary care provider concerns about management
637 of chronic pain in community clinic populations. *Journal of General Internal Medicine*. 2006;
638 216:652–5.
639

640 18. O'Rorke JE, Chen I, Genao I, Panda M, Cykert S. Physicians comfort in caring for
641 patients with chronic nonmalignant pain. *American Journal of Medical Science*.
642 2007;3332:93–100.
643

644 19. Synnott A, O'Keeffe M, Bunzli S, Dankaerts W, O'Sullivan P, O'Sullivan K.
645 Physiotherapists may stigmatise or feel unprepared to treat people with low back pain and
646 psychosocial factors that influence recovery: A systematic review. *Journal of Physiotherapy*.
647 2015;612:68–76.
648

649 20. Parsons S, Harding G, Breen A, Foster N, Pincus T, Vogel S, et al. The Influence of
650 Patients and Primary Care Practitioners Beliefs and Expectations About Chronic
651 Musculoskeletal Pain on the Process of Care; A Systematic Review of Qualitative Studies.
652 *Clinical Journal of Pain*. 2006;231:91–98.

- 653
654 21. Darlow B, Fullen BM, Dean S, Hurley DA, Baxter GD, Dowell A. The association
655 between health care professional attitudes and beliefs and the attitudes and beliefs, clinical
656 management, and outcomes of patients with low back pain: A systematic review. *European*
657 *Journal of Pain*. 2012;161:3–17.
658
659 22. Gardner T, Refshauge K, Smith L, McAuley J, Hübsher M, Goodall S. Physiotherapists'
660 beliefs and attitudes influence clinical practice in chronic low back pain: a systematic review
661 of quantitative and qualitative studies. *Journal of Physiotherapy*. 2017;633:132–143.
662
663 23. Gibbs M, Morrison N, Marshall P. Biomedical beliefs explain the clinical decisions made
664 by exercise-based practitioners for people with chronic low back pain. *SPINE*. 2021; doi:
665 10.1097/BRS.0000000000003698.
666
667 24. Mankelow J. Exploring healthcare students pain knowledge and attitudes towards
668 function in people with pain. PhD thesis. Teesside University. 2021.
669
670 25. Lin IB, O'Sullivan PB, Coffin JA, Mak DB, Toussaint S, Straker LM. Disabling chronic
671 low back pain as an iatrogenic disorder: A qualitative study in Aboriginal Australians. *BMJ*
672 *Open*. 2013; doi: 10.1136/bmjopen-2013-002654.
673
674
675 26. Setchell J, Costa N, Ferreira M, Makovey J, Nielsen M, Hodges PW. Individuals
676 explanations for their persistent or recurrent low back pain: A cross-sectional survey. *BMC*
677 *Musculoskeletal Disorders*. 2017; doi: 10.1186/s12891-017-1831-7.
678
679 27. Thompson K, Johnson MI, Milligan J, Briggs M. Twenty-five years of pain education
680 research-what have we learned? Findings from a comprehensive scoping review of research
681 into pre-registration pain education for health professionals. *Pain*. 2018;15911:2146–21.
682
683 28. Hogans BB, Watt-Watson J, Wilkinson P, Carr ECJ, Gordon DB. Perspective: update on
684 pain education. *Pain*. 2019;1599:1681–1682.
685
686 29. Polacek C, Christopher R, Mann M, Udall M, Craig T, Deminski M, et al. Healthcare
687 professionals perceptions of challenges to chronic pain management. *American Journal of*
688 *Managed Care*. 2020;264:E135–E139.
689
690 30. Ajzen I. The theory of planned behavior: Frequently asked questions. *Human Behavior*
691 *and Emerging Technologies*. 2020; doi:10.1002/hbe2.195.
692
693 31. Chief Allied Health Professions Officers Team. Allied Health Professions into Action:
694 Using Allied Health Professionals to transform health, care and wellbeing. NHS England.
695 2017. [Accessed on 1 March 2021]. Available from: [https://www.england.nhs.uk/wp-](https://www.england.nhs.uk/wp-content/uploads/2017/01/ahp-action-transform-hlth.pdf)
696 [content/uploads/2017/01/ahp-action-transform-hlth.pdf](https://www.england.nhs.uk/wp-content/uploads/2017/01/ahp-action-transform-hlth.pdf).
697
698 32. Briggs E, Carr E, Whittaker M. Survey of undergraduate pain curricula for healthcare
699 professionals in the United Kingdom, *European Journal of Pain*. 2011: [Accessed on 1 March
700 2021]. Available from: www.britishpainsociety.org.
701
702 33. European Federation of IASP Chapters. The Pain Management Core Curriculum for
703 Undergraduate Medical Education. 2013. [Accessed on 1 Mar 2021] Available from:
704 [https://europeanpainfederation.org/wp-](https://europeanpainfederation.org/wp-content/uploads/2018/10/CoreCurriculumPainManagement-EFIC-June-2013_FINAL1.pdf)
705 [content/uploads/2018/10/CoreCurriculumPainManagement-EFIC-June-2013_FINAL1.pdf](https://europeanpainfederation.org/wp-content/uploads/2018/10/CoreCurriculumPainManagement-EFIC-June-2013_FINAL1.pdf).
706
707 34. Briggs EV, Battelli D, Gordon D, Kopf A, Ribeiro S, Puig MM, Kress HG. Current pain

- 708 education within undergraduate medical studies across Europe: Advancing the Provision of
709 Pain Education and Learning APPEAL study. *BMJ Open*. 2015; doi: 10.1136/bmjopen-2014-
710 006984.
- 711
- 712 34. Shipton EE, Bate F, Garrick R, Steketee C, Shipton EA, Visser EJ. Pain medicine content,
713 teaching and assessment in medical school curricula in Australia and New Zealand. *BMC*
714 *Medical Education*, 2018;181:139–161.
- 715
- 716 35. Ali N, Thomson D. A comparison of the knowledge of chronic pain and its management
717 between final year physiotherapy and medical students. *European Journal of Pain*.
718 2009;131:38–50.
- 719
- 720 36. Morris H, Ryan C, Lauchlan D, Field M. Do medical student attitudes towards patients
721 with chronic low back pain improve during training? A cross-sectional study. *BMC Medical*
722 *Education*. 2012;121:2–7.
- 723
- 724 37. Eyob T, Mulatu A, Abrha H. Knowledge And Attitude Towards Pain Management
725 Among Medical And Paramedical Students Of An Ethiopian University. *Journal of Pain &*
726 *Relief*. 2014; doi: 10.4172/2167-0846.1000127.
- 727
- 728 38. Adillón C, Lozano È, Salvat I. Comparison of pain neurophysiology knowledge among
729 health sciences students: a cross-sectional study. *BMC Research Notes*.
730 2015;doi:10.1186/s13104-015-1585-y.
- 731
- 732 39. Carroll SP, Augeard N, Tennant J, Seenan C. How do the attitudes, confidence,
733 knowledge and understanding differ in pre-registration healthcare students towards treating
734 people with chronic pain: an observational, cross-sectional study. *European Journal of*
735 *Physiotherapy*. 2020; doi: 10.1080/21679169.2020.1746830.
- 736
- 737 40. Catley MJ, O'Connell NE, Moseley GL. How good is the neurophysiology of pain
738 questionnaire? A rasch analysis of psychometric properties, *Journal of Pain*. 2013;148:818–
739 827.
- 740
- 741 41. Houben RMA, Vlaeyen JWS, Peters M, Ostelo RWJG, Wolters PMJC, Stomp-van den
742 Berg SGM. Health care providers attitudes and beliefs towards common low back pain: factor
743 structure and psychometric properties of the HC-PAIRS. *Clinical Journal of Pain*.
744 2004;201:37–44.
- 745
- 746 42. Moseley L. Unraveling the barriers to reconceptualization of the problem in chronic pain:
747 The actual and perceived ability of patients and health professionals to understand the
748 neurophysiology. *Journal of Pain*. 2003;44:184–189.
- 749
- 750 43. Colleary G, O'Sullivan K, Griffin D, Ryan CG, Martin DJ. Effect of pain neurophysiology
751 education on physiotherapy students understanding of chronic pain, clinical recommendations
752 and attitudes towards people with chronic pain: a randomised controlled trial, *Physiotherapy*.
753 2017;1034:423–429.
- 754
- 755 44. Alodaibi F, Alhowimel A, Alsobayel H. Pain neurophysiology knowledge among
756 physical therapy students in Saudi Arabia: A cross-sectional study, *BMC Medical Education*.
757 2018; doi: 10.1186/s12909-018-1329-5.
- 758
- 759 45. Louw A, Podalak J, Zimney K, Schmidt S, Puentedura EJ. Can pain beliefs change in
760 middle school students? A study of the effectiveness of pain neuroscience education.
761 *Physiotherapy Theory and Practice*. 2018;347:542–550.
- 762

- 763 46. Maguire N, Chesterton P, Ryan C. The effect of pain neuroscience education on sports
764 therapy and rehabilitation students knowledge, attitudes, and clinical recommendations
765 toward athletes with chronic pain. *Journal of Sport Rehabilitation*. 2019;285:438–443.
766
- 767 47. Mankelov J, Ryan CG, Taylor PC, Martin DJ. The effect of pain neurophysiology
768 education on healthcare students knowledge, attitudes and behaviours towards pain: A mixed-
769 methods randomised controlled trial. *Musculoskeletal Science and Practice*. 2020; doi:
770 10.1016/j.msksp.2020.102249.
771
- 772 48. Cohen J. *Statistical power analysis for the behavioural sciences*. 2nd Ed. Hillsdale NJ:
773 Lawrence Earlbaum Associates. 1988.
774
- 775 49. Norman GR, Sloan JA, Wyrwich KW. Interpretation of changes in health-related quality
776 of life: The remarkable universality of half a standard deviation. *Medical Care*.
777 2003;415:582–592.
778
- 779 50. Copay AG, Glassman SD, Subach BR, Berven S, Schuler TC, Carreon LY. Minimum
780 clinically important difference in lumbar spine surgery patients: a choice of methods using the
781 Oswestry Disability Index, Medical Outcomes Study questionnaire Short Form 36, and Pain
782 Scales. *Spine Journal*. 2008;86: 968–974.
783
- 784 51. Dworkin RH, Turk, DC, Wyrwich KW, Beaton D, Cleeland CS, Farrar JT, et al.
785 Interpreting the Clinical Importance of Treatment Outcomes in Chronic Pain Clinical Trials:
786 IMMPACT Recommendations. *Journal of Pain*. 2008;92:105–121.
787
- 788 52. Moran RW, Rushworth WM, Mason J. Investigation of four self-report instruments
789 FABT, TSK-HC, Back-PAQ, HC-PAIRS to measure healthcare practitioners attitudes and
790 beliefs toward low back pain: Reliability, convergent validity and survey of New Zealand
791 osteopaths and manipulative physio. *Musculoskeletal Science and Practice*. 2017;32:44–50.
792
- 793 53. Latimer, J, Maher C, Refshauge K. The Attitudes and Beliefs of Physiotherapy Students
794 to Chronic Back Pain. *Clinical Journal of Pain*. 2004; 201:45–50.
795
- 796 54. Evans DW, Breen AC, Pincus T, Sim J, Underwood M, Vogel S, Foster NE. The
797 effectiveness of a posted information package on the beliefs and behavior of musculoskeletal
798 practitioners: The UK chiropractors, osteopaths, and musculoskeletal physiotherapists low
799 back pain management COMPLEMENT randomized trial. *Spine*. 2010;358:858–866.
800
- 801 55. Briggs AM, Slater H, Smith AJ, Parkin-Smith GF, Watkins K, Chua J. Low Back Pain-
802 related Beliefs and Likely Practice Behaviours Among Final-year Cross-discipline Health
803 Students, *European Journal of Pain*. 2013;175:766–775.
804
- 805 56. Macdonald RJD, Vaucher P, Esteves JE. The beliefs and attitudes of UK registered
806 osteopaths towards chronic pain and the management of chronic pain sufferers - A cross-
807 sectional questionnaire based survey. *International Journal of Osteopathic Medicine*. 2018;
808 doi: 10.1016/j.ijosm.2018.07.003.
809
- 810 57. Domenech J, Sánchez-Zuriaga D, Segura-Ortí E, Espejo-Tort B, Lisón JF. Impact of
811 biomedical and biopsychosocial training sessions on the attitudes, beliefs, and
812 recommendations of health care providers about low back pain: A randomised clinical trial.
813 *Pain*. 2011;15211:2557–2563.
814
- 815 58. Chalmers KJ, Madden VJ. Shifting beliefs across society would lay the foundation for
816 truly biopsychosocial care. *Journal of Physiotherapy*. 2019;653:121–122.
817

- 818 60. Quinn T, Ryan C, Jones D. Physiotherapy students attitudes towards the functional ability
819 of patients with chronic low back pain. *Pain and Rehabilitation*. 2014;3737:20–23.
820
- 821 61. Ryan C, Murphy D, Clark M, Lee A. The effect of a physiotherapy education compared
822 with a non-healthcare education on the attitudes and beliefs of students towards functioning in
823 individuals with back pain: An observational, cross-sectional study. *Physiotherapy*.
824 2010;962:144–150.
825
- 826 62. Amponsah AK, Kyei-Dompim J, Kyei EF, Oduro E, Afaya RA, Aphoto CK. Final Year
827 Nursing Students Knowledge and Attitudes regarding Childrens Pain, Pain Research and
828 Management. 2020; doi: 10.1155/2020/7283473.
829
- 830 63. Leahy A, O'Keeffe, M, Robinson K, O'Sullivan K. The beliefs of healthcare students
831 about the harmfulness of daily activities for their back: a cross-sectional study. *European*
832 *Journal of Physiotherapy*. 2019; doi: 10.1080/21679169.2019.1630854.
833
- 834 64. Mukoka G, Olivier B, Ravat S. Level of knowledge, attitudes and beliefs towards patients
835 with chronic low back pain among final year school of therapeutic sciences students at the
836 university of the Witwatersrand – A cross-sectional study. *South African Journal of*
837 *Physiotherapy*. 2019; doi: 10.4102/sajp.v75i1.683.
838
- 839 65. Goodrich C. Students and faculty members knowledge and attitudes regarding pain
840 management: a descriptive survey. *Journal of Nursing Education*. 2005;453:140–142.
841
- 842 66. Springer S, Gleicher H, Hababou H. Attitudes and beliefs about musculoskeletal pain and
843 its association with pain neuroscience knowledge among physiotherapy students in Israel.
844 *Israel Journal of Health Policy Research*. 2018; doi: 10.1186/s13584-018-0266-4.
845
- 846 67. Karran EL, Hillier SL, Yau Y-H, McAuley JH, Moseley GL. A quasi-randomised,
847 controlled, feasibility trial of GLITtER Green Light Imaging Interpretation to Enhance
848 Recovery-a psychoeducational intervention for adults with low back pain attending secondary
849 care. *PeerJ*. 2018; doi: 10.7717/peerj.4301.
850
- 851 68. Jarvik JG, Meier EN, James KT, Gold LS, Tan KW, Kessler LG. The Effect of Including
852 Benchmark Prevalence Data of Common Imaging Findings in Spine Image Reports on Health
853 Care Utilization Among Adults Undergoing Spine Imaging: A Stepped-Wedge Randomized
854 Clinical Trial. *JAMA network open*. 2020; doi: 10.1001/jamanetworkopen.2020.15713.
855
- 856 69. Kyei KA, Antwi WK, Opoku SY, Hemans S, Anim-Sampong S, Engel-Hills P.
857 Radiographers knowledge, attitude and challenges on pain management. *The South African*
858 *Radiographer*. 2014;521:22–25.
859
- 860 70. Carragee EJ, Todd AF, Miller JL, Carragee JM. Discographic, MRI and psychosocial
861 determinants of low back pain disability and remission: a prospective study in subjects with
862 benign persistent back pain. *The Spine Journal*. 2005; doi: 10.1016/j.spinee.2004.05.250.
863
- 864 71. Suri P, Boyko EJ, Goldberg J, Forsberg CW, Jarvik JG. Longitudinal associations
865 between incident lumbar spine MRI findings and chronic low back pain or radicular
866 symptoms: Retrospective analysis of data from the longitudinal assessment of imaging and
867 disability of the back LAIDBACK. *BMC Musculoskeletal Disorders*. 2014; doi:
868 10.1186/1471-2474-15-152.
869
- 870 72. Brinjikji W, Diehn FE, Jarvik JG, Carr CM, Kallmes DF, Murad MH, Luetmer PH. MRI
871 findings of disc degeneration are more prevalent in adults with low back pain than in
872 asymptomatic controls: A systematic review and meta-analysis. *American Journal of*

873 Neuroradiology. 2015;3612:2394–2399.

874

875 73. Brinjikji W, Luetmer PH, Comstock B, Bresnahan BW, Chen LE, Deyo RA, Halabi S,
876 Turner JA, Avins AL, James K, Wald JT, Kallmes DF, Jarvik JG. Systematic Literature
877 Review of Imaging Features of Spinal Degeneration in Asymptomatic Populations. American
878 Journal of Neuroradiology. 2015;364:811–816.

879

880 74. Mankelov J, Ryan C, Taylor P, Atkinson G, Martin D. A Systematic Review and Meta-
881 Analysis of the Effects of Biopsychosocial Pain Education upon Health Care Professional
882 Pain Attitudes, Knowledge, Behavior and Patient Outcomes. Journal of Pain. 2022;23:1-24.
883 doi: 10.1016/j.jpain.2021.06.010.

884

885

886