



Modelling and quantifying tomorrow's risks from natural hazards

Gemma Cremen^{a,*}, Carmine Galasso^{a,b}, John McCloskey^c

^a Department of Civil, Environmental and Geomatic Engineering, University College London (UCL), London, UK

^b Scuola Universitaria Superiore (IUSS) Pavia, Pavia, Italy


^c School of GeoSciences, The University of Edinburgh, Edinburgh, UK



HIGHLIGHTS

- We explore work that has modelled and quantified future risks from natural hazards.
- We focus on individual components of risk (hazard, exposure, and vulnerability).
- We discuss risk-modelling challenges, and their effect on relevant decision making.

GRAPHICAL ABSTRACT



ARTICLE INFO

Article history:

Received 21 October 2021

Received in revised form 15 December 2021

Accepted 15 December 2021

Available online 22 December 2021

Editor: Jay Gan

Keywords:

Natural hazards
Risks
Decision making
Climate change
Population increase
Ageing infrastructure

ABSTRACT

Understanding and modelling future risks from natural hazards is becoming increasingly crucial as the climate changes, human population grows, asset wealth accumulates, and societies become more urbanised and interconnected. This need is recognised by the 2015–2030 Sendai Framework for Disaster Risk Reduction, which emphasises the importance of preparing for the disasters that our world may face tomorrow through strategies/policies that aim to minimise uncontrolled development in hazardous areas. While the vast majority of natural-hazard risk-assessment frameworks have so far focused on static impacts associated with current conditions and/or are influenced by historical context, some authors have sought to provide decision makers with risk-quantification approaches that can be used to cultivate a sustainable future. This Review documents these latter efforts, explicitly examining work that has modelled and quantified the individual components that comprise tomorrow's risk, i.e., future natural hazards affected by climate change, future exposure (e.g., in terms of population, land use, and the built environment), and the evolving physical vulnerabilities of the world's infrastructure. We end with a discussion on the challenges faced by modellers in determining the risks that tomorrow's world may face from natural hazards, and the constraints these place on the decision-making abilities of relevant stakeholders.

Contents

1.	Introduction	2
2.	Future (natural) hazard	2
2.1.	Climate modelling	2
2.2.	Emission scenarios	3

* Corresponding author.

E-mail address: g.cremen@ucl.ac.uk (G. Cremen).

2.3.	Additional dynamic hazard features	3
2.4.	Modelling of future non-climate-related hazards	3
3.	Future exposure	4
3.1.	Modelling population and socioeconomic growth.	4
3.2.	Land-use change and urbanisation	4
4.	Future (physical) vulnerability	5
4.1.	Assessing increasing vulnerability	5
4.2.	Accounting for adapted vulnerability	6
4.3.	Non-physical vulnerability	6
5.	Risk metrics	6
6.	Modelling future risk	7
6.1.	Challenges	7
6.2.	Perspectives.	8
7.	Conclusions	8
	Data availability	8
	Declaration of competing interest	8
	Acknowledgments	8
	References	8

1. Introduction

Global population is projected to increase to almost 11 billion by 2100 (United Nations, 2019). By this time, 75% or more of people are expected to be living in cities (Angel et al., 2012) and there is general consensus in the scientific community that climactic conditions will be significantly different from those currently experienced (Hallegatte, 2009). Even by 2030, the total amount of urban land located in high-frequency flood zones is predicted to increase by 33% compared to 2015 (Güneralp et al., 2015). These changes are set against a backdrop of ever-increasing wealth disparities between the world's rich and poor (Brian, 2015), which can act to either amplify or mitigate the impacts of natural-hazard disasters. All of these facts point to the critical importance of evaluating risks to natural hazards through a future-focused lens, so that properly informed decisions are taken today on the spatial planning and risk-prevention measures/policies that will shape the coming decades (e.g., Neumann et al., 2015; Lavell et al., 2012; Burby et al., 2000; Ferrier and Haque, 2003; Huppert and Sparks, 2006).

The need to consider the long-term implications of natural hazards is recognised by well-established intergovernmental organisations, such as the United Nations (UN), the European Union (EU), and the World Bank, among others. The UN Sendai Framework for Disaster Risk Reduction (UNISDR, 2015) inherently focuses on the future, through its aim to substantially reduce disaster loss by 2030; the goal to mitigate forthcoming disasters is further underscored by Article 8 of the UN Framework Convention on Climate Change “Paris Agreement” (Savaresi, 2016). The EU encourages a forward-looking approach to assessing disaster risks, particularly highlighting the importance of accounting for climate change impacts (European Commission, 2020). The World Bank underlines the obligation of disaster risk managers to “move instead towards risk assessments that can guide decision makers towards a resilient future” (Fraser et al., 2016). Furthermore, the Intergovernmental Panel on Climate Change (IPCC) state that “effective risk reduction and adaptation strategies consider the dynamics of vulnerability and exposure and their linkages with socioeconomic processes, sustainable development, and climate change” (IPCC, 2014).

Despite the evident necessity, most studies of natural-hazard risks do not consider its long-term evolutionary nature, relying instead on past or static observations of society within current environmental conditions (Gallina et al., 2016). A somewhat recent examination of 80 open-source risk assessment tools for natural hazards (GFDRR, 2014) found that none accounted for future risk in an explicit sense. Even catastrophe loss models used by the insurance industry tend to disregard the dynamic relationship between built-environment portfolios and the natural hazards that affect them (Fraser et al., 2016).

The primary purpose of this paper is to provide an overview of state-of-practice methods and approaches that feature in efforts of the literature to

quantify and model risk from natural hazards in tomorrow's world – rather than to conduct a systematic review of all work that has been published on future natural-hazard risk; interested readers are referred to Ward et al. (2020) for the latter type of assessment at the global scale. We define the term ‘risk’ as the convolution of hazard, exposure, and vulnerability, in line with many previous studies of natural hazards like flooding and earthquakes (e.g., Birkmann and Welle, 2015; Erdik, 2017). All three of these components are changing over time under natural and human influences and impacting future risk estimates. We explicitly examine attempts to model each of these separate components into the future, primarily from a physical sciences and engineering perspective (this means that social vulnerability is broadly outside the scope of this study, for example). We also review the metrics that are used to define future risk, and end with a discussion on the challenges associated with realising and leveraging future natural-hazard risk assessments.

2. Future (natural) hazard

A hazard is defined according to the UN General Assembly as “a process, phenomenon or human activity that may cause loss of life, injury or other health impacts, property damage, social and economic disruption or environmental degradation” (UNDRR, 2020). Natural hazards are classified under a number of categories (IFRC, 2020), (1) geophysical (which covers earthquakes, landslides, tsunamis, and volcanic eruptions); (2) hydrological (which includes avalanches and floods); (3) climatological (which accounts for extreme temperatures, drought, and wildfires); (4) meteorological (which incorporates cyclones/hurricanes and storm surges); and (5) biological (consisting of disease epidemics and insect/animal plagues). They may also be categorised as (i) rapid onset (e.g., earthquakes, flash floods, and landslides); or (ii) slow onset (e.g., sea level rise, increasing temperatures, and droughts) (Tosun and Howlett, 2021). This Review specifically relates to the aforementioned classifications (1) to (4).

2.1. Climate modelling

The quantification of tomorrow's natural hazards commonly involves accounting for future climactic conditions (see Fig. 1). Climate change raises sea levels globally, affects the intensity/frequency of the strongest storms, increases extreme temperatures, and modifies precipitation patterns (e.g., Dessler, 2021). Therefore, climate change projections are frequently used as input to models for heat-, drought-, wind-, fire-, and flood-related risk assessments (e.g., Zscheischler et al., 2018; Kwadijk et al., 2010; Dowdy et al., 2019). These data are derived from so-called climate models that are combined with information on possible future scenarios of emissions. Note that the term climate model used here incorporates a variety of approaches for quantifying climate dynamics (in line with Van



Fig. 1. A typical methodological approach to modelling future climate-related hazards. Adapted from Fig. 8 of [Salman and Li \(2018\)](#). Information on possible scenarios of future emissions are combined with climate models that are then appropriately downscaled to provide finer resolution data for hazard analysis in the region of interest. The downscaled climate models are used to simulate values for dynamic weather variables, from which hazardous events can be derived. Then, local intensities associated with these events are calculated. The last three steps of the process can be repeated for many temporal instants in the future, to generate frequency exceedance curves for the hazard intensity measures of interest.

[Vuuren et al., 2011](#)) including (in order of increasing complexity) Simple Climate Models (SCMs), Earth-System Models of Intermediate Complexity (ESMIC), and Atmosphere-Ocean General Circulation Models (AOGCMs). SCMs (such as the Model for the Assessment of Greenhouse-Gas Induced Climate Change; e.g., [Wigley and Raper, 2001](#)) describe the ocean-atmosphere system as a collection of global or hemispherical boxes, and may be used to provide dynamic estimates of global mean temperature and sea-level rise. AOGCMs are the main tool used for climate projections, and comprehensively capture dynamics of the atmosphere-ocean system as well as other physical phenomena (e.g., land-surface processes). Examples of commonly used AOGCMs include the HadCM3 (Hadley Centre Coupled Model version 3; e.g., [Gordon et al., 2000](#)) and the HadGEM2-ES (Hadley Centre Global Environment Model version 2 - Earth-System; e.g., [Bellouin et al., 2011](#)), which were developed by the UK Met Office. ESMIC operate at a lower spatial resolution than AOGCMs, but account for some dynamics of ocean-atmosphere circulations and can be leveraged to measure continental-scale climate change. Frequently used ESMIC include CLIMBER-2 (CLIMate-BiosPHERE model; [Petoukhov et al., 2000](#)). More details on each of the climate model categories is available in [IPCC \(2007\)](#).

It is important to note that, due to recognised biases, the outputs of these large-scale climate models should be appropriately downscaled to a higher resolution for the context of interest (unless it is of global scale), before they are directly integrated into risk assessments ([Fowler et al., 2007](#)). This can be achieved in two main ways, i.e., empirical (statistical) or dynamic downscaling ([Fowler et al., 2007](#)). Statistical downscaling relies on empirical relationships between global and regional climate models. It may involve the use of scaling factors (e.g., [Arnell, 2003](#)), regression models (e.g., [Salathé, 2005](#)), weather generators (i.e., stochastic models that combine historical data and the outputs of large-scale climate models; e.g., [Eum et al., 2011](#)), or weather typing (whereby particular weather classes are related with local weather conditions; e.g., [Bermúdez et al., 2020](#)). Dynamic downscaling refers to the development of higher resolution regional climate models that leverage boundary conditions from the larger scale models (e.g., [Ozturk et al., 2017](#)). Further correction of biases may also be necessary at fine resolutions, to account for residual discrepancies between simulations and observations (e.g., [Hempel et al., 2013](#); [Chen et al., 2021](#)). Numerous bias-correction techniques have been proposed in the literature, including the linear scaling of mean simulated values and non-linear adjustments that also address variance inaccuracies. A detailed review of these and other approaches can be found in [Teutschbein and Seibert \(2012\)](#).

Uncertainties are typically incorporated in climate change predictions via ensemble modelling ([Stewart et al., 2014](#)), to provide more informative risk assessments than those that are produced using discrete scenarios ([New et al., 2007](#)). Ensemble modelling may consist of using a series of data from a variety of climate models (e.g., [Mortin et al., 2014](#)), simulating data from sets of different internal parameters of a single climate model (e.g., [Murphy et al., 2007](#)) or generating data using a range of input parameters for the same model (e.g., [Murphy et al., 2004](#)). The first two methods capture modelling uncertainty, while the latter is used to quantify climate variability (which can be natural or otherwise, depending on the range of input parameters used). Examples of future risk modelling efforts that have leveraged climate ensemble modelling include [Rojas et al. \(2013\)](#), [Ward et al. \(2014\)](#), and [Muis et al. \(2015\)](#).

2.2. Emission scenarios

Information on future emissions can be obtained from scenario storylines, such as those provided in the Special Report on Emission Scenarios ([Nakicenovic et al., 2000](#), SRES; see “Future exposure” section for more details). For example, emission data from the SRES storylines scenarios were combined with a set of climate models to determine a series of future sea-level rise (SLR) scenarios in the IPCC Fourth Assessment Report ([IPCC, 2007](#)). These SLR scenarios have been used to quantify future flooding risk in a number of subsequent studies, including [Hallegatte et al. \(2011\)](#) and [Hinkel et al. \(2012\)](#). Further future risk assessments that have integrated SRES emissions data with various climate models include [Pardaens et al. \(2011\)](#), [Nicholls and Tol \(2006\)](#), [Rojas et al. \(2013\)](#), [Te Linde et al. \(2011\)](#), [Schuster et al. \(2012\)](#), and [Balbi et al. \(2013\)](#). SRES emission information was replaced with emission data from the Representative Concentration Pathways (RCPs) in the IPCC Fifth Assessment report ([Pachauri et al., 2015](#)). RCPs consist of four future pathways on emissions and land use ([Van Vuuren et al., 2011](#)) that are consistent with a large set of scenarios provided by the literature. Examples of future risk studies that have incorporated emission-related information from RCPs include [Hinkel et al. \(2014\)](#), who combined these data with four AOGCMs to determine coastal flooding damage in the 21st century on a global scale. Additional risk quantification work that has used RCPs include [Huang et al. \(2019\)](#), [Dottori et al. \(2018\)](#), [Winsemius et al. \(2015\)](#), and [Arnell and Lloyd-Hughes \(2014\)](#). Emission scenarios are usually treated in a discrete manner, such that climate change predictions are produced for each one individually. However, some recent work has leveraged variance-based sensitivity analysis to quantify their uncertainties ([Xu et al., 2019](#)).

2.3. Additional dynamic hazard features

The impact-model used to represent the physical process that directly results in the hazard may also have dynamic features, to align with the previously discussed climate projections. For example, the annual maximum discharges and windspeeds that are respectively used to compute future flood and storm hazards may be non-stationary (e.g., [Rojas et al., 2013](#); [Byun and Hamlet, 2020](#); [Li and Stewart, 2011](#); [Condon et al., 2015](#); [Xu et al., 2020](#)). In addition, dynamic modelling of ground-cover change (e.g., replacement of natural ground with impermeable surfaces, changes to drainage or irrigation systems, and deforestation) can influence future projections of flooding and landslide hazards (e.g., [Fraser et al., 2016](#); [Guo et al., 2020](#); [Avand et al., 2021](#); [Pisano et al., 2017](#)).

2.4. Modelling of future non-climate-related hazards

Time-dependent approaches are also available for projecting future hazards that are not directly related to climate change, such as earthquakes. In this case, long-term time-dependent hazard assessment procedures condition the expected duration until a fault’s next event on the amount of time that has elapsed since its last earthquake (or the occurrence of an event on a neighbouring fault), to account for changes in elastic strain energy. A comprehensive review of these methods is provided in [Iacopetti et al. \(2021\)](#). Short-term time-dependent seismic hazard assessments

account for the spatial and temporal clustering of aftershock events that follow the occurrence of a mainshock earthquake (e.g., Papadopoulos et al., 2020).

3. Future exposure

Exposure is defined as “the people, property, systems, or other elements present in hazard zones that are thereby subject to potential losses” (UNISDR, 2009). Future exposure to natural hazards will be driven by increases in population (such as urbanisation, i.e., the movement of people from rural to urban areas), socioeconomic growth (e.g., Angel et al., 2011), as well as choices on land use (e.g., Santini and Valentini, 2011) (Fig. 2).

3.1. Modelling population and socioeconomic growth

Future population and economic growth are commonly modelled on the basis of socioeconomic scenario storylines (e.g., Hammond, 1998; De Vries et al., 1994). These narratives detail alternative characterisations of how the world may evolve, depending on political, economic, technical and social developments at global and regional levels. The first set of socioeconomic scenario storylines adopted for widespread use in research were presented in SRES, which were developed based on an extensive review of the literature. SRES comprises four scenario families for describing future emission-driven impacts on the world, based on population, economy, technology, energy, land use and agriculture. For example, one of these storylines (A1) describes a future of rapid economic growth, global population that is maximised in the middle of the 21st century and decreases afterward, and the rapid introduction of new technologies. Six different emission modelling approaches are used to quantify each storyline in a number of scenarios, which are designed to capture the full range of (known) uncertainties in the underlying factors. Studies that have adopted the socioeconomic component of the SRES scenarios at the global scale include Nicholls (2004), who examined changes in flooding by storm surges and Nicholls and Tol (2006), who quantified the impacts of sea-level rise. Rojas et al. (2013), which examined the future risk of flooding in Europe, is an example of a continent-level study that has leveraged socioeconomic data from the SRES scenarios. Socioeconomic information from SRES scenarios has also been incorporated into smaller-scale analyses at country or regional level (e.g., Hall et al., 2005; Kebede et al., 2012; Penning-Rowsell et al., 2013).

O'Neill et al. (2014) developed the shared socioeconomic pathways (SSPs) framework as an update to the SRES storylines. This contains five alternative plausible means by which society and ecosystems will evolve in the 21st century (in both narrative form and using a quantified set of socioeconomic metrics that define scenarios), and is more focused on the challenges of climate change than SRES. Examples of studies that have leveraged SSPs are Arnell and Lloyd-Hughes (2014); Dottori et al. (2018); Huang et al. (2019), as well as the most recent IPCC report (IPCC, 2021). Other socioeconomic storylines include the UN Environment Programme's Global Environmental Outlook, the Millennium Ecosystem Assessment, and

the World Energy Outlook; a comprehensive review of these and other scenario families is provided by Van Vuuren et al. (2012).

Socioeconomic scenario storylines are presented in a highly aggregated manner and are therefore often downscaled to obtain finer resolution data. Various downscaling approaches have been applied in the literature. Population and economic growth forecasts can be proportionally downscaled to national or regional level, which assumes that the growth rates are consistent at both scales (e.g., Gaffin et al., 2004; van Vuuren et al., 2007). Other methods for population downscaling explicitly account for urbanisation trends, prioritizing growth in urban regions (e.g., Grübler et al., 2007; Nicholls, 2004). Economic growth downscaling can also be determined based on convergence approaches, which ensure consistency with local average values (e.g., van Vuuren et al., 2007), and more finer downscaling may differentiate between urban and rural gross domestic product (GDP) (e.g., Grübler et al., 2007) or involve stakeholder input (e.g., Carter et al., 2004). The aforementioned methods may be broadly categorised as top-down downscaling approaches; a comprehensive review of these types of procedures is provided in van Vuuren et al. (2010). Bottom-up approaches to downscaling have additionally been developed in the literature, which leverage participatory input on site-specific conditions (e.g., Nilsson et al., 2017) or are based on shared policy assumptions for the local context (e.g., Frame et al., 2018). While socioeconomic storylines are typically implemented in future risk assessments as discrete (deterministic) scenarios, some work has translated their GDP information into probabilistic distributions (Güneralp and Seto, 2013).

Population projections can also be quantified or obtained in other ways. Single “best-guess” forecasts of country-wide populations are available from agencies such as the World Bank and the Population Reference Bureau, for example (Lutz and KC, 2010). The UN produces high, medium, and low scenarios of future population at national and international scale (United Nations, 2019), which are characterised by corresponding variations in fertility, mortality, and migration. These scenarios can be combined with uncertainty estimates produced by the U.S. National Research Council (Council and Population, 2000), to construct future population probability distribution functions (Seto et al., 2012; Muis et al., 2015). Fully probabilistic population projections may also be obtained for 13 regions of the world from modelling work of the International Institute for Applied Systems Analysis (Lutz et al., 1997, 2001, 2008).

3.2. Land-use change and urbanisation

It is important to note that the term land-use change can cover a multitude of transitions, such as the abandonment or adaptation of agricultural land (Acosta et al., 2014; Acosta-Michlik and Espaldon, 2008). For the purposes of natural-hazard risk assessments, land-use change is typically interpreted as urban growth (e.g., Penning-Rowsell et al., 2013), which has been modelled in the literature in a variety of ways. Bottom-up procedures can include the use of cellular automata (CA) or agent-based modelling (ABM). An example of this kind of approach is the SLEUTH (Slope, Land cover, Excluded regions, Urban land cover, Transportation, and Hillshade) model (Chaudhuri and Clarke, 2013). SLEUTH uses previous




Fig. 2. Future exposure to natural hazards will be influenced by increases in urbanisation, socioeconomic growth, and fluctuations in land-use choices.

maps of historical growth (containing the first five features of its acronym) to probabilistically determine future land use according to relatively simple simulation rules that are founded on spatial auto-correlation and neighbourhood effects. It has recently been applied to simulate urban growth in the city of Istanbul (e.g., [Sarica et al., 2020](#)), the state of California ([Clarke and Johnson, 2020](#)), and at a global scale ([Zhou et al., 2019](#)). ABM and CA approaches are sometimes integrated with other tools to model urban growth. For example, ABM (or multi-agent systems) has been coupled with weighted geographical regression ([Calderón and Silva, 2021](#)), logistic regression ([Pandey and Joshi, 2015](#)), and Bayesian networks ([Kocabas and Dragicevic, 2013](#)), whereas CA has been combined with Markov chain methods ([Lallemant, 2015](#); [Rimal et al., 2019](#)). Urban growth may also be simulated using empirical (e.g., [Hu and Lo, 2007](#)) or semi-empirical (e.g., [Santini and Valentini, 2011](#)) methods, landscape metrics (e.g., [Taubenböck et al., 2009](#)), machine learning approaches (such as artificial neural networks, e.g., [Tayyebi et al., 2011](#)), coupled Markov chain-genetic algorithm models (e.g., [Tang et al., 2007](#)), weight of evidence ([Soares-Filho et al., 2004](#)), socioeconomic storylines (e.g., [Te Linde et al., 2011](#)), as well as integrated top-down, bottom-up approaches that capture interactions at various scales ([Verburg and Overmars, 2009](#); [Promper et al., 2014](#)).

Some work has leveraged population and economic growth projections to better constrain urban expansion models. For example, [Seto et al. \(2012\)](#) and [Muis et al. \(2015\)](#) informed their urban growth model with per capita land approximations, which were empirically computed from per capita GDP estimates (that were determined based on forecasted population and economic growth values). [Calderón and Silva \(2021\)](#) adjusted their pre-calibrated land-use model based on estimates of built-up area, which were empirically derived from future population projections. On a smaller scale, population increases have been used as a proxy for estimating annual increases in housing number ([Bjarnadottir et al., 2014](#); [Stewart and Li, 2010](#)).

4. Future (physical) vulnerability

Vulnerability is broadly defined as “the characteristics and circumstances of a community, system or asset that make it susceptible to the damaging effects of a hazard” ([UNISDR, 2009](#)). We particularly focus on physical vulnerability in this Review, which may be interpreted in this context as the tendency of exposed elements (assets) to suffer adverse effects when impacted by natural hazards ([Cardona et al., 2012](#)). Risk assessment methods account for the future vulnerability of physical infrastructure to natural hazards from two main perspectives. The first of these outlooks assumes that vulnerability will increase in time as a result of unplanned/informal modifications or maintenance/degradation challenges, whereas the second focuses on the reduction in vulnerability that can be achieved by adapting infrastructure to future conditions (see [Fig. 3](#)). These divergent standpoints reflect the two main drivers of vulnerability, i.e., socioeconomic conditions and policy decisions ([Fraser et al., 2016](#)).

4.1. Assessing increasing vulnerability

One of the foremost attempts to quantify dynamically increasing building-level physical vulnerability is the work of [Lallemant et al. \(2017\)](#). This study developed a framework for modelling the time-dependent earthquake fragility (i.e., the probability of collapse) of incrementally expanding construction, which is a commonly adopted strategy for accommodating ever-increasing urban growth in developing countries (e.g., [Amoako and Boamah, 2017](#)). Markov chains were leveraged to quantify the probability of residential buildings transitioning to a new state with more stories, and the framework was demonstrated by quantifying dynamic projections of building collapses in Kathmandu valley, Nepal. Other work has characterised the short-term increase in physical vulnerability that may occur after or during a series of natural-hazard events. For instance, attempts have been made to model the accumulation of damage in structures that takes place after a mainshock earthquake and during the subsequent




Fig. 3. Sample changes in the vulnerability of a structure through time, from T_1 to T_3 . Dynamic increases in structural vulnerability (as shown in the figure) may be driven by informal structural modifications, environment-induced deterioration, and accumulation of damage due to cascading hazard events (e.g., aftershocks). These increases can be offset by vulnerability adaptation measures (shifting the vulnerability curves to the right), many of which may be broadly categorised as “climate adaptation engineering” approaches.

aftershock sequence; very recent examples include [Aljawhari et al. \(2020\)](#), [Gentile and Galasso \(2020\)](#), and [Papadopoulos et al. \(2020\)](#).

A vast array of studies have focused on modelling the time-dependent environment-induced deteriorating vulnerability of infrastructure. We only discuss here the most timely advances in the field for brevity, but we note that several reviews on approaches for modelling/quantifying the environmental degradation of physical assets have been released in recent times (e.g., [Kashani et al., 2019](#); [Amaya-Gómez et al., 2019](#); [Guo et al., 2019](#)). Most recent state-of-the-art developments in the field include (1) the study of [Zamanian et al. \(2020\)](#), which simulated the impacts of corrosion on buried concrete sewer pipes using a novel high-fidelity finite element model that captured the nonlinear behaviours of concrete, soil, and combined pipe-soil interactions; (2) the work of [Soltani et al. \(2021\)](#), which developed a new statistical (multiple linear regression) model for quantifying the reduced shear capacity of corroded reinforced concrete beams; and (3) research by [Zanini et al. \(2020\)](#), which leveraged systems reliability theory to compute original time-dependent fragility functions and corresponding dynamic seismic reliability profiles for environmentally degraded reinforced concrete bridges.

Research on the effects of environmentally-driven deterioration to the built environment is beginning to account for future climatic conditions ([Bastidas-Arteaga and Stewart, 2015, 2016](#); [Wang et al., 2012](#); [Stewart et al., 2012](#); [El Hassan et al., 2010](#); [Yang and Frangopol, 2020](#); [Sevieri and Galasso, 2021](#)). This is important, given that rising carbon dioxide levels associated with global warming will increase the likelihood of carbonation-induced corrosion ([Stewart et al., 2011](#)). One of the first attempts to incorporate future weather in the assessment of corrosion was the work of [Bastidas-Arteaga et al. \(2010\)](#), which specifically focused on reinforced concrete structures. This study leveraged a model that coupled the effects of convection, chloride binding, and concrete ageing with temperature and humidity data that were informed by three scenarios of global warming. [Stewart et al. \(2011\)](#) provided a probabilistic and reliability-based approach for predicting the probability of corrosion initiation and damage to concrete infrastructure, deriving temperature and humidity information from nine AOGCMs combined with three emission scenarios. [Peng and Stewart \(2016\)](#) used a series of climate projections to probabilistically determine future carbonation-induced damage to reinforced concrete structures in different Chinese cities. Lately, the work of [Yang and Frangopol \(2019a\)](#) leveraged various general circulation models to determine future scour-induced failure probabilities of bridges (and the associated long-term risks).

4.2. Accounting for adapted vulnerability

Work that has focused on a potential reduction in the future vulnerability of the built environment is often framed within the context of dynamically increasing climate risks and termed “climate adaptation engineering” (Stewart et al., 2014; Stewart and Deng, 2015; Mondoro et al., 2018). A significant motivation for this body of research is that the changing climate is generally likely to inflict more damage on engineering assets in the future, even if design standards remain unchanged (e.g., Stewart, 2015). Many of these studies have particularly focused on Australia. For example, Stewart (2015) assessed the risk, costs, and benefits of climate adaptation strategies for reducing the vulnerability of new housing to evolving wind-induced risk for three cities in Australia, under three emission scenarios. Li and Stewart et al. (2011) studied the economic viability of various mitigation strategies (such as building retrofitting and an improvement in new building design) against future cyclone hazards in Northern Australia. Qin and Stewart (2020) probabilistically examined two adaptation options for reducing the vulnerability of Australian housing to non-cyclonic windstorms in a changing climate, i.e., reinforcing the building envelope or increasing the water resistance of the building interior. In addition, Ryan et al. (2016); Ryan and Stewart (2017, 2020) considered the feasibility of reducing the vulnerability of Australian power distribution networks to future climate impacts. Further important contributions to the field of climate adaptation engineering are the work of Dong and Frangopol (2017), which proposed a genetic algorithm-based optimisation approach for the process of adapting residential buildings to the future impacts of hurricanes; the study of Liu et al. (2020), which presented optimal adaptation schedules for bridge networks vulnerable to abutment scour that considered the cost of both adaptation and network-level failure; and the investigation by Du et al. (2020) of various building vulnerability adaptation strategies (including dry-proofing, wet-proofing, and the construction of raised floors) on the effect of future flood risk in Shanghai.

Other studies have examined the potential implementation of physical or natural protective measures to reduce the vulnerability of regions to future flooding events. For example, the study of Dawson et al. (2005) quantified the reduction in inundation area associated with raising the Thames Estuary flood defences and installing additional tidal flood barriers in London, under different sea-level rise scenarios. The construction of dikes has been assessed as an alternative flood-related vulnerability reduction measure in a number of studies (e.g., Hinkel et al., 2013; Tol, 2007), including the work of Ward et al. (2017), which demonstrated that dyke investment could reduce flood risk below current levels in some areas of the world (despite climate change and dynamic economic conditions). The level of future flood-vulnerability mitigation offered by various nature-based solutions (i.e., actions inspired and supported by nature; e.g., Faivre et al., 2017) is an active area of research. For example, Zölch et al. (2017) measured the reduction in future urban surface runoff due to trees and green roofs in Munich, while Reguero et al. (2014) investigated the effectiveness of wetland restoration/conservation, beach nourishment, and oyster reef restoration in lowering flood vulnerability along the U.S. Gulf Coast. Additional research (Prudhomme et al., 2010) has taken a more conceptual, reliability-based approach to the concept of future flood vulnerability by developing an analytical framework for quantifying the fraction of climate model scenarios that would be facilitated by a prescribed adaptation measure.

4.3. Non-physical vulnerability

While quantitative natural-hazard risk assessments centre predominantly on the future vulnerability of the built environment, we briefly mention here efforts that have focused elsewhere. In fact, socioeconomic vulnerability also changes over time, impacted by the occurrence of disasters (disrupting lives and livelihoods) and by the effects of climate change. Bjarnadottir et al. (2011) quantified social vulnerability dynamics due to climate change, using a time-dependent index that accounted for the most relevant factors (e.g., race, age, gender and socioeconomic status) (Cutter

et al., 2008). A weighted socioeconomic vulnerability index for climate-change-affected regions in Bangladesh was developed in Ahsan and Warner (2014), to guide future adaptation policies, while adapted vulnerability to floods was defined in terms of expected annual fatalities by Jongman et al. (2015). Finally, some work on future drought events (Naumann et al., 2021) has accounted for adapted socioeconomic vulnerability on the basis of GDP growth, using pre-determined empirical relations (Formetta and Feyen, 2019).

5. Risk metrics

Risk metrics act as a crucial bridge between the quantitative risk assessment and the decision-making process that they are used to inform (Johansen and Rausand, 2014). High-level preliminary assessments have used relative risk indices to identify potential future natural-hazard risk hotspots (e.g., Hawchar et al., 2020). Most future-focused approaches to quantifying natural-hazard risk have concentrated on measuring risk in terms of conventional asset losses. For example, Stewart et al. (2018) examined the consequences of climate-induced extreme wind loading on residential roofs in terms of its percentage of house value. Both Forzieri et al. (2018) and Dawson et al. (2009) quantified the effects of future natural-hazard events as the expected annual cost of damage to physical infrastructure and/or agricultural land, while Schuster et al. (2012) expressed the impacts of flooding to a facility as the expected cumulative damage cost over its lifetime. Salmanidou et al. (2021) measured risks from future tsunamis in terms of their effect on the value of household-level assets. Other authors (e.g., Gaslikova et al., 2011; Schwierz et al., 2010) have focused exclusively on insured property losses.

Tomorrow's risk from natural hazards may also be presented in terms of its direct effect on the human population. For example, Maaskant et al. (2009), Nagai et al. (2020), and Stone et al. (2014) examined the potential mortality associated with different future natural-hazard events, while Nicholls (2004) measured the number of people likely to be affected by sea-level rise. The study of Darwin and Tol (2001) explicitly accounted for welfare changes associated with sea-level rise, using the equivalent variation metric to express losses.

Wider perspectives on risk have been offered by some authors, who have quantified the impacts of future natural-hazard events in terms of multiple metrics and/or based on a broader network-level perspective. Stewart (2016) examined structural, non-structural and business interruption losses associated with roof cover damage to industrial and commercial buildings in future wind events. Stewart et al. (2014) considered both the direct and indirect losses (in the form of, for example, residential clean-up, alternative household accommodation, disaster response and relief, injuries and fatalities, and business and economic disruption) associated with projected wind-related damage to residential buildings. Yang and Frangopol (2019b) quantified the societal risk of the impacts of climate change on transportation networks, which accounted for the costs incurred by users due to necessary detours and the loss of time. Both Liu et al. (2020) and Yang and Frangopol (2019a) considered the societal risk metric in addition to the direct economic cost associated with transportation-network bridge failure. Verschuur et al. (2020) measured the risk associated with future Bangladesh flooding events using a Composite Development Index, which incorporated information on welfare losses, the relative extent to which policy interventions reduce these losses for poor households, and resilience (i.e., the ratio of asset to welfare losses).

The Dynamic Interactive Vulnerability Assessment (DIVA) tool has been leveraged to determine future risk in a number of studies (e.g., Hinkel et al., 2013, 2012; Mcleod et al., 2010). In addition to quantifying direct economic losses, this model describes the consequences of sea-level rise in terms of three socially-focused indicators (Hinkel et al., 2010), 1) the number of people who live in the coastal floodplain (i.e., below the 1000-year storm-surge level); 2) the expected annual number of people subjected to flooding; and 3) the number of people forced to migrate as a result of erosion. The work of Bosello et al. (2012) used DIVA in combination with GTAP (Global Trade Analysis Project) – EF, a computable general

equilibrium model for Europe, to also quantify wider indirect economic effects of sea-level rise (e.g., impacts on GDP, investment, and international trade flows).

6. Modelling future risk

6.1. Challenges

Decision makers of today have a notable opportunity to positively influence the risks of tomorrow, through their choices on implementing strategies/policies that control future risk drivers. However, the selection of appropriate risk-management interventions strongly relies on our ability to model tomorrow's risks from natural hazards, explicitly accounting for the dynamic nature of hazard, exposure, and vulnerability. One of the main challenges to modelling future risk that has been identified in the literature is the significant inherent uncertainty in each of these components of tomorrow's risk (e.g., Hemmati et al., 2020). Some of the unknowns related to future risk, such as natural climate variability, cannot be (completely) reduced with new knowledge. Epistemic (i.e., modelling) uncertainties are captured by leveraging a combination (ensemble) of different models in a given risk assessment, which poses its own difficulties. For example, while general circulation models for climate have different levels of skill in modelling climate variables, there is no generally accepted protocol for ranking/aggregating models (e.g., Stewart et al., 2011). This means that current best practice is to average climate projections across all models in an ensemble, which may not be optimal (Gleckler et al., 2008). In addition, these ensembles may not capture all potential versions of the future. For instance, it has recently been argued that the IPCC scenarios for global sea-level rise under strong warming are concentrated on merely the lower end of possible outcomes (Siegert et al., 2020). Furthermore, the necessary downscaling of climate projections amplifies their uncertainties because subgrid parameterisations (i.e., submodels) are only approximate, idealistic representations of complex physical phenomena (Hourdin et al., 2017). Even if climate projections were well constrained, quantifying their impact on future hazard would be hampered by a lack of data on extreme events (Hallegatte, 2007). We have focused our discussion here on hazard uncertainty in particular, but note that this is even

further exacerbated by comparable unknowns in terms of future population (Cohen, 2004), the point in time at which an updated vulnerability measure may be introduced (Kwadijk et al., 2010), and the eventual success of any adopted adaptation measures (Dilling et al., 2015), for example. The combined effects of these uncertainties can inhibit the utility of future risk assessments in decision-making processes (e.g., Wilby and Dessai, 2010), which may help to explain why disaster risk managers tend to frame their risk perceptions in terms of today's - rather than tomorrow's - hazards (Dilling et al., 2015).

Despite clear and numerous efforts to model individual risk components associated with future natural-hazard events, there remains a lack of a commonly accepted analytical framework for complete end-to-end risk quantification. This is reflected in the wide range of inconsistent approaches found in the literature for calculating future loss projections, which incorporate various degrees of complexity (e.g., Jurgilevich et al., 2017). To illustrate this point, we take risk modelling for future tropical cyclones (hurricanes) as an example. Some work in this area (e.g., Webersik et al., 2010) quantified cyclone losses under the assumption that population distribution, economic growth, and vulnerability remain unchanged from 2010 conditions. Other studies (e.g., Bjarnadottir et al., 2014) accounted for the dynamic variability of both hazard and exposure, but leveraged a static representation of vulnerability. A more comprehensive approach was taken by Mendelsohn et al. (2012), who incorporated projected changes to all components of risk (hazard, exposure, and vulnerability) into their predictions of future tropical cyclone damage. Robust methodological frameworks and protocols are required to overcome these types of modelling inconsistencies (Bouwer, 2013).

Some work related to this need has already been accomplished. For example, Bouwer (2013) presented a framework for modelling future natural-hazard risk (see Fig. 4), which accounted for the dynamic nature of hazard (via climate change and natural weather variability), vulnerability (via socioeconomic change and adaptation measures), and exposure (via temporal adjustments to population and economic growth). Muis et al. (2015) developed a similarly structured framework with a specific focus on flood hazard and the corresponding expected annual damage. Verschuur et al. (2020) provided a more detailed framework for specifically analysing future welfare risks that accompany asset losses caused by Bangladesh flooding




Fig. 4. End-to-end framework for quantifying future natural-hazard risk, as proposed by Bouwer (2013). Note that the term “sensitivity” in this context refers to the empirical relationship between the hazard and the loss. Adapted from Fig. 1 of the aforementioned paper.

events, through a pro-poor lens. Finally, Lee et al. (2018) proposed an adaptive decision-making methodology to determine optimal strategies for mitigating seismic risks in the face of evolving hazard, exposure, and vulnerability.

6.2. Perspectives

While the aforementioned frameworks help to facilitate comprehensive assessments of future risk, they have some limitations that may hinder decision-making related to tomorrow's natural hazards. Firstly, each centres on just one hazard. This narrow-focused approach may lead to increased risks from other overlooked hazards (Gill and Malamud, 2016) as part of a failure to account for "as synergies" in disaster risk management (de Ruiter et al., 2021), is not effective for spatial planning purposes (Greiving et al., 2006), and is not aligned with the "all-hazards" perspective adopted in conventional emergency management (Ferrier and Haque, 2003). A multi-hazard (or multi-risk) outlook would significantly improve the risk-reduction potential of these frameworks (Gill and Malamud, 2014; Kappes et al., 2012). This could be achieved by making use of the multi-risk mathematical environment developed by Selva (2013), for example, which accounts for interactions between hazardous events across all three components of risk. However, it is important to mention here that modelling these interactions can currently be inhibited by shortages in computational and storage capacities (Bevacqua et al., 2019), data constraints that can further exacerbate the uncertainties associated with single-hazard assessments (Ciscar et al., 2019), and a lack of innovative risk assessment methods that capture the complexity of joint hazard occurrences within a single unifying framework (Mignan et al., 2014).

Secondly, these frameworks incorporate a limited set of risk metrics (e.g., direct economic costs and household resilience) that do not explicitly account for disruptions to socio-economic and technical networks, such as supply-chain interruption (e.g., Jabbarzadeh et al., 2016) and accessibility reduction (e.g., Miller and Baker, 2016). Interconnected risks often form the main source of crisis in the aftermath of a natural-hazard event (Pescaroli and Alexander, 2018), thus consideration of their effect should play a central role in future risk models. Quantification of these types of risks could benefit from more detailed datasets of future socio-economic projections than those typically leveraged in current models. For instance, many studies measure the effects of future natural hazards using aggregated population predictions (Lutz and KC, 2010), whereas more refined age-based projections could be useful for deriving time-dependent estimates of user numbers in a transportation network or predicting the perceived habitability of a neighbourhood after an event.

A further disadvantage of the aforementioned approaches is that they do not explicitly facilitate formal decision-making processes on future risk-management and mitigation measures. Cost-benefit analyses (CBA), which have already been successfully leveraged to appraise different climate adaptation policies (Ryan and Stewart, 2017) could be integrated into risk assessment procedures to overcome this issue. However, CBA methodologies fail to account for the different risk perspectives of various stakeholders, which are crucial for effective risk governance (Komendantova et al., 2014) and the successful adoption of any risk-management strategies (Carpignano et al., 2009). While the Composite Development Index metric output of the framework developed in Verschuur et al. (2020) allows for differently weighted policy objectives to reflect diverse priorities and preferences, the considered objectives (reducing disaster losses, increasing resilience, and promoting pro-poor solutions) are specifically tailored to the Bangladesh case study carried out and the framework is only demonstrated for the scenario of equal weights. One way of better promoting a participatory approach to future natural-hazard risk quantification would be to explicitly include an additional decision-making (or impact assessment) module in the future risk-assessment framework (see Fig. 5), representing a process in which the stakeholder would rank the importance of various risk metrics of interest and a final evaluation

of the weighted risk (or impact) would be carried out. This could be achieved using a similar methodology to that developed for selecting optimal earthquake early warning actions in Cremen and Galasso (2021), where risk outputs of a traditional framework would act as an input to a multi-criteria decision-making process that accounted for stakeholder preferences. Note that the impact assessment module could also feed back to the risk calculations (including the exposure inputs - which may also affect the multi-hazard simulations - and the vulnerability component), to allow stakeholder perspectives, management plans, and priorities to be accounted for in the final quantification and selection of risk metrics.

7. Conclusions

Modelling and quantifying tomorrow's risks from natural hazards is becoming ever more critical as we face an uncertain future of increasing human populations, relentless urbanisation, and significantly different climatic conditions. This paper has reviewed state-of-the-art approaches to future natural-hazard risk quantification, including various methodologies that have been developed to model tomorrow's hazards, exposure, and (physical) vulnerability. We have also documented diverse ways in which future risk has been measured, such as in terms of asset damage, business interruption, welfare losses, and travel-time delays.

Our Review has revealed that while there have been numerous efforts to model the individual components of future natural-hazard risk, some significant obstacles to quantifying this risk for management and mitigation purposes still remain. These include the substantial uncertainty associated with future risk projections, some of which (including natural climate variability) cannot be (completely) reduced with new information. There is also a clear lack of a standard framework for end-to-end future risk modelling that (1) captures multiple hazards; (2) accounts for network-level disruptions in our increasingly interconnected societies; (3) incorporates a broad set of risk metrics that, for instance, account for socioeconomic disparities; and (4) facilitates the diverse perspectives of a wide set of stakeholders involved in risk governance.

In summary, it is clear that there is some work to do before future risks from natural hazards can be quantified in a manner that is meaningful for the important decisions that need to be made as our world keeps changing. Our efforts will inevitably be enhanced as more data becomes available to constrain our models, but we first need to develop consistent risk-quantification procedures. It is also important to remember that we can readily improve our current assessments of tomorrow's natural-hazard risks, by leveraging existing tools (e.g., related to multi-criteria decision-making and multi-hazard theories) that have been successfully adopted for risk quantification in other contexts. The Tomorrow's Cities Hub of the United Kingdom Research and Innovation (UKRI) Global Challenge Research Fund (GCRF), which is aiming to develop a comprehensive methodology for facilitating the risk-sensitive design of future cities that is co-produced with local stakeholders, accounts for multiple hazards, and incorporates both social and physical vulnerabilities (Galasso et al., 2021), represents a promising step in the right direction towards addressing some of the current challenges.

Data availability

No data are created or directly analysed as part of this review.

Declaration of competing interest

The authors declare no competing interests.

Acknowledgments

The authors acknowledge funding from UKRI GCRF under grant NE/S009000/1, Tomorrow's Cities Hub. No new data were created as part of this study.




Fig. 5. Outline of a proposed idealised stakeholder-centred end-to-end risk modelling framework for future natural hazards. The conventional risk assessment approach (that consists of combining a single hazard, exposure, and vulnerability to produce consequence values typically related to economic and/or building-level loss) is advanced to account for: (1) multiple hazards; (2) a flexible suite of risk metrics that can account for both socio-economic and technical networks; and (3) stakeholder perspectives and priorities towards different dimensions of risk. Feedback loops to the risk calculations (including exposure inputs and the vulnerability component) enable stakeholder perspectives, plans, and priorities to be accounted for in the final quantification and selection of risk metrics.

References

- Acosta, L., Rounsevell, M.D., Bakker, M., Van Doorn, A., Gómez-Delgado, M., Delgado, M., 2014. An agent-based assessment of land use and ecosystem changes in traditional agricultural landscape of Portugal. *Intell. Inf. Manag.* 6, 55–80.
- Acosta-Michlik, L., Espaldon, V., 2008. Assessing vulnerability of selected farming communities in the Philippines based on a behavioural model of agent's adaptation to global environmental change. *Glob. Environ. Chang.* 18 (4), 554–563.
- Ahsan, M.N., Warner, J., 2014. The socioeconomic vulnerability index: a pragmatic approach for assessing climate change led risks—a case study in the south-western coastal Bangladesh. *Int. J. Disaster Risk Reduct.* 8, 32–49.
- Aljawhari, K., Gentile, R., Freddi, F., Galasso, C., 2020. Effects of ground-motion sequences on fragility and vulnerability of case-study reinforced concrete frames. *Bull. Earthq. Eng.* 1–31.
- Amaya-Gómez, R., Sanchez-Silva, M., Bastidas-Arteaga, E., Schoefs, F., Munoz, F., 2019. Reliability assessments of corroded pipelines based on internal pressure—a review. *Eng. Fail. Anal.* 98, 190–214.
- Amoako, C., Boamah, E.F., 2017. Build as you earn and learn: informal urbanism and incremental housing financing in Kumasi, Ghana. *J. Housing Built Environ.* 32 (3), 429–448.
- Angel, S., Parent, J., Civco, D.L., Blei, A., Potere, D., 2011. The dimensions of global urban expansion: estimates and projections for all countries, 2000–2050. *Prog. Plan.* 75 (2), 53–107.
- Angel, S., Blei, A.M., Civco, D.L., Parent, J., 2012. *Atlas of Urban Expansion*. Lincoln Institute of Land Policy Cambridge, MA.
- Arnell, N.W., 2003. Relative effects of multi-decadal climatic variability and changes in the mean and variability of climate due to global warming: future streamflows in Britain. *J. Hydrol.* 270 (3–4), 195–213.
- Arnell, N.W., Lloyd-Hughes, B., 2014. The global-scale impacts of climate change on water resources and flooding under new climate and socio-economic scenarios. *Clim. Chang.* 122 (1), 127–140.
- Avand, M., Moradi, H., et al., 2021. Using machine learning models, remote sensing, and GIS to investigate the effects of changing climates and land uses on flood probability. *J. Hydrol.* 595, 125663.
- Balbi, S., Giupponi, C., Perez, P., Alberti, M., 2013. A spatial agent-based model for assessing strategies of adaptation to climate and tourism demand changes in an alpine tourism destination. *Environ. Model Softw.* 45, 29–51.
- Bastidas-Arteaga, E., Stewart, M.G., 2015. Damage risks and economic assessment of climate adaptation strategies for design of new concrete structures subject to chloride-induced corrosion. *Struct. Saf.* 52, 40–53.
- Bastidas-Arteaga, E., Stewart, M.G., 2016. Economic assessment of climate adaptation strategies for existing reinforced concrete structures subjected to chloride-induced corrosion. *Struct. Infrastruct. Eng.* 12 (4), 432–449.
- Bastidas-Arteaga, E., Chateaufort, A., Sánchez-Silva, M., Bressolette, P., Schoefs, F., 2010. Influence of weather and global warming in chloride ingress into concrete: a stochastic approach. *Struct. Saf.* 32 (4), 238–249.
- Bellouin, N., Collins, W., Culverwell, I., Halloran, P., Hardiman, S., Hinton, T., Jones, C., McDonald, R., McLaren, A., O'Connor, F., et al., 2011. The HADGEM2 family of Met Office Unified Model climate configurations. *Geosci. Model Dev.* 4 (3), 723–757.
- Bermúdez, M., Cea, L., Van Uytven, E., Willems, P., Farfán, J., Puertas, J., 2020. A robust method to update local river inundation maps using global climate model output and weather typing based statistical downscaling. *Water Resour. Manag.* 34 (14), 4345–4362.
- Bevacqua, E., Maraun, D., Voudoukas, M.I., Voukouvalas, E., Vrac, M., Mentaschi, L., Widmann, M., 2019. Higher probability of compound flooding from precipitation and storm surge in Europe under anthropogenic climate change. *Sci. Adv.* 5 (9), eaaw5531.
- Birkmann, J., Welle, T., 2015. Assessing the risk of loss and damage: exposure, vulnerability and risk to climate-related hazards for different country classifications. *Int. J. Glob. Warm.* 8 (2), 191–212.
- Bjarnadottir, S., Li, Y., Stewart, M.G., 2011. Social vulnerability index for coastal communities at risk to hurricane hazard and a changing climate. *Nat. Hazards* 59 (2), 1055–1075.

- Bjarnadottir, S., Li, Y., Stewart, M.G., 2014. Regional loss estimation due to hurricane wind and hurricane-induced surge considering climate variability. *Struct. Infrastruct. Eng.* 10 (11), 1369–1384.
- Bosello, F., Nicholls, R.J., Richards, J., Roson, R., Tol, R.S., 2012. Economic impacts of climate change in Europe: sea-level rise. *Clim. Chang.* 112 (1), 63–81.
- Bouwer, L.M., 2013. Projections of future extreme weather losses under changes in climate and exposure. *Risk Anal.* 33 (5), 915–930.
- Brian, K., 2015. OECD Insights Income Inequality the Gap between Rich and Poor: The Gap between Rich and Poor. OECD Publishing.
- Burby, R.J., Deyle, R.E., Godschalk, D.R., Olshansky, R.B., 2000. Creating hazard resilient communities through land-use planning. *Nat. Hazards Rev.* 1 (2), 99–106.
- Byun, K., Hamlet, A.F., 2020. A risk-based analytical framework for quantifying non-stationary flood risks and establishing infrastructure design standards in a changing environment. *J. Hydrol.* 584, 124575.
- Calderón, A., Silva, V., 2021. Exposure forecasting for seismic risk estimation: application to Costa Rica. *Earthq. Spectra* 37 (3), 1806–1826.
- Cardona, O.D., Van Aalst, M.K., Birkmann, J., Fordham, M., Mc Gregor, G., Rosa, P., Pulwarty, R.S., Schipper, E.L.F., Sinh, B.T., Décamps, H., et al., 2012. Determinants of risk: exposure and vulnerability. Managing the Risks of Extreme Events and Disasters to Advance Climate Change Adaptation: Special Report of the Intergovernmental Panel on Climate Change. Cambridge University Press, pp. 65–108.
- Carpignano, A., Golia, E., Mauro, C.D., Bouchon, S., Nordvik, J., 2009. A methodological approach for the definition of multi-risk maps at regional level: first application. *J. Risk Res.* 12 (3–4), 513–534.
- Carter, T.R., Fronzek, S., Bärlund, I., 2004. Finsken: a framework for developing consistent global change scenarios for Finland in the 21st century. *Boreal Environ. Res.* 9 (2), 91–107.
- Chaudhuri, G., Clarke, K., 2013. The SLEUTH land use change model: a review. *Environ. Resour. Res.* 1 (1), 88–105.
- Chen, J., Arsenault, R., Brissette, F.P., Zhang, S., 2021. Climate change impact studies: should we bias correct climate model outputs or post-process impact model outputs? *Water Resour. Res.* 57 (5), e2020WR028638.
- Ciscar, J.-C., Rising, J., Kopp, R.E., Feyen, L., 2019. Assessing future climate change impacts in the EU and the USA: insights and lessons from two continental-scale projects. *Environ. Res. Lett.* 14 (8), 084010.
- Clarke, K.C., Johnson, J.M., 2020. Calibrating SLEUTH with big data: projecting California's land use to 2100. *Comput. Environ. Urban. Syst.* 83, 101525.
- Cohen, B., 2004. Urban growth in developing countries: a review of current trends and a caution regarding existing forecasts. *World Dev.* 32 (1), 23–51.
- Condon, L., Gangopadhyay, S., Pruitt, T., 2015. Climate change and non-stationary flood risk for the upper Truckee river basin. *Hydrol. Earth Syst. Sci.* 19 (1), 159–175.
- Council, N.R., Population, C., 2000. Beyond Six Billion: Forecasting the World's Population. National Academies Press.
- Cremen, G., Galasso, C., 2021. A decision-making methodology for risk-informed earthquake early warning. *Comput. Aided Civil Infrastruct. Eng.* 36 (6), 747–761.
- Cutter, S.L., Barnes, L., Berry, M., Burton, C., Evans, E., Tate, E., Webb, J., 2008. A place-based model for understanding community resilience to natural disasters. *Glob. Environ. Chang.* 18 (4), 598–606.
- Darwin, R.F., Tol, R.S., 2001. Estimates of the economic effects of sea level rise. *Environ. Resour. Econ.* 19 (2), 113–129.
- Dawson, R.J., Hall, J., Bates, P., Nicholls, R., 2005. Quantified analysis of the probability of flooding in the Thames estuary under imaginable worst-case sea level rise scenarios. *Water Resour. Dev.* 21 (4), 577–591.
- Dawson, R.J., Dickson, M.E., Nicholls, R.J., Hall, J.W., Walkden, M.J., Stansby, P.K., Mokrech, M., Richards, J., Zhou, J., Milligan, J., et al., 2009. Integrated analysis of risks of coastal flooding and cliff erosion under scenarios of long term change. *Clim. Chang.* 95 (1), 249–288.
- De Vries, H., Olivier, J., Van den Wijngaert, R., Kreileman, G., Toet, A., 1994. Model for calculating regional energy use, industrial production and greenhouse gas emissions for evaluating global climate scenarios. *Water Air Soil Pollut.* 76 (1), 79–131.
- Dessler, A.E., 2021. Introduction to Modern Climate Change. Cambridge University Press.
- Dilling, L., Daly, M.E., Travis, W.R., Wilhelm, O.V., Klein, R.A., 2015. The dynamics of vulnerability: why adapting to climate variability will not always prepare us for climate change. *Wiley Interdiscip. Rev. Clim. Chang.* 6 (4), 413–425.
- Dong, Y., Frangopol, D.M., 2017. Adaptation optimization of residential buildings under hurricane threat considering climate change in a lifecycle context. *J. Perform. Constr. Facil.* 31 (6), 04017099.
- Dottori, F., Szewczyk, W., Ciscar, J.-C., Zhao, F., Alfieri, L., Hirabayashi, Y., Bianchi, A., Mongelli, I., Frieler, K., Betts, R.A., et al., 2018. Increased human and economic losses from river flooding with anthropogenic warming. *Nat. Clim. Chang.* 8 (9), 781–786.
- Dowdy, A.J., Ye, H., Pepler, A., Thatcher, M., Osbrough, S.L., Evans, J.P., Di Virgilio, G., McCarthy, N., 2019. Future changes in extreme weather and pyroconvection risk factors for Australian wildfires. *Sci. Rep.* 9 (1), 1–11.
- Du, S., Scussolini, P., Ward, P.J., Zhang, M., Wen, J., Wang, L., Koks, E., Diaz-Loaiza, A., Gao, J., Ke, Q., et al., 2020. Hard or soft flood adaptation? Advantages of a hybrid strategy for Shanghai. *Glob. Environ. Chang.* 61, 102037.
- El Hassan, J., Bressolette, P., Chateauneuf, A., El Tawil, K., 2010. Reliability-based assessment of the effect of climatic conditions on the corrosion of RC structures subject to chloride ingress. *Eng. Struct.* 32 (10), 3279–3287.
- Erdik, M., 2017. Earthquake risk assessment. *Bull. Earthq. Eng.* 15 (12), 5055–5092.
- Eum, H.-I., Sredojevic, D., Simonovic, S.P., 2011. Engineering procedure for the climate change flood risk assessment in the upper Thames river basin. *J. Hydrol. Eng.* 16 (7), 608–612.
- European Commission, 2020. Overview of Natural and Man-made Disaster Risks the European Union May Face. European Commission.
- Faivre, N., Fritz, M., Freitas, T., de Boissezon, B., Vandewoestijne, S., 2017. Nature-based solutions in the EU: innovating with nature to address social, economic and environmental challenges. *Environ. Res.* 159, 509–518.
- Ferrier, N., Haque, C.E., 2003. Hazards risk assessment methodology for emergency managers: a standardized framework for application. *Nat. Hazards* 28 (2), 271–290.
- Formetta, G., Feyen, L., 2019. Empirical evidence of declining global vulnerability to climate-related hazards. *Glob. Environ. Chang.* 57, 101920.
- Forzieri, G., Bianchi, A., Silva, F.B., Herrera, M.A.M., Leblois, A., Lavalle, C., Aerts, J.C., Feyen, L., 2018. Escalating impacts of climate extremes on critical infrastructures in Europe. *Glob. Environ. Chang.* 48, 97–107.
- Fowler, H.J., Blenkinsop, S., Tebaldi, C., 2007. Linking climate change modelling to impacts studies: recent advances in downscaling techniques for hydrological modelling. *Int. J. Climatol.* 27 (12), 1547–1578.
- Frame, B., Lawrence, J., Ausseil, A.-G., Reisinger, A., Daigneault, A., 2018. Adapting global shared socio-economic pathways for national and local scenarios. *Clim. Risk Manag.* 21, 39–51.
- Fraser, S., Jongman, B., Balog, S., Simpson, A., Saito, K., Himmelfarb, A., 2016. The making of a riskier future: how our decisions are shaping future disaster risk. Global Facility for Disaster Reduction and Recovery.
- Gaffin, S.R., Rosenzweig, C., Xing, X., Yetman, G., 2004. Downscaling and geo-spatial gridding of socio-economic projections from the IPCC special report on emissions scenarios (SRES). *Glob. Environ. Chang.* 14 (2), 105–123.
- Galasso, C., McCloskey, J., Pelling, M., Hope, M., Bean, C., Cremen, G., Guragain, R., Hancilar, U., Menocal, J., Mwanga, K., 2021. Risk-based, pro-poor urban design and planning for tomorrow's cities. *Int. J. Disaster Risk Reduct.* 102158.
- Gallina, V., Torresan, S., Critto, A., Sperotto, A., Glade, T., Marcomini, A., 2016. A review of multi-risk methodologies for natural hazards: consequences and challenges for a climate change impact assessment. *J. Environ. Manag.* 168, 123–132.
- Gaslikova, L., Schwarzmann, A., Raible, C., Stocker, T., 2011. Future storm surge impacts on vulnerable coasts for the North Sea region. *Nat. Hazards Earth Syst. Sci.* 11 (4), 1205–1216.
- Gentile, R., Galasso, C., 2020. Hysteretic energy-based state-dependent fragility for ground-motion sequences. *Earthq. Eng. Struct. Dyn.* 50 (4), 1187–1203.
- GFDRR, 2014. Understanding Risk: Review of Open Source and Access Software Packages Available to Quantify Risk from Natural Hazards. World Bank Publications.
- Gill, J.C., Malamud, B.D., 2014. Reviewing and visualizing the interactions of natural hazards. *Rev. Geophys.* 52 (4), 680–722.
- Gill, J.C., Malamud, B.D., 2016. Hazard interactions and interaction networks (cascades) within multi-hazard methodologies. *Earth Syst. Dyn.* 7 (3), 659–679.
- Gleckler, P.J., Taylor, K.E., Doutriaux, C., 2008. Performance metrics for climate models. *J. Geophys. Res. Atmos.* 113 (D6).
- Gordon, C., Cooper, C., Senior, C.A., Banks, H., Gregory, J.M., Johns, T.C., Mitchell, J.F., Wood, R.A., 2000. The simulation of SST, sea ice extents and ocean heat transports in a version of the Hadley Centre coupled model without flux adjustments. *Clim. Dyn.* 16 (2), 147–168.
- Greiving, S., Fleischhauer, M., Lückenköter, J., 2006. A methodology for an integrated risk assessment of spatially relevant hazards. *J. Environ. Plan. Manag.* 49 (1), 1–19.
- Grübler, A., O'Neill, B., Riahi, K., Chirkov, V., Goujon, A., Kolp, P., Prommer, I., Scherbov, S., Slentee, E., 2007. Regional, national, and spatially explicit scenarios of demographic and economic change based on SRES. *Technol. Forecast. Soc. Chang.* 74 (7), 980–1029.
- Güneralp, B., Seto, K., 2013. Futures of global urban expansion: uncertainties and implications for biodiversity conservation. *Environ. Res. Lett.* 8 (1), 014025.
- Güneralp, B., Güneralp, İ., Liu, Y., 2015. Changing global patterns of urban exposure to flood and drought hazards. *Glob. Environ. Chang.* 31, 217–225.
- Guo, S., Si, R., Dai, Q., You, Z., Ma, Y., Wang, J., 2019. A critical review of corrosion development and rust removal techniques on the structural/environmental performance of corroded steel bridges. *J. Clean. Prod.* 233, 126–146.
- Guo, Y., Fang, G., Xu, Y.-P., Tian, X., Xie, J., 2020. Identifying how future climate and land use/cover changes impact streamflow in Xinjiang basin, East China. *Sci. Total Environ.* 710, 136275.
- Hall, J.W., Sayers, P.B., Dawson, R.J., 2005. National-scale assessment of current and future flood risk in England and Wales. *Nat. Hazards* 36 (1–2), 147–164.
- Hallegatte, S., 2007. The use of synthetic hurricane tracks in risk analysis and climate change damage assessment. *J. Appl. Meteorol. Climatol.* 46 (11), 1956–1966.
- Hallegatte, S., 2009. Strategies to adapt to an uncertain climate change. *Glob. Environ. Chang.* 19 (2), 240–247.
- Hallegatte, S., Ranger, N., Mestre, O., Dumas, P., Corfee-Morlot, J., Herweijer, C., Wood, R.M., 2011. Assessing climate change impacts, sea level rise and storm surge risk in port cities: a case study on Copenhagen. *Clim. Chang.* 104 (1), 113–137.
- Hammond, A., 1998. Which World?: Scenarios for the 21st Century: Global Destinies, Regional Choices. Earthscan.
- Hawchar, L., Naughton, O., Nolan, P., Stewart, M.G., Ryan, P.C., 2020. A GIS-based framework for high-level climate change risk assessment of critical infrastructure. *Clim. Risk Manag.* 29, 100235.
- Hemmati, M., Ellingwood, B.R., Mahmoud, H.N., 2020. The role of urban growth in resilience of communities under flood risk. *Earth's Future* 8 (3), e2019EF001382.
- Hempel, S., Frieler, K., Warszawski, L., Schewe, J., Piontek, F., 2013. A trend-preserving bias correction—the ISI-MIP approach. *Earth Syst. Dyn.* 4 (2), 219–236.
- Hinkel, J., Nicholls, R.J., Vafeidis, A.T., Tol, R.S., Avagianou, T., 2010. Assessing risk of and adaptation to sea-level rise in the European Union: an application of DIVA. *Mitig. Adapt. Strateg. Glob. Chang.* 15 (7), 703–719.
- Hinkel, J., Brown, S., Exner, L., Nicholls, R.J., Vafeidis, A.T., Kebede, A.S., 2012. Sea-level rise impacts on Africa and the effects of mitigation and adaptation: an application of DIVA. *Reg. Environ. Chang.* 12 (1), 207–224.
- Hinkel, J., van Vuuren, D.P., Nicholls, R.J., Klein, R.J., 2013. The effects of adaptation and mitigation on coastal flood impacts during the 21st century. An application of the DIVA and IMAGE models. *Clim. Chang.* 117 (4), 783–794.

- Hinkel, J., Lincke, D., Vafeidis, A.T., Perrette, M., Nicholls, R.J., Tol, R.S., Marzeion, B., Fettweis, X., Ionescu, C., Levermann, A., 2014. Coastal flood damage and adaptation costs under 21st century sea-level rise. *Proc. Natl. Acad. Sci.* 111 (9), 3292–3297.
- Hourdin, F., Mauritsen, T., Gettelman, A., Golaz, J.-C., Balaji, V., Duan, Q., Folini, D., Ji, D., Klocke, D., Qian, Y., et al., 2017. The art and science of climate model tuning. *Bull. Am. Meteorol. Soc.* 98 (3), 589–602.
- Hu, Z., Lo, C., 2007. Modeling urban growth in Atlanta using logistic regression. *Comput. Environ. Urban. Syst.* 31 (6), 667–688.
- Huang, K., Li, X., Liu, X., Seto, K.C., 2019. Projecting global urban land expansion and heat island intensification through 2050. *Environ. Res. Lett.* 14 (11), 114037.
- Huppert, H.E., Sparks, R.S.J., 2006. Extreme natural hazards: population growth, globalization and environmental change. *Philos. Trans. R. Soc. A Math. Phys. Eng. Sci.* 364 (1845), 1875–1888.
- Iacoletti, S., Cremen, G., Galasso, C., 2021. Advancements in multi-rupture time-dependent seismic hazard modeling, including fault interaction. *Earth Sci. Rev.* 220, 103650 in press.
- IFRC, 2020. World Disasters Report 2020: Come Heat or High Water. International Federation of Red Cross and Red Crescent Societies, Geneva.
- IPCC, 2007. Climate Change 2007: The Physical Science Basis. In: Solomon, S., Qin, D., Manning, M., Chen, Z., Marquis, M., Averyt, K.B., Tignor, M., Miller, H.L. (Eds.), Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, p. 996.
- IPCC, 2014. Summary for policymakers. In: Field, C.B., Barros, V.R., Dokken, D.J., Mach, K.J., Mastrandrea, M.D., Bilir, T.E., Chatterjee, M., Ebi, K.L., Estrada, Y.O., Genova, R.C., Girma, B., Kissel, E.S., Levy, A.N., MacCracken, S., Mastrandrea, P.R., White, L.L. (Eds.), Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, pp. 1–32.
- IPCC, 2021. Summary for Policymakers. In: Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S.L., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M.I., Huang, M., Leitzell, K., Lonnoy, E., Matthews, J.B.R., Maycock, T.K., Waterfield, T., Yelekçi, O., Yu, R., Zhou, B. (Eds.), Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge University Press In Press.
- Jabbarzadeh, A., Fahimnia, B., Sheu, J.-B., Moghadam, H.S., 2016. Designing a supply chain resilient to major disruptions and supply/demand interruptions. *Transp. Res. B Methodol.* 94, 121–149.
- Johansen, I.L., Rausand, M., 2014. Foundations and choice of risk metrics. *Saf. Sci.* 62, 386–399.
- Jongman, B., Winsemius, H.C., Aerts, J.C., De Perez, E.C., Van Aalst, M.K., Kron, W., Ward, P.J., 2015. Declining vulnerability to river floods and the global benefits of adaptation. *Proc. Natl. Acad. Sci.* 112 (18), E2271–E2280.
- Jurgilevich, A., Räsänen, A., Groundstroem, F., Juhola, S., 2017. A systematic review of dynamics in climate risk and vulnerability assessments. *Environ. Res. Lett.* 12 (1), 013002.
- Kappes, M.S., Papathoma-Koehle, M., Keiler, M., 2012. Assessing physical vulnerability for multi-hazards using an indicator-based methodology. *Appl. Geogr.* 32 (2), 577–590.
- Kashani, M.M., Maddocks, J., Dizaj, E.A., 2019. Residual capacity of corroded reinforced concrete bridge components: state-of-the-art review. *J. Bridge. Eng.* 24 (7), 03119001.
- Kebede, A.S., Nicholls, R.J., Hanson, S., Mokrech, M., 2012. Impacts of climate change and sea-level rise: a preliminary case study of Mombasa, Kenya. *J. Coast. Res.* 28 (1A), 8–19.
- Kocabas, V., Dragicevic, S., 2013. Bayesian networks and agent-based modeling approach for urban land-use and population density change: a BNAS model. *J. Geogr. Syst.* 15 (4), 403–426.
- Komendantova, N., Mrzyglocki, R., Mignán, A., Khazai, B., Wenzel, F., Patt, A., Fleming, K., 2014. Multi-hazard and multi-risk decision-support tools as a part of participatory risk governance: feedback from civil protection stakeholders. *Int. J. Disaster Risk Reduct.* 8, 50–67.
- Kwadijk, J.C., Haasnoot, M., Mulder, J.P., Hoogvliet, M.M., Jeuken, A.B., van der Krogt, R.A., van Oostrom, N.G., Schelfhout, H.A., van Velzen, E.H., van Waveren, H., et al., 2010. Using adaptation tipping points to prepare for climate change and sea level rise: a case study in the Netherlands. *Wiley Interdiscip. Rev. Clim. Chang.* 1 (5), 729–740.
- Lallemant, D., 2015. Modeling the Future Disaster Risk of Cities to Envision Paths Towards Their Future Resilience. PhD thesisStanford University.
- Lallemant, D., Burton, H., Ceferino, L., Bullock, Z., Kiremidjian, A., 2017. A framework and case study for earthquake vulnerability assessment of incrementally expanding buildings. *Earthq. Spectra* 33 (4), 1369–1384.
- Lavell, A., Oppenheimer, M., Diop, C., Hess, J., Lempert, R., Li, J., Muir-Wood, R., Myeong, S., Moser, S., Takeuchi, K., 2012. Climate change: new dimensions in disaster risk, exposure, vulnerability, and resilience. Managing the Risks of Extreme Events and Disasters to Advance Climate Change Adaptation: Special Report of the Intergovernmental Panel on Climate Change. Cambridge University Press, pp. 25–64.
- Lee, J.Y., Burton, H.V., Lallemant, D., 2018. Adaptive decision-making for civil infrastructure systems and communities exposed to evolving risks. *Struct. Saf.* 75, 1–12.
- Li, Y., Stewart, M.G., 2011. Cyclone damage risks caused by enhanced greenhouse conditions and economic viability of strengthened residential construction. *Nat. Hazards Rev.* 12 (1), 9–18.
- Liu, L., Yang, D.Y., Frangopol, D.M., 2020. Network-level risk-based framework for optimal bridge adaptation management considering scour and climate change. *J. Infrastruct. Syst.* 26 (1), 04019037.
- Lutz, W., KC, S., 2010. Dimensions of global population projections: what do we know about future population trends and structures? *Philos. Trans. R. Soc. B Biol. Sci.* 365 (1554), 2779–2791.
- Lutz, W., Sanderson, W., Scherbov, S., 1997. Doubling of world population unlikely. *Nature* 387 (6635), 803–805.
- Lutz, W., Sanderson, W., Scherbov, S., 2001. The end of world population growth. *Nature* 412 (6846), 543–545.
- Lutz, W., Sanderson, W., Scherbov, S., 2008. The coming acceleration of global population ageing. *Nature* 451 (7179), 716–719.
- Maaskant, B., Jonkman, S.N., Bouwer, L.M., 2009. Future risk of flooding: an analysis of changes in potential loss of life in South Holland (The Netherlands). *Environ. Sci. Policy* 12 (2), 157–169.
- McLeod, E., Hinkel, J., Vafeidis, A.T., Nicholls, R.J., Harvey, N., Salm, R., 2010. Sea-level rise vulnerability in the countries of the Coral Triangle. *Sustain. Sci.* 5 (2), 207–222.
- Mendelsohn, R., Emanuel, K., Chonabayashi, S., Bakkensen, L., 2012. The impact of climate change on global tropical cyclone damage. *Nat. Clim. Chang.* 2 (3), 205–209.
- Mignán, A., Wiemer, S., Giardini, D., 2014. The quantification of low-probability-high-consequences events: part I. A generic multi-risk approach. *Nat. Hazards* 73 (3), 1999–2022.
- Miller, M., Baker, J.W., 2016. Coupling mode-destination accessibility with seismic risk assessment to identify at-risk communities. *Reliabil. Eng. Syst. Saf.* 147, 60–71.
- Mondoro, A., Frangopol, D.M., Liu, L., 2018. Bridge adaptation and management under climate change uncertainties: a review. *Nat. Hazards Rev.* 19 (1), 04017023.
- Mortin, J., Graversen, R.G., Svensson, G., 2014. Evaluation of pan-Arctic melt-freeze onset in CMIP5 climate models and reanalyses using surface observations. *Clim. Dyn.* 42 (7), 2239–2257.
- Muis, S., Güneralp, B., Jongman, B., Aerts, J.C., Ward, P.J., 2015. Flood risk and adaptation strategies under climate change and urban expansion: a probabilistic analysis using global data. *Sci. Total Environ.* 538, 445–457.
- Murphy, J.M., Sexton, D.M., Barnett, D.N., Jones, G.S., Webb, M.J., Collins, M., Stainforth, D.A., 2004. Quantification of modelling uncertainties in a large ensemble of climate change simulations. *Nature* 430 (7001), 768–772.
- Murphy, J.M., Booth, B.B., Collins, M., Harris, G.R., Sexton, D.M., Webb, M.J., 2007. A methodology for probabilistic predictions of regional climate change from perturbed physics ensembles. *Philos. Trans. R. Soc. A Math. Phys. Eng. Sci.* 365 (1857), 1993–2028.
- Nagai, R., Takabatake, T., Esteban, M., Ishii, H., Shibayama, T., 2020. Tsunami risk hazard in Tokyo bay: the challenge of future sea level rise. *Int. J. Disaster Risk Reduct.* 45, 101321.
- Nakicenovic, N., Alcamo, J., Davis, G., Vries, B.D., Fenhann, J., Gaffin, S., Gregory, K., Grubler, A., Jung, T.Y., Kram, T., 2000. Special report on emissions scenarios. A Special Report of IPCC Working Group III.
- Naumann, G., Cammalleri, C., Mentaschi, L., Feyen, L., 2021. Increased economic drought impacts in Europe with anthropogenic warming. *Nat. Clim. Chang.* 11 (6), 485–491.
- Neumann, B., Vafeidis, A.T., Zimmermann, J., Nicholls, R.J., 2015. Future coastal population growth and exposure to sea-level rise and coastal flooding—a global assessment. *PLoS One* 10 (3), e0118571.
- New, M., Lopez, A., Dessai, S., Wilby, R., 2007. Challenges in using probabilistic climate change information for impact assessments: an example from the water sector. *Philos. Trans. R. Soc. A Math. Phys. Eng. Sci.* 365 (1857), 2117–2131.
- Nicholls, R.J., 2004. Coastal flooding and wetland loss in the 21st century: changes under the SRES climate and socio-economic scenarios. *Glob. Environ. Chang.* 14 (1), 69–86.
- Nicholls, R.J., Tol, R.S., 2006. Impacts and responses to sea-level rise: a global analysis of the SRES scenarios over the twenty-first century. *Philos. Trans. R. Soc. A Math. Phys. Eng. Sci.* 364 (1841), 1073–1095.
- Nilsson, A.E., Bay-Larsen, I., Carlsen, H., van Oort, B., Bjørkan, M., Jylhä, K., Klyuchnikova, E., Masloboev, V., van der Watt, L.-M., 2017. Towards extended shared socioeconomic pathways: a combined participatory bottom-up and top-down methodology with results from the Barents region. *Glob. Environ. Chang.* 45, 124–132.
- O'Neill, B., Krieger, E., Riahi, K., Ebi, K.L., Hallegatte, S., Carter, T.R., Mathur, R., van Vuuren, D.P., 2014. A new scenario framework for climate change research: the concept of shared socioeconomic pathways. *Clim. Chang.* 122 (3).
- Ozturk, T., Turp, M.T., Türkeş, M., Kurnaz, M.L., 2017. Projected changes in temperature and precipitation climatology of Central Asia CORDEX Region 8 by using REGCM4.3.5. *Atmos. Res.* 183, 296–307.
- Pachauri, R.K., Allen, M.R., Barros, V.R., Broome, J.R., Cramer, W., Christ, R., Church, J.A., Clarke, L., Dahe, Q., Dasgupta, P., Dubash, N.K., Edenhofer, O., Elgizouli, I., Field, C.B., Forster, P., Friedlingstein, P., Fuglestedt, J., Gomez-Echeverri, L., Hallegatte, S., Hegerl, G., Howden, M., Jiang, K., Jimenez Cisneros, B., Kattsov, V., Lee, H., Mach, K.J., Marotzke, J., Mastrandrea, M.D., Meyer, L., Minx, J., Muluetta, Y., O'Brien, K., Oppenheimer, M., Pereira, J.J., Pichs-Madruga, R., Plattner, G.K., Pörtner, H.O., Power, S.B., Preston, B., Ravindranath, N.H., Reisinger, A., Riahi, K., Rusticucci, M., Scholes, R., Seyboth, K., Sokona, Y., Stavins, R., Stocker, T.F., Tschakert, P., van Vuuren, D., van Ypersele, J.P., 2014. In: Pachauri, R., Meyer, L. (Eds.), Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. IPCC, Geneva, Switzerland, p. 151 ISBN: 978-92-9169-143-2.
- Pandey, B., Joshi, P., 2015. Numerical modelling spatial patterns of urban growth in Chandigarh and surrounding region (India) using multi-agent systems. *Model. Earth Syst. Environ.* 1 (3), 1–14.
- Papadopoulos, A.N., Bazzurro, P., Marzocchi, W., 2020. Exploring probabilistic seismic risk assessment accounting for seismicity clustering and damage accumulation: part I. Hazard analysis. *Earthq. Spectra* 37 (2), 803–826 page 8755293020957338.
- Pardaens, A., Lowe, J., Brown, S., Nicholls, R., de Gusmão, D.D., 2011. Sea-level rise and impacts projections under a future scenario with large greenhouse gas emission reductions. *Geophys. Res. Lett.* 38 (12).
- Peng, L., Stewart, M.G., 2016. Climate change and corrosion damage risks for reinforced concrete infrastructure in China. *Struct. Infrastruct. Eng.* 12 (4), 499–516.
- Penning-Rowsell, E., Yanyan, W., Watkinson, A., Jiang, J., Thorne, C., 2013. Socioeconomic scenarios and flood damage assessment methodologies for the Taihu Basin, China. *J. Flood Risk Manag.* 6 (1), 23–32.
- Pescaroli, G., Alexander, D., 2018. Understanding compound, interconnected, interacting, and cascading risks: a holistic framework. *Risk Anal.* 38 (11), 2245–2257.

- Petoukhov, V., Ganopolski, A., Brovkin, V., Claussen, M., Eliseev, A., Kubatzki, C., Rahmstorf, S., 2000. Climber-2: a climate system model of intermediate complexity. Part I: model description and performance for present climate. *Clim. Dyn.* 16 (1), 1–17.
- Pisano, L., Zumpano, V., Malek, Z., Rosskopf, C.M., Parise, M., 2017. Variations in the susceptibility to landslides, as a consequence of land cover changes: a look to the past, and another towards the future. *Sci. Total Environ.* 601, 1147–1159.
- Promper, C., Puissant, A., Malet, J.-P., Glade, T., 2014. Analysis of land cover changes in the past and the future as contribution to landslide risk scenarios. *Appl. Geogr.* 53, 11–19.
- Prudhomme, C., Wilby, R., Crooks, S., Kay, A.L., Reynard, N.S., 2010. Scenario-neutral approach to climate change impact studies: application to flood risk. *J. Hydrol.* 390 (3–4), 198–209.
- Qin, H., Stewart, M.G., 2020. Risk-based cost-benefit analysis of climate adaptation measures for Australian contemporary houses under extreme winds. *J. Infrastruct. Preserv. Resil.* 1 (1), 1–19.
- Reguero, B.G., Bresch, D.N., Beck, M., Calil, J., Meliane, I., 2014. Coastal risks, nature-based defenses and the economics of adaptation: an application in the Gulf of Mexico, USA. *Coast. Eng. Proc.* 1 (34), 25.
- Rimal, B., Keshthkar, H., Sharma, R., Stork, N., Rijal, S., Kunwar, R., 2019. Simulating urban expansion in a rapidly changing landscape in eastern Tarai, Nepal. *Environ. Monit. Assess.* 191 (4), 1–14.
- Rojas, R., Feyen, L., Watkiss, P., 2013. Climate change and river floods in the European Union: socio-economic consequences and the costs and benefits of adaptation. *Glob. Environ. Chang.* 23 (6), 1737–1751.
- de Ruiter, M.C., De Bruijn, J.A., Enghardt, J., Daniell, J.E., de Moel, H., Ward, P.J., 2021. The asynergies of structural disaster risk reduction measures: comparing floods and earthquakes. *Earth's Future* 9 (1), e2020EF001531.
- Ryan, P.C., Stewart, M.G., 2017. Cost-benefit analysis of climate change adaptation for power pole networks. *Clim. Chang.* 143 (3), 519–533.
- Ryan, P.C., Stewart, M.G., 2020. Regional variability of climate change adaptation feasibility for timber power poles. *Struct. Infrastruct. Eng.* 1–11.
- Ryan, P.C., Stewart, M.G., Spencer, N., Li, Y., 2016. Probabilistic analysis of climate change impacts on timber power pole networks. *Int. J. Electr. Power Energy Syst.* 78, 513–523.
- Salathé Jr., E.P., 2005. Downscaling simulations of future global climate with application to hydrologic modelling. *Int. J. Climatol.* 25 (4), 419–436.
- Salman, A.M., Li, Y., 2018. Flood risk assessment, future trend modeling, and risk communication: a review of ongoing research. *Nat. Hazards Rev.* 19 (3), 04018011.
- Salmanidou, D.M., Ehara, A., Himaz, R., Heidarzadeh, M., Guillas, S., 2021. Impact of future tsunamis from the Java trench on household welfare: merging geophysics and economics through catastrophe modelling. *Int. J. Disaster Risk Reduct.* 61, 102291.
- Santini, M., Valentini, R., 2011. Predicting hot-spots of land use changes in Italy by ensemble forecasting. *Reg. Environ. Chang.* 11 (3), 483–502.
- Sarica, G.M., Zhu, T., Pan, T.-C., 2020. Spatio-temporal dynamics in seismic exposure of Asian megacities: past, present and future. *Environ. Res. Lett.* 15 (9), 094092.
- Savarese, A., 2016. The Paris agreement: a new beginning? *J. Energ. Nat. Resour. Law* 34 (1), 16–26.
- Schuster, Z.T., Potter, K.W., Liebl, D.S., 2012. Assessing the effects of climate change on precipitation and flood damage in Wisconsin. *J. Hydrol. Eng.* 17 (8), 888–894.
- Schwierz, C., Köllner-Heck, P., Mutter, E.Z., Bresch, D.N., Vidale, P.-L., Wild, M., Schär, C., 2010. Modelling European winter wind storm losses in current and future climate. *Clim. Chang.* 101 (3), 485–514.
- Selva, J., 2013. Long-term multi-risk assessment: statistical treatment of interaction among risks. *Nat. Hazards* 67 (2), 701–722.
- Seto, K.C., Güneralp, B., Hutrya, L.R., 2012. Global forecasts of urban expansion to 2030 and direct impacts on biodiversity and carbon pools. *Proc. Natl. Acad. Sci.* 109 (40), 16083–16088.
- Sevieri, G., Galasso, C., 2021. Typhoon risk and climate-change impact assessment for cultural heritage asset roofs. *Struct. Saf.* 91, 102065.
- Siegert, M., Alley, R.B., Rignot, E., Englander, J., Corell, R., 2020. Twenty-first century sea-level rise could exceed IPCC projections for strong-warming futures. *One Earth* 3 (6), 691–703.
- Soares-Filho, B., Alencar, A., Nepstad, D., Cerqueira, G., Vera Diaz, M.D.C., Rivero, S., Solórzano, L., Voll, E., 2004. Simulating the response of land-cover changes to road paving and governance along a major Amazon highway: the Santarém-Cuiabá corridor. *Glob. Change Biol.* 10 (5), 745–764.
- Soltani, M., Abu-Abaleh, A., Scott Rowe, B., 2021. Statistical approach to modeling reduced shear capacity of corrosion-damaged reinforced concrete beams. *Pract. Period. Struct. Des. Constr.* 26 (2), 04020073.
- Stewart, M.G., 2015. Risk and economic viability of housing climate adaptation strategies for wind hazards in Southeast Australia. *Mitig. Adapt. Strateg. Glob. Chang.* 20 (4), 601–622.
- Stewart, M.G., 2016. Climate change impact assessment of metal-clad buildings subject to extreme wind loading in non-cyclonic regions. *Sustain. Resil. Infrastruct.* 1 (1–2), 32–45.
- Stewart, M.G., Deng, X., 2015. Climate impact risks and climate adaptation engineering for built infrastructure. *ASCE-ASME J. Risk Uncertain. Eng. Syst. Part A* 1 (1), 04014001.
- Stewart, M.G., Li, Y., 2010. Methodologies for economic impact and adaptation assessment of cyclone damage risks due to climate change. *Aust. J. Struct. Eng.* 10 (2), 121–135.
- Stewart, M.G., Wang, X., Nguyen, M.N., 2011. Climate change impact and risks of concrete infrastructure deterioration. *Eng. Struct.* 33 (4), 1326–1337.
- Stewart, M.G., Wang, X., Nguyen, M.N., 2012. Climate change adaptation for corrosion control of concrete infrastructure. *Struct. Saf.* 35, 29–39.
- Stewart, M.G., Val, D.V., Bastidas-Arteaga, E., O'Connor, A.J., Wang, X., 2014. Climate Adaptation Engineering and Risk-based Design and Management of Infrastructure.
- Stewart, M.G., Ginger, J.D., Henderson, D.J., Ryan, P.C., 2018. Fragility and climate impact assessment of contemporary housing roof sheeting failure due to extreme wind. *Eng. Struct.* 171, 464–475.
- Stone Jr., B., Vargo, J., Liu, P., Habeeb, D., DeLucia, A., Trail, M., Hu, Y., Russell, A., 2014. Avoided heat-related mortality through climate adaptation strategies in three US cities. *PLoS one* 9 (6), e100852.
- Tang, J., Wang, L., Yao, Z., 2007. Spatio-temporal urban landscape change analysis using the Markov chain model and a modified genetic algorithm. *Int. J. Remote Sens.* 28 (15), 3255–3271.
- Taubenböck, H., Wegmann, M., Roth, A., Mehl, H., Dech, S., 2009. Urbanization in India—spatiotemporal analysis using remote sensing data. *Comput. Environ. Urban. Syst.* 33 (3), 179–188.
- Tayyebi, A., Pijanowski, B.C., Tayyebi, A.H., 2011. An urban growth boundary model using neural networks, GIS and radial parameterization: an application to Tehran, Iran. *Lands. Urban Plan.* 100 (1–2), 35–44.
- Te Linde, A., Bubeck, P., Dekkers, J., Moel, H.D., Aerts, J., 2011. Future flood risk estimates along the river Rhine. *Nat. Hazards and Earth System Sciences* 11 (2), 459–473.
- Teutschbein, C., Seibert, J., 2012. Bias correction of regional climate model simulations for hydrological climate-change impact studies: review and evaluation of different methods. *J. Hydrol.* 456, 12–29.
- Tol, R.S., 2007. The double trade-off between adaptation and mitigation for sea level rise: an application of FUND. *Mitig. Adapt. Strateg. Glob. Chang.* 12 (5), 741–753.
- Tosun, J., Howlett, M., 2021. Managing slow onset events related to climate change: the role of public bureaucracy. *Curr. Opin. Environ. Sustain.* 50, 43–53.
- UNDRR, 2020. Hazard Definition & Classification Review. United Nations Office for Disaster Risk Reduction, Geneva.
- UNISDR, 2009. Terminology on Disaster Risk Reduction. United Nations International Strategy for Disaster Reduction, Geneva.
- UNISDR, 2015. Sendai Framework for Disaster Risk Reduction 2015–30. United Nations International Strategy for Disaster Reduction, Geneva.
- United Nations, 2019. World population prospects: the 2019 revision. Population Division of the Department of Economic and Social Affairs of the United Nations Secretariat, New York.
- Van Vuuren, D.P., Edmonds, J., Kainuma, M., Riahi, K., Thomson, A., Hibbard, K., Hurtt, G.C., Kram, T., Krey, V., Lamarque, J.-F., et al., 2011. The representative concentration pathways: an overview. *Clim. Chang.* 109 (1), 5–31.
- Van Vuuren, D.P., Kok, M.T., Girod, B., Lucas, P.L., de Vries, B., 2012. Scenarios in global environmental assessments: key characteristics and lessons for future use. *Glob. Environ. Chang.* 22 (4), 884–895.
- Verburg, P.H., Overmars, K.P., 2009. Combining top-down and bottom-up dynamics in land use modeling: exploring the future of abandoned farmlands in Europe with the Dyna-CLUE model. *Lands. Ecol.* 24 (9), 1167–1181.
- Verschuur, J., Koks, E., Haque, A., Hall, J., 2020. Prioritising resilience policies to reduce welfare losses from natural disasters: a case study for coastal Bangladesh. *Glob. Environ. Chang.* 65, 102179.
- van Vuuren, D.P., Lucas, P.L., Hilderink, H., 2007. Downscaling drivers of global environmental change: enabling use of global sres scenarios at the national and grid levels. *Glob. Environ. Chang.* 17 (1), 114–130.
- van Vuuren, D.P., Smith, S.J., Riahi, K., 2010. Downscaling socioeconomic and emissions scenarios for global environmental change research: a review. *Wiley Interdiscip. Rev. Clim. Chang.* 1 (3), 393–404.
- Wang, X., Stewart, M.G., Nguyen, M., 2012. Impact of climate change on corrosion and damage to concrete infrastructure in Australia. *Clim. Chang.* 110 (3), 941–957.
- Ward, P., Van Pelt, S., De Keizer, O., Aerts, J., Beersma, J., Van den Hurk, B., Te Linde, A., 2014. Including climate change projections in probabilistic flood risk assessment. *J. Flood Risk Manag.* 7 (2), 141–151.
- Ward, P.J., Jongman, B., Aerts, J.C., Bates, P.D., Botzen, W.J., Loaiza, A.D., Hallegatte, S., Kind, J.M., Kwadijck, J., Scussolini, P., et al., 2017. A global framework for future costs and benefits of river-flood protection in urban areas. *Nat. Clim. Chang.* 7 (9), 642–646.
- Ward, P.J., Blauhut, V., Bloemendaal, N., Daniell, J.E., Ruitter, M.C.D., Duncan, M.J., Emberson, R., Jenkins, S.F., Kirschbaum, D., Kunz, M., 2020. Natural hazard risk assessments at the global scale. *Nat. Hazards Earth Syst. Sci.* 20 (4), 1069–1096.
- Webersik, C., Esteban, M., Shibayama, T., 2010. The economic impact of future increase in tropical cyclones in Japan. *Nat. Hazards* 55 (2), 233–250.
- Wigley, T.M., Raper, S.C., 2001. Interpretation of high projections for global-mean warming. *Science* 293 (5529), 451–454.
- Wilby, R.L., Dessai, S., 2010. Robust adaptation to climate change. *Weather* 65 (7), 180–185.
- Winsemius, H.C., Jongman, B., Veldkamp, T.I., Hallegatte, S., Bangalore, M., Ward, P.J., 2015. Disaster Risk, Climate Change, and Poverty: Assessing the Global Exposure of Poor People to Floods and Droughts.
- Xu, K., Xu, B., Ju, J., Wu, C., Dai, H., Hu, B.X., 2019. Projection and uncertainty of precipitation extremes in the crip5 multimodel ensembles over nine major basins in China. *Atmos. Res.* 226, 122–137.
- Xu, H., Lin, N., Huang, M., Lou, W., 2020. Design tropical cyclone wind speed when considering climate change. *J. Struct. Eng.* 146 (5), 04020063.
- Yang, D.Y., Frangopol, D.M., 2019a. Physics-based assessment of climate change impact on long-term regional bridge scour risk using hydrologic modeling: application to Lehigh river watershed. *J. Bridg. Eng.* 24 (11), 04019099.
- Yang, D.Y., Frangopol, D.M., 2019b. Societal risk assessment of transportation networks under uncertainties due to climate change and population growth. *Struct. Saf.* 78, 33–47.
- Yang, D.Y., Frangopol, D.M., 2020. Risk-based portfolio management of civil infrastructure assets under deep uncertainties associated with climate change: a robust optimisation approach. *Struct. Infrastruct. Eng.* 16 (4), 531–546.
- Zamanian, S., Hur, J., Shafieezadeh, A., 2020. A high-fidelity computational investigation of buried concrete sewer pipes exposed to truckloads and corrosion deterioration. *Eng. Struct.* 221, 111043.

- Zanini, M.A., Toska, K., Faleschini, F., Pellegrino, C., 2020. Seismic reliability of reinforced concrete bridges subject to environmental deterioration and strengthened with FRCM composites. *Soil Dyn. Earthq. Eng.* 136, 106224.
- Zhou, Y., Varquez, A.C., Kanda, M., 2019. High-resolution global urban growth projection based on multiple applications of the SLEUTH urban growth model. *Sci. Data* 6 (1), 1–10.
- Zölch, T., Henze, L., Keilholz, P., Pauleit, S., 2017. Regulating urban surface runoff through nature-based solutions—an assessment at the micro-scale. *Environ. Res.* 157, 135–144.
- Zscheischler, J., Westra, S., Van Den Hurk, B.J., Seneviratne, S.I., Ward, P.J., Pitman, A., AghaKouchak, A., Bresch, D.N., Leonard, M., Wahl, T., et al., 2018. Future climate risk from compound events. *Nat. Clim. Chang.* 8 (6), 469–477.