



Drew, S., Fox, F. E., Gregson, C. L., Patel, R., Judge, A., Marques, E. M. R., Capelas Barbosa, E., Bradshaw, M. G. E. L., Whale, K., Ben-Shlomo, Y., & Gooberman-Hill, R. J. S. (2022). *Multiple Organisational Factors Improve Multi-Disciplinary Care Delivery to Patients with Hip Fractures: A Qualitative Study of Service Delivery*. Abstract from British Geriatrics Society Spring Meeting 2022.

Peer reviewed version

[Link to publication record in Explore Bristol Research](#)
PDF-document

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

British Geriatrics Society Spring Meeting 2022

Title: 127 characters (max 128)

Multiple Organisational Factors Improve Multi-Disciplinary Care Delivery to Patients with Hip Fractures: A Qualitative Study of Service Delivery

Authors: 70 words (max 70)

S Drew¹; F Fox¹; CL Gregson¹; R Patel¹; A Judge¹; A Johansen²; EMR Marques¹; E Capelas Barbosa¹; J Griffin³; M Bradshaw¹; K Whale¹, T Chesser⁴; XL Griffin⁵; MK Javaid⁶; Y Ben-Shlomo¹, Gooberman-Hill R ^{1, 7}

1. University of Bristol; 2. Cardiff University and University Hospital of Wales; 3. Royal Osteoporosis Society; 4. North Bristol NHS Trust; 5. Queen Mary University of London; 6. University of Oxford, 7. NIHR Bristol BRC

Abstract: 300 words (max 300)

Introduction

Hip fractures are devastating injuries which incur high healthcare costs. Despite national standards and guidelines, there is substantial variation in hospital delivery of hip fracture care and in patient outcomes. This study aimed to understand organisational processes that facilitate successful delivery of hip fracture services.

Methods

Forty qualitative interviews were conducted with healthcare professionals involved in delivering hip fracture care at four English hospitals. Interview data were supplemented with documentary analysis of 23 anonymised British Orthopaedic Association hospital-initiated peer-review reports of services. Data were analysed thematically, with themes transposed onto key components of the care pathway.

Results

We identified multiple aspects of service organisation that facilitated good care delivery. At admission, standardisation of training in nerve block administration impacted care delivery. During hospital stays, service delivery was improved by integrated, shared-care between orthopaedics and orthogeriatrics, and by strategies to improve trauma list efficiency. Adequately staffed orthogeriatric services and the 'right' skills and seniority mix were important to holistic care provision. Placing patients on designated hip fracture wards concentrated staff expertise. Collaborative working was achieved through multi-disciplinary team (MDT) meetings between key staff, protocols and care pathways that defined roles and responsibilities, MDT documentation, 'joined-up' IT systems within hospitals and with primary care, and shared working spaces such as shared offices and on wards. Trauma and hip fracture coordinators organised care processes and provided a valuable central point of contact within teams. Nominated leads, representing diverse specialties, worked together in MDT planning meetings to develop joint protocols, establish audit priorities, and agree shared goals.

Routine, comprehensive monitoring and evaluation of service delivery, with findings shared throughout the MDT, was beneficial.

Conclusions

Our study has characterised potentially modifiable elements of successful hip fracture service delivery. Findings are intended to help services overcome organisational barriers towards delivery of high-quality hip fracture services.