

Samir Al Mohammady; Sid Ahmed Ould Beinane; Sid Ahmed Ould Ahmed Mahmoud
On (n, m) - A -normal and (n, m) - A -quasinormal semi-Hilbertian space operators

Mathematica Bohemica, Vol. 147 (2022), No. 2, 169–186

Persistent URL: <http://dml.cz/dmlcz/150326>

Terms of use:

© Institute of Mathematics AS CR, 2022

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

ON (n, m) - A -NORMAL AND (n, m) - A -QUASINORMAL
SEMI-HILBERTIAN SPACE OPERATORS

SAMIR AL MOHAMMADY, SID AHMED OULD BEINANE,
SID AHMED OULD AHMED MAHMOUD, Sakaka

Received November 30, 2019. Published online June 4, 2021.
Communicated by Laurian Suciu

Abstract. The purpose of the paper is to introduce and study a new class of operators on semi-Hilbertian spaces, i.e. spaces generated by positive semi-definite sesquilinear forms. Let \mathcal{H} be a Hilbert space and let A be a positive bounded operator on \mathcal{H} . The semi-inner product $\langle h | k \rangle_A := \langle Ah | k \rangle$, $h, k \in \mathcal{H}$, induces a semi-norm $\|\cdot\|_A$. This makes \mathcal{H} into a semi-Hilbertian space. An operator $T \in \mathcal{B}_A(\mathcal{H})$ is said to be (n, m) - A -normal if $[T^n, (T^{\sharp_A})^m] := T^n(T^{\sharp_A})^m - (T^{\sharp_A})^m T^n = 0$ for some positive integers n and m .

Keywords: semi-Hilbertian space; A -normal operator; (n, m) -normal operator; (n, m) -quasinormal operator

MSC 2020: 54E40, 47B99

1. INTRODUCTION AND PRELIMINARIES

Throughout this paper, let $(\mathcal{H}, \langle \cdot | \cdot \rangle)$ be a complex Hilbert space equipped with the norm $\|\cdot\|$. Let $\mathcal{B}(\mathcal{H})$ denote the C^* -algebra of all bounded linear operators on \mathcal{H} and let $\mathcal{B}(\mathcal{H})^+$ be the *cone of positive operators* of $\mathcal{B}(\mathcal{H})$ defined as

$$\mathcal{B}(\mathcal{H})^+ := \{A \in \mathcal{B}(\mathcal{H}) : \langle Ah | h \rangle \geq 0 \ \forall h \in \mathcal{H}\}.$$

For every $T \in \mathcal{B}(\mathcal{H})$ its range is denoted by $\mathcal{R}(T)$, its null space by $\mathcal{N}(T)$, and its adjoint by T^* . If \mathcal{M} is a linear subspace of \mathcal{H} , then $\overline{\mathcal{M}}$ stands for its closure in the norm topology of \mathcal{H} . We denote the orthogonal projection onto a closed linear subspace \mathcal{M} of \mathcal{H} by $P_{\mathcal{M}}$. The positive operator $A \in \mathcal{B}(\mathcal{H})$ defines a positive semi-definite sesquilinear form $\langle \cdot | \cdot \rangle_A : \mathcal{H} \times \mathcal{H} \rightarrow \mathbb{C}$ given by $\langle h | k \rangle_A = \langle Ah | k \rangle$. Note that $\langle \cdot | \cdot \rangle_A$ defines a semi-inner product on \mathcal{H} , and the semi-norm induced by it is

given by $\|h\|_A = \sqrt{\langle h | h \rangle_A}$ for every $h \in \mathcal{H}$. Observe that $\|h\|_A = 0$ if and only if $h \in \mathcal{N}(A)$. Then $\|\cdot\|_A$ is a norm if and only if A is injective, and the semi-normed space $(\mathcal{H}, \|\cdot\|_A)$ is a complete space if and only if $\mathcal{R}(A)$ is closed.

The above semi-norm induces a semi-norm on the subspace $\mathcal{B}^A(\mathcal{H})$ of $\mathcal{B}(\mathcal{H})$ consisting of all $T \in \mathcal{B}(\mathcal{H})$ so that for some $c > 0$ and for all $h \in \mathcal{H}$, $\|Th\|_A \leq c\|h\|_A$. Indeed, if $T \in \mathcal{B}^A(\mathcal{H})$, then

$$\|T\|_A := \sup \left\{ \frac{\|Th\|_A}{\|h\|_A}, h \notin \mathcal{N}(A) \right\}.$$

For $T \in \mathcal{B}(\mathcal{H})$, an operator $S \in \mathcal{B}(\mathcal{H})$ is called an *A-adjoint operator* of T if for every $h, k \in \mathcal{H}$ we have $\langle Th | k \rangle_A = \langle h | Sk \rangle_A$, that is, $AS = T^*A$. If T is an *A-adjoint* of itself, then T is called an *A-selfadjoint operator*.

Generally, the existence of an *A-adjoint operator* is not guaranteed. The set of all operators that admit *A-adjoints* is denoted by $\mathcal{B}_A(\mathcal{H})$. An application of the Douglas theorem (see [13]) shows that

$$\begin{aligned} \mathcal{B}_A(\mathcal{H}) &= \{T \in \mathcal{B}(\mathcal{H}) : \mathcal{R}(T^*A) \subseteq \mathcal{R}(A)\} \\ &= \{T \in \mathcal{B}(\mathcal{H}) : \exists c > 0 : \|ATx\| \leq c\|Ax\| \ \forall x \in \mathcal{H}\}. \end{aligned}$$

Note that $\mathcal{B}_A(\mathcal{H})$ is a subalgebra of $\mathcal{B}(\mathcal{H})$, which is neither closed nor dense in $\mathcal{B}(\mathcal{H})$. Moreover, the inclusions $\mathcal{B}_A(\mathcal{H}) \subseteq \mathcal{B}^A(\mathcal{H}) \subseteq \mathcal{B}(\mathcal{H})$ hold with equality if A is one-to-one and has a closed range. If $T \in \mathcal{B}_A(\mathcal{H})$, the reduced solution of the equation $AX = T^*A$ is a distinguished *A-adjoint operator* of T , which is denoted by $T^{\sharp A}$. Note that $T^{\sharp A} = A^\dagger T^*A$ in which A^\dagger is the Moore-Penrose inverse of A . It was observed that the *A-adjoint operator* $T^{\sharp A}$ satisfies

$$AT^{\sharp A} = T^*A, \quad \mathcal{R}(T^{\sharp A}) \subseteq \overline{\mathcal{R}(A)}$$

and

$$\mathcal{N}(T^{\sharp A}) = \mathcal{N}(T^*A).$$

For $T, S \in \mathcal{B}_A(\mathcal{H})$, it is easy to see that $\|TS\|_A \leq \|T\|_A\|S\|_A$ and $(TS)^{\sharp A} = S^{\sharp A}T^{\sharp A}$.

Notice that if $T \in \mathcal{B}_A(\mathcal{H})$, then $T^{\sharp A} \in \mathcal{B}_A(\mathcal{H})$, $(T^{\sharp A})^{\sharp A} = P_{\overline{\mathcal{R}(A)}} T P_{\overline{\mathcal{R}(A)}}$ and $((T^{\sharp A})^{\sharp A})^{\sharp A} = T^{\sharp A}$. (For more detail on the concepts cited above see [5], [4], [6].)

In [17] it was observed that if $T \in \mathcal{B}_A(\mathcal{H})$ is such that $TA = AT$, then $T^{\sharp A} = PT^*$. For an arbitrary operator $T \in \mathcal{B}_A(\mathcal{H})$, we can write

$$\operatorname{Re}_A(T) := \frac{1}{2}(T + T^{\sharp A}) \quad \text{and} \quad \operatorname{Im}_A(T) := \frac{1}{2i}(T - T^{\sharp A}).$$

The concept of n -normal operators as a generalization of normal operators on Hilbert spaces has been introduced and studied by Jibril (see [15]) and Alzuraiqi et al. (see [3]). The class of n -power normal operators is denoted by $[n\mathbf{N}]$. An operator T is called n -power normal if $[T^n, T^*] = 0$ (equivalently $T^n T^* = T^* T^n$). Very recently, several papers have appeared on n -normal operators. We refer the interested reader to [12], [11], [16] for the complete details.

In [1] and [2], the authors introduced and studied the classes of (n, m) -normal powers and (n, m) -power quasinormal operators as follows: An operator $T \in \mathcal{B}(\mathcal{H})$ is said to be (n, m) -power normal if $T^n (T^m)^* = (T^m)^* T^n$ and it is said to be (n, m) -power quasinormal if $T^n (T^*)^m T = (T^*)^m T T^n$ where n, m are two nonnegative integers. We refer the interested reader to [11] for the complete details on (n, m) -power normal operators.

The classes of normal, (α, β) -normal, and n -power quasinormal operators, isometries, partial isometries, unitary operators etc. on Hilbert spaces have been generalized to semi-Hilbertian spaces by many authors in many papers. (See, for more details, [5]–[7], [9], [10], [14], [17], [18], [21].)

An operator $T \in \mathcal{B}_A(\mathcal{H})$ is said to be

- (1) A -normal if $T^{\sharp_A} T = T T^{\sharp_A}$ (see [17]),
- (2) (α, β) - A -normal if $\beta^2 T^{\sharp_A} T \geq_A T T^{\sharp_A} \geq_A \alpha^2 T^{\sharp_A} T$ for $0 \leq \alpha \leq 1 \leq \beta$ (see [9]),
- (3) (A, n) -power-quasinormal if $T^n (T^{\sharp_A} T) = (T T^{\sharp_A}) T^n$ (see [14]),
- (4) an A -isometry if $T^{\sharp_A} T = P_{\overline{\mathcal{R}(A)}}$ (see [5]),
- (5) A -unitary if $T^{\sharp_A} T = (T^{\sharp_A})^{\sharp_A} T^{\sharp_A} = P_{\overline{\mathcal{R}(A)}}$, i.e. T and T^{\sharp_A} are A -isometries (see [5]).

From now on, A denotes a positive operator on \mathcal{H} , i.e. $A \in \mathcal{B}(\mathcal{H})^+$.

This paper is devoted to the study of some new classes of operators on semi-Hilbertian spaces called (n, m) - A -normal operators and (n, m) - A -quasinormal operators. Some properties of these classes are investigated.

2. (n, m) - A -NORMAL OPERATORS

In this section, the class of (n, m) - A -normal operators as a generalization of the classes of A -normal operators is introduced. In addition, we study several properties of members of this class of operators.

Definition 2.1. Let $T \in \mathcal{B}_A(\mathcal{H})$. We say that T is (n, m) - A -normal if

$$(2.1) \quad [T^n, (T^{\sharp_A})^m] := T^n (T^{\sharp_A})^m - (T^{\sharp_A})^m T^n = 0$$

for some positive integers n and m . The set of all operators which are (n, m) - A -normal is denoted by $[(n, m)\mathbf{N}]_A$.

Remark 2.1. We make the following observations:

- (1) Every A -normal operator is an (n, m) - A -normal for all $n, m \in \mathbb{N}$.
- (2) If $n = m = 1$, every $(1, 1)$ - A -normal operator is an A -normal operator.
- (3) If $T \in [(1, m)\mathbf{N}]_A$ then $T \in [(n, m)\mathbf{N}]_A$ and if $T \in [(n, 1)\mathbf{N}]_A$ then $T \in [(n, m)\mathbf{N}]_A$.
- (4) If $T \in [(n, m)\mathbf{N}]_A$ then $T \in [(2n, m)\mathbf{N}]_A \cap [(n, 2m)\mathbf{N}]_A \cap [(2n, 2m)\mathbf{N}]_A$.

Remark 2.2. In the following example we present an operator that is (n, m) - A -normal for some positive integers n and m but is not an A -normal operator.

Example 2.1. Let $T = \begin{pmatrix} 2 & 0 \\ 1 & -2 \end{pmatrix}$ and $A = \begin{pmatrix} 1 & 0 \\ 0 & 2 \end{pmatrix}$ be operators acting on two-dimensional Hilbert space \mathbb{C}^2 . A simple calculation shows that $T^{\sharp A} = \begin{pmatrix} 2 & 2 \\ 0 & -2 \end{pmatrix}$. Moreover, $T^{\sharp A}T \neq TT^{\sharp A}$ and $T^{\sharp A}T^2 = T^2T^{\sharp A}$. Therefore T is a $(2, 1)$ - A -normal but not an A -normal operator.

In [17], Theorem 2.1 it was observed that if $T \in \mathcal{B}_A(\mathcal{H})$ then T is A -normal if and only if

$$\|Th\|_A = \|T^{\sharp A}h\|_A \quad \forall h \in \mathcal{H} \quad \text{and} \quad \mathcal{R}(TT^{\sharp A}) \subseteq \overline{\mathcal{R}(A)}.$$

In the following theorem, we generalize this characterization to (n, m) - A -normal operators.

Theorem 2.1. *Let $T \in \mathcal{B}_A(\mathcal{H})$. Then T is an (n, m) - A -normal operator for some positive integers n and m if and only if T satisfies the conditions:*

- (1) $\langle (T^{\sharp A})^m h \mid (T^{\sharp A})^n h \rangle_A = \langle (T^n h \mid T^m h)_A \quad \forall h \in \mathcal{H},$
- (2) $\mathcal{R}(T^n (T^{\sharp A})^m) \subseteq \overline{\mathcal{R}(A)}.$

Proof. Assume that T is an (n, m) - A -normal operator and we need to proof that T satisfies the conditions (1) and (2). In fact, we have

$$\begin{aligned} \langle [T^n, (T^{\sharp A})^m]h \mid h \rangle_A = 0 &\Rightarrow \langle T^n (T^{\sharp A})^m h \mid h \rangle_A - \langle (T^{\sharp A})^m T^n h \mid h \rangle_A = 0 \\ &\Rightarrow \langle (T^{\sharp A})^m h \mid T^{*n} A h \rangle - \langle A (T^{\sharp A})^m T^n h \mid h \rangle = 0 \\ &\Rightarrow \langle (T^{\sharp A})^m h \mid (T^{\sharp A})^n h \rangle_A - \langle T^n h \mid T^m h \rangle_A = 0 \\ &\Rightarrow \langle (T^{\sharp A})^m h \mid (T^{\sharp A})^n h \rangle_A = \langle T^n h \mid T^m h \rangle_A. \end{aligned}$$

Moreover, the condition $[T^n, (T^{\sharp A})^m] = 0$ implies that $T^n (T^{\sharp A})^m = (T^{\sharp A})^m T^n$. Therefore

$$\mathcal{R}(T^n (T^{\sharp A})^m) = \mathcal{R}((T^{\sharp A})^m T^n) \subseteq \mathcal{R}(T^{\sharp A}) \subseteq \overline{\mathcal{R}(A)}.$$

Conversely, assume that T satisfies the conditions (1) and (2) and we prove that T is an (n, m) - A -normal operator. From the condition (1), a simple computation shows that

$$\begin{aligned} \langle (T^{\sharp A})^m h \mid (T^{\sharp A})^n h \rangle_A - \langle T^n h \mid T^m h \rangle_A &= 0 \\ \Rightarrow \langle T^n (T^{\sharp A})^m h \mid h \rangle_A - \langle (T^{\sharp A})^m T^n h \mid h \rangle_A &= 0 \\ \Rightarrow \langle [T^n, (T^{\sharp A})^m] h \mid h \rangle_A &= 0, \end{aligned}$$

which implies that $\mathcal{R}([T^n, (T^{\sharp A})^m]) \subseteq \mathcal{N}(A)$.

On the other hand, since the condition (2) holds, it follows that

$$\mathcal{R}([T^n, (T^{\sharp A})^m]) \subseteq \overline{\mathcal{R}(A)} = \mathcal{N}(A)^\perp.$$

We deduce that $[T^n, (T^{\sharp A})^m] = 0$ which means that the operator T is (n, m) - A -normal. \square

Remark 2.3. If $n = m = 1$, then Theorem 2.1 coincides with Theorem 2.1 of [17].

The following proposition discusses the relation between (n, m) - A -normal operators and (m, n) - A -normal operators.

Proposition 2.1. *Let $T \in \mathcal{B}_A(\mathcal{H})$ be such that $\mathcal{N}(A)^\perp$ is an invariant subspace of T . Then the following statements are equivalent.*

- (1) T is an (n, m) - A -normal operator.
- (2) T is an (m, n) - A -normal operator.

Proof. (1) \Rightarrow (2) Assume that T is an (n, m) - A -normal operator. It follows that

$$T^n (T^{\sharp A})^m - (T^{\sharp A})^m T^n = 0.$$

Then

$$\begin{aligned} T^n (T^{\sharp A})^m - (T^{\sharp A})^m T^n &= 0 \\ \Rightarrow [(T^{\sharp A})^{\sharp A}]^m (T^n)^{\sharp A} - (T^n)^{\sharp A} [(T^{\sharp A})^{\sharp A}]^m &= 0 \\ \Rightarrow (P_{\overline{\mathcal{R}(A)}} T P_{\overline{\mathcal{R}(A)}})^m (T^n)^{\sharp A} - (T^n)^{\sharp A} (P_{\overline{\mathcal{R}(A)}} T P_{\overline{\mathcal{R}(A)}})^m &= 0 \\ \Rightarrow P_{\overline{\mathcal{R}(A)}} (T^m (T^n)^{\sharp A} - (T^n)^{\sharp A} T^m) &= 0. \end{aligned}$$

This means that $(T^m (T^n)^{\sharp A} - (T^n)^{\sharp A} T^m)h \in \mathcal{N}(A)$ for all $h \in \mathcal{H}$.

On the other hand, this fact and $\mathcal{R}(T^{\sharp A n}) \subset \mathcal{R}(T^{\sharp A}) \subset \overline{\mathcal{R}(A)}$ and the assumption that $\mathcal{N}(A)^\perp$ is an invariant subspace for T imply that $(T^m (T^n)^{\sharp A} - (T^n)^{\sharp A} T^m)h \in \overline{\mathcal{R}(A)}$ for all $h \in \mathcal{H}$. Consequently, $(T^m (T^n)^{\sharp A} - (T^n)^{\sharp A} T^m)h = 0$ for all $h \in \mathcal{H}$. Therefore $[T^m, (T^{\sharp A})^n] = 0$. Hence $T^{\sharp A}$ is an (m, n) - A -normal operator.

(2) \Rightarrow (1) By the same way hence we omit it. \square

It is well known that if $T \in \mathcal{B}_A(\mathcal{H})$ is A -normal, then T^n is A -normal. In the following theorem, we extend this result to an (n, m) - A -normal operator as follows.

Theorem 2.2. *Let $T \in \mathcal{B}_A(\mathcal{H})$. If T is an (n, m) - A -normal operator then the following statements hold:*

- (i) T^j is A -normal where j is the least common multiple of n and m , i.e. $j = LCM(n, m)$,
- (ii) T^{nm} is an A -normal operator.

Proof. (i) Assume that T is (n, m) - A -normal that is $T^n(T^{\sharp_A})^m = (T^{\sharp_A})^m T^n$. Let $j = pn$ and $j = qm$. By computation we get

$$\begin{aligned}
 T^j (T^j)^{\sharp_A} &= T^{pn} ((T^{\sharp_A})^{qm}) = (T^n)^p ((T^{\sharp_A})^m)^q \\
 &= \underbrace{T^n \dots T^n}_{p\text{-times}} \underbrace{(T^{\sharp_A})^m \dots (T^{\sharp_A})^m}_{q\text{-times}} \\
 &= \underbrace{(T^{\sharp_A})^m \dots (T^{\sharp_A})^m}_{q\text{-times}} \underbrace{T^n \dots T^n}_{p\text{-times}} \\
 &= (T^{\sharp_A})^{qm} T^{np} = (T^{qm})^{\sharp_A} T^{np} = (T^j)^{\sharp_A} T^j,
 \end{aligned}$$

which means that T^j is A -normal.

- (ii) This statement is proved in the same way as the statement (i). □

Proposition 2.2. *Let $T \in \mathcal{B}_A(\mathcal{H})$, $X = T^n + (T^{\sharp_A})^m$, $Y = T^n - (T^{\sharp_A})^m$ and $Z = T^n (T^{\sharp_A})^m$. The following statements hold:*

- (1) T is (n, m) - A -normal if and only if $[X, Y] = 0$.
- (2) If $T \in [(n, m)\mathbf{N}]_A$, then $[Z, X] = [Z, Y] = 0$.
- (3) $T \in [(n, m)\mathbf{N}]_A$ if and only if $[T^n, X] = 0$.
- (4) $T \in [(n, m)\mathbf{N}]_A$ if and only if $[T^n, Y] = 0$.

Proof. (1)

$$\begin{aligned}
 [X, Y] = XY - YX = 0 &\Leftrightarrow ((T^n + (T^{\sharp_A})^m)(T^n - (T^{\sharp_A})^m)) \\
 &\quad - ((T^n - (T^{\sharp_A})^m)(T^n + (T^{\sharp_A})^m)) = 0 \\
 &\Leftrightarrow T^{2n} - T^n (T^{\sharp_A})^m + (T^{\sharp_A})^m T^n - (T^{\sharp_A})^{2m} \\
 &\quad - T^{2n} - T^n (T^{\sharp_A})^m + (T^{\sharp_A})^m T^n - (T^{\sharp_A})^{2m} = 0 \\
 &\Leftrightarrow T^n (T^{\sharp_A})^m - (T^{\sharp_A})^m T^n = 0 \\
 &\Leftrightarrow [T^n, (T^{\sharp_A})^m] = 0.
 \end{aligned}$$

Hence $[X, Y] = 0$ if and only if T is (n, m) - A -normal.

Proofs of the statements (2), (3) and (4) are straightforward. □

Proposition 2.3. *Let $T, V \in \mathcal{B}_A(\mathcal{H})$ be such that $\mathcal{N}(A)^\perp$ is an invariant subspace for both T and V . If T is an (n, m) - A -normal operator and V is an A -isometry, then VTV^{\sharp_A} is an (n, m) - A -normal operator.*

Proof. Since V is an A -isometry then $V^{\sharp_A}V = P_{\overline{\mathcal{R}(A)}}$. Moreover from the fact that $\mathcal{N}(A)^\perp$ is an invariant subspace for T we have $P_{\overline{\mathcal{R}(A)}}T = TP_{\overline{\mathcal{R}(A)}}$ which implies that $T^{\sharp_A}P_{\overline{\mathcal{R}(A)}} = P_{\overline{\mathcal{R}(A)}}T^{\sharp_A}$ since $P_{\overline{\mathcal{R}(A)}}^{\sharp_A} = P_{\overline{\mathcal{R}(A)}}$. In a similar way we have

$$VP_{\overline{\mathcal{R}(A)}} = P_{\overline{\mathcal{R}(A)}}V \quad \text{and} \quad V^{\sharp_A}P_{\overline{\mathcal{R}(A)}} = P_{\overline{\mathcal{R}(A)}}V^{\sharp_A}.$$

It is easy to check that

$$\begin{aligned} (VTV^{\sharp_A})^j &= \underbrace{(VTV^{\sharp_A})(VTV^{\sharp_A}) \dots (VTV^{\sharp_A})}_{j\text{-times}} \\ &= (VTP_{\overline{\mathcal{R}(A)}}TV^{\sharp_A}) \dots (VTV^{\sharp_A}) \\ &= P_{\overline{\mathcal{R}(A)}}VT^2V^{\sharp_A} \dots (VTV^{\sharp_A}) \\ &\quad \vdots \\ &= P_{\overline{\mathcal{R}(A)}}VT^jV^{\sharp_A}. \end{aligned}$$

The same arguments yield

$$\begin{aligned} (VTV^{\sharp_A})^{\sharp_A j} &= \underbrace{(VTV^{\sharp_A})^{\sharp_A} (VTV^{\sharp_A})^{\sharp_A} \dots (VTV^{\sharp_A})^{\sharp_A}}_{j\text{-times}} \\ &= (P_{\overline{\mathcal{R}(A)}}VP_{\overline{\mathcal{R}(A)}}T^{\sharp_A}V^{\sharp_A}) \dots (P_{\overline{\mathcal{R}(A)}}VP_{\overline{\mathcal{R}(A)}}T^{\sharp_A}V^{\sharp_A}) \\ &\quad \vdots \\ &= P_{\overline{\mathcal{R}(A)}}V(T^{\sharp_A})^jV^{\sharp_A}. \end{aligned}$$

From the above calculation, we deduce that

$$\begin{aligned} (2.2) \quad \langle \{(VTV^{\sharp_A})^{\sharp_A}\}^m h \mid \{(VTV^{\sharp_A})^{\sharp_A}\}^n h \rangle_A & \\ &= \langle P_{\overline{\mathcal{R}(A)}}V(T^{\sharp_A})^m V^{\sharp_A} h \mid P_{\overline{\mathcal{R}(A)}}V(T^{\sharp_A})^n V^{\sharp_A} h \rangle_A \\ &= \langle (T^{\sharp_A})^m V^{\sharp_A} h \mid (T^{\sharp_A})^n V^{\sharp_A} h \rangle_A. \end{aligned}$$

It is also easy to show that

$$\begin{aligned} (2.3) \quad \langle (VTV^{\sharp_A})^n h \mid (VTV^{\sharp_A})^m h \rangle_A &= \langle P_{\overline{\mathcal{R}(A)}}VT^n V^{\sharp_A} h \mid P_{\overline{\mathcal{R}(A)}}VT^m V^{\sharp_A} h \rangle_A \\ &= \langle T^n V^{\sharp_A} h \mid T^m V^{\sharp_A} h \rangle_A. \end{aligned}$$

Since T is (n, m) - A -normal, by combining (2.2) and (2.3) we have

$$\langle \{(VTV^{\sharp_A})^{\sharp_A}\}^m h \mid \{(VTV^{\sharp_A})^{\sharp_A}\}^n h \rangle_A = \langle (VTV^{\sharp_A})^n h \mid (VTV^{\sharp_A})^m h \rangle_A \quad \forall h \in \mathcal{H}.$$

On the other hand, we have

$$\begin{aligned} \mathcal{R}((VTV^{\sharp_A})^n \{(VTV^{\sharp_A})^{\sharp_A}\}^m) &= \mathcal{R}(P_{\overline{\mathcal{R}(A)}} V T^n V^{\sharp_A} P_{\overline{\mathcal{R}(A)}} V (T^{\sharp_A})^m V^{\sharp_A}) \\ &= \mathcal{R}(P_{\overline{\mathcal{R}(A)}} V T^n (T^{\sharp_A})^m V^{\sharp_A}) \\ &\subseteq \mathcal{R}(P_{\overline{\mathcal{R}(A)}}) \subseteq \overline{\mathcal{R}(A)}. \end{aligned}$$

In view of Theorem 2.1, it follows that VTV^{\sharp_A} is (n, m) - A -normal operator. \square

Proposition 2.4. *Let $T \in \mathcal{B}_A(\mathcal{H})$ and $S \in \mathcal{B}_A(\mathcal{H})$ be such that $TS = ST$ and $ST^{\sharp_A} = T^{\sharp_A}S$. If T is (n, n) - A -normal, the following statements hold:*

- (1) *If S is an A -self adjoint, then TS is an (n, n) - A -normal operator.*
- (2) *If S is an A -normal operator, then TS is an (n, n) - A -normal operator.*

Proof. (1) Let $h \in \mathcal{H}$, under the assumption that S is A -self-adjoint ($AS = S^*A$) and the statement (1) of Theorem 2.1 we have

$$\begin{aligned} \langle (TS)^{\sharp_{A^n}} h \mid (TS)^{\sharp_{A^n}} h \rangle_A &= \langle (S)^{\sharp_{A^n}} (T)^{\sharp_{A^n}} h \mid (S)^{\sharp_{A^n}} (T)^{\sharp_{A^n}} h \rangle_A \\ &= \langle A(S)^{\sharp_{A^n}} (T)^{\sharp_{A^n}} h \mid (S)^{\sharp_{A^n}} (T)^{\sharp_{A^n}} h \rangle \\ &= \langle (S^*)^n A(T)^{\sharp_{A^n}} h \mid (S)^{\sharp_{A^n}} (T)^{\sharp_{A^n}} h \rangle \\ &= \langle A(S)^n (T)^{\sharp_{A^n}} h \mid (S)^{\sharp_{A^n}} (T)^{\sharp_{A^n}} h \rangle \\ &= \langle A(T)^{\sharp_{A^n}} S^n h \mid (S)^{\sharp_{A^n}} (T)^{\sharp_{A^n}} h \rangle \\ &= \langle (T)^{\sharp_{A^n}} S^n h \mid A(S)^{\sharp_{A^n}} (T)^{\sharp_{A^n}} h \rangle \\ &= \langle (T)^{\sharp_{A^n}} S^n h \mid (T)^{\sharp_{A^n}} S^n h \rangle_A \\ &= \langle T^n S^n h \mid T^n S^n h \rangle_A \\ &= \langle (TS)^n h \mid (TS)^n h \rangle_A. \end{aligned}$$

On the other hand, we have

$$\mathcal{R}((TS)^n (TS)^{\sharp_{A^n}}) = \mathcal{R}(T^n T^{\sharp_{A^n}} S^n S^{\sharp_{A^n}}) \subseteq \overline{\mathcal{R}(A)}.$$

This means that TS is an (n, n) - A -normal operator by Theorem 2.1.

(2) Let S be an A -normal operator then $SS^{\sharp A} = S^{\sharp A}S$ and because T is an (n, n) - A -normal operator we get the relations

$$\begin{aligned}
\langle (ST)^{\sharp A^n} h \mid (ST)^{\sharp A^n} h \rangle_A &= \langle S^{\sharp A^n} T^{\sharp A^n} h \mid S^{\sharp A^n} T^{\sharp A^n} h \rangle_A \\
&= \langle AS^{\sharp A^n} T^{\sharp A^n} h \mid S^{\sharp A^n} T^{\sharp A^n} h \rangle \\
&= \langle S^{*n} AT^{\sharp A^n} h \mid S^{\sharp A^n} T^{\sharp A^n} h \rangle \\
&= \langle T^{\sharp A^n} h \mid S^n S^{\sharp A^n} T^{\sharp A^n} h \rangle_A \\
&= \langle T^{\sharp A^n} h \mid (S^{\sharp A})^n S^n T^{\sharp A^n} h \rangle_A \\
&= \langle S^n T^{\sharp A^n} h \mid S^n T^{\sharp A^n} h \rangle_A \\
&= \langle T^{\sharp A^n} S^n h \mid T^{\sharp A^n} S^n h \rangle_A \\
&= \langle T^n S^n h \mid T^n S^n h \rangle_A \quad (\text{since } T \text{ is } (n, n)\text{-}A\text{-normal}) \\
&= \langle (TS)^n h \mid (TS)^n h \rangle_A.
\end{aligned}$$

On the other hand, based on the (n, n) - A -normality of T we get the inclusion

$$\mathcal{R}((TS)^n (TS)^{\sharp A^n}) = \mathcal{R}(T^n S^n T^{\sharp A^n} S^{\sharp A^n}) \subseteq \mathcal{R}(T^n T^{\sharp A^n}) \subseteq \overline{\mathcal{R}(A)}.$$

From the items (1) and (2) of Theorem 2.1, the operator TS is an (n, n) - A -normal operator. \square

In the following proposition, we study the relation between the classes $[(2, m)\mathbf{N}]_A$ and $[(3, m)\mathbf{N}]_A$.

Proposition 2.5. *Let $T \in \mathcal{B}_A(\mathcal{H})$ be such that $T \in [(2, m)\mathbf{N}]_A \cap [(3, m)\mathbf{N}]_A$ for some positive integer m , then $T \in [(n, m)\mathbf{N}]_A$ for all positive integers $n \geq 4$.*

Proof. It is obvious from Definition 2.1 that if $T \in [(2, m)\mathbf{N}]_A$ then $T \in [(4, m)\mathbf{N}]_A$. However, $T \in [(2, m)\mathbf{N}]_A \cap [(3, m)\mathbf{N}]_A$ implies that $T \in [(5, m)\mathbf{N}]_A$.

Assume that $T \in [(n, m)\mathbf{N}]_A$ for $n \geq 5$, that is,

$$T^n (T^{\sharp A})^m = (T^{\sharp A})^m T^n.$$

Then we have

$$\begin{aligned}
[T^{n+1}, (T^{\sharp A})^m] &= T^{n+1} (T^{\sharp A})^m - (T^{\sharp A})^m T^{n+1} \\
&= T (T^{\sharp A})^m T^n - (T^{\sharp A})^m T^{n+1} \\
&= T (T^{\sharp A})^m T^2 T^{n-2} - (T^{\sharp A})^m T^{n+1} \\
&= T^3 (T^{\sharp A})^m T^{n-2} - (T^{\sharp A})^m T^{n+1} \\
&= (T^{\sharp A})^m T^{n+1} - (T^{\sharp A})^m T^{n+1} = 0.
\end{aligned}$$

This means that $T \in [(n+1, m)\mathbf{N}]_A$. The proof is complete. \square

Proposition 2.6. *Let $T \in \mathcal{B}_A(\mathcal{H})$. If $T \in [(n, m)\mathbf{N}]_A \cap [(n + 1, m)\mathbf{N}]_A$, then $T \in [(n + 2, m)\mathbf{N}]_A$ for some positive integers n and m . In particular $T \in [(j, m)\mathbf{N}]_A$ for all $j \geq n$.*

Proof. Let $T \in [(n, m)\mathbf{N}]_A \cap [(n + 1, m)\mathbf{N}]_A$, then it follows that

$$T^n(T^{\sharp_A})^m - (T^{\sharp_A})^m T^n = 0 \quad \text{and} \quad T^{n+1}(T^{\sharp_A})^m - (T^{\sharp_A})^m T^{n+1} = 0.$$

Note that

$$\begin{aligned} [T^{n+2}, (T^{\sharp_A})^m] &= T^{n+2}(T^{\sharp_A})^m - (T^{\sharp_A})^m T^{n+2} \\ &= T T^{n+1}(T^{\sharp_A})^m - (T^{\sharp_A})^m T^{n+2} \\ &= T(T^{\sharp_A})^m T^{n+1} - (T^{\sharp_A})^m T^{n+2} \\ &= T T^n (T^{\sharp_A})^m T - (T^{\sharp_A})^m T^{n+2} \\ &= (T^{\sharp_A})^m T^{n+2} - (T^{\sharp_A})^m T^{n+2} = 0. \end{aligned}$$

Hence $T \in [(n+2, m)\mathbf{N}]_A$. By repeating this process we can prove that $T \in [(j, m)\mathbf{N}]_A$ for all $j \geq n$. \square

Proposition 2.7. *Let $T \in \mathcal{B}_A(\mathcal{H})$. If $T \in [(n, m)\mathbf{N}]_A \cap [(n + 1, m)\mathbf{N}]_A$ is one-to-one, then $T \in [(1, m)\mathbf{N}]_A$.*

Proof. Let $T \in [(n, m)\mathbf{N}]_A \cap [(n + 1, m)\mathbf{N}]_A$, then it follows that,

$$T^n(T(T^{\sharp_A})^m - (T^{\sharp_A})^m T) = 0.$$

Since T is one-to-one, then so is T^n and it follows that $T(T^{\sharp_A})^m - (T^{\sharp_A})^m T = 0$. Therefore $T \in [(1, m)\mathbf{N}]_A$. \square

Proposition 2.8. *Let $T \in \mathcal{B}_A(\mathcal{H})$. The following statements are equivalent.*

- (1) *If $T \in [(n, 2)\mathbf{N}]_A \cap [(n, 3)\mathbf{N}]_A$ for some positive integer n , then $T \in [(n, m)\mathbf{N}]_A$ for all positive integers $m \geq 4$.*
- (2) *If $T \in [(n, m)\mathbf{N}]_A \cap [(n, m + 1)\mathbf{N}]_A$, then $T \in [(n, m + 2)\mathbf{N}]_A$ for some positive integers n, m . In particular $T \in [(n, j)\mathbf{N}]_A$ for all $j \geq m$.*

Proof. The proof follows by applying Proposition 2.1 and Proposition 2.5. \square

Proposition 2.9. *Let $T \in \mathcal{B}_A(\mathcal{H})$. If $T \in [(n, m)\mathbf{N}]_A \cap [(n, m + 1)\mathbf{N}]_A$ is such that T^{\sharp_A} is one-to one, then $T \in [(n, 1)\mathbf{N}]_A = [n\mathbf{N}]_A$.*

P r o o f. Since $T \in [(n, m)\mathbf{N}]_A \cap [(n, m + 1)\mathbf{N}]_A$, it follows that

$$(T^{\sharp_A})^m(T^n T^{\sharp_A} - T^{\sharp_A} T^n) = 0.$$

If T^{\sharp_A} is one-to-one, then so is $(T^{\sharp_A})^m$ and we obtain $T^n T^{\sharp_A} - T^{\sharp_A} T^n = 0$. Consequently $T \in [(n, 1)\mathbf{N}]_A$. \square

In [19], Theorem 2.4 it was proved that if T is (n, m) -power normal such that T^m is a partial isometry, then T is $(n + m, m)$ -power normal. In the following theorem we extend this result to (n, m) - A -normal operators.

Theorem 2.3. *Let $T \in \mathcal{B}_A(\mathcal{H})$ be (n, m) - A -normal for some positive integers n and m . The following statements hold:*

- (1) *If $n \geq m$ and $T^m(T^{\sharp_A})^m T^m = T^m$, then $T \in [(n + m, m)\mathbf{N}]_A$.*
- (2) *If $m \geq n$ and $(T^{\sharp_A})^n T^n (T^{\sharp_A})^n = (T^{\sharp_A})^n$, then $T \in [(n, m + n)\mathbf{N}]_A$.*

P r o o f. (1) Under the assumption that $T^m(T^{\sharp_A})^m T^m = T^m$, it follows that

$$T^m(T^{\sharp_A})^m T^n = T^n \quad \text{and} \quad T^n(T^{\sharp_A})^m T^m = T^n \quad \text{for } n \geq m,$$

which means that $T^n(T^{\sharp_A})^m T^m = T^m(T^{\sharp_A})^m T^n$. Since T is (n, m) - A normal, we get

$$(T^{\sharp})^m T^{n+m} = T^{n+m} (T^{\sharp_A})^m.$$

So, $T \in [(m + n, m)\mathbf{N}]_A$.

(2) In same way, under the assumption $(T^{\sharp_A})^n T^n (T^{\sharp_A})^n = (T^{\sharp_A})^n$, it follows that

$$(T^{\sharp_A})^n T^n (T^{\sharp_A})^m = (T^{\sharp_A})^m \quad \text{and} \quad (T^{\sharp_A})^m T^n (T^{\sharp_A})^n = (T^{\sharp_A})^m \quad \text{for } m \geq n,$$

which means that $(T^{\sharp_A})^n T^n (T^{\sharp_A})^m = (T^{\sharp_A})^m T^n (T^{\sharp_A})^n$. Since T is (n, m) - A normal, we get

$$(T^{\sharp})^{m+n} T^n = T^n (T^{\sharp_A})^{n+m}.$$

So, $T \in [(n, m + n)\mathbf{N}]_A$ and the proof is complete. \square

Proposition 2.10. *Let $T \in \mathcal{B}_A(\mathcal{H})$ be an (n, m) - A -normal operator for some positive integers n and m . Then T satisfies the relation $T^{2n}(T^{\sharp_A})^{2m} = (T^n(T^{\sharp_A})^m)^2$.*

P r o o f. Since T is an (n, m) - A -normal operator, it follows that

$$T^{2n}(T^{\sharp_A})^{2m} = T^n T^n (T^{\sharp_A})^m (T^{\sharp_A})^m = \underbrace{T^n (T^{\sharp_A})^m}_{\text{}} \underbrace{T^n (T^{\sharp_A})^m}_{\text{}} = (T^n (T^{\sharp_A})^m)^2.$$

\square

The idea of the following proposition is inspired by [20].

Proposition 2.11. *Let $T \in \mathcal{B}_A(\mathcal{H})$ be such that $AT = TA$. If T is an n -normal operator, then T is an (n, m) - A -normal operator for $m \in \mathbb{N}$.*

Proof. Indeed, since T^n is normal and $T^m T^n = T^n T^m$, it follows from the Fuglede theorem (see [14]) that $T^{*m} T^n = T^n T^{*m}$. Taking in consideration that under the assumptions we have $P_{\mathcal{R}(A)} T = T P_{\mathcal{R}(A)}$ and $T^{\sharp A} = P_{\mathcal{R}(A)} T^*$. Then

$$\begin{aligned} [T^n, (T^{\sharp A})^m] &= T^n (T^{\sharp A})^m - (T^{\sharp A})^m T^n \\ &= T^n (P_{\mathcal{R}(A)} T^*)^m - (P_{\mathcal{R}(A)} T^*)^m T^n \\ &= P_{\mathcal{R}(A)} [T^n, T^{*m}] = 0. \end{aligned}$$

Therefore T is (n, m) - A -normal. \square

Corollary 2.1. *Let $T \in \mathcal{B}_A(\mathcal{H})$ be such that $AT = TA$. If T is an (n, m) -normal operator, then T is a (j, r) - A -normal operator where $r \in \mathbb{N}$ and j is the least common multiple of n and m .*

Proof. Since T is (n, m) -normal, it was observed in [11], Lemma 4.4 that T^j is a normal operator where $j = LCM(n, m)$. By applying Proposition 2.11 we get that T is a (j, r) - A -normal operator. \square

3. (n, m) - A -QUASINORMAL OPERATORS

In [8] the author has introduced the class of (n, m) - A -quasinormal operators as follows. An operator $T \in \mathcal{B}_A(\mathcal{H})$ is said to be (n, m) - A -quasinormal if T satisfies

$$[T^n, (T^{\sharp A})^m T] := T^n (T^{\sharp A})^m T - (T^{\sharp A})^m T T^n = 0$$

for some positive integers n and m . This class of operators is denoted by $[(n, m)\mathbf{QN}]_A$.

Remark 3.1. Clearly, the class of (n, m) - A -quasinormal operators includes the class of (n, m) - A -normal one, i.e. the following inclusion holds

$$[(n, m)\mathbf{N}]_A \subset [(n, m)\mathbf{QN}]_A.$$

We give an example to show that there exists an (n, m) - A -quasinormal operator which is not (n, m) - A -normal for some positive integers n and m .

Example 3.1. Let T be a unilateral shift, that is, if $\mathcal{H} = l^2$, the matrix

$$T = \begin{pmatrix} 0 & 0 & 0 & \dots \\ 1 & 0 & 0 & \dots \\ 0 & 1 & 0 & \dots \\ \dots & \dots & \dots & \dots \end{pmatrix} \quad \text{and} \quad A = I_{l^2} \text{ (the identity operator).}$$

It is easily verified that $[T^2, T^{\sharp A}] \neq 0$ and $[T^2, T^{\sharp A}T] = 0$. So that T is not a $(2, 1)$ - A -normal operator but it is a $(2, 1)$ - A -quasinormal operator.

The following theorem gives a characterization of (n, m) - A -quasinormal operators.

Theorem 3.1. *Let $T \in \mathcal{B}_A(\mathcal{H})$. Then T is an (n, m) - A -quasinormal operator for some positive integers n and m if and only if T satisfies the following conditions:*

- (1) $\langle (T^{\sharp A})^m T h \mid (T^{\sharp A})^n h \rangle_A = \langle T^n T h \mid T^m h \rangle_A \quad \forall h \in \mathcal{H},$
- (2) $\mathcal{R}(T^n (T^{\sharp A})^m T) \subseteq \overline{\mathcal{R}(A)}.$

Proof. We omit the proof, since the techniques are similar to those of Theorem 2.1. □

Remark 3.2. Theorem 3.1 is an improved version of [8], Lemma 4.4.

Proposition 3.1. *Let $T \in \mathcal{B}_A(\mathcal{H})$ and $S \in \mathcal{B}_A(\mathcal{H})$ be (n, m) - A -normal operators. Then their product ST is an (n, m) - A -normal operator if the conditions $ST = TS$, $ST^{\sharp A} = T^{\sharp A}S$ and $TS^{\sharp A} = S^{\sharp A}T$ are satisfied.*

Proof. It is

$$\begin{aligned} (TS)^n ((TS)^{\sharp A})^m (TS) &= T^n S^n (T^{\sharp A})^m (S^{\sharp A})^m TS = T^n (T^{\sharp A})^m T S^n (S^{\sharp A})^m S \\ &= (T^{\sharp A})^m T T^n (S^{\sharp A})^m S S^n = ((TS)^{\sharp A})^m (TS) (TS)^n. \end{aligned}$$

Therefore TS is an (n, m) - A -quasinormal operator. □

Remark 3.3. Proposition 3.1 is an improved version of [8], Proposition 4.5.

Proposition 3.2. *Let $T \in \mathcal{B}_A(\mathcal{H})$. If $T \in [(n, m)\mathbf{QN}]_A \cap [(n + 1, m)\mathbf{QN}]_A$, then $T \in [(n + 2, m)\mathbf{QN}]_A$.*

Proof. Assume that $T \in [(n, m)\mathbf{QN}]_A \cap [(n+1, m)\mathbf{QN}]_A$, it follows that

$$T^{n+1}(T^{\sharp_A})^m T - (T^{\sharp_A})^m T T^{n+1} = 0 \quad \text{and} \quad T^n (T^{\sharp_A})^m T - (T^{\sharp_A})^m T T^n = 0.$$

On the other hand, we have

$$\begin{aligned} T^{n+2}(T^{\sharp_A})^m T - (T^{\sharp_A})^m T T^{n+2} &= T(T^{\sharp_A})^m T T^{n+1} - (T^{\sharp_A})^m T T^{n+2} \\ &= T^{n+1}(T^{\sharp_A})^m T T - (T^{\sharp_A})^m T T^{n+2} \\ &= (T^{\sharp_A})^m T T^{n+2} - (T^{\sharp_A})^m T T^{n+2} = 0. \end{aligned}$$

□

In [19] it was proved that if $T \in [(n, m)\mathbf{QN}]$ such that T^m is a partial isometry, then $T \in [(n+m, m)\mathbf{QN}]$ for $n \geq m$. We extend this result to the class of $[(n, m)\mathbf{QN}]_A$ as follows.

Theorem 3.2. *Let $T \in \mathcal{B}_A(\mathcal{H})$ be such that $T \in [(n, m)\mathbf{QN}]$ for some positive integers n and m . If $T^m (T^{\sharp_A})^m T^m = T^m$ for $n \geq m$, then $T \in [(n+m, m)\mathbf{QN}]_A$.*

Proof. (1) Assume that T^m satisfies $T^m (T^{\sharp_A})^m T^m = T^m$ for $m \geq n$, then we have

$$(3.1) \quad T^m (T^{\sharp_A})^m T T^{m-1} = T^m.$$

Multiplying (3.1) from the left by T^{n-m} and from the right by T we get

$$(3.2) \quad T^n ((T^{\sharp_A})^m T) T^m = T^{n+1}.$$

Multiplying (3.1) from the right by T^{n-m+1} we get

$$(3.3) \quad T^m ((T^{\sharp_A})^m T) T^n = T^{n+1}.$$

Combining (3.2), (3.3) and using the fact that $T \in [(n, m)\mathbf{QN}]$ we obtain

$$T^{n+m} ((T^{\sharp_A})^m T) = ((T^{\sharp_A})^m T) T^{n+m}.$$

Therefore $T \in [(n+m, m)\mathbf{QN}]_A$ as required. □

Proposition 3.3. *Let $T \in \mathcal{B}_A(\mathcal{H})$, n and m positive integers. The following statements hold:*

- (1) *If $T \in [(n, m)\mathbf{QN}]_A \cap [(n+1, m)\mathbf{QN}]_A$ such that T is one-to-one, then $T \in [(1, m)\mathbf{QN}]_A$.*
- (2) *If $T \in [(n, m)\mathbf{QN}]_A \cap [(n, m+1)\mathbf{QN}]_A$ such that T^* is one-to-one and $\mathcal{R}(T^{\sharp_A})^m T = \mathcal{R}(A)$, then $T \in [(n, 1)\mathbf{N}]_A$.*

Proof. (1) Under the assumption $T \in [(n, m)\mathbf{QN}]_A \cap [(n+1, m)\mathbf{QN}]_A$, it follows that

$$T^n(T(T^{\sharp_A})^m T - (T^{\sharp_A})^m T T) = 0.$$

If T is injective, then so is T^n and we have $T(T^{\sharp_A})^m T - (T^{\sharp_A})^m T T = 0$. Hence, $T \in [(1, m)\mathbf{QN}]_A$.

(2) Since $T \in [(n, m)\mathbf{QN}]_A \cap [(n, m+1)\mathbf{QN}]_A$, we have

$$\begin{aligned} T^n(T^{\sharp_A})^{m+1} T - (T^{\sharp_A})^{m+1} T T^n &= 0 \\ \Rightarrow T^n T^{\sharp_A} (T^{\sharp_A})^m T - T^{\sharp_A} (T^{\sharp_A})^m T T^n &= 0 \\ \Rightarrow (T^n T^{\sharp_A} - T^{\sharp_A} T^n) (T^{\sharp_A})^m T &= 0 \\ \Rightarrow (T^n T^{\sharp_A} - T^{\sharp_A} T^n) \equiv 0 \quad \text{on } \overline{\mathcal{R}((T^{\sharp_A})^m T)} &= \overline{\mathcal{R}(A)}. \end{aligned}$$

On the other hand, since $T \in \mathcal{B}_A(\mathcal{H})$, we have $T(\mathcal{N}(A)) \subseteq \mathcal{N}(A)$. Moreover, by the assumption that T^* is injective we obtain $\mathcal{N}(T^{\sharp_A}) = \mathcal{N}(A)$. If $h \in \mathcal{N}(A)$ it follows from the above observation that

$$(T^n T^{\sharp_A} - T^{\sharp_A} T^n)h = T^n T^{\sharp_A} h - T^{\sharp_A} T^n h = 0.$$

Consequently, $(T^n T^{\sharp_A} - T^{\sharp_A} T^n) = 0$ on \mathcal{H} . Therefore $T \in [(n, 1)\mathbf{N}]_A$. \square

Proposition 3.4. *Let $T \in \mathcal{B}_A(\mathcal{H})$ be such that $T \in [(2, m)\mathbf{QN}]_A \cap [(3, m)\mathbf{QN}]_A$ for some positive integer m , then $T \in [(n, m)\mathbf{QN}]_A$ for all positive integers $n \geq 4$.*

Proof. We prove the assertion by using the mathematical induction. Since $T \in [(2, m)\mathbf{QN}]_A \cap [(3, m)\mathbf{QN}]_A$, it follows immediately that

$$T^4(T^{\sharp_A})^m T - (T^{\sharp_A})^m T T^4 = 0 \quad \text{and} \quad T^5(T^{\sharp_A})^m T - (T^{\sharp_A})^m T T^5 = 0.$$

Now assume that the result is true for $n \geq 5$, that is,

$$T^n(T^{\sharp_A})^m T - (T^{\sharp_A})^m T T^n = 0,$$

then

$$\begin{aligned} T^{n+1}(T^{\sharp_A})^m T - (T^{\sharp_A})^m T T^{n+1} &= T(T^{\sharp_A})^m T T^n - (T^{\sharp_A})^m T T^{n+1} \\ &= T^3(T^{\sharp_A})^m T T^{n-2} - (T^{\sharp_A})^m T T^{n+1} \\ &= (T^{\sharp_A})^m T T^{n+1} - (T^{\sharp_A})^m T T^{n+1} = 0. \end{aligned}$$

Therefore $T \in [(n+1, m)\mathbf{QN}]_A$. The proof is complete. \square

Now we discuss the (n, m) - A -quasinormality of an operator under some commutation conditions on its real and imaginary part.

Theorem 3.3. *Let $T \in \mathcal{B}_A(\mathcal{H})$ be such that $\mathcal{R}(T^{m-1})$ is dense. If $TA = AT$. Then the following statements are equivalent.*

- (1) $T \in [(n, m)\mathbf{QN}]_A$.
- (2) $C_{m,A}$ commutes with $\operatorname{Re}_A(T^n)$ and $\operatorname{Im}_A(T^n)$, where $C_{m,A} = \sqrt{(T^{\sharp_A})^m T^m}$.

Proof. Since T is (n, m) - A -quasinormal, it follows that

$$T^n (T^{\sharp_A})^m T = (T^{\sharp_A})^m T T^n.$$

Hence,

$$T^n (T^{\sharp_A})^m T^m = (T^{\sharp_A})^m T^m T^n.$$

From the conditions that $TA = AT$ and $\mathcal{N}(A)^\perp$ is an invariant subspace for T , we observe that

$$T P_{\overline{\mathcal{R}(A)}} = T P_{\overline{\mathcal{R}(A)}}, \quad T^{\sharp_A} P_{\overline{\mathcal{R}(A)}} = T^{\sharp_A} P_{\overline{\mathcal{R}(A)}} \quad \text{and} \quad T^{\sharp_A} = P_{\overline{\mathcal{R}(A)}} T^*.$$

Therefore, $C_{m,A}$ is a nonnegative definite operator and by elementary calculation we get

$$C_{m,A}^2 \operatorname{Re}_A(T^n) = \operatorname{Re}_A(T^n) C_{m,A}^2.$$

Consequently,

$$C_{m,A} \operatorname{Re}_A(T^n) = \operatorname{Re}_A(T^n) C_{m,A}.$$

In a similar way we can prove that $C_{m,A} \operatorname{Im}_A(T^n) = \operatorname{Im}_A(T^n) C_{m,A}$. Conversely, assume that $C_{m,A} \operatorname{Re}_A(T^n) = \operatorname{Re}_A(T^n) C_{m,A}$ and $C_{m,A} \operatorname{Im}_A(T^n) = \operatorname{Im}_A(T^n) C_{m,A}$. Then

$$C_{m,A}^2 \operatorname{Re}_A(T^n) = \operatorname{Re}_A(T^n) C_{m,A}^2 \quad \text{and} \quad C_{m,A}^2 \operatorname{Im}_A(T^n) = \operatorname{Im}_A(T^n) C_{m,A}^2.$$

Hence

$$C_{m,A}^2 (\operatorname{Re}_A(T^n) + i \operatorname{Im}_A(T^n)) = (\operatorname{Re}_A(T^n) + i \operatorname{Im}_A(T^n)) C_{m,A}^2,$$

and therefore

$$C_{m,A}^2 T^n = T^n C_{m,A}^2.$$

On the other hand, we have

$$\begin{aligned} C_{m,A}^2 T^n = T^n C_{m,A}^2 &\Leftrightarrow (T^{\sharp_A})^m T^m T^n - T^n (T^{\sharp_A})^m T^m = 0 \\ &\Leftrightarrow ((T^{\sharp_A})^m T T^n - T^n (T^{\sharp_A})^m T) T^{m-1} = 0 \\ &\Leftrightarrow (T^{\sharp_A})^m T T^n - T^n (T^{\sharp_A})^m T = 0 \quad (\overline{\mathcal{R}(T^{m-1})} = \mathcal{H}). \end{aligned}$$

Therefore $T \in [(n, m)\mathbf{QN}]_A$. □

Theorem 3.4. Let $T \in \mathcal{B}_A(\mathcal{H})$ be such that $\mathcal{R}(T^{m-1})$ is dense and $TA = AT$.

If T satisfies the conditions

- (i) $B_{m,A}$ commutes with $\text{Re}_A(T^m)$ and $\text{Im}_A(T^m)$,
- (ii) $C_{m,A}^2 T^n = T^n B_{m,A}^2$, where $B_{m,A} = \sqrt{T^m (T^{\sharp_A})^m}$.

Then T is an (m, m) - A -quasinormal operator.

Proof. Since

$$B_{m,A} \text{Re}_A(T^m) = \text{Re}_A(T^m) B_{m,A} \quad \text{and} \quad B_{m,A} \text{Im}_A(T^m) = \text{Im}_A(T^m) B_{m,A},$$

it follows that

$$\begin{cases} B_{m,A}^2 T^m + B^2(T^m)^{\sharp_A} = T^m B_{m,A}^2 + (T^m)^{\sharp_A} B_{m,A}^2, \\ B_{m,A}^2 T^m - B_{m,A}^2 (T^m)^{\sharp_A} = T^m B_{m,A}^2 - (T^m)^{\sharp_A} B_{m,A}^2. \end{cases}$$

This gives

$$B_{m,A}^2 T^m = T^m B_{m,A}^2 = C_{m,A}^2 T^m.$$

On the other hand, we have

$$\begin{aligned} B_{m,A}^2 T^m = C_{m,A}^2 T^m &\Rightarrow T^m (T^{\sharp_A})^m T^m - (T^{\sharp_A})^m T^m T^m = 0 \\ &\Rightarrow (T^m (T^{\sharp_A})^m T - (T^{\sharp_A})^m T T^m) T^{m-1} = 0 \\ &\Rightarrow T^m (T^{\sharp_A})^m T - (T^{\sharp_A})^m T T^m = 0 \quad \text{on } \overline{\mathcal{R}(T^{m-1})} = \mathcal{H}. \end{aligned}$$

Therefore $T^m (T^{\sharp_A})^m T - (T^{\sharp_A})^m T T^m = 0$ and T is an (m, m) - A -quasinormal operator. \square

Proposition 3.5. Let $T \in \mathcal{B}_A(\mathcal{H})$ be (n, m) - A -quasinormal, then

$$(T^{\sharp_A})^{2m} T^{2n} = ((T^{\sharp_A})^m T^n)^2.$$

Proof. Since T is (n, m) - A -quasinormal, it follows that

$$T^n (T^{\sharp_A})^m T = (T^{\sharp_A})^m T T^n.$$

On the other hand, we have

$$\begin{aligned} (T^{\sharp_A})^{2m} T^{2n} &= (T^{\sharp_A})^m (T^{\sharp_A})^m T^n T^n = (T^{\sharp_A})^m (T^{\sharp_A})^m T T^n T^{n-1} \\ &= (T^{\sharp_A})^m T^n (T^{\sharp_A})^m T^n = ((T^{\sharp_A})^m T^n)^2. \end{aligned}$$

\square

References

- [1] *E. H. Abood, M. A. Al-loz*: On some generalization of normal operators on Hilbert space. *Iraqi J. Sci.* *56* (2015), 1786–1794.
- [2] *E. H. Abood, M. A. Al-loz*: On some generalizations of (n, m) -normal powers operators on Hilbert space. *J. Progressive Res. Math. (JPRM)* *7* (2016), 1063–1070.
- [3] *S. A. Alzuraiqi, A. B. Patel*: On n -normal operators. *Gen. Math. Notes* *1* (2010), 61–73. [zbl](#)
- [4] *M. L. Arias, G. Corach, M. C. Gonzalez*: Metric properties of projections in semi-Hilbertian spaces. *Integral Equations Oper. Theory* *62* (2008), 11–28. [zbl](#) [MR](#) [doi](#)
- [5] *M. L. Arias, G. Corach, M. C. Gonzalez*: Partial isometries in semi-Hilbertian spaces. *Linear Algebra Appl.* *428* (2008), 1460–1475. [zbl](#) [MR](#) [doi](#)
- [6] *M. L. Arias, G. Corach, M. C. Gonzalez*: Lifting properties in operator ranges. *Acta Sci. Math.* *75* (2009), 635–653. [zbl](#) [MR](#)
- [7] *H. Baklouti, K. Feki, O. A. M. Sid Ahmed*: Joint normality of operators in semi-Hilbertian spaces. *Linear Multilinear Algebra* *68* (2020), 845–866. [zbl](#) [MR](#) [doi](#)
- [8] *V. Bavithra*: (n, m) -power quasi normal operators in semi-Hilbertian spaces. *J. Math. Informatics* *11* (2017), 125–129.
- [9] *A. Benali, O. A. M. Sid Ahmed*: (α, β) - A -normal operators in semi-Hilbertian spaces. *Afr. Mat.* *30* (2019), 903–920. [zbl](#) [MR](#) [doi](#)
- [10] *C. Chellali, A. Benali*: Class of (A, n) -power-hyponormal operators in semi-Hilbertian space. *Func. Anal. Approx. Comput.* *11* (2019), 13–21. [zbl](#) [MR](#)
- [11] *M. Chō, J. E. Lee, K. Tanahashic, A. Uchiyamad*: Remarks on n -normal operators. *Filomat* *32* (2018), 5441–5451. [MR](#) [doi](#)
- [12] *M. Chō, B. Načevska*: Spectral properties of n -normal operators. *Filomat* *32* (2018), 5063–5069. [MR](#) [doi](#)
- [13] *R. G. Douglas*: On majorization, factorization, and range inclusion of operators in Hilbert space. *Proc. Am. Math. Soc.* *17* (1966), 413–415. [zbl](#) [MR](#) [doi](#)
- [14] *S. H. Jah*: Class of (A, n) -power quasi-normal operators in semi-Hilbertian spaces. *Int. J. Pure Appl. Math.* *93* (2014), 61–83. [zbl](#) [doi](#)
- [15] *A. A. S. Jibril*: On n -power normal operators. *Arab. J. Sci. Eng., Sect. A, Sci.* *33* (2008), 247–251. [zbl](#) [MR](#)
- [16] *J. S. I. Mary, P. Vijaylakshmi*: Fuglede-Putnam theorem and quasi-nilpotent part of n -normal operators. *Tamkang J. Math.* *46* (151–165). [zbl](#) [MR](#) [doi](#)
- [17] *A. Saddi*: A -normal operators in semi-Hilbertian spaces. *Aust. J. Math. Anal. Appl.* *9* (2012), Article ID 5, 12 pages. [zbl](#) [MR](#)
- [18] *O. A. M. Sid Ahmed, A. Benali*: Hyponormal and k -quasi-hyponormal operators on semi-Hilbertian spaces. *Aust. J. Math. Anal. Appl.* *13* (2016), Article ID 7, 22 pages. [zbl](#) [MR](#)
- [19] *O. A. M. Sid Ahmed, O. B. Sid Ahmed*: On the classes (n, m) -power D -normal and (n, m) -power D -quasi-normal operators. *Oper. Matrices* *13* (2019), 705–732. [zbl](#) [MR](#) [doi](#)
- [20] *O. B. Sid Ahmed, O. A. M. Sid Ahmed*: On the class of n -power D - m -quasi-normal operators on Hilbert spaces. *Oper. Matrices* *14* (2020), 159–174. [zbl](#) [MR](#) [doi](#)
- [21] *L. Suci*: Orthogonal decompositions induced by generalized contractions. *Acta Sci. Math.* *70* (2004), 751–765. [zbl](#) [MR](#)

Authors' address: Samir Al Mohammady, Sid Ahmed Ould Beinane, Sid Ahmed Ould Ahmed Mahmoud, Jouf University, Mathematics Department, College of Science, P.O.Box 2014, Sakaka, Saudi Arabia, e-mail: senssar@ju.edu.sa; beinane06@gmail.com, sabeinane@ju.edu.sa; sidahmed@ju.edu.sa, sidahmed.sidha@gmail.com.