

Insights into the chemical diversity of selected fungi from the Tza Itzá cenote of the Yucatan peninsula

By: Carlos A. Fajardo-Hernández, [Firoz Shah Tuglak Khan](#), Laura Flores-Bocanegra, Alejandra Prieto-Davó, Baojie Wan, Rui Ma, Mallique Qader, Rodrigo Villanueva-Silva, Anahí Martínez-Cárdenas, Marian A. López-Lobato, [Shabnam Hematian](#), Scott G. Franzblau, [Huzefa A. Raja](#), Rodolfo García-Contreras, and [Mario Figueroa](#)

Carlos A. Fajardo-Hernández, Firoz Shah Tuglak Khan, Laura Flores-Bocanegra, Alejandra Prieto-Davó, Baojie Wan, Rui Ma, Mallique Qader, Rodrigo Villanueva-Silva, Anahí Martínez-Cárdenas, Marian A. López-Lobato, Shabnam Hematian, Scott G. Franzblau, Huzefa A. Raja, Rodolfo García-Contreras, and Mario Figueroa. Insights into the Chemical Diversity of Selected Fungi from the Tza Itzá Cenote of the Yucatan Peninsula. *ACS Omega* 2022, 7, 14, 12171–12185. <https://doi.org/10.1021/acsomega.2c00544>



© 2022 The Authors. This work is licensed under a [Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License](#).

Abstract:

Cenotes are habitats with unique physical, chemical, and biological features. Unexplored microorganisms from these sinkholes represent a potential source of bioactive molecules. Thus, a series of cultivable fungi (*Aspergillus* spp. NCA257, NCA264, and NCA276, *Stachybotrys* sp. NCA252, and *Cladosporium* sp. NCA273) isolated from the cenote Tza Itzá were subjected to chemical, coculture, and metabolomic analyses. Nineteen compounds were obtained and tested for their antimicrobial potential against ESKAPE pathogens, *Mycobacterium tuberculosis*, and nontuberculous mycobacteria. In particular, phenylspirodrimanes from *Stachybotrys* sp. NCA252 showed significant activity against MRSA, MSSA, and mycobacterial strains. On the other hand, the absolute configuration of the new compound 17-deoxy-aspergillin PZ (**1**) isolated from *Aspergillus* sp. NCA276 was established via single-crystal X-ray crystallography. Also, the chemical analysis of the cocultures between *Aspergillus* and *Cladosporium* strains revealed the production of metabolites that were not present or were barely detected in the monocultures. Finally, molecular networking analysis of the LC-MS-MS/MS data for each fungus was used as a tool for the annotation of additional compounds, increasing the chemical knowledge on the corresponding fungal strains. Overall, this is the first systematic chemical study on fungi isolated from a sinkhole in Mexico.

Keywords: fungi | Tza Itzá cenote | cenote diversity

Article:

***Note: Full text of article below

Insights into the Chemical Diversity of Selected Fungi from the Tza Itzá Cenote of the Yucatan Peninsula

Carlos A. Fajardo-Hernández, Firoz Shah Tuglak Khan, Laura Flores-Bocanegra, Alejandra Prieto-Davó, Baojie Wan, Rui Ma, Mallique Qader, Rodrigo Villanueva-Silva, Anahí Martínez-Cárdenas, Marian A. López-Lobato, Shabnam Hematian, Scott G. Franzblau, Huzefa A. Raja, Rodolfo García-Contreras, and Mario Figueroa*



Cite This: *ACS Omega* 2022, 7, 12171–12185



Read Online

ACCESS |



Metrics & More




Article Recommendations



Supporting Information

ABSTRACT: Cenotes are habitats with unique physical, chemical, and biological features. Unexplored microorganisms from these sinkholes represent a potential source of bioactive molecules. Thus, a series of cultivable fungi (*Aspergillus* spp. NCA257, NCA264, and NCA276, *Stachybotrys* sp. NCA252, and *Cladosporium* sp. NCA273) isolated from the cenote Tza Itzá were subjected to chemical, coculture, and metabolomic analyses. Nineteen compounds were obtained and tested for their antimicrobial potential against ESKAPE pathogens, *Mycobacterium tuberculosis*, and nontuberculous mycobacteria. In particular, phenylspirodrimanones from *Stachybotrys* sp. NCA252 showed significant activity against MRSA, MSSA, and mycobacterial strains. On the other hand, the absolute configuration of the new compound 17-deoxy-aspergillin PZ (**1**) isolated from *Aspergillus* sp. NCA276 was established via single-crystal X-ray crystallography. Also, the chemical analysis of the cocultures between *Aspergillus* and *Cladosporium* strains revealed the production of metabolites that were not present or were barely detected in the monocultures. Finally, molecular networking analysis of the LC-MS-MS/MS data for each fungus was used as a tool for the annotation of additional compounds, increasing the chemical knowledge on the corresponding fungal strains. Overall, this is the first systematic chemical study on fungi isolated from a sinkhole in Mexico.



INTRODUCTION

One of the most famous geological features in the Yucatan peninsula is the ring of cenotes (from the Maya word *ts'otnot*), which is a group of sinkholes formed by the process of karstification.¹ The origin of cenotes is related to the impact of the Chicxulub asteroid around 66 millions of years ago, which caused changes in the karst features.^{2,3} These habitats are physically and chemically unique due to the phototrophic activity of its microbial communities. In particular, the carbon and sulfur cycles completed by different microbes affect the sulfate reduction into aqueous sulfide, accelerating the limestone dissolution, and increasing alkalinity.^{4,5} In addition, most of the cenotes have hydraulic connections or networks across the region, and the microbial communities, especially freshwater fungi and bacteria, play a key role in keeping the balance of the entire ecosystem.⁶

Metagenomic analysis of soil samples from freshwater environments in the Yucatan peninsula revealed a remarkable microbial diversity, where fungi (ascomycetes) prevail over bacteria (actinobacteria) in terms of biomass production and enzymatic substrate degradation.⁷ Furthermore, the biosynthetic potential of these communities of microbes was assessed using bioinformatics tools coupled with direct amplification of environmental DNA.^{8–10} The number of studies demonstrat-

ing the potential of microorganisms from cenotes for the discovery of bioactive natural products is scarce. Such studies have focused on the biological activity of extracts, exhibiting a high percentage of hits (~81%). About half of the extracts showed varied activities, including insecticidal and nematotoxic.^{11–13} To date, the only formal chemical study reported from a fungal strain isolated from plant litter submerged in a sinkhole in Merida, Yucatan, led to the discovery of the novel hexahydroacremointriol. This compound showed moderate insecticidal activity against phytophagous *Myzus persicae*, and *Rhopalosiphum padi*.¹⁴

The number of microorganisms from this region seems to be underexplored. The Yucatan peninsula thus remains an invaluable source of biological diversity for bioprospecting purposes.^{15,16} In this study, the chemical diversity and antimicrobial properties of a series of cultivable fungi isolated

Received: January 26, 2022

Accepted: March 14, 2022

Published: March 28, 2022





Figure 1. Location of cenote Tza Itzá ($20^{\circ}43'50.27''$ N, $89^{\circ}27'56.82''$ W) in the Yucatan peninsula.




Figure 2. Fungal isolates from cenote Tza Itzá on PDA medium: *Aspergillus* spp. NCA257, NCA264, and NCA276; *Stachybotrys* sp. NCA252; and *Cladosporium* sp. NCA273.

from the Tza Itzá cenote were explored by combining conventional chemical studies, cocultures analysis, and metabolomics (Figure 1). The isolation of several compounds from single-strain cultures demonstrated their potential to produce interesting chemistry. In addition, the analysis of cocultures led to the identification of compounds that were not present, or were barely detected, in the monocultures. Moreover, molecular networking (MN) analysis of the liquid chromatography coupled to high-resolution mass spectrometry (LC-HRMS) data of each fungus, allowed the annotation of additional compounds beyond to the isolated. Finally, several

isolated compounds showed significant antibacterial activity against ESKAPE pathogens, *Mycobacterium tuberculosis*, and nontuberculous mycobacterias.

RESULTS AND DISCUSSION

Fungal Strains Isolation and Identification. Five culturable fungal strains were isolated from sediments samples collected in the cenote Tza Itzá. The taxonomic identity of the strains was determined by molecular sequencing of the ITS rDNA, followed by BLAST search and maximum likelihood analysis.^{17,18} The strains were identified as *Aspergillus*




Figure 3. Isolated compounds from *Aspergillus* spp. NCA257, NCA264, and NCA276.

(NCA257, NCA264, and NCA276), *Stachybotrys* (NCA252), and *Cladosporium* (NCA273) (Figure 2 and Table S1, Supporting Information). Next, and as part of a program to explore the chemical diversity and antimicrobial potential of fungal species isolated from unexplored areas of Mexico, isolated strains were subjected to chemical, biological, and metabolomic analyses.

Chemical Study and Molecular Networking Analysis of *Aspergillus* spp. The chemical study of the defatted extracts from moist rice cultures of the *Aspergillus* spp. yielded a new compound, 17-deoxy-aspergillin PZ (1),¹⁹ and the known aspergillin PZ (2),²⁰ aspochalasin D (3),²¹ asperphenamate (4),²² *N*-benzoyl-*L*-phenylalaninol (5),²³ and 2-*O*-methylbutyrolactone II (6)²⁴ from strain NCA276; diorcinol (7),²⁵ sydonic acid (8),²⁶ and 11-dehydroxy sydonic acid (9)²⁷ from NCA264; and asterriquinol D dimethyl ether (10)²⁸ from NCA257 (Figure 3). All known compounds were elucidated by comparison with reported spectroscopic data (Supporting Information). Compound 1 was originally isolated from a jar fermentation of a presumably *Aspergillus* species, and its planar structure was established by NMR analysis.¹⁹ However, its absolute configuration had not been determined before. Suitable colorless crystals (Figure S1, Supporting Information) of 1 were obtained for X-ray structural determination and its absolute configuration was established as 3*S*,3*a**R*,4*S*,6*a*-*S*,8*a**S*,9*R*,13*S*,13*a**R*,13*b**R*-1 (Figure 4).

Next, the organic extracts of these strains were analyzed by ultrahigh-performance liquid chromatography tandem high-resolution electrospray mass spectrometry (UPLC-HRESIMS-MS/MS). Then, metabolomic analyses were performed using an in-house dereplication procedure^{29,30} and the Global Natural Products Social (GNPS) platform to perform feature-based molecular networking (FBMN) and spectral library search.^{31–33} The strain NCA276 showed the largest and most chemically diverse MN. Its metabolite features were




Figure 4. Displacement ellipsoid plot (50% probability level) of 1 at 100(2) K.

grouped into 800 nodes arranged in 22 clusters or chemical families with >3 nodes per cluster, 29 with two nodes, and 506 singletons (Figure 5). Chemical ontology analysis by the MolNetEnhancer GNPS tool classified the molecular features in six classes of compounds (Figure 5). Detailed analysis of the aspochalasins (cytochalasins) cluster allowed to manually annotate the isolated compounds 1–3 using their HRMS-MS/MS data along with a series of related compounds (aspochalasins I and K, and flavichalasin N)^{34,35} annotated by GNPS (Figure 5 and Table 1). In the phenylalanine and derivatives cluster, 4 and its precursor 5 were annotated (Figure 5). Lastly, emeriphenolicin F³⁶ and erythrokyrin³⁷ were found by GNPS in the glutamic and derivatives and furofurans nodes (Figure 5). For strain NCA264, FBMN analysis grouped the molecular features (591 nodes) into 9 classes of compounds clustered in 29 chemical families with >3 nodes per cluster, 29 with two nodes, and 542 singletons (Figure 6). The biggest family in the network belongs to the sesquiterpenoids, where 8, 9, and its derivative (*S*)-10-hydroxysydonic acid³⁸ were annotated (Figure 6). In addition, 7 was successfully annotated by GNPS in the phenyl-




Figure 5. Metabolomic analysis of *Aspergillus* sp. NCA276. (A) FBMN (>2 nodes per cluster) and (B–E) selected clusters with nodes showing the compounds manually annotated (red circles) and by GNPS library search. Singletons (no molecular relatives) are not shown in the network.

propanoids and polyketides family (Table 1). Finally, MN of strain NCA257 showed 154 nodes grouped in 11 clusters with >3 nodes per cluster, 3 with two nodes, and 99 singletons. Interestingly, FBMN analysis did not show any structural families, but the annotation of **10** was possible using its HRMS-MS/MS data (Figure 7 and Table 1).

Chemical Study and Molecular Networking Analysis of *Stachybotrys* sp. NCA252 and *Cladosporium* sp. NCA273. From the organic extract of *Stachybotrys* sp. NCA252 culture, the isocoumarin, *O*-methylmellein (**11**),³⁹ the dolabellane-type diterpenoids, atranones A and B (**12** and **13**),⁴⁰ and the phenylspirodrimanones, stachybotrolide acetate (**14**), stachybotryol acetate (**15**), and stachybotryolide (**16**),^{41,42} were isolated (Figure 8 and Supporting Informa-

tion). Compounds **12** and **13** are commonly observed in the chemotype A of *Stachybotrys* spp., and no compounds from chemotype S (e.g., macrocyclic trichothecenes such as satratoxins and roridins) were detected in the NCA252 strain.^{43,44} In addition, the interconversion of **15** into lactone **14** was observed by NMR⁴⁵ (Figure S44). Finally, this is the first report of the isolation of **11** in a fungus of the *Stachybotrys* genera.

GNPS analysis of this strain grouped the metabolite features into 325 nodes arranged in 12 clusters with >3 nodes per cluster, 17 with two nodes, and 103 singletons. Chemical ontology analysis showed the presence of seven classes of compounds (Figure 9). Detailed analysis of the benzofurans and phthalides nodes allowed the manual annotation of **14** and

Table 1. Chemical Annotation of Metabolites from *Aspergillus* spp

compound	observed ion (m/z) ^d	adduct	molecular formula	exact mass (m/z) ^e	mass accuracy (ppm)
<i>Aspergillus</i> sp. NCA276					
17-deoxyaspergillin PZ (1) ^{a,b}	386.268	[M+H] ⁺	C ₂₄ H ₃₆ NO ₃	386.2682	-2.0
aspergillin PZ (2) ^{a,b}	402.263	[M+H] ⁺	C ₂₄ H ₃₆ NO ₄	402.2630	-2.2
aspochalasin D (3) ^{a,b}	384.253	[M-H ₂ O+H] ⁺	C ₂₄ H ₃₄ NO ₃	384.2523	-2.7
asperphenamate (4) ^{a,c}	507.228	[M+H] ⁺	C ₃₂ H ₃₁ N ₂ O ₄	507.2279	+0.1
<i>N</i> -benzoyl-L-phenylalaninol (5) ^{a,c}	256.133	[M+H] ⁺	C ₁₆ H ₁₈ NO ₂	256.1328	+1.6
aspochalasin I ^c	416.240	[M+H] ⁺	C ₂₄ H ₃₄ NO ₅	416.2437	+1.3
aspochalasin K ^c	434.289	[M+H] ⁺	C ₂₅ H ₄₀ NO ₅	434.2906	+1.2
flavichalasin N ^c	434.253	[M+H] ⁺	C ₂₅ H ₄₀ NO ₅	434.2906	+1.2
emeriphenolicin F ^c	734.459	[M+H] ⁺	C ₄₄ H ₆₄ NO ₈	734.4632	+0.8
erythrokyrin ^c	456.235	[M+H] ⁺	C ₂₆ H ₃₄ NO ₆	456.2386	+1.2
<i>Aspergillus</i> sp. NCA264					
diorcinol (7) ^{a,c}	229.086	[M-H] ⁻	C ₁₄ H ₁₃ O ₃	229.0863	-3.1
sydonic acid (8) ^{a,c}	265.144	[M-H] ⁻	C ₁₅ H ₂₁ O ₄	265.1442	-1.3
11-dehydrosydonic acid (9) ^{a,c}	263.129	[M-H] ⁻	C ₁₅ H ₁₉ O ₄	263.1286	-1.1
(<i>S</i>)-10-hydroxysydonic acid ^c	281.139	[M-H] ⁻	C ₁₅ H ₂₁ O ₅	281.1389	-1.9
<i>Aspergillus</i> sp. NCA257					
Asterriquinol D dimethyl ether (10) ^{a,b}	429.180	[M+H] ⁺	C ₂₆ H ₂₅ N ₂ O ₄	429.1800	-2.1

^aIsolated compound. ^bManual annotation. ^cGNPS annotation. ^dValues from GNPS. ^eValues from UPLC-HRESIMS-MS/MS analysis.




Figure 6. Metabolomic analysis of *Aspergillus* sp. NCA264. (A) FBMN and (B,C) selected clusters with nodes showing the compounds annotated by GNPS library search.

16, and by GNPS the annotation of a series of related phenylspirodrimanes, including 15, asperugin,⁴⁶ and stachy-bysbin A⁴⁷ (Figure 9 and Table 2).

The chemical study of the culture of *Cladosporium* sp. NCA273 did not yield any pure compound due to the scarcity

of the organic extract obtained from the solid culture (poor growth in this medium). In addition, the MN analysis of this strain grouped the metabolite features into 77 nodes arranged in 6 clusters with >3 nodes per cluster, 8 with two nodes, and 63 singletons. No matches were found using the in-house




Figure 7. Metabolomic analysis of *Aspergillus* sp. NCA257 and selected node showing manually annotated 9 (red circle).




Figure 8. Isolated compounds from *Stachybotrys* sp. NCA252.

dereplication procedure nor the spectral library search in the GNPS, and only two chemical families (O-glycosyl compounds and glycerophosphocholines) were observed by FBMN (Figure S45).

***Aspergillus* spp. NCA257 and NCA276 and *Cladosporium* sp. NCA273 Coculture Analysis.** During the isolation process of the fungal strains from the cenote sediment samples, a series of antagonistic interactions between *Aspergillus* spp. NCA257 and NCA276 and *Cladosporium* sp. NCA273 were observed (Figure 10). It is known that coculture fermentation can trigger the expression of silent biosynthetic gene clusters to produce new secondary metabolites, increase the amounts of others, or in some cases inhibit the production of some compounds.^{48–50} This strategy was thus employed to analyze

the chemical variation of these strains when cocultivated in solid media (Supporting Information).

High-performance liquid chromatography (HPLC) analysis of the organic extract of the coculture of *Aspergillus* sp. NCA257 and *Cladosporium* sp. NCA273 revealed the presence of penicillic acid (17),⁵¹ 5,6-dihydropenicillic acid (18),⁵² and 3-isobutyl-6-(1-hydroxy-2-methylpropyl)-pyrazin-2(1H)-one (19)⁵³ (Figure 11). Compound 17 was not isolated from the monoculture of *Aspergillus* sp. NCA257 due to its low abundance. However, the induction of 17–19 by coculture allowed its isolation and structural elucidation (Supporting Information). Interestingly, 17 abundance (AUC by HPLC analysis) was 250% times higher in the coculture compared to the monoculture of *Aspergillus* sp. NCA257, whereas 10 was totally suppressed. On the other hand, the analysis of the *Aspergillus* spp. NCA257 and NCA276 coculture revealed similar results than those of NCA257 and NCA273 coculture, that is, 17–19 were induced in the coculture, and the aspochalasins 1 and 3 were suppressed (Figure 12).

The MN analysis by GNPS of the cocultures of NCA257/NCA273 and NCA257/NCA276, and the single-cultures of all three strains consisted of 1380 nodes, which were grouped into 114 clusters with >2 nodes per cluster (Figure S46). Interestingly, the MN showed a major contribution from strain NCA257 (*Aspergillus* sp.) in the clusters of the chemical features (Figure S46). This agrees with the HPLC profiles (Figures 11 and 12), where the production of the penicillic acids 17 and 18, and the pyrazinone 19 suppresses the biosynthesis of the main metabolites from the other strains.⁵⁴ Finally, aspochalasin J, aspochalasinol A, and alterporriol B were annotated by GNPS in MN (Figure S46 and Table S4, Supporting Information).

Biological Activity of Pure Compounds. The antibacterial activity of the isolated compounds was assessed against a panel of ESKAPE pathogens using the microdilution assay^{55,56} at 100 μg/mL or 10 μg/mL (Table 3). From these, the aspochalasin 3, the amino acid ester 4, the bis-indolyl




Figure 9. Metabolomic analysis of *Stachybotrys* sp. NCA252. (A) FBMN and (B–F) selected clusters with nodes showing the compounds manually annotated (red circles) and by GNPS library search.

benzenoid 10, and the phenylspirodrimanes 14–16 showed high activity (>50% inhibition) against MSSA and MRSA strains with 14 as the most potent metabolite (>81% inhibition at test concentration of 10 $\mu\text{g}/\text{mL}$). None of the tested

compounds were active against *Pseudomonas aeruginosa* ATCC 27853, *Klebsiella aerogenes* ATCC 13048, *K. pneumoniae* ATCC 700603, *Acinetobacter baumannii* ATCC 17978, and the multidrug-resistant *A. baumannii* clinical strain A564.⁵⁷

Table 2. Chemical Annotation of Metabolites from *Stachybotrys* sp. NCA252

compound	observed ion (m/z) ^d	adduct	molecular formula	exact mass (m/z) ^e	mass accuracy (ppm)
O-methylmellein (11) ^{a,b}	215.068	[M+Na] ⁺	C ₁₁ H ₁₂ O ₃ Na	215.0675	+1.7
atranone A (12) ^{a,b}	461.218	[M+HCOO] ⁻	C ₂₅ H ₃₃ O ₈	461.2182	+0.2
atranone B (13) ^{a,b}	445.223	[M-H] ⁻	C ₂₅ H ₃₃ O ₇	445.2226	-1.3
stachybotrolide acetate (14) ^{a,b}	429.227	[M+H] ⁺	C ₂₅ H ₃₃ O ₆	429.2272	+0.1
stachybotrydial acetate (15) ^a	369.206	[M-AcO+H] ⁺	C ₂₃ H ₂₉ O ₄	369.2061	+0.2
stachybotrolide (16) ^{a,b}	387.217	[M+H] ⁺	C ₂₃ H ₃₁ O ₅	387.2160	-1.6
stachybotramide ^c	430.258	[M+H] ⁺	C ₂₅ H ₃₆ NO ₅	430.2593	+1.2
K-76 diacetate ^c	487.233	[M+H] ⁺	C ₂₇ H ₃₅ O ₈	487.2332	+1.1
stachybotrysin H ^c	471.238	[M-H ₂ O+H] ⁺	C ₂₇ H ₃₅ O ₇	471.2383	+1.2
myrothecisin D ^c	371.222	[M-H ₂ O+H] ⁺	C ₂₃ H ₃₁ O ₄	371.2222	+1.4
F1839E ^c	446.254	[M+H] ⁺	C ₂₅ H ₃₆ NO ₆	446.2542	+1.1
asperugin ^c	401.232	[M+H] ⁺	C ₂₄ H ₃₃ O ₅	401.2328	+1.4
stachybybin A ^c	385.201	[M+H] ⁺	C ₂₃ H ₂₉ O ₅	385.2015	+1.4

^aIsolated compound. ^bManually annotation. ^cGNPS annotation. ^dValues from GNPS. ^eValues from UPLC-HRESIMS-MS/MS analysis.




Figure 10. Fungal interaction between *Aspergillus* spp. NCA257 and NCA276 and *Cladosporium* sp. NCA273 in PDA plates.

Finally, 7, 9, and 19 were not tested due to the scarcity of the isolated material.

In addition, the anti-mycobacteria activity of the isolated compounds was assessed using the microplate Alamar blue (MABA) and low oxygen recovery (LORA) assays^{58,59} against tuberculous (TB) and nontuberculous mycobacterias (NTM), respectively, and a cytotoxicity assay against Vero cell line (ATCC CCL-81)⁶⁰ (Table 4). Compounds 3, 4, 7, 14, 16, and 17 showed >98% growth inhibition of TB at 50 μ g/mL and CC₅₀ values >35 μ M against Vero cells. Moreover, 3, 4, 7, 14, and 16, displayed anti-*M. avium* activity with MIC values of 22.6, 37.0, 20.0, 11.6, and 17.0 μ M, respectively, and 7, 14, and 16 against *M. marinum* with MIC values of 47.2, 23.1, and 47.0 μ M, respectively. Compounds 2 and 9 were not tested in these assays due to the paucity of the isolated material.

CONCLUSIONS

In summary, this work represents the first contribution to the chemical diversity and biology of fungi from sediments in a

cenote of the Yucatan peninsula. Strains *Aspergillus* (NCA257, NCA264, and NCA276), *Stachybotrys* (NCA252), and *Cladosporium* (NCA273) were isolated from sediments' samples collected in the cenote Tza Itzá and taxonomically identified by molecular sequencing of the ITS rDNA. From this, several isolated compounds showed significant activity against MSSA, MRSA, other ESKAPE pathogens and mycobacterial strains and could be further studied for the development of potential drug leads. The absolute configuration of new 17-deoxy-aspergillin PZ (1) was established via single-crystal X-ray crystallography. Finally, cocultures and metabolomics analysis revealed that communication between the strains is needed to produce metabolites that were not present in the monocultures. Overall, this work uncovers the chemical and biological potential of understudied fungal strains to produce biosynthetic bioactive natural products.

EXPERIMENTAL SECTION

General Experimental Procedures. Optical rotations, and UV data were measured using a Rudolph Research Autopol III polarimeter (Rudolph Research Analytical), and a Varian Cary 100 Bio UV-vis spectrophotometer (Varian Inc.), respectively. NMR experiments were conducted in CDCl₃ or methanol-*d*₄, using a JEOL ECA-600 spectrometer (JEOL Ltd.), a Varian VNMRS 400 (Varian Inc.), or a Bruker Avance III 400 MHz (Bruker BioSpin Corp.). HRESIMS data were acquired using a Q Exactive Plus system (Thermo Fisher Scientific), equipped with an electrospray ionization source with an HCD cell. Data were collected in both positive and negative modes via direct injection or through an Acquity UPLC system (Waters Corp.) using a BEH C₁₈ column (50 \times 2.1 mm i.d., 1.7 μ m; Waters Corp.) with a gradient solvent system from 15:85 to 100:0 CH₃CN-0.1% aqueous formic acid for 10 min at a flow rate of 0.3 mL/min. Analytical and preparative HPLC were carried out on a Waters HPLC system equipped with a 2535 quaternary pump, a 2707 autosampler, and 2998 PDA and 2424 ELSD detectors, using Gemini C₁₈ and Kinetex C₁₈ columns (5 μ m, 110 Å, 250 \times 4.6 mm i.d. and 5 μ m, 110 Å, 250 \times 21.2 mm i.d.; Phenomenex) for analytical and preparative runs, respectively. Data acquisition and management were performed with the Empower 3 software (Waters Corp.). Flash chromatography was conducted on a CombiFlash Rf+ Lumen system (Teledyne Technologies Inc.) equipped with PDA and ELSD detectors using RediSep Rf Gold Si-gel columns (Teledyne Technologies Inc.). Reagent-




Figure 11. HPLC traces (PDA UV $\lambda = 254$ nm) of monoculture extracts of *Aspergillus* sp. NCA257 and *Cladosporium* sp. NCA273 and the coculture extract.

grade chloroform, *n*-hexane, and methanol, and HPLC- and MS-grade acetonitrile, methanol, and water were purchased from J.T. Baker (Avantor Performance Materials). Deuterated NMR solvents were acquired from Cambridge Isotope Laboratories, Inc.

Fungal Strains Isolation and Identification. The fungi of interest were isolated from sediment samples collected from cenote *Tza Itzá* (20°43'50.27" N, 89°27'56.82" W), in the state of Yucatán, Mexico in 2018. The cenote dimensions are a maximum depth of 40 m, water mirror of 7.6 × 10.5 m, and dry cave of 30 × 8 × 5 m (L × W × H). The isolation of the strains was made in A1 10% media with peptone (0.4 g), starch (1.0 g), yeast extract (0.2 g), supplemented with rifampicin (5 μ g/mL) and cycloheximide (0.1 mg/mL) to reduce the growth of Gram-negative bacteria as previously described.¹⁰ The fungal cultures are maintained at -80 °C at the microbial culture collection from Unidad de Química en Sisal, Facultad de Química, UNAM, in Yucatan, Mexico. A copy of each strain used for the chemical studies is also maintained at the Departamento de Farmacia, Facultad de Química, UNAM, Mexico. Axenic cultures of strains NCA252, NCA257, NCA264, NCA273, and NCA276 were subjected to molecular identification by ITS sequencing and BLAST search with the RefSeq database using the methodologies reported.^{61,62} For designation of taxonomic names, the results of the ITS BLAST search using GenBank were interpreted with caution using modification of outlined criteria.⁶³ Since we sequenced only the ITS region, we chose a rather conservative approach and

made identifications only to the genus level even when BLAST sequence homology was $\geq 99\%$. Sequences for each strain were deposited in the National Center for Biotechnology Information (NCBI, <http://www.ncbi.nlm.nih.gov/BLAST/>) and given the GenBank Data Base accession numbers indicated in (Table S1 Supporting Information).

Fungal Cultivation, Extraction, and Isolation and Characterization of Pure Compounds. The methods utilized to culture the fungal strains (mono and cocultures) to prepare their organic extracts and to isolate and structurally elucidate compounds 1–19 from the fungi followed well-established protocols^{64–67} and are detailed in the Supporting Information.

17-Deoxy-aspergillin PZ (1). Colorless solid; $[\alpha]_D^{18.1} + 7.5$ (*c* 0.1, CHCl₃); UV (CHCl₃) λ_{\max} (log ϵ) 322 (3.28), 285 (3.34) nm; ¹H and ¹³C NMR, see Table S3 on Supporting Information; HRESIMS *m/z* 386.2682 [M + H]⁺ (calcd. for C₂₄H₃₆NO₃, 386.2695).

X-ray Crystallographic Analysis of 17-deoxy-aspergillin PZ (1). Suitable X-ray quality single crystals for 17-deoxy-aspergillin PZ (1) were successfully obtained from MeOH–CHCl₃–H₂O (20:1:1) recrystallization mixture. The crystallographic data and data collection parameters are mentioned in Table S2. All reflection intensities were measured at 100(2) K using a Gemini R diffractometer (equipped with Atlas detector) with CuK α radiation ($\lambda = 1.54178$ Å) under the program CrysAlisPro (Version CrysAlisPro 1.171.38.43f, Rigaku OD, 2015). The refinement




Figure 12. HPLC traces (PDA UV $\lambda = 254$ nm) of monoculture extracts of *Aspergillus* spp. NCA257 and NCA276 and the coculture extract.

Table 3. Antimicrobial Activity of Isolated Compounds against ESKAPE Bacteria at 100 $\mu\text{g/mL}$

compound	growth inhibition (%)				
	VSEF ^c	VREF ^c	MSSA ^c	MRSA ^c	EC ^c
17-deoxy-aspergillin PZ (1)			11.8	27.7	28.1
aspergillin PZ (2)				23.0	9.0
aspochalasin D (3)	37.2	23.8	81.4	87.2	
asperphenamate (4)			10.4	62.9	22.5
<i>N</i> -benzoyl- <i>L</i> -phenylalaninol (5)			13.3	50.7	
2- <i>O</i> -methylbutyrolactone II (6)					
sydonic acid (8)				34.0	11.9
asterriquinol D dimethyl ether (10)			50.9	62.3	48.2
<i>O</i> -methymllelin (11)					16.2
atranone A (12)	13.0				6.0
atranone B (13)					12.2
stachybotrolide acetate (14) ^d			81.0	92.9	21.5
stachybotrydial acetate (15)			82.7	89.3	18.3
stachybotrolide (16)			82.2	91.2	23.7
penicillic acid (17)			80.4	51.4	
5,6-dihydropenicillic acid (18)	38.3				
MIC positive control in $\mu\text{g/mL}$.	3.75 ^a	25.0 ^a	200.0 ^b	2.5 ^a	0.5 ^c

^aVancomycin - Inactive at 100 $\mu\text{g/mL}$. ^bAmpicillin - Inactive at 100 $\mu\text{g/mL}$. ^cGentamicin - Inactive at 100 $\mu\text{g/mL}$. ^dTested at 10 $\mu\text{g/mL}$. ^eVSEF, vancomycin-susceptible *E. faecalis* ATCC 29212; VREF, vancomycin-resistant *E. faecalis* ATCC 51299; MSSA, methicillin-susceptible *S. aureus* ATCC 25923; MRSA, methicillin-resistant *S. aureus* ATCC 43300; and EC, *E. cloacae* ATCC 700323.

of cell dimensions and data reduction were performed using the most recent version of the program (viz. CrysAlisPro 1.171.40.53, Rigaku OD, 2019). The structure was solved with the program SHELXT-2018/2 and was refined on F^2 by the

full-matrix least-squares method using SHELXL-2018/3.⁶⁸ An empirical absorption correction was applied using CrysAlisPro 1.171.40.53.⁶⁹ Non-hydrogen atoms were refined anisotropically. In the refinement, hydrogens attached to carbon, were

Table 4. Anti-mycobacteria and Cytotoxic Activities of Isolated Compounds

compound	<i>M. tuberculosis</i> H37Rv		<i>M. abscessus</i> ATCC 19977	<i>M. chelonae</i> ATCC 35752	<i>M. marinum</i> ATCC 927	<i>M. avium</i> ATCC 15769	<i>M. kansasii</i> ATCC 12478	Vero cell ATCC CCL- 81 ^d
	MABA ^a	LORA ^b			MABA ^c			
17-deoxy-aspergillin PZ (1)	19 (ND)	ND	ND	ND	ND	ND	ND	ND
aspochalasin D (3)	101 (39.5)	>50	>50 (0)	>50 (17)	>50 (31)	22.55	>50 (16)	37.5
asperphenamate (4)	101 (16.6)	>50	>50 (0)	>50 (6)	>50 (22)	37.00	>50 (0)	>50
<i>N</i> -benzoyl-L-phenylalaninol (5)	20 (ND)	ND	ND	ND	ND	ND	ND	ND
2- <i>O</i> -methylbutyrolactone II (6)	83 (ND)	ND	ND	ND	ND	ND	ND	ND
diorcinol (7)	100 (47.9)	>50	>50 (0)	>50 (12)	47.24	19.98	>50 (0)	39.9
sydonic acid (8)	30 (ND)	ND	ND	ND	ND	ND	ND	ND
asterriquinol D dimethyl ether (10)	13 (ND)	ND	ND	ND	ND	ND	ND	ND
<i>O</i> -methylmellein (11)	22 (ND)	ND	ND	ND	ND	ND	ND	ND
atranone A (12)	29 (ND)	ND	ND	ND	ND	ND	ND	ND
atranone B (13)	33 (ND)	ND	ND	ND	ND	ND	ND	ND
stachybotrolide acetate (14)	99 (48.3)	>50	>50 (0)	>50(0)	23.15	11.63	>50(0)	38.2
stachybotrydial acetate (15)	38 (ND)	ND	ND	ND	ND	ND	ND	ND
stachybotrolide (16)	103 (49.4)	>50	>50 (0)	>50 (0)	47.02	16.92	>50(0)	39.7
penicillic acid (17)	98 (49.5)	>50	ND	ND	ND	ND	ND	ND
5,6-dihydropenicillic acid (18)	42 (ND)	ND	>50 (0)	>50 (73)	>50 (58)	>50 (53)	>50(0)	35.4
3-isobutyl-6-(1-hydroxy-2-methylpropyl)- pyrazin-2(1H)-one (19)	23 (ND)	ND	ND	ND	ND	ND	ND	ND
MIC Rifampicin ($\mu\text{g/mL}$)	100 (0.03)	0.08	>8.0	4.25	0.09	0.05	0.41	>8.0

^a% Inhibition at 50 $\mu\text{g/mL}$ (MIC μM). ^bMIC μM . ^cMIC μM (% Inhibition). ^dCC₅₀, cytotoxic concentration to 50% inhibition of the cell line. ND, not determined since % inhibition of *M. tuberculosis* H37Rv using MABA was below 90%.

treated as riding atoms using SHELXL default parameters, while those attached to nitrogen were located with electron difference maps. The crystallographic data of **1** in CIF format was deposited in the Cambridge Structural Database (CSD) (CCDC no. 2142221).

Biological Activity Assays. Pure compounds were tested *in vitro* for antibacterial activity using the Clinical and Laboratory Standards Institute (CLSI) broth dilution standard method.^{55,56} Target bacteria used in the assays belong to the ESKAPE group: vancomycin-susceptible *Enterococcus faecalis* ATCC 29212 (VSEF), vancomycin-resistant *E. faecalis* ATCC 51299 (VREF), methicillin-susceptible *Staphylococcus aureus* ATCC 25923 (MSSA), methicillin-resistant *S. aureus* ATCC 43300 (MRSA), *Klebsiella aerogenes* ATCC 13048, *K. pneumoniae* ATCC 700603, *Acinetobacter baumannii* ATCC 17978, multidrug-resistant *A. baumannii* strain 564 (clinical isolated),⁵⁷ *Pseudomonas aeruginosa* ATCC 27853, and *Enterobacter cloacae* ATCC 700323. The compounds were dissolved in DMSO to obtain a stock solution and then tested at a final concentration of 100 $\mu\text{g/mL}$ or 10 $\mu\text{g/mL}$. The bioassays were carried out in 96-well plates in triplicate at concentrations of pure compound. In addition, pure compounds were also tested against *M. tuberculosis* H37Rv (MT) strain under both aerobic (replicating) and anaerobic (nonreplicating; NRMT) conditions using MABA and LORA assays.⁵⁸ Compounds with activity >90% of inhibition of the growth of MT in MABA were further tested against *M. abscessus* ATCC 19977, *M. chelonae* ATCC 35752, *M. marinum* ATCC 927, *M. avium* ATCC 15769, and *M. kansasii* ATCC 12478, using MABA⁵⁸ and a cytotoxicity assay against Vero cell line ATCC CCL-81.⁶⁰ For the latter, 0.6 mM of resazurin was used and the absorbance was recorded at 530 nm (excitation) and 590 nm (emission). Positive controls for all assays are indicated in Tables 3 and 4.

Molecular Networking and Metabolomics Analysis.

Each extract from the mono and cocultures was analyzed by LC-HRMS-MS/MS using a previously described methodology.^{66,67} Raw data were converted to mzML format using the ProteoWizard tool MsConvert (version 3.0.20239) and the resulting files uploaded to the Global Natural Products Social (GNPS; <https://gnps.ucsd.edu>) Web server using the FTP Server Version 5.17.16. Molecular networks were generated following the workflow previously published.^{31,32} Data was filtered by removing all MS/MS fragment ions within ± 17 Da of the precursor *m/z*. The precursor ion mass tolerance was set to 0.01 Da and a MS/MS fragment ion tolerance of 0.02 Da. A network was then created where edges were filtered to have a cosine score above 0.7 and more than five matched peaks. Furthermore, edges between two nodes were kept in the network if and only if each of the nodes appeared in each other's respective top 10 most similar nodes. Finally, the maximum size of a molecular family was set to 100, and the lowest scoring edges were removed from molecular families until the molecular family size was below this threshold. The spectra in the network were then searched against GNPS' spectral libraries. The library spectra were filtered in the same manner as the input data. All matches between network spectra and library spectra were required to have a score above 0.7 and at least five matched peaks. GNPS MolNetEnhancer and Dereplicator+ tools were applied for chemical classification.^{33,70} Molecular networks were visualized with Cytoscape 3.8.1.⁷¹ Finally, manual dereplication was assessed using UV-absorption maxima and HRMS-MS/MS data against MS/MS data of **1–19** and by comparison with those reported in the Dictionary of Natural Products,⁷² SciFinder,⁷³ and an in-house mycotoxins database. The annotation of isolated compounds was at confidence level 1, according to the metabolomics standards initiative and exact mass accuracy <5 ppm.⁷⁴

■ ASSOCIATED CONTENT

SI Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acsomega.2c00544>.

Methods for fungal cultivation, extraction, and isolation and characterization of pure compounds; identity of fungal isolates; NMR and MS spectra of 1–19; crystallographic data of 1; MN and chemical annotation of cocultures (PDF)

■ AUTHOR INFORMATION

Corresponding Author

Mario Figueroa – *Departamento de Farmacia, Facultad de Química, Universidad Nacional Autónoma de México, Ciudad de México 04510, Mexico*; orcid.org/0000-0001-7004-0591; Email: mafiquer@unam.mx

Authors

Carlos A. Fajardo-Hernández – *Departamento de Farmacia, Facultad de Química, Universidad Nacional Autónoma de México, Ciudad de México 04510, Mexico*; orcid.org/0000-0003-1487-1926

Firoz Shah Tuglak Khan – *Department of Chemistry and Biochemistry, University of North Carolina Greensboro, Greensboro, North Carolina 27402, United States*; orcid.org/0000-0002-3800-0423

Laura Flores-Bocanegra – *Department of Chemistry and Biochemistry, University of North Carolina Greensboro, Greensboro, North Carolina 27402, United States*; orcid.org/0000-0002-1393-7834

Alejandra Prieto-Davó – *Unidad de Química en Sisal, Facultad de Química, Universidad Nacional Autónoma de México, Sisal, Yucatán 97356, Mexico*

Baojie Wan – *Institute for Tuberculosis Research, College of Pharmacy, University of Illinois at Chicago, Chicago, Illinois 60612, United States*

Rui Ma – *Institute for Tuberculosis Research, College of Pharmacy, University of Illinois at Chicago, Chicago, Illinois 60612, United States*

Mallique Qader – *Institute for Tuberculosis Research, College of Pharmacy, University of Illinois at Chicago, Chicago, Illinois 60612, United States*

Rodrigo Villanueva-Silva – *Departamento de Farmacia, Facultad de Química, Universidad Nacional Autónoma de México, Ciudad de México 04510, Mexico*

Anahí Martínez-Cárdenas – *Departamento de Farmacia, Facultad de Química, Universidad Nacional Autónoma de México, Ciudad de México 04510, Mexico*

Marian A. López-Lobato – *Departamento de Farmacia, Facultad de Química, Universidad Nacional Autónoma de México, Ciudad de México 04510, Mexico*

Shabnam Hematian – *Department of Chemistry and Biochemistry, University of North Carolina Greensboro, Greensboro, North Carolina 27402, United States*; orcid.org/0000-0002-0788-7615

Scott G. Franzblau – *Institute for Tuberculosis Research, College of Pharmacy, University of Illinois at Chicago, Chicago, Illinois 60612, United States*; orcid.org/0000-0002-8698-0243

Huzefa A. Raja – *Department of Chemistry and Biochemistry, University of North Carolina Greensboro, Greensboro, North*

Carolina 27402, United States; orcid.org/0000-0002-0824-9463

Rodolfo García-Contreras – *Departamento de Microbiología y Parasitología, Facultad de Medicina, Universidad Nacional Autónoma de México, Ciudad de México 04510, Mexico*; orcid.org/0000-0001-8475-2282

Complete contact information is available at:

<https://pubs.acs.org/10.1021/acsomega.2c00544>

Author Contributions

This work was part for the Ph.D. thesis of C.A.F.-H. from the Posgrado en Ciencias Químicas, UNAM. C.A.F.-H. and M.F. designed the experiments. C.A.F.-H., F.S.T.K., L.F.-B., A.P.-D., B.W., R.M., M.Q., S.G.F., R.V.-S., A.M.-C., M.A.L.-L., and H.A.R. performed the experiments. C.A.F.-H., A.P.-D., S.H., S.G.F., H.A.R., R.G.-C., and M.F. analyzed the data and revised the manuscript. C.A.F.-H. and M.F. wrote, reviewed, and edited the manuscript. All authors have read and agreed to the revised version of the manuscript.

Notes

The authors declare no competing financial interest. LC-MS/MS data can be accessed at MassIVE (accession no. MSV000088691; accessed on January 26, 2022). The molecular network data set of fungal coculture species *Aspergillus* spp. NCA257, NCA276 and *Cladosporium* sp. NCA273 can be accessed at MassIVE (accession no. MSV000088690; accessed on January 26, 2022). The molecular network of *Stachybotrys* sp. NCA252 at <https://gnps.ucsd.edu/ProteoSAFe/status.jsp?task=2e6ed82df2854ee481908c5e7be883e0> (MolNetEnhancer analysis; accessed on January 26, 2022), molecular network of *Aspergillus* sp. NCA257 at <https://gnps.ucsd.edu/ProteoSAFe/status.jsp?task=2aea96c6e47746e28349aa3241c8d065> (MolNetEnhancer analysis; accessed on January 26, 2022), molecular network of *Aspergillus* sp. NCA264 at <https://gnps.ucsd.edu/ProteoSAFe/status.jsp?task=63f8482f83c94d3ea4b34910a972f170> (MolNetEnhancer analysis; accessed on January 26, 2022), molecular network of *Cladosporium* sp. NCA273 at <https://gnps.ucsd.edu/ProteoSAFe/status.jsp?task=5f13c0f6339742879130611c389af5da> (MolNetEnhancer analysis; accessed on January 26, 2022) and molecular network of *Aspergillus* sp. NCA276 at <https://gnps.ucsd.edu/ProteoSAFe/status.jsp?task=ee9f321cb43d4e1da6f38fde4cfa696b> (MolNetEnhancer analysis; accessed on January 26, 2022). Molecular network of fungal coculture species *Aspergillus* spp. NCA257, NCA276 and *Cladosporium* sp. NCA273 at <https://gnps.ucsd.edu/ProteoSAFe/status.jsp?task=66c153a4faa4418485b252df22c99023> (accessed on January 26, 2022).

■ ACKNOWLEDGMENTS

This work was supported by grants from UNAM-DGAPA-PAPIIT IN222220 (M.F.) and FQ-PAIP 5000-9145 (M.F.). A.P.-D. thanks to CONACyT, Ciencia Básica Project A1-S-10785 and the Rockefeller University for providing funds for sampling expeditions and to the divers Biol. Luis Arturo Liévano and M.S. Efraín Chávez Solís for collecting the samples from the cenote. C.A.F.-H. thanks to CONACyT for the fellowship (no. 596831) to pursue his Ph.D. studies. A.M.-C. acknowledges the postdoctoral fellowship from DGAPA, UNAM. S.H. and F.S.T.K. gratefully acknowledge the JONT

School of Nanoscience and Nanoengineering (JSNN) for providing access to the X-ray diffraction facility. M.F. thanks to Q.F.B. Alejandro Camacho Cruz (Cepario, FQ, UNAM), Dr. Isabel Rivero-Cruz and Ramiro del Carmen (FQ, UNAM), Dr. Norma Angélica Márquez-Velázquez (Unidad de Química en Sisal, FQ, UNAM), and M.S. Rosa Isela del Villar and M.S. Nayeli López-Balbiaux, (USAI, FQ, UNAM) for their valuable technical assistance. The authors are deeply grateful to Professor Nicholas H. Oberlies for providing access to the nuclear magnetic resonance (NMR) and mass spectrometry (MS) facilities at the University of North Carolina at Greensboro.

REFERENCES

- (1) Schmitter-Soto, J. J.; Comín, F. A.; Escobar-Briones, E.; Herrera-Silveira, J.; Alcocer, J.; Suárez-Morales, E.; Elías-Gutiérrez, M.; Díaz-Arce, V.; Marín, L. E.; Steinich, B. Hydrogeochemical and biological characteristics of cenotes in the Yucatan peninsula (SE Mexico). *Hydrobiol.* **2002**, *467* (1–3), 215–228.
- (2) Gómez-Nicolás, M.; Rebolledo-Vieyra, M.; Canto-Lugo, E.; Huerta-Quintanilla, R.; Ochoa-Sandoval, P. Connectivity in a karst system using electrical resistivity tomography and network theory. *Groundwater* **2018**, *56* (5), 732–741.
- (3) Hildebrand, A. R.; Pilkington, M.; Connors, M.; Ortiz-Aleman, C.; Chavez, R. E. Size and structure of the Chicxulub crater revealed by horizontal gravity gradients and cenotes. *Nature* **1995**, *376* (6539), 415–417.
- (4) Lian, B.; Yuan, D.; Liu, Z. Effect of microbes on karstification in karst ecosystems. *Chin. Sci. Bull.* **2011**, *56*, 3743–3747.
- (5) Stoessell, R. K.; Moore, Y. H.; Coke, J. G. The occurrence and effect of sulfate reduction and sulfide oxidation on coastal limestone dissolution in Yucatan cenotes. *Groundwater* **1993**, *31* (4), 566–575.
- (6) Hahn, M. W. The microbial diversity of inland waters. *Curr. Opin. Biotechnol.* **2006**, *17* (3), 256–261.
- (7) Krauss, G. J.; Sole, M.; Krauss, G.; Schlosser, D.; Wesenberg, D.; Baerlocher, F. Fungi in freshwaters: ecology, physiology and biochemical potential. *FEMS Microbiol. Rev.* **2011**, *35* (4), 620–651.
- (8) Estrada-Medina, H.; Canto-Canché, B. B.; De los Santos-Briones, C.; O'Connor-Sánchez, A. Yucatán in black and red: Linking edaphic analysis and pyrosequencing-based assessment of bacterial and fungal community structures in the two main kinds of soil of Yucatán State. *Microbiol. Res.* **2016**, *188*, 23–33.
- (9) Marfil-Santana, M. D.; O'Connor-Sánchez, A.; Ramírez-Prado, J. H.; De los Santos-Briones, C.; Lluvia, K.; Rojas-Herrera, R.; Lago-Lestón, A.; Prieto-Davó, A. A computationally simplistic poly-phasic approach to explore microbial communities from the Yucatan aquifer as a potential sources of novel natural products. *J. Microbiol.* **2016**, *54* (11), 774–781.
- (10) Parera-Valadez, Y.; Yam-Puc, A.; López-Aguilar, L. K.; Borges-Argáez, R.; Figueroa-Saldivar, M. A.; Cáceres-Farfán, M.; Márquez-Velázquez, N. A.; Prieto-Davó, A. Ecological strategies behind the selection of cultivable actinomycete strains from the Yucatan peninsula for the discovery of secondary metabolites with antibiotic activity. *Microb. Ecol.* **2019**, *77* (4), 839–851.
- (11) De la Rosa-García, S. C.; Muñoz-García, A. A.; Barahona-Pérez, L. F.; Gamboa-Angulo, M. M. Antimicrobial properties of moderately halotolerant bacteria from cenotes of the Yucatan peninsula. *Letts. Appl. Microbiol.* **2007**, *45* (3), 289–294.
- (12) Marcela Gamboa-Angulo. Antimicrobial screening of tropical microfungi isolated from sinkholes located in the Yucatan peninsula, Mexico. *Afr. J. Microbiol. Res.* **2012**, *6* (10), 2305–2312.
- (13) Ruiz-Jiménez, A. L.; González-Coloma, A.; Andrés-Yeves, M. F.; Ruiz-Sánchez, E.; Heredia, G.; Peraza-Sánchez, S. R.; Medina-Baizabal, I. L.; Reyes-Estebanez, M.; Canto-Caché, B.; Gamboa-Angulo, M. Insect deterrent and nematicidal screening of microfungi from Mexico and anti-aphid compounds from *Glomastix maseei*. *Rev. Argent. Microbiol.* **2017**, *49* (1), 83–92.
- (14) Ruiz-Jiménez, A. L.; Ruiz-Sánchez, E.; Heredia, G.; Tapia-Tussell, R.; González-Coloma, A.; Peraza-Jiménez, K.; Moo-Koh, F. A.; Medina-Baizabal, I. L.; Hernández-Romero, Y.; Mena-Rejón, G. J.; Quijano-Quiñones, R. F.; Gamboa-Angulo, M. Identification of insect-deterrent metabolites from *Acremonium maseei* strain CICY026, a saprophytic fungus from a sinkhole in Yucatán. *Microorganisms* **2019**, *7* (12), 712.
- (15) Pech-Puch, D.; Pérez-Povedano, M.; Lenis-Rojas, O. A.; Rodríguez, J.; Jiménez, C. Marine natural products from the Yucatan peninsula. *Mar. Drugs* **2020**, *18* (1), 59.
- (16) Hernández-Bolio, G. I.; Ruiz-Vargas, J. A.; Peña-Rodríguez, L. M. Natural products from the yucatecan flora: Structural diversity and biological activity. *J. Nat. Prod.* **2019**, *82* (3), 647–656.
- (17) Gardes, M.; Bruns, T. D. ITS primers with enhanced specificity for basidiomycetes-application to the identification of mycorrhizae and rusts. *Mol. Ecol.* **1993**, *2* (2), 113–118.
- (18) White, T. J.; Bruns, T.; Lee, S. J. W. T.; Taylor, J. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. *PCR protocols: a guide to methods and applications*; Academic Press: San Diego, 1990; Vol. 18, pp 315–322.
- (19) Ding, X.; Terui, Y.; Chen, Y. Separation and structural confirmation of three new compounds with antibacterial activity. *Huaxue Yanjiu Yu Yingyong* **2006**, *18* (9), 1026–1028.
- (20) Chen, L.; Liu, Y. T.; Song, B.; Zhang, H. W.; Ding, G.; Liu, X. Z.; Gu, Y. C.; Zou, Z. M. Stereochemical determination of new cytochalasins from the plant endophytic fungus *Trichoderma gamsii*. *Fitoterapia* **2014**, *96*, 115–122.
- (21) Tomikawa, T.; Shin-Ya, K.; Kinoshita, T.; Miyajima, A.; Seto, H.; Hayakawa, Y. Selective cytotoxicity and stereochemistry of aspochalasin D. *J. Antibiot.* **2001**, *54* (4), 379–381.
- (22) Hou, X. M.; Zhang, Y. H.; Hai, Y.; Zheng, J. Y.; Gu, Y. C.; Wang, C. Y.; Shao, C. L. Aspersymmetide A, a new centrosymmetric cyclohexapeptide from the marine-derived fungus *Aspergillus versicolor*. *Mar. drugs* **2017**, *15* (11), 363.
- (23) Li, W.; Fan, A.; Wang, L.; Zhang, P.; Liu, Z.; An, Z.; Yin, W. B. Asperphenamate biosynthesis reveals a novel two-module NRPS system to synthesize amino acid esters in fungi. *Chem. Sci.* **2018**, *9* (9), 2589–2594.
- (24) Chen, M.; Wang, K. L.; Liu, M.; She, Z. G.; Wang, C. Y. Bioactive steroid derivatives and butyrolactone derivatives from a Gorgonian-derived *Aspergillus* sp. fungus. *Chem. Biodivers.* **2015**, *12* (9), 1398–1406.
- (25) Sanchez, J. F.; Chiang, Y. M.; Szewczyk, E.; Davidson, A. D.; Ahuja, M.; Oakley, C. E.; Bok, J. W.; Keller, N.; Oakley, B. R.; Wang, C. C. Molecular genetic analysis of the orsellinic acid/F9775 gene cluster of *Aspergillus nidulans*. *Mol. Biosyst.* **2010**, *6* (3), 587–593.
- (26) Li, D.; Xu, Y.; Shao, C. L.; Yang, R. Y.; Zheng, C. J.; Chen, Y. Y.; Fu, X. M.; Qian, P. Y.; She, Z. G.; de Voogd, N. J.; Wang, C. Y. Antibacterial bisabolane-type sesquiterpenoids from the sponge-derived fungus *Aspergillus* sp. *Mar. drugs* **2012**, *10* (12), 234–241.
- (27) Lu, Z.; Zhu, H.; Fu, P.; Wang, Y.; Zhang, Z.; Lin, H.; Liu, P.; Zhuang, Y.; Hong, K.; Zhu, W. Cytotoxic polyphenols from the marine-derived fungus *Penicillium expansum*. *J. Nat. Prod.* **2010**, *73* (5), 911–914.
- (28) Lacey, H. J.; Vuong, D.; Pitt, J. I.; Lacey, E.; Piggott, A. M. Kumbicins A–D: bis-indolyl benzenoids and benzoquinones from an Australian soil fungus, *Aspergillus kumbius*. *Aust. J. Chem.* **2016**, *69* (2), 152–160.
- (29) El-Elimat, T.; Figueroa, M.; Ehrmann, B. M.; Cech, N. B.; Pearce, C. J.; Oberlies, N. H. High-resolution MS, MS/MS, and UV database of fungal secondary metabolites as a dereplication protocol for bioactive natural products. *J. Nat. Prod.* **2013**, *76* (9), 1709–1716.
- (30) Paguigan, N. D.; El-Elimat, T.; Kao, D.; Raja, H. A.; Pearce, C. J.; Oberlies, N. H. Enhanced dereplication of fungal cultures via use of mass defect filtering. *J. Antibiot.* **2017**, *70*, 553–561.
- (31) Aron, A. T.; Gentry, E. C.; McPhail, K. L.; Nothias, L. F.; Nothias-Espósito, M.; Bouslimani, A.; Petras, D.; Gauglitz, J. M.; Sikora, N.; Vargas, F.; et al. Reproducible molecular networking of

- untargeted mass spectrometry data using GNPS. *Nat. Protoc.* **2020**, *15*, 1954–1991.
- (32) Wang, M.; Carver, J. J.; Phelan, V. V.; Sanchez, L. M.; Garg, N.; Peng, Y.; Nguyen, D. D.; Watrous, J.; Kaponov, C. A.; Luzzatto-Knaan, T.; et al. Sharing and community curation of mass spectrometry data with Global Natural Products Social Molecular Networking. *Nat. Biotechnol.* **2016**, *34*, 828–837.
- (33) Ernst, M.; Kang, K. B.; Caraballo-Rodríguez, A. M.; Nothias, L. F.; Wandy, J.; Chen, C.; Wang, M.; Rogers, S.; Medema, M. H.; Dorrestein, P. C.; Van Der Hoof, J. J. MolNetEnhancer: Enhanced molecular networks by integrating metabolome mining and annotation tools. *Metabolites* **2019**, *9* (7), 144.
- (34) Zhou, G. X.; Wijeratne, E. K.; Bigelow, D.; Pierson, L. S.; VanEtten, H. D.; Gunatilaka, A. L. Aspochalasins I, J, and K: three new cytotoxic cytochalasins of *Aspergillus flavipes* from the rhizosphere of *Ericameria laricifolia* of the Sonoran Desert. *J. Nat. Prod.* **2004**, *67* (3), 328–332.
- (35) Si, Y.; Tang, M.; Lin, S.; Chen, G.; Feng, Q.; Wang, Y.; Hua, H.; Bai, J.; Wang, H.; Pei, Y. Cytotoxic cytochalasins from *Aspergillus flavipes* PJ03–11 by OSMAC method. *Tetrahedron Lett.* **2018**, *59* (18), 1767–1771.
- (36) Zhou, H.; Sun, X.; Li, N.; Che, Q.; Zhu, T.; Gu, Q.; Li, D. Isoindolone-containing meroperpenoids from the endophytic fungus *Emericella nidulans* HDN12–249. *Org. Lett.* **2016**, *18* (18), 4670–4673.
- (37) Shibata, S.; Sankawa, U.; Taguchi, H.; Yamasaki, K. Biosynthesis of Natural Products. III. Biosynthesis of erythrokyrine, a coloring matter of *Penicillium islandicum* SOPP. *Chem. Pharm. Bull.* **1966**, *14* (5), 474–478.
- (38) Li, X. D.; Li, X. M.; Xu, G. M.; Zhang, P.; Wang, B. G. Antimicrobial phenolic bisabolanes and related derivatives from *Penicillium aculeatum* SD-321, a deep sea sediment-derived fungus. *J. Nat. Prod.* **2015**, *78* (4), 844–849.
- (39) Glauser, G.; Gindro, K.; Fringeli, J.; De Joffrey, J. P.; Rudaz, S.; Wolfender, J. L. Differential analysis of mycoalexins in confrontation zones of grapevine fungal pathogens by ultrahigh pressure liquid chromatography/time-of-flight mass spectrometry and capillary nuclear magnetic resonance. *J. Agr. Food Chem.* **2009**, *57* (4), 1127–1134.
- (40) Hinkley, S. F.; Jiang, J.; Mazzola, E. P.; Jarvis, B. B. Atranes: Novel diterpenoids from the toxigenic mold *Stachybotrys atra*. *Tetrahedron Lett.* **1999**, *40* (14), 2725–2728.
- (41) Jarvis, B. B.; Salemme, J.; Morales, A. *Stachybotrys* toxins. I. *Nat. Toxins* **1995**, *3* (1), 10–16.
- (42) Ayer, W. A.; Miao, S. Secondary metabolites of the aspen fungus *Stachybotrys cylindrospora*. *Can. J. Chem.* **1993**, *71* (4), 487–493.
- (43) Andersen, B.; Nielsen, K. F.; Thrane, U.; Szaro, T.; Taylor, J. W.; Jarvis, B. B. Molecular and phenotypic descriptions of *Stachybotrys chlorohalonata* sp. nov. and two chemotypes of *Stachybotrys chartarum* found in water-damaged buildings. *Mycologia* **2003**, *95* (6), 1227–1238.
- (44) Semeiks, J.; Borek, D.; Otwinowski, Z.; Grishin, N. V. Comparative genome sequencing reveals chemotype-specific gene clusters in the toxigenic black mold *Stachybotrys*. *BMC Genomics* **2014**, *15* (1), 590.
- (45) Jagels, A.; Lindemann, V.; Ulrich, S.; Gottschalk, C.; Cramer, B.; Hübner, F.; Gareis, M.; Humpf, H. U. Exploring secondary metabolite profiles of *Stachybotrys* spp. by LC-MS/MS. *Toxins* **2019**, *11* (3), 133.
- (46) Ballantine, J. A.; Hassall, C. H.; Jones, G. Asperugin, a metabolite associated with abnormal morphology of *Aspergillus rugulosus*. *Tetrahedron Lett.* **1964**, *5* (49), 3739–3740.
- (47) Bao, Y. R.; Chen, G. D.; Wu, Y. H.; Li, X. X.; Hu, D.; Liu, X. Z.; Li, Y.; Yao, X. S.; Gao, H. Stachybisbins A and B, the first cases of seco-bisabosquals from *Stachybotrys bisbyi*. *Fitoterapia* **2015**, *105*, 151–155.
- (48) Arora, D.; Gupta, P.; Jaglan, S.; Roullier, C.; Grovel, O.; Bertrand, S. Expanding the chemical diversity through microorganisms co-culture: Current status and outlook. *Biotechnol. Adv.* **2020**, *40*, 107521.
- (49) Knowles, S. L.; Raja, H. A.; Roberts, C. D.; Oberlies, N. H. Fungal–fungal co-culture: a primer for generating chemical diversity. *Nat. Prod. Rep.* **2022**, DOI: 10.1039/D1NP00070E.
- (50) Li, H. L.; Li, X. M.; Yang, S. Q.; Cao, J.; Li, Y. H.; Wang, B. G. Induced terreins production from marine red algal-derived endophytic fungus *Aspergillus terreus* EN-539 co-cultured with symbiotic fungus *Paecilomyces lilacinus* EN-531. *J. Antibiot.* **2020**, *73* (2), 108–111.
- (51) Nonaka, K.; Chiba, T.; Suga, T.; Asami, Y.; Iwatsuki, M.; Masuma, R.; Ōmura, S.; Shiomi, K. Coculnol, a new penicillic acid produced by a coculture of *Fusarium solani* FKI-6853 and *Talaromyces* sp. FKA-65. *J. Antibiot.* **2015**, *68* (8), 530–532.
- (52) Sassa, T.; Hayakari, S.; Ikeda, M.; Miura, Y. Plant growth inhibitors produced by fungi Part I. Isolation and identification of penicillic acid and dihydropenicillic acid. *Agr. Biol. Chem.* **1971**, *35* (13), 2130–2131.
- (53) Li, H. J.; Cai, Y. T.; Chen, Y. Y.; Lam, C. K.; Lan, W. J. Metabolites of marine fungus *Aspergillus* sp. collected from soft coral *Sarcophyton tortuosum*. *Chem. Res. Chin. Univ.* **2010**, *26* (3), 415–419.
- (54) Frisvad, J. C. A critical review of producers of small lactone mycotoxins: patulin, penicillic acid and moniliformin. *World Mycotoxin J.* **2018**, *11* (1), 73–100.
- (55) Balouiri, M.; Sadiki, M.; Ibsouda, S. K. Methods for in vitro evaluating antimicrobial activity: A review. *J. Pharm. Anal.* **2016**, *6* (2), 71–79.
- (56) Clinical and Laboratory Standards Institute, <https://clsi.org/standards/?page=1&sort=date&sortdir=desc&subcat=&area=>.
- (57) Cruz-Muñoz, M. Y.; López-Jacome, L. E.; Hernández-Durán, M.; Franco-Cendejas, R.; Licona-Limón, P.; Ramos-Balderas, J. L.; Martínez-Vázquez, M.; Belmont-Díaz, J. A.; Wood, T. K.; García-Contreras, R. Repurposing the anticancer drug mitomycin C for the treatment of persistent *Acinetobacter baumannii* infections. *Int. J. Antimicrob. Agents* **2017**, *49* (1), 88–92.
- (58) Cho, S.; Lee, H. S.; Franzblau, S. Microplate Alamar Blue Assay (MABA) and Low Oxygen Recovery Assay (LORA) for *Mycobacterium tuberculosis*. In *Mycobacteria Protocols*; Parish, T., Roberts, D. M., Eds.; Springer: New York, 2015; pp 281–292.
- (59) Gao, W.; Kim, J.-Y.; Anderson, J. R.; Akopian, T.; Hong, S.; Jin, Y.-Y.; Kandror, O.; Kim, J.-W.; Lee, I.-A.; Lee, S.-Y.; et al. The cyclic peptide ecumicin targeting ClpC1 is active against *Mycobacterium tuberculosis* in vivo. *Antimicrob. Agents Chemother.* **2015**, *59*, 880–889.
- (60) Nandikolla, A.; Srinivasarao, S.; Khetmalis, Y. M.; Kumar, B. K.; Murugesan, S.; Shetye, G.; Ma, R.; Franzblau, S. G.; Sekhar, K. V. G. C. Design, synthesis and biological evaluation of novel 1, 2, 3-triazole analogues of Imidazo-[1, 2-a]-pyridine-3-carboxamide against *Mycobacterium tuberculosis*. *Toxicol. In Vitro* **2021**, *74*, 105137.
- (61) Raja, H. A.; Miller, A. N.; Pearce, C. J.; Oberlies, N. H. Fungal identification using molecular tools: a primer for the natural products research community. *J. Nat. Prod.* **2017**, *80* (3), 756–770.
- (62) Schoch, C. L.; Seifert, K. A.; Huhndorf, S.; Robert, V.; Spouge, J. L.; Levesque, C. A.; Chen, W.; Bolchacova, E.; Voigt, K.; Crous, P. W.; Miller, A. N.; Wingfield, M. J.; Aime, M. C.; An, K.-D.; Bai, F.-Y.; Barreto, R. W.; Begerow, D.; Bergeron, M.-J.; Blackwell, M.; Boekhout, T.; Bogale, M.; Boonyuen, N.; Burgaz, A. R.; Buyck, B.; Cai, L.; Cai, Q.; Cardinali, G.; Chaverri, P.; Coppins, B. J.; Crespo, A.; Cubas, P.; Cummings, C.; Damm, U.; de Beer, Z. W.; de Hoog, G. S.; Del-Prado, R.; Dentinger, B.; Dieguez-Urbeondo, J.; Divakar, P. K.; Douglas, B.; Duenas, M.; Duong, T. A.; Eberhardt, U.; Edwards, J. E.; Elshahed, M. S.; Fliiegerova, K.; Furtado, M.; Garcia, M. A.; Ge, Z.-W.; Griffith, G. W.; Griffiths, K.; Groenewald, J. Z.; Groenewald, M.; Grube, M.; Gryzenhout, M.; Guo, L.-D.; Hagen, F.; Hambleton, S.; Hamelin, R. C.; Hansen, K.; Harrold, P.; Heller, G.; Herrera, C.; Hirayama, K.; Hirooka, Y.; Ho, H.-M.; Hoffmann, K.; Hofstetter, V.; Hognabba, F.; Hollingsworth, P. M.; Hong, S.-B.; Hosaka, K.; Houbaken, J.; Hughes, K.; Huhtinen, S.; Hyde, K. D.; James, T.; Johnson, E. M.; Johnson, J. E.; Johnston, P. R.; Jones, E.B. G.; Kelly, L. J.; Kirk, P. M.; Knapp, D. G.; Koljalg, U.; Kovacs, G. M.; Kurtzman, C. P.; Landvik, S.; Leavitt, S. D.; Ligenstoffer, A. S.; Liimatainen, K.

Lombard, L.; Luangsa-ard, J. J.; Lumbsch, H. T.; Maganti, H.; Maharachchikumbura, S. S. N.; Martin, M. P.; May, T. W.; McTaggart, A. R.; Methven, A. S.; Meyer, W.; Moncalvo, J.-M.; Mongkolsamrit, S.; Nagy, L. G.; Nilsson, R. H.; Niskanen, T.; Nyilasi, I.; Okada, G.; Okane, I.; Olariaga, I.; Otte, J.; Papp, T.; Park, D.; Petkovits, T.; Pino-Bodas, R.; Quaedvlieg, W.; Raja, H. A.; Redecker, D.; Rintoul, T. L.; Ruibal, C.; Sarmiento-Ramirez, J. M.; Schmitt, I.; Schußler, A.; Shearer, C.; Sotome, K.; Stefani, F. O.P.; Stenroos, S.; Stielow, B.; Stockinger, H.; Suetrong, S.; Suh, S.-O.; Sung, G.-H.; Suzuki, M.; Tanaka, K.; Tedersoo, L.; Telleria, M. T.; Tretter, E.; Untereiner, W. A.; Urbina, H.; Vagvolgyi, C.; Vialle, A.; Vu, T. D.; Walther, G.; Wang, Q.-M.; Wang, Y.; Weir, B. S.; Weiß, M.; White, M. M.; Xu, J.; Yahr, R.; Yang, Z. L.; Yurkov, A.; Zamora, J.-C.; Zhang, N.; Zhuang, W.-Y.; Schindel, D. Nuclear ribosomal internal transcribed spacer (ITS) region as a universal DNA barcode marker for Fungi. *Proc. Natl. Acad. Sci. U. S. A.* **2012**, *109* (16), 6241–6246.

(63) Goncalves, V. N.; Vaz, A. B. M.; Rosa, C. A.; Rosa, L. H. Diversity and distribution of fungal communities in lakes of Antarctica. *FEMS Microbiol. Ecol.* **2012**, *82* (2), 459–471.

(64) Aparicio-Cuevas, M. A.; Rivero-Cruz, I.; Sánchez-Castellanos, M.; Menéndez, D.; Raja, H. A.; Joseph-Nathan, P.; González, M. C.; Figueroa, M. Dioxomorpholines and derivatives from a marine-facultative *Aspergillus* species. *J. Nat. Prod.* **2017**, *80* (8), 2311–2318.

(65) Aparicio-Cuevas, M. A.; González, M. C.; Raja, H. A.; Rivero-Cruz, I.; Kurina, S. J.; Burdette, J. E.; Oberlies, N. H.; Figueroa, M. Metabolites from the marine-facultative *Aspergillus* sp. MEXU 27854 and *Gymnoascus hyalinosporus* MEXU 29901 from Caleta Bay, Mexico. *Tetrahedron Lett.* **2019**, *60* (25), 1649–1652.

(66) Martínez-Cárdenas, A.; Cruz-Zamora, Y.; Fajardo-Hernández, C. A.; Villanueva-Silva, R.; Cruz-García, F.; Raja, H. A.; Figueroa, M. Genome Mining and Molecular Networking-Based Metabolomics of the Marine Facultative *Aspergillus* sp. MEXU 27854. *Molecules* **2021**, *26* (17), 5362.

(67) Villanueva-Silva, R.; Velez, P.; Riquelme, M.; Fajardo-Hernández, C. A.; Martínez-Cárdenas, A.; Arista-Romero, A.; Wan, B.; Ma, R.; Qader, M.; Franzblau, S. G.; Figueroa, M. Chemical Diversity and Antimicrobial Potential of Cultivable Fungi from Deep-Sea Sediments of the Gulf of Mexico. *Molecules* **2021**, *26* (23), 7328.

(68) Sheldrick, G. M. *SHELXL-2018: Program for Crystal Structure Refinement*; University of Göttingen: Göttingen, Germany, 2018.

(69) Rigaku Oxford Diffraction. *CrysAlisPro: Empirical absorption correction using spherical harmonics, implemented in SCALE3 ABSPACK scaling algorithm*; Agilent Technologies Ltd.: Yarnton, U.K., 2019.

(70) Mohimani, H.; Gurevich, A.; Shlemov, A.; Mikheenko, A.; Korobeynikov, A.; Cao, L.; Shcherbin, E.; Nothias, L. F.; Dorrestein, P. C.; Pevzner, P. A. Dereplication of microbial metabolites through database search of mass spectra. *Nat. Commun.* **2018**, *9*, 4035.

(71) Shannon, P.; Markiel, A.; Ozier, O.; Baliga, N. S.; Wang, J. T.; Ramage, D.; Amin, N.; Schwikowski, B.; Ideker, T. Cytoscape: a software environment for integrated models of biomolecular interaction networks. *Genome Res.* **2003**, *13* (11), 2498–2504.

(72) Dictionary of Natural Products 30.1. Available online at <https://dnp.chemnetbase.com/faces/chemical/ChemicalSearch.xhtml> (accessed on 26 November 2021).

(73) SciFinder—CAS. Available online at <https://scifinder.cas.org> (accessed on 26 November 2021).

(74) Sumner, L. W.; Amberg, A.; Barrett, D.; Beale, M. H.; Beger, R.; Daykin, C. A.; Fan, T.W.-M.; Fiehn, O.; Goodacre, R.; Griffin, J. L.; et al. Proposed minimum reporting standards for chemical analysis: Chemical Analysis Working Group (CAWG) Metabolomics Standards Initiative (MSI). *Metabolomics* **2007**, *3*, 211–221.