

MAIT cells are imprinted by the microbiota in early life and promote tissue repair

Michael G. Constantinides, Verena M. Link, Samira Tamoutounour, Andrea C. Wong, P. Juliana Perez-Chaparro, Seong-Ji Han, Y. Erin Chen, Kelin Li, Sepideh Farhat, Antonin Weckel, Siddharth R. Krishnamurthy, Ivan Vujkovic-Cvijin, Jonathan L. Linehan, Nicolas Bouladoux, E. Dean Merrill, Sobhan Roy, Daniel J. Cua, Erin J. Adams, Avinash Bhandoola, Tiffany C. Scharschmidt, Jeffrey Aubé, Michael A. Fischbach, Yasmine Belkaid*

INTRODUCTION: The microbiota promotes the maturation and homeostasis of the immune system, in part through the release of microbial products. Early-life microbial colonization has been shown to play a fundamental role in the development of the immune system and imparts long-lasting effects on host fitness. However, despite the importance of this early host-microbiota dialogue, little is known about the microbial-derived signals and antigens involved. This question is of particular importance for mucosal-associated invariant T (MAIT) cells, which are predominantly located in tissues colonized by the microbiota and characterized by their recognition of microbial-derived intermediates of vitamin B2 (riboflavin) synthesis. Because riboflavin synthesis is broadly conserved among bacteria and fungi, MAIT cells are


thought to be particularly dependent on the microbiota.

RATIONALE: Although MAIT cells are the predominant innate-like lymphocyte subset in humans, there is remarkable variability in their abundance between individuals. How commensal-derived antigens contribute to the variability in MAIT cell abundance and their function has not been established. Furthermore, the extent to which MAIT cells promote tissue physiology remains to be determined.

RESULTS: MAIT cells were highly abundant within human and mouse skin. However, genetically identical mice housed in distinct cages showed striking variability in the proportion of MAIT cells. By contrast, animals in the same cage had similar frequencies of MAIT cells,

Early-life microbial exposure imprints the abundance of MAIT cells, and subsequent interactions with the microbiota modulate their function.

(Top) A lack of microbial exposure in early life, results in an absence of MAIT cells in adults. (Middle) Early colonization with a low abundance of vitamin B2-synthesizing (synth.) commensals induces minimal MAIT cell development, whereas (Bottom) a high abundance of these microbes imprints a high frequency of tissue-resident MAIT cells in adults. MAIT cells are selected on double-positive (DP) thymocytes and require IL-23 for their development and/or accumulation. Subsequent recognition of commensals requires MR1-mediated presentation of riboflavin derivatives and IL-18. Microbial stimulation of MAIT cells prompts IL-1-dependent IL-17A production and reinforces a tissue repair program inherent to MAIT cells.



supporting the hypothesis that these differences were associated with distinct microbiota. MAIT cells accumulated in barrier tissues between 2 and 3 weeks of age, bolstering the idea that MAIT cells develop during a very specific temporal window and in response to defined microbial exposure. The isolation of early-life intestinal commensals and subsequent colonization of neonatal germ-free mice with defined bacteria induced MAIT cell development. Conversely, colonization later in life failed to promote their development within tissues, indicating that microbial exposure

ON OUR WEBSITE
Read the full article at <http://dx.doi.org/10.1126/science.aax6624>

must occur during an early-life window, imprinting MAIT cell abundance for life. Commensals that induced MAIT cell development were capable of synthesizing riboflavin, demonstrating antigen necessity, whereas the development of MAIT cells after treatment with a riboflavin derivative indicated sufficiency. After their development in response to early-life commensals, MAIT cells represented a dominant type-17 effector subset in the skin, and cutaneous MAIT cells distinctly expressed a transcriptional program associated with tissue repair. Cutaneous MAIT cells were tissue-resident and required the cytokine interleukin-23 (IL-23) for their homeostasis. These lymphocytes were capable of responding locally to skin commensals in a manner that required IL-1 and IL-18 as well as antigen presentation mediated by the major histocompatibility complex molecule MR1. MR1-mediated presentation of riboflavin metabolites was necessary and sufficient for MAIT cell recognition of skin commensals and further enhanced the tissue-repair program of these lymphocytes. Within the skin, MAIT cells were distinctly localized at the interface of the dermis and epidermis, in close proximity to the basal layer. Topical application of a riboflavin derivative selectively increased MAIT cells in the skin and was sufficient to promote cutaneous wound healing, demonstrating that MAIT cells contribute to skin physiology.

CONCLUSION: Our work demonstrates how early microbial encounters have long-term effects on the composition of the immune system. We show that MAIT cells are induced during a specific early-life window in response to riboflavin-synthesizing commensals. This phenomenon permanently imprints the abundance of this subset in tissues, controlling tissue repair and homeostasis.

The list of author affiliations is available in the full article online.
*Corresponding author. Email: ybelkaid@niaid.nih.gov
Cite this article as M. Constantinides et al., *Science* 366, eaax6624 (2019). DOI: 10.1126/science.aax6624

MAIT cells are imprinted by the microbiota in early life and promote tissue repair

Michael G. Constantinides¹, Verena M. Link¹, Samira Tamoutounour^{1*}, Andrea C. Wong², P. Juliana Perez-Chaparro³, Seong-Ji Han¹, Y. Erin Chen⁴, Kelin Li⁵, Sepideh Farhat⁶, Antonin Weckel⁶, Siddharth R. Krishnamurthy¹, Ivan Vujkovic-Cvijin¹, Jonathan L. Linehan^{1,†}, Nicolas Bouladoux^{1,3}, E. Dean Merrill¹, Sobhan Roy⁷, Daniel J. Cua^{8,‡}, Erin J. Adams⁷, Avinash Bhandoola⁹, Tiffany C. Scharschmidt⁶, Jeffrey Aubé⁵, Michael A. Fischbach⁴, Yasmine Belkaid^{1,3§}

How early-life colonization and subsequent exposure to the microbiota affect long-term tissue immunity remains poorly understood. Here, we show that the development of mucosal-associated invariant T (MAIT) cells relies on a specific temporal window, after which MAIT cell development is permanently impaired. This imprinting depends on early-life exposure to defined microbes that synthesize riboflavin-derived antigens. In adults, cutaneous MAIT cells are a dominant population of interleukin-17A (IL-17A)-producing lymphocytes, which display a distinct transcriptional signature and can subsequently respond to skin commensals in an IL-1-, IL-18-, and antigen-dependent manner. Consequently, local activation of cutaneous MAIT cells promotes wound healing. Together, our work uncovers a privileged interaction between defined members of the microbiota and MAIT cells, which sequentially controls both tissue-imprinting and subsequent responses to injury.

Every barrier site harbors a distinct community of commensal microbes, known as the microbiota, that controls host physiology (1). These microbes promote the maturation and homeostasis of the immune system, in part through the release of microbial products, including metabolites such as short-chain fatty acids, aryl hydrocarbon receptor (AhR) ligands, and polyamines (2). In turn, the immune system modulates the composition of the microbiota and maintains the segregation of commensals by sustaining barrier tissue function (1).

Early-life microbial colonization has been shown to play a fundamental role in the development of the immune system and imparts

long-lasting effects on host fitness. For example, neonatal colonization of the lungs promotes tolerance to allergens in adult mice through the induction of regulatory T cells (T_{reg} cells) (3), whereas colonization of neonatal skin similarly induces T_{reg} cell-mediated tolerance to commensal microbes (4). During weaning, the microbiota induces a transient up-regulation of phosphorylated signal transducer and activator of transcription 3 (STAT3) in epithelial cells and innate lymphoid cells (ILCs) (5), as well as an increase in colonic T_{reg} cells (6), which can affect host metabolism and susceptibility to inflammatory disorders. High microbial diversity within the intestine during early life is necessary to impede immunoglobulin E (IgE) class switching of mucosal B cells, reducing susceptibility to orally induced anaphylaxis (7). Furthermore, early colonization with the intestinal commensal *Bacteroides fragilis* inhibits the proliferation of natural killer T (NKT) cells, which prevents oxazolone-induced colitis in adult mice (8, 9). However, despite the importance of this early host-microbiota dialogue, little is known about the microbially derived signals and antigens involved. This question is of particular importance for unconventional T cells, which are predominantly located in tissues colonized by the microbiota and characterized by their recognition of conserved antigenic motifs.

Exhibiting characteristics of both innate and adaptive immunity, innate-like lymphocytes such as mucosal-associated invariant T (MAIT) cells, NKT cells, and $\gamma\delta$ T cells acquire their effector functions during development, which

direct their localization to tissues and enable them to respond immediately upon primary antigen recognition (10). Consequently, within barrier sites, innate-like lymphocytes are likely poised to respond to the microbiota and play a dominant role in mediating host-commensal interactions. Both $\gamma\delta$ T cells and $CD8^+$ T cells restricted by the nonclassical major histocompatibility complex (MHC)-Ib molecule H2-M3 respond to skin commensals (11, 12), whereas $\gamma\delta$ T cells can mediate both beneficial and deleterious effects of the microbiota in the intestines and lungs (13, 14). MAIT cells express semi-invariant $\alpha\beta$ T cell receptors that recognize microbial-derived intermediates of vitamin B2 (riboflavin) synthesis presented by the MHC-Ib molecule MRI (15). Because riboflavin synthesis is broadly conserved among bacteria and fungi (16), MAIT cells are thought to be particularly dependent on the microbiota. Germ-free (GF) mice exhibit fewer MAIT cells than do animals housed in specific-pathogen-free (SPF) conditions (17, 18). Although these cells are the predominant innate-like lymphocyte subset in humans, where they compose up to 45% of hepatic lymphocytes (19), surprisingly little is known about them. Although the function of MAIT cells remains largely unclear, in humans these cells can be altered in defined inflammatory or infectious settings (20), and their frequencies show remarkable variability among individuals (21, 22). Although the paucity of MAIT cells in mice has rendered the exploration of their function challenging, MAIT cells have been proposed to promote inflammation and microbial defense (23–25). How commensal-derived antigens contribute to the abundance and function of MAIT cells has not been established. Furthermore, the extent to which MAIT cells promote tissue physiology remains to be determined.

Here, we show that the development of MAIT cells and their long-term frequencies within tissues depend on a very specific developmental window and, in particular, early-life exposure to defined microbial communities enriched in riboflavin-synthesizing bacteria, such as Enterobacteriaceae. This defined microbial exposure must occur during the first few weeks of life, after which MAIT cell development is permanently impaired. After their development, MAIT cells become a dominant, tissue-resident population of interleukin-17A (IL-17A)-producing lymphocytes within the skin and display a transcriptional profile that is distinct from MAIT cells in other tissues. Cutaneous MAIT cells can subsequently respond to defined skin commensals or commensal-derived metabolites in an antigen-dependent manner. Consequently, the local activation of cutaneous MAIT cells promotes wound healing. This work therefore uncovers a privileged interaction between defined members of the microbiota and MAIT cells, resulting in a sequential

¹Metaorganism Immunity Section, Laboratory of Immune System Biology, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Bethesda, MD 20892, USA. ²Immunology Graduate Group, University of Pennsylvania, Philadelphia, PA 19104, USA. ³NIAID Microbiome Program, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Bethesda, MD 20892, USA. ⁴Department of Bioengineering and ChEM-H, Stanford University, Stanford, CA 94305, USA. ⁵Division of Chemical Biology and Medicinal Chemistry, UNC Eshelman School of Pharmacy, University of North Carolina, Chapel Hill, NC 27599, USA. ⁶Department of Dermatology, University of California, San Francisco, CA 94143, USA. ⁷Department of Biochemistry and Molecular Biology, University of Chicago, Chicago, IL 60637, USA. ⁸Merck & Co., Merck Research Laboratories, Palo Alto, CA 94304, USA. ⁹Laboratory of Genome Integrity, Center for Cancer Research, National Cancer Institute, National Institutes of Health, Bethesda, MD 20892, USA.

*Present address: L'Oréal Research and Innovation, Aulnay-sous-Bois, 93600, France. †Present address: Department of Cancer Immunology, Genentech, South San Francisco, CA 94080, USA. ‡Present address: Janssen Research and Development, Spring House, PA 19477, USA.

§Corresponding author. Email: ybelkaid@niaid.nih.gov

dialogue that controls both tissue-imprinting and subsequent responses to injury.

Early-life exposure to riboflavin-synthesizing commensals is required for MAIT cell development

Although MAIT cells are abundant in human tissues (19), they are comparatively infrequent in mouse organs, where they typically represent less than 1% of $\alpha\beta$ T cells (26). Because the skin is characterized by an unusually high frequency of unconventional T cells (Fig. 1A), we assessed the frequency of MAIT cells at this barrier site using an MRI tetramer loaded with the riboflavin derivative 5-(2-oxopropylideneamino)-6-D-ribitylaminouracil (5-OP-RU) (27). Although nonspecific MRI tetramer staining was minimal, as demonstrated by use of a MRI tetramer loaded with 6-formylpterin (6-FP) (fig. S1A), we used 5-OP-RU-loaded MRI tetramers conjugated to two different fluorochromes for increased specificity, when possible (fig. S1B). This analysis revealed that MAIT cells were highly enriched in the skin (Fig. 1, B and C), where they composed up to 40% of $\alpha\beta$ T cells (Fig. 1, D and E). The frequency of MAIT cells in the skin correlated highly with their frequency in other organs (fig. S1C). Furthermore, MAIT cells were also enriched in human skin compared with the reported ~1% in peripheral blood (20), averaging ~2% of CD3⁺ lymphocytes (Fig. 1F). Thus, MAIT cells represent a substantial population of skin lymphocytes, both in mice and humans.

A comparison of mice housed in conventional SPF conditions to GF animals confirmed previous observations that the microbiota is necessary for the presence of MAIT cells in the thymus, spleen, and gut (17, 18) and extended this requirement for MAIT cells in the skin, lungs, and liver (Fig. 1B). Although MAIT cells were highly abundant within the skin, wild-type (WT) SPF mice housed in distinct cages showed striking variability in the proportion of MAIT cells among genetically identical animals, ranging from <5 to 40% of $\alpha\beta$ T cells (Fig. 1, D and E). By contrast, animals in the same cage had similar frequencies of MAIT cells in the skin and other organs (Fig. 1E), supporting the hypothesis that these differences were associated with distinct microbiota.

To establish how the microbiota could modulate MAIT cell development and/or accumulation, we first assessed the kinetics of unconventional T cell accumulation within barrier tissues. Although $\gamma\delta$ T cells and NKT cells were present in barrier tissues of 2-week-old mice, there was a dearth of MAIT cells in the skin, lungs, and intestines (Fig. 1G). MAIT cells accumulated in all barrier tissues assessed by 3 weeks of age, supporting the idea that MAIT cell development occurs during a very specific developmental window and in response

to defined microbial exposure. Therefore, we longitudinally assessed the intestinal microbiota of WT mice during early life using 16S ribosomal RNA (rRNA) gene sequencing (Fig. 1G). At 2 weeks of age, the intestinal microbiota was predominantly composed of Lactobacillaceae, with a transient enrichment of Enterobacteriaceae (Fig. 1G), which is consistent with previous work that has identified an abundance of these two bacterial families in early life (28). Isolation of several bacterial strains from 2-week-old mice, which developed MAIT cells as adults (fig. S1, D and E), allowed us to generate an early-life microbial community composed of two Lactobacillaceae members (*Lactobacillus murinus* and *Lactobacillus johnsonii*), two Enterobacteriaceae members (*Proteus mirabilis* and *Klebsiella oxytoca*), and the common intestinal commensal *Enterococcus faecalis*. To determine whether this five-species community was sufficient to promote MAIT cell development, we colonized GF mice by administering the bacteria at weeks 1, 2, and 3 by means of oral gavage (Fig. 1H). The colonization of neonatal GF mice with our early-life microbial community induced MAIT cell development to the same extent observed in SPF mice harboring high MAIT cell frequencies (Fig. 1I). Thus, this limited community was sufficient to provide the necessary developmental and/or survival signals. Conversely, exposure later in life failed to promote MAIT cell development and accumulation within tissues (Fig. 1, H and I), while still inducing T helper 17 (T_H17) and IL-17A⁺ $\gamma\delta$ T cells (fig. S1F). Similarly, conventionalized mice (adult GF mice cohoused with SPF animals) remained deficient in MAIT cells (Fig. 1I). Thus, exposure to the microbiota during an early-life window has a profound impact on MAIT cell development, and exposure later in life cannot compensate for a lack of neonatal colonization.

Whole-genome sequencing of the five-species community indicated that *P. mirabilis* and *K. oxytoca* have the genes *ribA*, *ribD*, and *pyrP2*, which encode the riboflavin synthesis enzymes necessary to generate the MAIT cell antigen 5-OP-RU, whereas the *Lactobacillus* species and *E. faecalis* do not (Fig. 1J), suggesting that, within our defined community, early colonization with Enterobacteriaceae were responsible for the emergence of the MAIT cell lineage. The monocolonization of neonatal GF mice with *P. mirabilis*, but not *L. johnsonii*, was sufficient to drive MAIT cell development (Fig. 1K). Segmented filamentous bacteria (SFB), a known inducer of T_H17 cells that lacks riboflavin synthesis enzymes (Fig. 1J and fig. S1G) (29, 30), also failed to induce MAIT cells (Fig. 1K). As we observed with the five-species early-life community, the ability of *P. mirabilis* to promote MAIT cell development was dependent on neonatal colonization or exposure from birth, both in the skin and other tissues (fig.

S1H). *P. mirabilis* colonization of both GF neonates and adults was sufficient to induce mature CD24⁻ CD44⁺ MAIT cells in the thymus (fig. S1I), indicating that microbial exposure in early life is required for the development of tissue MAIT cells but not the thymic developmental stages. This corroborates recent work that demonstrates that microbially derived 5-OP-RU can promote the thymic development of MAIT cells in GF mice (31). Enterobacteriaceae, including both *Proteus* and *Klebsiella* species, are enriched in the gut microbiota of both human and mouse neonates and decrease over time (28, 32, 33). Thus, the early abundance of Enterobacteriaceae may play an essential role in the long-term imprinting of MAIT cells.

We next assessed whether exposure to microbial antigens alone during early life was sufficient to induce MAIT cell development and whether such exposure could occur at other barrier sites, such as the skin. To this end, we administered a mixture of 5-amino-6-D-ribitylaminouracil (5-A-RU) and methylglyoxal, which react to form the unstable MAIT cell ligand 5-OP-RU (34), to the skin of GF mice on a weekly basis, beginning at 1 week of age. Topical application of 5-OP-RU was sufficient to induce a significant increase in the number of MAIT cells within the skin (Fig. 1L), without promoting the accumulation of other T cell subsets (fig. S1J). This demonstrated that antigen recognition in the absence of microbial-induced proinflammatory cytokines was adequate for MAIT cell development. In agreement with our observation that adult microbial colonization was insufficient to promote MAIT cell development (Fig. 1I), topical application of 5-OP-RU to the skin of adult GF mice failed to induce MAIT cells (Fig. 1L). Thus, early-life exposure to commensals that produce defined metabolites, or to microbially derived metabolites alone, is required for MAIT cell development and their presence later in life. Although this exposure to microbial antigens can occur at multiple barrier sites, it must take place during a defined period in early life. If not, MAIT cell development in tissues is permanently impaired.

Cutaneous MAIT cells express a type-17 transcriptional program and require homeostatic IL-23

MAIT cells have previously been shown to produce either type-1- or type-17-associated cytokines (18, 26). After stimulation, skin MAIT cells robustly produced IL-17A but not interferon- γ (IFN γ) (Fig. 2A), suggesting that within the skin, these cells are exclusively type-17 lymphocytes. As such, in mice with high MAIT cell frequencies, MAIT cells accounted for the largest proportion of IL-17A⁺ $\alpha\beta$ T cells (Fig. 2B), indicating that these innate-like lymphocytes can represent a dominant type-17 effector subset in the skin.




Fig. 1. Early-life exposure to riboflavin-synthesizing commensals is required for MAIT cell development. (A) Percentage of T cells ($\text{TCR}\beta^+$ or $\text{TCR}\gamma\delta^+$) that are MAIT, NKT, or $\gamma\delta$ T cells in WT mice. SI LP, small intestine lamina propria. (B) Percentage of MAIT cells among $\alpha\beta$ T cells in SPF and GF WT mice. (C) Flow cytometry of $\text{TCR}\beta^+$ lymphocytes from the skin of SPF and GF WT mice. (D and E) Analysis of SPF mice housed in different cages (denoted a to g). (D) Flow cytometry of $\text{TCR}\beta^+$ lymphocytes from the skin of mice housed in the indicated cages. (E) The percentage of MAIT cells among $\alpha\beta$ T cells, with each dot representing an individual mouse. (F) Flow cytometry of $\text{CD}3^+$ lymphocytes from a human skin biopsy. (G) 16S rRNA gene sequencing of feces from WT SPF mice longitudinally sampled from 2 to 6 weeks of age and the number of MAIT, NKT, and $\gamma\delta$ T cells present in WT SPF mice at the corresponding ages. Asterisks denote statistically significant changes in cell number from 2 weeks of age. (H) Neonatal GF mice received oral gavages at 1, 2, and 3 weeks after birth, whereas adult GF mice received oral gavages at 7, 8, and 9 weeks of age. Both were analyzed 5 weeks after the initial gavage. (I) Number of MAIT cells in the skin of GF mice administered oral gavages

of the five-species (5-spp.) community (*P. mirabilis*, *K. oxytoca*, *E. faecalis*, *L. johnsonii*, and *L. murinus*) either as neonates (Neo.) or adults (Ad.) as depicted in (H) or conventionalized (conv.) by cohousing with SPF mice for 3 weeks as adults. (J) Presence of riboflavin synthesis genes in bacteria of the five-species community and SFB denoted with the appropriate color. 5-A-RU, which reacts with methylglyoxal to form the MAIT cell antigen 5-OP-RU, and 6,7-dimethyl-8-(1-D-ribityl)lumazine (RL-6,7-diMe) are denoted in red. (K) Number of MAIT cells in the skin of GF mice monocolonized as neonates, with the indicated bacterial species compared with GF and SPF controls. (L) PBS solution of 1 mM 5-A-RU and 25 mM methylglyoxal (referred to as 5-OP-RU) was topically applied once per week to the skin of GF mice beginning either at 1 week of age (Neo.) or 7 weeks of age (Ad.). Number of MAIT cells in the skin was assessed 5 weeks after the initial application. Flow cytometry gate frequencies and graphs indicate means \pm SEM. Data represent at least two experiments with four or more mice per group. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$, and **** $P < 0.0001$ as calculated with Student's *t* test. "ns" denotes that comparison was not significant.




Fig. 2. Cutaneous MAIT cells express a type-17 transcriptional program and require homeostatic IL-23. (A) Flow cytometry of cytokine production by MAIT cells from the skin of WT mice. (B) Percentage of IL-17A⁺ $\alpha\beta$ T cells within WT mouse skin that were CD4⁺, CD8⁺, NKT, or MAIT cells. (C and D) RNA-seq data of cutaneous MAIT cells and CD4⁺ CD25⁻ T cells. (C) Top 20 GO terms that were enriched in MAIT cells and, for the indicated GO terms, the top 30 genes up-regulated in MAIT cells. Positive (pos.), negative (neg.), and regulation (reg.) are abbreviated. (D) Expression plot normalized to transcripts per million (TPM), with a minimum fold-change of 2 and FDR < 5%. Type-17 genes are denoted in red, and genes associated with Type-1, Type-2, T regulatory, and T helper programs are highlighted in blue. (E) Representative flow cytometry plot of TCR β ⁺ lymphocytes from the skin of *Il23r^{GFP/GFP}* mice. (F) Percentage change of the indicated T cells in the skin of *Il23r^{GFP/GFP}* mice compared with WT controls. Statistics denote

whether the percentage differs significantly from the WT mean (100%). (G) Number of cutaneous MAIT cells in *Il23r^{GFP/GFP}* mice and WT controls. (H) Percentage of MAIT cells in the indicated organs that expressed CD4, CD8, or neither coreceptor [double negative (DN)]. (I and J) scRNA-seq data of MAIT cells sorted from mouse skin, spleen, lung, and liver. Clusters were assigned to "Skin" or "Spleen/Lung/Liver" on the basis of the presence of hashtag oligonucleotides (HTOs) from those tissues. Clusters that did not have a predominance of HTOs from either skin or the other tissues were not assigned ("NA"). (I) UMAP plot displaying the distribution of MAIT cell clusters assigned to skin and the other tissues. (J) UMAP plots depicting expression of the indicated transcripts. Flow cytometry gate frequencies and graphs indicate means \pm SEM. Data represent at least two experiments with four or more mice per group. ***P* < 0.01, ****P* < 0.001, and *****P* < 0.0001 as calculated with Student's *t* test. "ns" denotes that comparison was not significant.

To establish the gene expression profile of cutaneous MAIT cells, we compared them with conventional CD4⁺ T cells from the skin using RNA-sequencing (RNA-seq) (fig. S2A). Gene Ontology (GO) enrichment analysis revealed

that cutaneous MAIT cells express a transcriptional profile consistent with T cell activation, including leukocyte differentiation, migration and cell adhesion, cytokine production, and the extracellular signal-regulated kinase-1 and -2

(ERK1/2) cascade, which is downstream of T cell receptor (TCR) signaling (Fig. 2C). MAIT cells were also enriched in GO terms associated with tissue repair (Fig. 2C), suggesting that these lymphocytes may contribute

to the maintenance or restoration of tissue homeostasis.

As expected, cutaneous MAIT cells expressed the gene *Zbtb16* (Fig. 2D), which encodes the transcription factor promyelocytic leukemia zinc finger (PLZF) and is necessary for their development (18, 26). Consistent with their IL-17A potential, MAIT cells expressed type-17-associated transcripts, including *Il22* and *Rorc*, the gene for the transcription factor retinoic acid receptor-related orphan receptor- γ t (ROR γ t), as well as *Il1r1* and *Il23r*, which encode receptors for IL-1 and IL-23, respectively (Fig. 2D). Analysis of *Il1r1*^{-/-} mice indicated that cutaneous MAIT cells do not require IL-1 signaling for their development (fig. S2, B and C). MAIT cells expressed the highest levels of IL-23R among $\alpha\beta$ T cells (Fig. 2E), and IL-23 receptor-deficient mice (*Il23r*^{gfp/gfp}) exhibited fewer MAIT cells than did WT animals (Fig. 2, F and G, and fig. S2, D and E), supporting the idea that homeostatic IL-23 contributes to MAIT cell development and/or accumulation within tissues. Although SFB is known to promote IL-23 expression (35), SFB did not induce MAIT cells in GF mice (Fig. 1K), indicating that IL-23 is insufficient to promote MAIT cell development in the absence of microbial-derived metabolites. Thus, MAIT cells constitute a substantial IL-17A-producing population within the skin that requires IL-23 signaling for development and/or survival.

Although the MAIT cell TCR does not require the CD4 or CD8 coreceptors to facilitate recognition of antigens presented by MR1, CD4⁺, CD8⁺, and CD4⁻ CD8⁻ double-negative (DN) MAIT cells have been described in mice (26), whereas CD8⁺ and DN subsets exist in humans (36). We found that MAIT cells from the skin and other barrier sites, such as the lungs and gut, were almost exclusively DN (Fig. 2H). Conversely, a large proportion of MAIT cells in the thymus, spleen, and liver expressed either CD4 or CD8. In humans, DN MAIT cells produce more IL-17 and can arise from CD8⁺ MAIT cells in vitro (36), suggesting that the mouse DN MAIT cells at barrier tissues may represent the most mature developmental stage.

We next compared the transcriptional profile of MAIT cells from the skin to other tissues by means of single-cell RNA-sequencing (scRNA-seq). Cutaneous MAIT cells displayed a markedly different transcriptional profile (Fig. 2, I and J, and fig. S2, F and G). In concordance with our previous data (Fig. 2, D and H), skin MAIT cells expressed *Rorc*, *Il1r1*, and *Il23r* and predominantly lacked expression of *Cd4* and *Cd8a* (Fig. 2J). Type-1 MAIT cells expressing *Tbx21*, which encodes T-bet, and *Cxcr3* were present in nonskin tissues, whereas the *Rorc*⁺ MAIT cells expressed *Ccr6* (Fig. 2J). Additionally, scRNA-seq confirmed that cutaneous MAIT cells express multiple genes associated with tissue repair, including *Lgals3*, *F2r*,

Sdc1, and *Npnt* (Fig. 2J). MAIT cells expressing either coreceptor did not cluster distinctly from DN MAIT cells from the same tissue (Fig. 2J), suggesting that the transcriptional differences between the CD4⁺, CD8⁺, and DN subsets did not outweigh their tissue-specific programs.

Skin-resident MAIT cells respond to cutaneous microbes in an IL-1- and IL-18-dependent manner

The skin is readily exposed to the environment and colonized by a diverse microbial community that is second in size only to the intestinal microbiome (37). We have previously shown that topical application of the skin commensal *Staphylococcus epidermidis* promotes the accumulation of conventional CD4⁺ T cells, H2-M3-restricted CD8⁺ T cells, and $\gamma\delta$ T cells within the skin without causing inflammation (11, 12, 38). Although the relative abundance of MAIT cells within tissues is imprinted through early exposure to defined microbes, MAIT cells may also subsequently respond to alterations in the abundance or composition of tissue-resident microbiota—specifically, those capable of synthesizing riboflavin. Using strains of *S. epidermidis* that either induce nonclassical CD8⁺ T cells or not (fig. S3A) (12), we observed that in both cases, topical association expanded the cutaneous DN MAIT cell population and increased their production of IL-17A (Fig. 3, A to D, and fig. S3, B to D). Furthermore, there was a strong positive correlation between the abundance of MAIT cells before and after association with *S. epidermidis* (fig. S3, E and F), indicating that SPF animals that were developmentally imprinted with a low MAIT cell frequency will retain a lower level of MAIT cells after subsequent interactions with the microbiota.

To determine whether the MAIT cell response to *S. epidermidis* resulted from local expansion or required priming in the lymph node, we associated *Lta*^{-/-} mice (39). Although the number of CD4⁺, CD8⁺, and $\gamma\delta$ T cells after *S. epidermidis* association were significantly decreased in the absence of lymph nodes as expected (Fig. 3E), *S. epidermidis* still promoted an expansion of MAIT cells in the skin of *Lta*-deficient mice. This suggested that MAIT cells respond to skin commensals locally, either through antigen recognition and/or cytokine stimulation. To further interrogate whether MAIT cells remain within the skin tissue after stimulation, we associated congenically labeled mice with *S. epidermidis* and performed parabiosis (Fig. 3F). As previously described for NKT cells, subsets of $\gamma\delta$ T cells, and MAIT cells in other tissues (40–43), unconventional T cells in the skin, including MAIT cells, were host-derived (Fig. 3G). Thus, MAIT cells persist in the skin as a tissue-resident population and respond locally to skin commensals.

MAIT cells have been shown to respond to cytokine stimulation in an antigen-independent

manner (44). More generally, the relative contribution of antigen versus cytokines in MAIT cell activation in vivo remains poorly understood. Although homeostatic IL-23 contributed to early MAIT cell development (Fig. 2, F and G, and fig. S2E), blocking the IL-23 receptor during *S. epidermidis* association did not reduce MAIT cell numbers (Fig. 3H). Because MAIT cells are known to up-regulate IL-18R α during development (18), we confirmed that cutaneous MAIT cells expressed IL-18R α (Fig. 3I). Neutralization of IL-18 significantly reduced the MAIT cell response to *S. epidermidis* association (Fig. 3, J to L, and fig. S3G), indicating that IL-18 signaling was required for optimal MAIT cell expansion. We have previously shown that topical application of *S. epidermidis* elicits the release of IL-1 α , which promotes IL-17 responses by $\gamma\delta$ and conventional $\alpha\beta$ T cells (45). Although IL-1 receptor-deficient animals (*Il1r1*^{-/-}) had equivalent numbers of cutaneous MAIT cells as WT mice at steady state and upon *S. epidermidis* application (Fig. 3M), their ability to produce IL-17A was significantly decreased (Fig. 3, N and O). Thus, IL-18 was necessary for the expansion of MAIT cells in response to *S. epidermidis*, and local IL-1 signaling was required for the licensing of IL-17A production.

We next assessed how *S. epidermidis* association affects the transcriptional profile of tissue-resident MAIT cells. RNA-seq analysis revealed that application of *S. epidermidis* led to an enrichment in GO terms associated with leukocyte activation and tissue repair (Fig. 3, P and Q), indicating that skin commensals may reinforce the functional capabilities of cutaneous MAIT cells. In addition to inducing the expression of genes downstream of TCR signaling, such as *Egr1* and *Egr2* (46), *S. epidermidis* association promoted the expression of genes associated specifically with angiogenesis, including angiogenin (*Ang*) and *Hgf* (47, 48), as well as genes that more broadly promote tissue repair, such as *Ceacam1*, *Grn*, *Hmoa1*, *Igfl1*, and *Pdgfa* (Fig. 3R) (49–53). Thus, metabolites produced by defined skin commensals may promote the local expansion and activation of cutaneous MAIT cells and enhance their tissue repair program.

MR1-mediated presentation of riboflavin metabolites is necessary and sufficient for MAIT cell recognition of skin commensals

Although the intestine harbors multiple bacterial families that are unable to generate riboflavin, riboflavin synthesis is highly conserved among skin commensals (Fig. 4A). Nearly all *Staphylococcaceae* species, including our *S. epidermidis* isolates, express riboflavin synthesis genes (Fig. 4, A and B). To assess whether recognition of riboflavin metabolites was necessary for MAIT cell responses within the skin, we generated a mutant strain of *S. epidermidis*




Fig. 3. Skin-resident MAIT cells respond to cutaneous microbes in an IL-1- and IL-18-dependent manner. (A to C) *S. epidermidis* (*S. epi*) LM061 was topically applied to the skin of WT mice on days 0, 2, 4, and 6. Animals were compared with unassociated (control) mice 14 days after the initial application. (A) Flow cytometry of TCRβ⁺ lymphocytes (top), IL-17A production by MAIT cells (middle), and coreceptor expression of MAIT cells (bottom) within the skin. (B) Number of cutaneous MAIT cells. (C) Number of cutaneous IL-17A⁺ MAIT cells. (D) Number of cutaneous MAIT cells after topical association with CD8⁺ T cell-inducing (LM087) and -noninducing (O5001 and LM061) strains of *S. epidermidis*. (E) Both *Lta*^{-/-} and WT mice were topically associated with *S. epidermidis* LM087, and the percentage change of T cells in the skin of *Lta*^{-/-} mice compared with WT controls is depicted. Statistics denote whether the percentage differs significantly from the WT mean (100%). (F) CD45.1 (5.1) and CD45.2 (5.2) mice were topically associated with *S. epidermidis* LM087, conjoined 7 weeks later, and analyzed 13 weeks after parabiosis. (G) Frequency of T cells in the indicated tissues of parabiotic mice that are host-derived. (H) WT mice were injected subcutaneously with 1 mg of either antibody to IL-23R or mlgG1 isotype control 2 days before the initial application of *S. epidermidis* LM061 on day 0 and again on day 6. Number of cutaneous MAIT

cells is depicted. (I) Flow cytometry of TCRβ⁺ lymphocytes from the skin of WT mice. (J to L) WT mice were injected intraperitoneally with either 1 mg of antibody to IL-18 or saline (vehicle) 2 days before the initial application of *S. epidermidis* LM061 on day 0 and again on days 1, 5, 8, and 11. (J) Number of cutaneous MAIT cells and (K) IL-17A⁺ MAIT cells and (L) percentage of MAIT cells that are IL-17A⁺ in mice treated with antibody to IL-18 and control mice (vehicle) that were associated with *S. epidermidis* LM061. (M and N) Number of cutaneous (M) MAIT cells and (N) IL-17A⁺ MAIT cells in *Il1r1*^{-/-} mice and WT controls associated with *S. epidermidis* LM061. (O) Flow cytometry of IL-17A production by MAIT cells within the skin of *S. epidermidis*-associated mice. (P to R) RNA-seq data of cutaneous MAIT cells from mice associated with *S. epidermidis* LM061 and unassociated controls. GO terms enriched in MAIT cells from *S. epidermidis*-associated mice that are related to (P) leukocyte activation and (Q) tissue repair. Positive (pos.), negative (neg.), and regulation (reg.) are abbreviated. (R) Heatmap depicting relative expression (exp.) of genes associated with tissue repair and leukocyte activation (Act.). Flow cytometry gate frequencies and graphs indicate means ± SEM. Data represent at least two experiments with four or more mice per group. **P* < 0.05, ***P* < 0.01, ****P* < 0.001, and *****P* < 0.0001 as calculated with Student's *t* test. "ns" denotes that comparison was not significant.




Fig. 4. MR1-mediated presentation of riboflavin metabolites is necessary and sufficient for MAIT cell recognition of skin commensals. (A) Heatmap displaying the abundance of riboflavin synthesis genes among species in the indicated bacterial families. (B) Presence of riboflavin synthesis genes in strains of *S. epidermidis* denoted with the appropriate color. (C) Number of cutaneous MAIT cells in WT mice associated with either *S. epidermidis* LM087 or mutant *S. epidermidis* LM087 Δ ribD. (D to H) 5-OP-RU was topically applied to the skin of WT mice on days 0, 2, 4, and 6. Animals were compared with untreated (control) mice 14 days after the initial application. (D) Flow cytometry of TCR β^+ lymphocytes (top) and IL-17A production by MAIT cells (bottom) within the skin of 5-OP-RU-treated and control mice. Shown in (E) and (G) are the percentage change of the indicated (E) T cells and (G) IL-17A⁺ T cells in the skin

of mice treated with 5-OP-RU compared with untreated controls. Statistics denote whether the percentage differs significantly from the control mean (100%). Shown in (F) and (H) are the number of cutaneous (F) MAIT cells and (H) IL-17A⁺ MAIT cells in 5-OP-RU-treated and untreated control mice. (I and J) *Mr1^{fl/fl} R26^{creERT2}* and *Mr1^{fl/fl}* littermate controls were injected intraperitoneally with 3 mg of tamoxifen 8, 6, 4, and 2 days before the initial association with *S. epidermidis* LM061 and were analyzed 14 days later. (I) Flow cytometry of TCR β^+ lymphocytes and cytokine production by MAIT cells. (J) Number of cutaneous MAIT cells. Flow cytometry gate frequencies and graphs indicate means \pm SEM. Data represent at least two experiments with four or more mice per group. * $P < 0.05$, ** $P < 0.01$, and *** $P < 0.001$ as calculated with Student's *t* test. "ns" denotes that comparison was not significant.

with a deletion of the *ribD* gene, which is required for the second step of the riboflavin synthesis pathway, preceding generation of 5-OP-RU (27). Although this mutation prevented riboflavin synthesis from guanosine-5'-triphosphate (GTP), the *S. epidermidis* Δ ribD mutant retained the *ribB*, *ribH*, and *ribE* genes necessary to produce MAIT cell antigens from

ribose 5-phosphate (Fig. 4B). WT and Δ ribD mutant strains colonized the skin equivalently (fig. S4A). However, in contrast with the WT *S. epidermidis* strain, association with the *S. epidermidis* Δ ribD mutant failed to induce a MAIT cell response (Fig. 4C), even though the CD8⁺ and $\gamma\delta$ T cell responses were conserved (fig. S4, B and C). Thus, these results

support the idea that production of riboflavin intermediates by *S. epidermidis* is required for local MAIT cell activation and that *Staphylococcus* species generate riboflavin exclusively from GTP. To assess whether exposure to microbial antigens was sufficient to activate MAIT cells within the skin, we topically applied 5-OP-RU to SPF mice. Application of




Fig. 5. MAIT cells promote tissue repair. (A) Representative confocal microscopy images of the skin of *Il17a^{cre} R26-STOP-YFP* mice that received topical 5-OP-RU and untreated controls. Shown is a projection along the z axis (top) or x axis (bottom). Colocalization of yellow fluorescent protein (YFP) and TCRβ is depicted in green, whereas the colocalization of YFP and either CD4 or CD8 is in red. White dashed lines denote the dermal-epidermal interface. Scale bars, 50 μm. (B) Flow cytometry of Thy1.2⁺ lymphocytes within the dermis and epidermis of WT mice. (C) Flow cytometry of TCRβ⁺ lymphocytes from the skin of *Tcrd*^{-/-} and WT mice either associated with *S. epidermidis* LM061 or left untreated. The TCRβ^{hi} cells in the *Tcrd*^{-/-} samples are DETCs that express αβ TCRs. (D) Number of cutaneous T cells in *Tcrd*^{-/-} and WT mice after association with *S. epidermidis* LM061. (E and F) *S. epidermidis* LM061 was topically applied to the backs of MAIT cell-deficient *Mr1*^{-/-} *Tcrd*^{-/-} mice and *Tcrd*^{-/-} littermates on days 0, 2, 4, and 6. Punch biopsies were taken through the back skin 12 days after

the initial application, and animals were assessed 5 days later. (E) Representative immunofluorescence images of back skin wounds 5 days after punch biopsies. Tissue sections were immunolabeled with Keratin 14 (red), which stains the advancing epidermal tongues (demarcated with white dashed lines) during re-epithelialization of the wounds. Scale bars, 1 mm. (F) Quantification of the epidermal tongue length 5 days after wounding, with each dot representing the measured length of an individual epidermal tongue. (G) 5-OP-RU was topically applied to the backs of WT mice on days 0, 2, 4, and 6. Punch biopsies were taken through the back skin 12 days after the initial application, and animals were assessed 5 days later, when epidermal tongue lengths were quantified. Flow cytometry gate frequencies and graphs indicate means ± SEM. Data represent at least two experiments with four or more mice per group. **P* < 0.05 and ***P* < 0.01 as calculated with Student's *t* test. "ns" denotes that comparison was not significant.

antigen alone resulted in a significant increase of cutaneous IL-17A⁺ MAIT cells, with minimal effects on other T cells subsets (Fig. 4, D to H, and fig. S4, D and E). As with *S. epidermidis*, the MAIT cell response to 5-OP-RU resulted from local expansion (fig. S4, F and G), without an accompanying expansion of MAIT cells in other tissues (figs. S4, H to M).

To further interrogate whether antigen presentation is necessary for the MAIT cell response to skin commensals, we generated *Mr1^{fl/fl}* conditional knock-out mice by inserting LoxP sites flanking exon 3 (fig. S4N), resulting in the elimination of the α2 helix, which together with the α1 helix composes the ligand-binding cleft of MR1 (15). Although previous work has suggested that MAIT cells are selected on double-positive (DP) thymocytes similarly to NKT cells (54), this study used TCR transgenic T cells, which have been shown to display an unusual PLZF⁺ CD44⁻ phenotype (26). Therefore, whether thymocytes are necessary for the selection of WT MAIT cells remained to be determined. Analysis of *Mr1^{fl/fl} Cd4^{cre}* mice revealed a pro-

found reduction in MAIT cells compared with littermate controls, whereas other lineages were unaltered (fig. S4, O and P), indicating that MAIT cells are indeed selected on thymocytes.

Previous studies have suggested that B cells promote MAIT cell development and may present commensal antigens (17, 55). However, *Mr1^{fl/fl} Cd19^{cre}* mice and *Mr1^{fl/fl}* littermates had similar numbers of MAIT cells before and after association with *S. epidermidis* (fig. S4Q). To determine whether the MAIT cell response to *S. epidermidis* required antigen recognition, we used a tamoxifen-inducible Cre recombinase to enable ablation of MR1 after MAIT cell development (56). Treating the resulting *Mr1^{fl/fl} Rosa26^{creERT2}* mice and *Mr1^{fl/fl}* littermates with tamoxifen did not affect MAIT cell numbers 3 weeks later, suggesting that continuous antigen presentation is not necessary to maintain MAIT cells in the skin under steady-state conditions (Fig. 4J). Tamoxifen treatment a week before the *S. epidermidis* association revealed that MR1-mediated antigen presentation is required for the MAIT cell response (Fig. 4, I and

J, and fig. S4, R and S). Thus, the production of riboflavin metabolites by defined skin commensals and presentation of these antigens promotes local expansion and activation of cutaneous MAIT cells. Although cytokines have been shown to be sufficient to activate MAIT cells (44), our results support the idea that in the context of homeostatic responses to the microbiota, the MAIT cell response is strictly TCR-dependent.

MAIT cells promote tissue repair

To explore a potential role for MAIT cells in skin physiology, we first determined their localization within the skin. Because MAIT cells account for the vast majority of IL-17A-producing αβ T cells after topical application of 5-OP-RU (fig. S5A), we imaged the skin of *Il17a^{cre} R26-STOP-YFP* fate-mapping mice by means of confocal microscopy and found that MAIT cells predominantly resided near the interface of the dermis and epidermis, in close proximity to the basal layer (Fig. 5A). Dissociation of the dermis and epidermis confirmed that although

few MAIT cells are present in the epidermal layer, the majority are localized in the dermis (Fig. 5B). This localization contrasted with that of conventional CD4⁺ or CD8⁺ T cells, which localized close to hair follicles and within the epidermis, respectively (38). Together with the tissue remodeling signature of these cells (Figs. 2, C and J, and 3, Q and R), this supported the idea that MAIT cells may regulate tissue responses to injury.

Although alterations in MAIT cell frequencies have been associated with various pathologies, including skin conditions (57, 58), the functions of these cells remain largely unclear. The previous difficulty assessing MAIT cell function in mice may have been due to the low frequency of MAIT cells in mouse tissues other than skin, as well as the potentially overlapping functions and compensatory responses of unconventional T cell subsets. NKT cell-deficient *Cd1d*^{-/-} mice harbor greater numbers of thymic and splenic MAIT cells (18), suggesting that this subset of innate-like lymphocytes competes with the MAIT cell population. However, such a compensatory response was not observed in the skin of *Cd1d*^{-/-} mice (fig. S5, B and C), likely because of the relatively small number of NKT cells in the skin compared with $\gamma\delta$ T cells. Conversely, in the absence of $\gamma\delta$ T cells, MAIT and NKT cell numbers were significantly increased at steady state and after *S. epidermidis* association (Fig. 5, C and D, and fig. S5, D to F). Therefore, $\gamma\delta$ T cells restrict the MAIT cell population at steady state and constrain their response to commensals. This observation provided us with an approach to test the function of MAIT cells when competition with other innate-like lymphocytes is limited, a setting that is relevant to the lymphocyte composition of human skin, where $\gamma\delta$ T cells are far less abundant than in mice (59).

Previous studies from our laboratory and others have indicated that innate-like lymphocytes can promote tissue repair. For instance, CD8⁺ T cells that recognize commensal-derived N-formylated peptides presented by the MHC-Ib molecule H2-M3 accelerate wound closure in mice (12), whereas mouse dendritic epidermal T cells (DETCs) promote epidermal thickening and keratinocyte proliferation (60). We used an experimental model that involved taking a punch biopsy through the mouse back skin 12 days after the initial association with *S. epidermidis* and assessing the re-epithelialization of the wound 5 days later by measuring the epidermal tongue of proliferating keratinocytes (fig. S5G) (12, 61). To minimize confounding effects of H2-M3-restricted CD8⁺ T cells, we topically associated *Tcrd*^{-/-} mice and MAIT cell-deficient *Mr1*^{-/-} *Tcrd*^{-/-} littermates with a strain of *S. epidermidis* that does not induce CD8⁺ T cells (12). After wounding, the epidermal tongue lengths of MAIT cell-sufficient *Tcrd*^{-/-} mice were significantly longer than

those observed in MAIT cell-deficient *Mr1*^{-/-} *Tcrd*^{-/-} animals (Fig. 5, E and F), indicating that MAIT cells promote tissue repair in the absence of compensatory responses by $\gamma\delta$ T cells.

To assess whether an increase in MAIT cells was sufficient to affect skin physiology and response to injury, we directly applied the 5-OP-RU antigen to the skin of WT mice before wounding to selectively increase MAIT cells (Fig. 4E). The topical application of 5-OP-RU significantly increased the epidermal tongue length (Fig. 5G), demonstrating that the microbially derived antigen 5-OP-RU was sufficient to promote wound healing in an intact host.

Discussion

After development, the skin harbors tissue-resident IL-17A⁺ MAIT cells, and exposure to defined skin microbes or microbially derived antigens can further MAIT cell accumulation and strengthen the tissue repair functions of these cells in an antigen-dependent manner. The majority of skin commensals were capable of synthesizing riboflavin-derived metabolites (Fig. 4A), suggesting that these antigens contribute to skin health and homeostasis and may represent new therapeutic approaches for promoting tissue repair.

We also uncovered how early microbial encounters have long-term effects on the composition of the immune system. We show that MAIT cells are induced during a specific early-life window in response to riboflavin-synthesizing commensals, a phenomenon that permanently imprints the abundance of this subset in tissues. Previous work uncovered a role for early-life exposure to epithelial cell-derived butyrophilin-like proteins for V γ 7⁺ $\gamma\delta$ intraepithelial lymphocyte development, gestational colonization for the induction of ILCs, and early colonization with the intestinal commensal *B. fragilis* to constrain the abundance of NKT cells in adults (8, 62, 63). Together, these findings imply that temporal windows of development and permanent imprinting of tissues may represent a general phenomenon for nonclassically restricted T cells. Our work also provides a possible explanation for the remarkable variation in MAIT cell abundance observed in humans (21, 22). In contrast to mice, human T cells develop in utero, which supports the hypothesis that such early MAIT cell imprinting may occur before birth. Functionally mature MAIT cells have been found in second-trimester human fetuses (64). How, in humans, microbiota-derived riboflavin intermediates before and after birth contribute to the final maturation and long-term tissue imprinting of MAIT cells remains to be addressed. Our work proposes that a central strategy used by the immune system to interact with the microbiota may be through the recognition of canonical microbially derived antigens by unconventional T cells, in-

cluding MAIT cells, a dialogue that may play a dominant role in the control of host physiology.

Materials and Methods

Mice

Il17a^{cre} (65), *Il23*^{gfp/gfp} (66), and *Mr1*^{-/-} (17) mice have been previously described and were generously provided by B. Stockinger (Francis Crick Institute), M. Oukka, (Seattle Children's Research Institute), and S. Gillfillan (Washington University School of Medicine), respectively. Wild-type (WT) C57BL/6NTac and BALB/c SPF mice were purchased from Taconic Biosciences. CD45.1 (B6.SJL-Ptpr^c Pepc^b/BoyJ) and *Il17r1*^{-/-} (C57BL/6-[KO]IL1r1) were obtained through the NIAID-Taconic exchange program. GF C57BL/6 mice were bred and maintained in the NIAID Microbiome Program gnotobiotic animal facility. *Cd1d*^{-/-} (B6.129S6-Del(3Cd1d2-Cd1d1)1Sbp/J), *Cd4*^{cre} (Tg(Cd4-cre)1Cwi/Bflu/J), *Cd19*^{cre} (B6.129P2(C)-Cd19^{tm1(cre)Cgn}/J), *Lta*^{-/-} (B6.129S2-Lta^{tm1Dch}/J), *R26-STOP-YFP* (B6.129X1-Gt(ROSA)26Sor^{tm1(EYFP)Cos}/J), *R26*^{creERT2} (B6.129-Gt(ROSA)26Sor^{tm1(cre/ERT2)Tyji}/J), and *Tcrd*^{-/-} (B6.129P2-Tcrd^{tm1Mom}/J) mice were purchased from The Jackson Laboratory. All mice were bred and maintained at an American Association for the Accreditation of Laboratory Animal Care (AAALAC)-accredited animal facility at the NIAID and housed in accordance with the procedures outlined in the Guide for the Care and Use of Laboratory Animals. All experiments were performed at NIAID under an animal study proposal (LISB-19E or LISB-20E) approved by the NIAID Animal Care and Use Committee. Unless otherwise noted, age- and sex-matched mice between 6 and 12 weeks of age were used for each experiment.

Generation of Mr1^{fl/fl} mice

CRISPR guide RNAs (gRNAs) were synthesized following a published protocol (67). The following oligos were used to target the second and third introns of *Mr1*: 5'-TAGGTTAAAGC-CATCTCCCAT-3' and 5'-AAACATGGGAGGATGGCTTTAA-3' (second) and 5'-TAGGCTAGGCATGTTAAGAATG-3' and 5'-AAACCATTCTTAACATGCCTAG-3' (third). Briefly, oligos were annealed and ligated into the pT7-gRNA vector (Addgene). These constructs were linearized and used as templates for in vitro transcription (IVT) using the MEGAscript T7 Transcription Kit (Thermo Fisher Scientific), which yielded the gRNAs. Cas9 mRNA was generated by using linearized pT3TS-nCas9n (Addgene) as the template for IVT with the mMACHINE T3 Transcription Kit (Thermo Fisher Scientific). Both the Cas9 mRNA and the gRNAs were purified using the MEGAclear Transcription Clean-Up Kit (Thermo Fisher Scientific) and eluted in RNase-free water. C57BL/6Ncr embryos were microinjected with Cas9 mRNA (100 ng/ μ L), both gRNAs (100 ng/ μ L), and the following two PAGE-purified ssDNA

oligos (300 ng/ μ L; Integrated DNA Technologies), which contain the LoxP sequences flanked by 75 bases of homology: 5'-CAAAAGTGCTTT-CAGCCATGTCTCCAGTTGTACCACTTGAGAA-GTCTCTCTGTCTCTGTGTTTAAAGCCATCC-TCCATAAATTCGTATAATGTATGCTATACGA-AGTTATCATAGGCTGTATATGTTTCAGGGC-TTCACACCTACCAGAGAATGATTGGCTGT-GAGTTGCTAGAAGATGGCAGCA-3' and 5'-TGGCTAAAGAGGTTCTTGGAAATATGGAAGA-GATACCCTAGAAAGAACAGGTAATGGGAAG-AGAGAACACCTCATATAAATTCGTATAATGT-ATGCTATACGAAGTTATTTAAACATGCCA-GGAAAACCTGGCTTTGTGTCTGACTTCTCTT-CTGCAGGTGATATTCGCCCTGTGCTGAAAT-3' and surgically transferred to recipient females. Progeny were screened by polymerase chain reaction (PCR) amplification with primers 5'-TTTCTGGCTCCTGCGTATCT-3' and 5'-ATTT-CAGCACAGGGCGAATA-3', followed by Sanger sequencing with the primer 5'-CTAATTC-CAAGGCAAGC-3'. The founder was bred to C57BL/6NTac mice to propagate the strain.

Inducible deletion of MR1

Mr1^{fl/fl} R26^{creERT2} and *Mr1^{fl/fl}* littermate controls were injected intraperitoneally with 3 mg of tamoxifen (Sigma-Aldrich) in a corn oil-ethanol (95:5) mixture 8, 6, 4, and 2 days before association with *S. epidermidis* NIHLM061.

Commensal culture and colonization

For the isolation of intestinal commensals, a stool pellet was homogenized in 200 μ L of BHI and 10 μ L of the homogenate was streaked on MacConkey (BD Biosciences) and de Man, Rogosa, and Sharpe (MRS; Sigma-Aldrich) agar plates and incubated at 37°C under aerobic and microaerophilic conditions, respectively. After 48 hours of incubation, colonies bearing distinct morphologies were sub-cultured to obtain pure cultures. Additionally, reiterative 1:10 dilutions in Brain Heart Infusion (BHI) broth allowed isolation of single bacteria in 96-well plates under aerobic conditions. Taxonomic identification of pure isolates was determined with Sanger sequencing. To prepare bacterial cultures for oral gavage, 10 μ L of each frozen strain was propagated on Columbia agar with 5% sheep blood (Thermo Scientific) and incubated under microaerophilic conditions overnight at 36°C. Two to three colonies of each strain were inoculated in Peptone Yeast Glucose (PYG) broth and incubated at 37°C under microaerophilic conditions for 12 hours, except for *Lactobacillus* strains, which were incubated for 24 hours. All cultures were normalized to an optical density (OD) of 1 at 600 nm, mixed in equal volumes when required (5-species community), centrifuged, and resuspended in 10% the volume of PYG broth. Neonates received 20 μ L of the bacterial culture by oral gavage, whereas adult animals received 200 μ L. For the topical association with *S. epidermidis*

NIH05001, NIHLM061, and NIHLM087 (referred to as 05001, LM061, and LM087 in the text) (68), strains were cultured for 18 hours in Tryptic Soy Broth (TSB) at 37°C, to a density of approximately 10⁹ colony-forming units (CFU)/mL. Mice were colonized by topically applying up to 5 mL of culture across the entire skin surface using a sterile swab. Association was repeated every other day, for a total of four times. To quantitate *S. epidermidis* colonization, the ear skin was swabbed with a sterile cotton swab soaked in TSB. Swabs were streaked on blood agar and plates were then incubated at 37°C under aerobic conditions for 18 hours before counting CFUs.

Generation of *S. epidermidis* mutant

The *S. epidermidis* NIHLM087 Δ *ribD* mutant was constructed by using a recently described method (Chen *et al.*, submitted). Approximately 1 kb up- and downstream of the gene *ribD* were amplified with the following primers: 5'-CGGTATCGATAAGCTTGATATCTGGAAT-AGTTTGAGCGTTAA-3' and 5'-GATACCTGTAAACATAGACATTCTACTCAATTGATCA-CCTC-3' and 5'-GAGGTGATCAATTGAGTA-GAATGTCTATGTTTACAGGTATC-3' and 5'-TATAGGGCGAATTGGAGCTCTTACGTGTTAA-CACACCATT-3'. These two flanking regions were assembled into the plasmid pIMAY (Addgene) (69) by using Gibson Assembly (New England Biolabs) (70). The resulting plasmid was transformed into *S. epidermidis* NIHLM087. Temperature shift to 37°C and anhydrotetracycline were then used to select for allelic recombination events that resulted in a *ribD* deletion without a genomic scar. Successful knockouts were confirmed by means of colony PCR and sequencing.

Parabiosis experiments and surgery

Age-matched CD45.1 and CD45.2 mice were topically associated with *S. epidermidis* NIHLM087 and 7 weeks after the initial colonization, parabiosis surgery was performed as described (7). Animals were treated with oral antibiotics for 2 weeks and remained conjoined for 13 weeks before analysis.

In vivo cytokine blockade

Mice were injected subcutaneously with 1 mg of either anti-IL-23R antibody (21A4; Merck) or mIgG1 isotype control (27F11; Merck) 2 d before the initial application of *S. epidermidis* NIHLM061 and again on day 6. Anti-IL-18 monoclonal antibody (SK1L3AE-4) (72) was purified from hybridoma supernatant using Protein G Sepharose beads (Abcam). Mice were injected intraperitoneally with either 1 mg of anti-IL-18 or saline (vehicle) 2 days before the initial application of *S. epidermidis* NIHLM061 and again on days 1, 5, 8, and 11.

Human skin specimens

Normal adult human skin was obtained from five patients at the University of California,

San Francisco, undergoing elective surgery, in which healthy skin was discarded as a routine procedure. The study was conducted in accordance with the Declaration of Helsinki principles.

Human skin processing

Skin samples were stored at 4°C in a sterile container with phosphate-buffered saline (PBS) and gauze until the time of digestion. Subcutaneous fat was removed, skin was minced finely with dissection scissors, and mixed in a six-well plate with 3 mL of digestion buffer consisting of 0.8 mg/mL collagenase type 4 (Worthington), 0.02 mg/mL DNase I (Sigma-Aldrich), 10% fetal bovine serum (FBS), 1% hydroxyethylpiperazine ethane sulfonic acid (HEPES), and 1% penicillin/streptavidin in RPMI medium. Samples were incubated overnight in 5% CO₂ and filtered through a 100- μ m filter with wash buffer (2% FBS, 1% penicillin/streptavidin in RPMI medium).

Mouse tissue processing

To isolate skin cells, ear pinnae were excised and separated into ventral and dorsal sheets, which were digested by placing them dermal side down in RPMI 1640 media supplemented with 2 mM L-glutamine, 1 mM sodium pyruvate, 1 mM nonessential amino acids, 50 mM β -mercaptoethanol, 20 mM HEPES, 100 U/mL penicillin, 100 mg/mL streptomycin, 0.5 mg/mL DNase I (Sigma-Aldrich), and 0.25 mg/mL of Liberase TL purified enzyme blend (Roche), and incubated for an hour and 45 min at 37°C. Digested skin sheets were homogenized by using the Medimachine tissue homogenizer system (BD Biosciences) and passed through 50- μ m filters. To separate the epidermis from the dermis, ear pinnae were first digested for 45 min with 500 CU Dispase (Corning Life Sciences), and curved forceps were used to remove the epidermis, before proceeding with the Liberase TL digestion as previously described. Lungs were diced and incubated in 2 mL of pre-warmed RPMI 1640 containing 1 mg/mL DNase I (Sigma-Aldrich) and 0.25 mg/mL of Liberase TL (Roche) for 45 min in a 37°C water bath. The digested tissue was passed through a 70- μ m filter, centrifuged, resuspended in 5 mL of 40% Percoll (Sigma-Aldrich), and centrifuged again. Thymus, spleen, and liver were dissociated through 70- μ m filters, after which 40% Percoll (Sigma-Aldrich) was used to remove hepatocytes from liver samples. Small intestine lamina propria were processed as previously described (73).

In vitro restimulation

Cells were cultured for 2.5 hours at 37°C in RPMI complete media (RPMI 1640 supplemented with 10% FBS, 2 mM L-glutamine, 1 mM sodium pyruvate, 1 mM nonessential amino acids, 50 mM β -mercaptoethanol, 20 mM HEPES, 100 U/mL penicillin, and 100 mg/mL

streptomycin) containing 50 ng/mL of phorbol myristate acetate (PMA; Sigma-Aldrich), 5 mg/mL ionomycin (Sigma-Aldrich), and a 1:1000 dilution of GolgiPlug (BD Biosciences).

Flow cytometric analysis

Fluorophore-conjugated antibodies that were used are listed in table S1. 5-OP-RU and 6-FP-loaded mMR1 and hMR1 tetramers and PBS57-loaded mCD1d tetramers were provided by the NIH Tetramer Core Facility and cells were stained in RPMI complete media for 1 hour at room temperature. For intracellular cytokine staining, cells were fixed and permeabilized using the Fixation/Permeabilization Solution Kit (BD Biosciences) and stained with fluorophore-conjugated antibodies for at least 60 min at 4°C. For transcription factor staining, cells were fixed and permeabilized with the Foxp3/Transcription Factor Staining Buffer Set (eBioscience) and stained with fluorophore-conjugated antibodies for at least 60 min at 4°C. All staining was performed in the presence of purified anti-mouse CD16/32 and purified rat gamma globulin (Jackson ImmunoResearch). Dead cells were excluded from live samples using 4',6-diamidino-2-phenylindol (DAPI; Sigma-Aldrich), whereas either LIVE/DEAD Fixable Blue Dead Cell Stain Kit (Invitrogen Life Technologies) or Ghost Dye Violet 510 (Tonbo Biosciences) was used in fixed samples. All antibodies were purchased from BD Biosciences, BioLegend, or eBioscience. Cells were acquired on an LSRII flow cytometer (BD Biosciences) running FACSDiva software (BD Biosciences), and data were analyzed by using FlowJo (TreeStar). Unless otherwise indicated, mouse MAIT cells were gated as MR1 tetramer⁺ TCRβ⁺ TCRγδ⁻ CD3⁺ CD90.2⁺ CD45⁺ (CD44⁺ as well in tissues other than skin); NKT cells as CD1d tetramer⁺ TCRβ⁺ TCRγδ⁻ CD3⁺ CD90.2⁺ CD45⁺; CD4⁺ T cells as CD4⁺ CD8β⁻ MR1 tetramer⁻ CD1d tetramer⁻ TCRβ⁺ TCRγδ⁻ CD3⁺ CD90.2⁺ CD45⁺; CD8⁺ T cells as CD8β⁺ CD4⁻ MR1 tetramer⁻ CD1d tetramer⁻ TCRβ⁺ TCRγδ⁻ CD3⁺ CD90.2⁺ CD45⁺; and γδ T cells as TCRγδ⁺ TCRβ⁻ CD3⁺ CD90.2⁺ CD45⁺.

Tetramer enrichment

For the enrichment of MAIT cells, samples were stained with phycoerythrin (PE)-conjugated MR1 tetramer, bound to anti-PE MicroBeads (Miltenyi Biotec), and enriched on an autoMACS Pro Separator (Miltenyi Biotec) using the POSSEL_S program.

Confocal microscopy

Ear pinnae were split with forceps, fixed in 1% paraformaldehyde (Electron Microscopy Sciences) overnight at 4°C, and blocked in PBS containing 1% bovine serum albumin (BSA) and 0.25% Triton X-100 for 2 hours at room temperature. Tissues were first stained with anti-CD4 (RM4-5, eBioscience), anti-CD8α (53-6.7, eBioscience), anti-CD49f (GoH3, eBioscience),

anti-GFP (A21311, Life Technologies), and anti-TCRβ (H57-597, eBioscience) antibodies overnight at 4°C, washed three times with PBS, and then stained with DAPI (Sigma-Aldrich) overnight before being mounted with ProLong Gold (Invitrogen). Ear pinnae images were captured on a Leica TCS SP8 confocal microscope equipped with HyD and PMT detectors and a 40X oil objective (HC PL APO 40X/1.3 oil). Images were analyzed using Imaris (Bitplane).

Wounding and epifluorescence microscopy of wound tissue

Wounding and quantitation of wound healing were performed as previously described (61). Male mice in the telogen phase of the hair cycle were anesthetized with ketamine/ xylazine and their back skin was then shaved using a ChroMini Pro (Moser). A 6-mm biopsy punch was first used to partially perforate the skin and then iris scissors were used to cut the epidermis and dermis along the punch outline, resulting in a circular, full-thickness wound. 5 d after wounding, the skin tissue was excised and fixed in 4% paraformaldehyde for 4 hours at 4°C, incubated overnight in 30% sucrose at 4°C, embedded in OCT compound (Tissue-Tek), frozen on dry ice, and cryo-sectioned (20-μm-thick sections). Sections were fixed in 4% paraformaldehyde for 10 min at room temperature, rinsed with PBS, permeabilized with 0.1% Triton X-100 (Sigma) for 10 min, and blocked for 1 hour at room temperature in blocking buffer (2.5% normal goat serum, 1% BSA, 0.3% Triton X-100 in PBS). Chicken anti-mouse Keratin 14 antibody (Poly9060, BioLegend) was diluted at 1:400 in blocking buffer containing rat gamma globulin and anti-CD16/32 and incubated overnight at 4°C. After washing with PBS, sections were stained with a polyclonal anti-chicken IgY-AlexaFluor647 secondary antibody (Jackson ImmunoResearch) at a 1:800 dilution for 1 hour at room temperature, washed with PBS, stained with DAPI, washed with PBS, and mounted with ProLong Gold (Invitrogen). Wound images were captured with a Leica DMI 6000 widefield epifluorescence microscope equipped with a Leica DFC360X monochrome camera. Tiled and stitched images of wounds were collected using a 20X/0.4 NA dry objective. Images were analyzed using Imaris (Bitplane).

16S rRNA gene sequencing and analysis

Fecal DNA was purified using the MagAttract PowerMicrobiome DNA/RNA kit (Qiagen). Amplification of the V4 hypervariable region of the bacterial 16S rRNA gene was performed using the 515f and 806r primers (515F: 5'-GTGCCAGCMGCCGCGGTAA-3'; 806R: 5'-GGA-CTACHVGGGTWTCTAAT-3'), followed by an additional PCR to append different barcodes to each sample. Amplicons were quantified using Kapa Library Quantification Complete Kit (ROX

Low) (Kapa Biosystems) and pooled at equimolar concentrations before being sequenced on a MiSeq (Illumina) using the v3 MiSeq Reagent kit (Illumina). The DADA2 algorithm (74) was used to denoise raw 16S reads after primer trimming and to tabulate sequence variants. Read counts were converted to relative abundance and taxonomy was assigned to each sequence variant using the RDP Training Set 16. Sequence variants were collapsed by family-level classifications, and families present in greater than 5% abundance in at least one time point were graphically represented, whereas remaining families were collapsed into the category "Other."

Bacterial whole-genome sequencing

For the *E. faecalis*, *K. oxytoca*, *L. johnsonii*, *L. murinus*, and *P. mirabilis* isolates, DNA libraries were prepared by using the paired-ended Nextera DNA Flex Library Prep Kit (Illumina). Sequencing was performed on a NextSeq 500 (Illumina) using the High Output v2 Kit (Illumina).

Assessing the presence of riboflavin synthesis genes

Protein sequences of representative riboflavin biosynthesis genes were downloaded from GenBank: *ribA* (NP_764994), *ribB* (CDOI2415), *ribD* (NP_764996), *ribE* (NP_764995), *ribH* (NP_764993), and *pyr2* (NP_764225). Isolates were either sequenced as described above or genomes were downloaded from National Center for Biotechnology Information (NCBI): *Candidatus Arthromitus* (BioProject PRJDA67835), *S. epidermidis* NIH05001 (BioProject PRJNA62391), *S. epidermidis* NIHLM061 (BioProject PRJNA62355), and *S. epidermidis* NIHLM087 (BioProject PRJNA62349). For taxa that are generally present in the intestinal and skin microbiota, genomes of different strains were downloaded from NCBI: Corynebacteriaceae (txid 1653), Propionibacteriaceae (txid 31957), Staphylococcaceae (txid 90964), Streptococcaceae (txid 1300), Bacteroidaceae (txid 815), Bifidobacteriaceae (txid 31953), Clostridiaceae (txid 31979), Enterobacteriaceae (txid 543), Enterococcaceae (txid 81852), Eubacteriaceae (txid 186806), Hungateiclostridiaceae (txid 2304686), Lachnospiraceae (txid 186803), Lactobacillaceae (txid 33958), Peptococcaceae (txid 186807), Peptostreptococcaceae (txid 186804), Ruminococcaceae (txid 541000), and Tannerellaceae (txid 2005525). These nucleotide scaffolds were used as subjects in a tBLASTn search using representative riboflavin biosynthesis genes, using an e-value <1×10⁻¹⁰. Alignments that met this threshold were considered as homologs, and thus those genomes were considered to contain that gene.

RNA-seq and analysis

The following populations were isolated by means of fluorescence-activated cell sorting

(FACS) from the ear pinnae of C57BL/6NTac mice associated with *S. epidermidis* NIHLM061 and untreated control animals: MAIT cells (MR1 tetramer⁺ TCRβ⁺ TCRγδ⁻ CD90.2⁺ CD45⁺ DAPI⁻) and CD4⁺ T cells (CD4⁺ CD25⁻ CD8β⁻ MR1 tetramer⁻ TCRβ⁺ TCRγδ⁻ CD90.2⁺ CD45⁺ DAPI⁻). RNA was isolated using the Arcturus PicoPure RNA Isolation Kit (Applied Biosystems), cDNA was synthesized using the SMART-Seq v4 Ultra Low Input RNA Kit (Clontech Laboratories), libraries were generated using Nextera XT DNA Library Preparation Kit (Illumina), and the samples were sequenced on a NextSeq 500 (Illumina) using the High Output v2 Kit (Illumina). Fastq files were mapped with STAR (75) to the mouse mm10 reference genome with default parameters. For analysis, the Hypergeometric Optimization of Motif Enrichment (HOMER) program was used (76). Tag directories were created with default parameters. To analyze differential gene expression, reads per gene were counted for two replicates per condition with analyzeRepeats rna with parameters -noadj and -condenseGenes. Subsequently, getDiffExpression.pl was used with default parameters [log2 fold change 1, false discovery rate (FDR) < 5%] using DESeq2 and rlog normalization. For GO term enrichment analysis Metascape (77) was used.

Single-cell RNA-seq and analysis

Lymphocytes from the ear pinnae (skin), spleens, lungs, and livers of pooled WT mice were stained with antibodies against surface markers and TotalSeq-A hashtag oligonucleotide (HTO) antibodies (BioLegend; hashtag 1 for skin, 2 for livers, 3 for lungs, and 4 for spleens). MAIT cells were sorted from each organ as MR1 tetramer⁺ TCRβ⁺ TCRγδ⁻ CD3⁺ CD90.2⁺ B220⁻ CD11b⁻ CD11c⁻ CD49f⁻ CD45⁺ (10,000 cells from skin, spleens, and lungs and 3500 cells from livers). Samples were pooled, encapsulated into droplets using the Chromium Single Cell Controller (10X Genomics), and libraries were prepared using Chromium Single Cell 3' Reagent Kits v2 (10X Genomics). The mRNA library was prepared following the 10X Genomics user guide, while the HTO library was prepared according to published guidelines (78). Libraries were sequenced on a NextSeq 500 with 10% of the lane occupied by the HTO library and the other 90% by the mRNA library.

Data was demultiplexed by using Cell Ranger 3.0.1 resulting in data for 6463 cells. HTO libraries were processed by using CITE-seq-Count (DOI: 10.5281/zenodo.2590196). Subsequently, data was analyzed using Seurat 3.0 (79). Cells with less than 900 or more than 3000 gene counts, more than 5% mitochondrial contamination, or more than 10,000 RNA counts were filtered out, leaving 5678 cells for downstream analysis. Data was normalized, scaled, principal components analysis was performed, and

neighbors were found using five dimensions. Uniform manifold approximation and projection (UMAP) reduction was performed on this dataset with 10 dimensions. HTO were read in and assigned to the single cells and only singlets were used for further analysis leaving 585 cells. To define origins of cells, number of HTOs per cluster were considered and classified as “Skin,” “Spleen/Lung/Liver,” or not assigned (“NA”). To analyze gene expression between tissues, all cells, including cells without HTO from the respective clusters were used.

Statistical analysis

Two-tailed unpaired Student's *t* test was performed by using Prism 7 (GraphPad) to determine statistical significance. **P* < 0.05, ***P* < 0.01, ****P* < 0.001, and *****P* < 0.0001. “ns” denotes that the comparison was not significant.

REFERENCES AND NOTES

1. Y. Belkaid, T. W. Hand, Role of the microbiota in immunity and inflammation. *Cell* **157**, 121–141 (2014). doi: [10.1016/j.cell.2014.03.011](https://doi.org/10.1016/j.cell.2014.03.011); pmid: 24679531
2. M. G. Rooks, W. S. Garrett, Gut microbiota, metabolites and host immunity. *Nat. Rev. Immunol.* **16**, 341–352 (2016). doi: [10.1038/nri.2016.42](https://doi.org/10.1038/nri.2016.42); pmid: 27231050
3. E. S. Gollwitzer *et al.*, Lung microbiota promotes tolerance to allergens in neonates via PD-L1. *Nat. Med.* **20**, 642–647 (2014). doi: [10.1038/nm.3568](https://doi.org/10.1038/nm.3568); pmid: 24813249
4. T. C. Scharnschmidt *et al.*, A wave of regulatory T cells into neonatal skin mediates tolerance to commensal microbes. *Immunity* **43**, 1011–1021 (2015). doi: [10.1016/j.immuni.2015.10.016](https://doi.org/10.1016/j.immuni.2015.10.016); pmid: 26588783
5. K. Mao *et al.*, Innate and adaptive lymphocytes sequentially shape the gut microbiota and lipid metabolism. *Nature* **554**, 255–259 (2018). doi: [10.1038/nature25437](https://doi.org/10.1038/nature25437); pmid: 29364878
6. Z. Al Nabhani *et al.*, A weaning reaction to microbiota is required for resistance to immunopathologies in the adult. *Immunity* **50**, 1276–1288.e5 (2019). doi: [10.1016/j.immuni.2019.02.014](https://doi.org/10.1016/j.immuni.2019.02.014); pmid: 30902637
7. J. Cahenzli, Y. Köller, M. Wyss, M. B. Geuking, K. D. McCoy, Intestinal microbial diversity during early-life colonization shapes long-term IgE levels. *Cell Host Microbe* **14**, 559–570 (2013). doi: [10.1016/j.chom.2013.10.004](https://doi.org/10.1016/j.chom.2013.10.004); pmid: 24237701
8. T. Olszak *et al.*, Microbial exposure during early life has persistent effects on natural killer T cell function. *Science* **336**, 489–493 (2012). doi: [10.1126/science.1219328](https://doi.org/10.1126/science.1219328); pmid: 22442383
9. D. An *et al.*, Sphingolipids from a symbiotic microbe regulate homeostasis of host intestinal natural killer T cells. *Cell* **156**, 123–133 (2014). doi: [10.1016/j.cell.2013.11.042](https://doi.org/10.1016/j.cell.2013.11.042); pmid: 24439373
10. M. G. Constantinides, Interactions between the microbiota and innate and innate-like lymphocytes. *J. Leukoc. Biol.* **103**, 409–419 (2018). doi: [10.1002/JLB.3RI0917-378R](https://doi.org/10.1002/JLB.3RI0917-378R); pmid: 29345366
11. V. K. Ridaura *et al.*, Contextual control of skin immunity and inflammation by *Corynebacterium*. *J. Exp. Med.* **215**, 785–799 (2018). doi: [10.1084/jem.20171079](https://doi.org/10.1084/jem.20171079); pmid: 29382696
12. J. L. Linehan *et al.*, Non-classical immunity controls microbiota impact on skin immunity and tissue repair. *Cell* **172**, 784–796.e18 (2018). doi: [10.1016/j.cell.2017.12.033](https://doi.org/10.1016/j.cell.2017.12.033); pmid: 29358051
13. A. S. Ismail *et al.*, Gammadelta intraepithelial lymphocytes are essential mediators of host-microbial homeostasis at the intestinal mucosal surface. *Proc. Natl. Acad. Sci. U.S.A.* **108**, 8743–8748 (2011). doi: [10.1073/pnas.1019574108](https://doi.org/10.1073/pnas.1019574108); pmid: 21555560
14. C. Jin *et al.*, Commensal microbiota promote lung cancer development via γδ T cells. *Cell* **176**, 998–1013.e16 (2019). doi: [10.1016/j.cell.2018.12.040](https://doi.org/10.1016/j.cell.2018.12.040); pmid: 30712876
15. L. Kjer-Nielsen *et al.*, MR1 presents microbial vitamin B metabolites to MAIT cells. *Nature* **491**, 717–723 (2012). doi: [10.1038/nature11605](https://doi.org/10.1038/nature11605); pmid: 23051753

16. L. Le Bourhis *et al.*, Antimicrobial activity of mucosal-associated invariant T cells. *Nat. Immunol.* **11**, 701–708 (2010). doi: [10.1038/ni.1890](https://doi.org/10.1038/ni.1890); pmid: 20581831
17. E. Treiner *et al.*, Selection of evolutionarily conserved mucosal-associated invariant T cells by MR1. *Nature* **422**, 164–169 (2003). doi: [10.1038/nature01433](https://doi.org/10.1038/nature01433); pmid: 12634786
18. H. F. Koay *et al.*, A three-stage intrathymic development pathway for the mucosal-associated invariant T cell lineage. *Nat. Immunol.* **17**, 1300–1311 (2016). doi: [10.1038/ni.3565](https://doi.org/10.1038/ni.3565); pmid: 27668799
19. M. Dusseaux *et al.*, Human MAIT cells are xenobiotic-resistant, tissue-targeted, CD161hi IL-17-secreting T cells. *Blood* **117**, 1250–1259 (2011). doi: [10.1182/blood-2010-08-303339](https://doi.org/10.1182/blood-2010-08-303339); pmid: 21084709
20. C. K. Vorkas *et al.*, Mucosal-associated invariant and γδ T cell subsets respond to initial *Mycobacterium tuberculosis* infection. *JCI Insight* **3**, e121899 (2018). doi: [10.1172/jci.insight.121899](https://doi.org/10.1172/jci.insight.121899); pmid: 30282828
21. G. Ben Youssef *et al.*, Ontogeny of human mucosal-associated invariant T cells and related T cell subsets. *J. Exp. Med.* **215**, 459–479 (2018). doi: [10.1084/jem.20171739](https://doi.org/10.1084/jem.20171739); pmid: 29339446
22. P. Chen *et al.*, Circulating mucosal-associated invariant T cells in a large cohort of healthy Chinese individuals from newborn to elderly. *Front. Immunol.* **10**, 260 (2019). doi: [10.3389/fimmu.2019.00260](https://doi.org/10.3389/fimmu.2019.00260); pmid: 30838000
23. J. E. Ussher, P. Klenerman, C. B. Willberg, Mucosal-associated invariant T-cells: New players in anti-bacterial immunity. *Front. Immunol.* **5**, 450 (2014). doi: [10.3389/fimmu.2014.00450](https://doi.org/10.3389/fimmu.2014.00450); pmid: 25339949
24. H. Wang *et al.*, MAIT cells protect against pulmonary Legionella longbeachae infection. *Nat. Commun.* **9**, 3350 (2018). doi: [10.1038/s41467-018-05202-8](https://doi.org/10.1038/s41467-018-05202-8); pmid: 30135490
25. B. van Wilgenburg *et al.*, MAIT cells contribute to protection against lethal influenza infection in vivo. *Nat. Commun.* **9**, 4706 (2018). doi: [10.1038/s41467-018-07207-9](https://doi.org/10.1038/s41467-018-07207-9); pmid: 30413689
26. A. Rahimpour *et al.*, Identification of phenotypically and functionally heterogeneous mouse mucosal-associated invariant T cells using MR1 tetramers. *J. Exp. Med.* **212**, 1095–1108 (2015). doi: [10.1084/jem.20142110](https://doi.org/10.1084/jem.20142110); pmid: 26101265
27. A. J. Corbett *et al.*, T-cell activation by transitory neo-antigens derived from distinct microbial pathways. *Nature* **509**, 361–365 (2014). doi: [10.1038/nature13160](https://doi.org/10.1038/nature13160); pmid: 24695216
28. Y. G. Kim *et al.*, Neonatal acquisition of *Clostridia* species protects against colonization by bacterial pathogens. *Science* **356**, 315–319 (2017). doi: [10.1126/science.aag2029](https://doi.org/10.1126/science.aag2029); pmid: 28428425
29. I. Ivanov *et al.*, Induction of intestinal Th17 cells by segmented filamentous bacteria. *Cell* **139**, 485–498 (2009). doi: [10.1016/j.cell.2009.09.033](https://doi.org/10.1016/j.cell.2009.09.033); pmid: 19836068
30. V. Gaboriau-Routhiau *et al.*, The key role of segmented filamentous bacteria in the coordinated maturation of gut helper T cell responses. *Immunity* **31**, 677–689 (2009). doi: [10.1016/j.immuni.2009.08.020](https://doi.org/10.1016/j.immuni.2009.08.020); pmid: 19833089
31. F. Legoux *et al.*, Microbial metabolites control the thymic development of mucosal-associated invariant T cells. *Science* **eaaw2719** (2019). doi: [10.1126/science.aaw2719](https://doi.org/10.1126/science.aaw2719); pmid: 31467190
32. D. M. Chu *et al.*, Maturation of the infant microbiome community structure and function across multiple body sites and in relation to mode of delivery. *Nat. Med.* **23**, 314–326 (2017). doi: [10.1038/nm.4272](https://doi.org/10.1038/nm.4272); pmid: 28112736
33. J. L. Combellick *et al.*, Differences in the fecal microbiota of neonates born at home or in the hospital. *Sci. Rep.* **8**, 15660 (2018). doi: [10.1038/s41598-018-33995-7](https://doi.org/10.1038/s41598-018-33995-7); pmid: 30353125
34. K. Li *et al.*, Synthesis, stabilization, and characterization of the MR1 ligand precursor 5-amino-6-D-ribitylaminouracil (5-A-RU). *PLOS ONE* **13**, e0191837 (2018). doi: [10.1371/journal.pone.0191837](https://doi.org/10.1371/journal.pone.0191837); pmid: 29401462
35. T. Sano *et al.*, An IL-23R/IL-22 circuit regulates epithelial serum amyloid A to promote local effector Th17 responses. *Cell* **163**, 381–393 (2015). doi: [10.1016/j.cell.2015.08.061](https://doi.org/10.1016/j.cell.2015.08.061); pmid: 26411290
36. J. Dias *et al.*, The CD4⁺CD8⁻ MAIT cell subpopulation is a functionally distinct subset developmentally related to the main CD8⁺ MAIT cell pool. *Proc. Natl. Acad. Sci. U.S.A.* **115**, E11513–E11522 (2018). doi: [10.1073/pnas.1812273115](https://doi.org/10.1073/pnas.1812273115); pmid: 30442667
37. A. L. Byrd, Y. Belkaid, J. A. Segre, The human skin microbiome. *Nat. Rev. Microbiol.* **16**, 143–155 (2018). doi: [10.1038/nrmicro.2017.157](https://doi.org/10.1038/nrmicro.2017.157); pmid: 29332945
38. S. Naik *et al.*, Commensal-dendritic-cell interaction specifies a unique protective skin immune signature. *Nature* **520**, 104–108 (2015). doi: [10.1038/nature14052](https://doi.org/10.1038/nature14052); pmid: 25539086

39. P. De Togni *et al.*, Abnormal development of peripheral lymphoid organs in mice deficient in lymphotoxin. *Science* **264**, 703–707 (1994). doi: [10.1126/science.8171322](https://doi.org/10.1126/science.8171322); pmid: [8171322](https://pubmed.ncbi.nlm.nih.gov/18171322/)
40. X. Jiang *et al.*, Dermal $\gamma\delta$ T cells do not freely re-circulate out of skin and produce IL-17 to promote neutrophil infiltration during primary contact hypersensitivity. *PLOS ONE* **12**, e0169397 (2017). doi: [10.1371/journal.pone.0169397](https://doi.org/10.1371/journal.pone.0169397); pmid: [28081153](https://pubmed.ncbi.nlm.nih.gov/28081153/)
41. S. Y. Thomas *et al.*, PLZF induces an intravascular surveillance program mediated by long-lived LFA-1-ICAM-1 interactions. *J. Exp. Med.* **208**, 1179–1188 (2011). doi: [10.1084/jem.20102630](https://doi.org/10.1084/jem.20102630); pmid: [21624939](https://pubmed.ncbi.nlm.nih.gov/21624939/)
42. M. Salou *et al.*, A common transcriptomic program acquired in the thymus defines tissue residency of MAIT and NKT subsets. *J. Exp. Med.* **216**, 133–151 (2019). doi: [10.1084/jem.20181483](https://doi.org/10.1084/jem.20181483); pmid: [30518599](https://pubmed.ncbi.nlm.nih.gov/30518599/)
43. M. J. Sobkowiak *et al.*, Tissue-resident MAIT cell populations in human oral mucosa exhibit an activated profile and produce IL-17. *Eur. J. Immunol.* **49**, 133–143 (2019). doi: [10.1002/eji.201847759](https://doi.org/10.1002/eji.201847759); pmid: [30372518](https://pubmed.ncbi.nlm.nih.gov/30372518/)
44. B. van Wilgenburg *et al.*, MAIT cells are activated during human viral infections. *Nat. Commun.* **7**, 11653 (2016). doi: [10.1038/ncomms11653](https://doi.org/10.1038/ncomms11653); pmid: [27337592](https://pubmed.ncbi.nlm.nih.gov/27337592/)
45. S. Naik *et al.*, Compartmentalized control of skin immunity by resident commensals. *Science* **337**, 1115–1119 (2012). doi: [10.1126/science.1225152](https://doi.org/10.1126/science.1225152); pmid: [22837383](https://pubmed.ncbi.nlm.nih.gov/22837383/)
46. M. P. Seiler *et al.*, Elevated and sustained expression of the transcription factors Egr1 and Egr2 controls NKT lineage differentiation in response to TCR signaling. *Nat. Immunol.* **13**, 264–271 (2012). doi: [10.1038/ni.2230](https://doi.org/10.1038/ni.2230); pmid: [22306690](https://pubmed.ncbi.nlm.nih.gov/22306690/)
47. F. Bussolino *et al.*, Hepatocyte growth factor is a potent angiogenic factor which stimulates endothelial cell motility and growth. *J. Cell Biol.* **119**, 629–641 (1992). doi: [10.1083/jcb.119.3.629](https://doi.org/10.1083/jcb.119.3.629); pmid: [1383237](https://pubmed.ncbi.nlm.nih.gov/1383237/)
48. M. Miyake, S. Goodison, A. Lawton, E. Gomes-Giacioia, C. J. Rosser, Angiogenin promotes tumoral growth and angiogenesis by regulating matrix metalloproteinase-2 expression via the ERK1/2 pathway. *Oncogene* **34**, 890–901 (2015). doi: [10.1038/nc.2014.2](https://doi.org/10.1038/nc.2014.2); pmid: [24561529](https://pubmed.ncbi.nlm.nih.gov/24561529/)
49. A. Toulon *et al.*, A role for human skin-resident T cells in wound healing. *J. Exp. Med.* **206**, 743–750 (2009). doi: [10.1084/jem.20081787](https://doi.org/10.1084/jem.20081787); pmid: [19307328](https://pubmed.ncbi.nlm.nih.gov/19307328/)
50. Z. He, C. H. Ong, J. Halper, A. Bateman, Progranulin is a mediator of the wound response. *Nat. Med.* **9**, 225–229 (2003). doi: [10.1038/nm816](https://doi.org/10.1038/nm816); pmid: [12524533](https://pubmed.ncbi.nlm.nih.gov/12524533/)
51. S. LeBlanc *et al.*, CEACAM1 deficiency delays important wound healing processes. *Wound Repair Regen.* **19**, 745–752 (2011). doi: [10.1111/j.1524-475X.2011.00742.x](https://doi.org/10.1111/j.1524-475X.2011.00742.x); pmid: [22092845](https://pubmed.ncbi.nlm.nih.gov/22092845/)
52. M. Demaria *et al.*, An essential role for senescent cells in optimal wound healing through secretion of PDGF-AA. *Dev. Cell* **31**, 722–733 (2014). doi: [10.1016/j.devcel.2014.11.012](https://doi.org/10.1016/j.devcel.2014.11.012); pmid: [25499914](https://pubmed.ncbi.nlm.nih.gov/25499914/)
53. A. Grochot-Przeczek *et al.*, Heme oxygenase-1 accelerates cutaneous wound healing in mice. *PLOS ONE* **4**, e5803 (2009). doi: [10.1371/journal.pone.0005803](https://doi.org/10.1371/journal.pone.0005803); pmid: [19495412](https://pubmed.ncbi.nlm.nih.gov/19495412/)
54. N. Seach *et al.*, Double-positive thymocytes select mucosal-associated invariant T cells. *J. Immunol.* **191**, 6002–6009 (2013). doi: [10.4049/jimmunol.1301212](https://doi.org/10.4049/jimmunol.1301212); pmid: [24244014](https://pubmed.ncbi.nlm.nih.gov/24244014/)
55. E. Martin *et al.*, Stepwise development of MAIT cells in mouse and human. *PLoS Biol.* **7**, e54 (2009). doi: [10.1371/journal.pbio.1000054](https://doi.org/10.1371/journal.pbio.1000054); pmid: [19278296](https://pubmed.ncbi.nlm.nih.gov/19278296/)
56. A. Ventura *et al.*, Restoration of p53 function leads to tumour regression in vivo. *Nature* **445**, 661–665 (2007). doi: [10.1038/nature05541](https://doi.org/10.1038/nature05541); pmid: [17251932](https://pubmed.ncbi.nlm.nih.gov/17251932/)
57. M. B. M. Teunissen *et al.*, The IL-17A-producing CD8⁺ T-cell population in psoriatic lesional skin comprises mucosa-associated invariant T cells and conventional T cells. *J. Invest. Dermatol.* **134**, 2898–2907 (2014). doi: [10.1038/jid.2014.261](https://doi.org/10.1038/jid.2014.261); pmid: [24945094](https://pubmed.ncbi.nlm.nih.gov/24945094/)
58. J. Li *et al.*, The frequency of mucosal-associated invariant T cells is selectively increased in dermatitis herpetiformis. *Australas. J. Dermatol.* **58**, 200–204 (2017). doi: [10.1111/ajd.12456](https://doi.org/10.1111/ajd.12456); pmid: [26940855](https://pubmed.ncbi.nlm.nih.gov/26940855/)
59. J. Swanpradid, Z. E. Holcomb, A. S. MacLeod, Emerging skin T-cell functions in response to environmental insults. *J. Invest. Dermatol.* **137**, 288–294 (2017). doi: [10.1016/j.jid.2016.08.013](https://doi.org/10.1016/j.jid.2016.08.013); pmid: [27784595](https://pubmed.ncbi.nlm.nih.gov/27784595/)
60. J. Jameson *et al.*, A role for skin $\gamma\delta$ T cells in wound repair. *Science* **296**, 747–749 (2002). doi: [10.1126/science.1069639](https://doi.org/10.1126/science.1069639); pmid: [11976459](https://pubmed.ncbi.nlm.nih.gov/11976459/)
61. B. E. Keyes *et al.*, Impaired epidermal to dendritic T cell signaling slows wound repair in aged skin. *Cell* **167**, 1323–1338. e14 (2016). doi: [10.1016/j.cell.2016.10.052](https://doi.org/10.1016/j.cell.2016.10.052); pmid: [27863246](https://pubmed.ncbi.nlm.nih.gov/27863246/)
62. R. Di Marco Barros *et al.*, Epithelia use butyrophilin-like molecules to shape organ-specific $\gamma\delta$ T cell compartments. *Cell* **167**, 203–218. e17 (2016). doi: [10.1016/j.cell.2016.08.030](https://doi.org/10.1016/j.cell.2016.08.030); pmid: [27641500](https://pubmed.ncbi.nlm.nih.gov/27641500/)
63. M. Gomez de Agüero *et al.*, The maternal microbiota drives early postnatal innate immune development. *Science* **351**, 1296–1302 (2016). doi: [10.1126/science.aad2571](https://doi.org/10.1126/science.aad2571); pmid: [26989247](https://pubmed.ncbi.nlm.nih.gov/26989247/)
64. E. Leeansyah, L. Loh, D. F. Nixon, J. K. Sandberg, Acquisition of innate-like microbial reactivity in mucosal tissues during human fetal MAIT-cell development. *Nat. Commun.* **5**, 3143 (2014). doi: [10.1038/ncomms4143](https://doi.org/10.1038/ncomms4143); pmid: [24452018](https://pubmed.ncbi.nlm.nih.gov/24452018/)
65. K. Hirota *et al.*, Fate mapping of IL-17-producing T cells in inflammatory responses. *Nat. Immunol.* **12**, 255–263 (2011). doi: [10.1038/ni.1993](https://doi.org/10.1038/ni.1993); pmid: [21278737](https://pubmed.ncbi.nlm.nih.gov/21278737/)
66. A. Awasthi *et al.*, Cutting edge: IL-23 receptor gp reporter mice reveal distinct populations of IL-17-producing cells. *J. Immunol.* **182**, 5904–5908 (2009). doi: [10.4049/jimmunol.0900732](https://doi.org/10.4049/jimmunol.0900732); pmid: [19414740](https://pubmed.ncbi.nlm.nih.gov/19414740/)
67. L. E. Jao, S. R. Wente, W. Chen, Efficient multiplex biallelic zebrafish genome editing using a CRISPR nuclease system. *Proc. Natl. Acad. Sci. U.S.A.* **110**, 13904–13909 (2013). doi: [10.1073/pnas.1308335110](https://doi.org/10.1073/pnas.1308335110); pmid: [23918387](https://pubmed.ncbi.nlm.nih.gov/23918387/)
68. S. Conlan *et al.*, *Staphylococcus epidermidis* pan-genome sequence analysis reveals diversity of skin commensal and hospital infection-associated isolates. *Genome Biol.* **13**, R64 (2012). doi: [10.1186/gb-2012-13-7-r64](https://doi.org/10.1186/gb-2012-13-7-r64); pmid: [22830599](https://pubmed.ncbi.nlm.nih.gov/22830599/)
69. I. R. Monk, I. M. Shah, M. Xu, M. W. Tan, T. J. Foster, Transforming the untransformable: Application of direct transformation to manipulate genetically *Staphylococcus aureus* and *Staphylococcus epidermidis*. *mBio* **3**, e00277-11 (2012). doi: [10.1128/mBio.00277-11](https://doi.org/10.1128/mBio.00277-11); pmid: [22434850](https://pubmed.ncbi.nlm.nih.gov/22434850/)
70. D. G. Gibson *et al.*, Enzymatic assembly of DNA molecules up to several hundred kilobases. *Nat. Methods* **6**, 343–345 (2009). doi: [10.1038/nmeth.1318](https://doi.org/10.1038/nmeth.1318); pmid: [19363495](https://pubmed.ncbi.nlm.nih.gov/19363495/)
71. D. E. Wright, A. J. Wagers, A. P. Gulati, F. L. Johnson, I. L. Weissman, Physiological migration of hematopoietic stem and progenitor cells. *Science* **294**, 1933–1936 (2001). doi: [10.1126/science.1064081](https://doi.org/10.1126/science.1064081); pmid: [11729320](https://pubmed.ncbi.nlm.nih.gov/11729320/)
72. M. Lochner, H. Wagner, M. Classen, I. Förster, Generation of neutralizing mouse anti-mouse IL-18 antibodies for inhibition of inflammatory responses in vivo. *J. Immunol. Methods* **259**, 149–157 (2002). doi: [10.1016/S0022-1759\(01\)00505-1](https://doi.org/10.1016/S0022-1759(01)00505-1); pmid: [11730850](https://pubmed.ncbi.nlm.nih.gov/11730850/)
73. J. A. Hall *et al.*, Essential role for retinoic acid in the promotion of CD4(+) T cell effector responses via retinoic acid receptor α . *Immunity* **34**, 435–447 (2011). doi: [10.1016/j.immuni.2011.03.003](https://doi.org/10.1016/j.immuni.2011.03.003); pmid: [21419664](https://pubmed.ncbi.nlm.nih.gov/21419664/)
74. B. J. Callahan *et al.*, DADA2: High-resolution sample inference from Illumina amplicon data. *Nat. Methods* **13**, 581–583 (2016). doi: [10.1038/nmeth.3869](https://doi.org/10.1038/nmeth.3869); pmid: [27214047](https://pubmed.ncbi.nlm.nih.gov/27214047/)
75. A. I. Dobin *et al.*, STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15–21 (2012). doi: [10.1093/bioinformatics/bts635](https://doi.org/10.1093/bioinformatics/bts635); pmid: [23104886](https://pubmed.ncbi.nlm.nih.gov/23104886/)
76. S. Heinz *et al.*, Simple combinations of lineage-determining transcription factors prime cis-regulatory elements required for macrophage and B cell identities. *Mol. Cell* **38**, 576–589 (2010). doi: [10.1016/j.molcel.2010.05.004](https://doi.org/10.1016/j.molcel.2010.05.004); pmid: [20513432](https://pubmed.ncbi.nlm.nih.gov/20513432/)
77. S. Tripathi *et al.*, Meta- and orthogonal integration of influenza “OMICs” data defines a role for UBR4 in virus budding. *Mol. Cell* **18**, 723–735 (2015). doi: [10.1016/j.chom.2015.11.002](https://doi.org/10.1016/j.chom.2015.11.002); pmid: [26651948](https://pubmed.ncbi.nlm.nih.gov/26651948/)
78. M. Stoeckius *et al.*, Cell Hashing with barcoded antibodies enables multiplexing and doublet detection for single cell genomics. *Genome Biol.* **19**, 224 (2018). doi: [10.1186/s13059-018-1603-1](https://doi.org/10.1186/s13059-018-1603-1); pmid: [30567574](https://pubmed.ncbi.nlm.nih.gov/30567574/)
79. T. Stuart *et al.*, Comprehensive integration of single-cell data. *Cell* **177**, 1888–1902. e21 (2019). doi: [10.1016/j.cell.2019.05.031](https://doi.org/10.1016/j.cell.2019.05.031); pmid: [31178118](https://pubmed.ncbi.nlm.nih.gov/31178118/)

ACKNOWLEDGMENTS

We thank National Institute of Allergy and Infectious Diseases (NIAID) animal facility staff—in particular, D. Trageser-Cesler and C. Acevedo (NIAID Microbiome Program Gnotobiotic Animal Facility); J. Chen, N. Modi, and S. Mistry (NIH Center for Human Immunology); M. Smelkinson (NIAID Biological Imaging Section); T. Hawley and E. Stregovsky (NIAID Flow Cytometry Section); R. Awasthi (NCI Transgenic Mouse Model Laboratory); and G. Trinchieri, R. Salcedo, J. McCulloch, S. Sen, K. Beacht, J. LeGrand, E. Lewis, and J. Kehr. MRI and CD1d tetramers were provided by the NIH Tetramer Core Facility. We acknowledge the NIAID Microbiome Program Sequencing Platform for performing the 16S rRNA and genomic sequencing. This study used the Office of Cyber Infrastructure and Computational Biology (OCICB) High Performance Computing (HPC) cluster at NIAID and the high-performance computational capabilities of the Biowulf Linux cluster at NIH. **Funding:** This research was supported in part by the Intramural Research Program of NIAID, NIH (ZIA-AI001115 and ZIA-AI001132 to Y.B.); the NIH Extramural Research Program (U19-AI111143 to J.A., R01-DK110174 and DP1-DK113598 to M.A.F., DP2-AI144968 to T.C.S.); the Cancer Research Institute Irvington Postdoctoral Fellowship Program (M.G.C. and I.V.-C.); the European Molecular Biology Organization (EMBO) Long-Term Fellowship Program and the ARC Foundation (S.T.); the National Institute of General Medical Sciences (NIGMS) Postdoctoral Research Associate Training (PRAT) Program (J.L.L.); Howard Hughes Medical Institute (HHMI) Hanna H. Gray Fellows Program (Y.E.C.); and the HHMI-Simons Faculty Scholars Program and the Chan Zuckerberg Biohub (M.A.F.). **Author contributions:** M.G.C. and Y.B. designed the study and experiments and wrote the manuscript; M.G.C. performed the experiments and analyzed the data; V.M.L., S.T., A.C.W., P.J.P.-C., S.-J.H., Y.E.C., K.L., S.F., A.W., S.R.K., I.V.-C., J.L.L., N.B., E.D.M., S.R., D.J.C., E.J.A., A.B., T.C.S., J.A., and M.A.F. participated in performing experiments, provided intellectual expertise, and/or helped to interpret experimental results; V.M.L. assisted with RNA-seq data analysis; S.T. and A.C.W. assisted with numerous experiments throughout; P.J.P.-C. isolated, cultured, and sequenced intestinal commensals; S.-J.H. performed confocal microscopy and parabiotic surgery; Y.E.C. and M.A.F. generated the *S. epidermidis* NIHLM087 Δ ribD mutant; K.L. and J.A. synthesized 5-A-RU; S.F., A.W., and T.C.S. analyzed human skin biopsies; S.R.K. analyzed the presence of riboflavin synthesis genes; I.V.-C. analyzed the 16S rRNA data; J.L.L. assisted with wounding studies; N.B. coordinated GF experiments; E.D.M. participated in the conventionalization of GF mice; S.R. and E.J.A. generated MRI tetramer used in initial experiments; D.J.C. provided antibody to IL-23R and isotype control; and A.B. coordinated the generation of *Mrl1^{+/+}* mice. **Competing interests:** The authors declare no competing interests. **Data and materials availability:** Reagents are available upon signing a materials transfer agreement. Antibody to IL-18 (SK113AE-4) is available from I. Förster under a materials transfer agreement with the University of Bonn. The RNA-seq data, 16S rRNA sequences, and bacterial genomes were deposited with the NCBI under accession nos. GSE128814 and PRJNA529261. All other data needed to evaluate the conclusions in this paper are present either in the main text or the supplementary materials.

SUPPLEMENTARY MATERIALS

science.sciencemag.org/content/366/6464/eaax6624/suppl/DC1
Figs. S1 to S5
Table S1

[View/request a protocol for this paper from Bio-protocol.](#)

11 April 2019; accepted 12 September 2019
10.1126/science.aax6624