



Published in final edited form as:

*Angew Chem Int Ed Engl.* 2008 ; 47(33): 6233–6235. doi:10.1002/anie.200801591.

## Highly Stereoselective Ring Expansion Reactions Mediated by Attractive, Non-Bonded Cation– $n$ Interactions†

**Timothy Ribelin,**

Department of Medicinal Chemistry, 1251 Wescoe Hall Drive, Malott Hall, Room 4070, University of Kansas, Lawrence, KS 66047

**Dr. Christopher E. Katz,**

Department of Medicinal Chemistry, 1251 Wescoe Hall Drive, Malott Hall, Room 4070, University of Kansas, Lawrence, KS 66047

**Donna Withrow,**

Department of Chemistry and Biochemistry, Old Dominion University, 4541 Hampton Boulevard, Norfolk, Virginia 23529

**Sherriel Smith,**

Department of Chemistry and Biochemistry, Old Dominion University, 4541 Hampton Boulevard, Norfolk, Virginia 23529

**Anna Manukyan,**

Department of Chemistry and Biochemistry, Old Dominion University, 4541 Hampton Boulevard, Norfolk, Virginia 23529

**Dr. Victor W. Day,**

Department of Medicinal Chemistry, 1251 Wescoe Hall Drive, Malott Hall, Room 4070, University of Kansas, Lawrence, KS 66047

**Benjamin Neuenswander,**

Department of Medicinal Chemistry, 1251 Wescoe Hall Drive, Malott Hall, Room 4070, University of Kansas, Lawrence, KS 66047

**Prof. Jennifer L. Poutsma, and**

Department of Chemistry and Biochemistry, Old Dominion University, 4541 Hampton Boulevard, Norfolk, Virginia 23529

**Prof. Jeffrey Aubé**

Department of Medicinal Chemistry, 1251 Wescoe Hall Drive, Malott Hall, Room 4070, University of Kansas, Lawrence, KS 66047

### Abstract

**Electrostatic control of leaving group stereochemistry** leads to superior diastereoselectivity in an asymmetric ring expansion reaction.

### Keywords

ring expansion; non-bonded interactions; azides; Schmidt reaction; stereoselectivity

†Supporting information for this article is available on the WWW under <http://www.angewandte.org> or from the author. CDCD

Correspondence to: Jennifer L. Poutsma; Jeffrey Aubé.

Most stereoselective reactions are ruled by steric effects. In particular, kinetically controlled asymmetric transformations utilizing chiral reagent, auxiliaries, or catalysts succeed due to energy differences in transition states that most often arise by the minimization of repulsive, non-bonded interactions. Stereoelectronic considerations, which arise when the alignment of particular orbitals are necessary for a successful reaction, can also play a role.<sup>[1]</sup> An iconic stereoelectronic effect in organic chemistry is the anomeric effect.<sup>[2]</sup> Reactions controlled by the anomeric effect, such as glycosidations, largely depend on the relative orientation of the non-bonding or *n* electrons of a nearby alkoxy group. In recent years, alkoxy group control of stereoselective reactions via electrostatic interactions has received renewed scrutiny, led by the Woerpel group.<sup>[3]</sup> In this communication, we report an alternative and highly effective approach to stereocontrol through the maximization of attractive non-bonded interactions between an alkoxy or alkylthio group and a positively charged leaving group.

The Lewis acid-promoted reaction of a symmetrically substituted cyclic ketone with a chiral hydroxyalkyl azide provides a stereoselective route to lactams (Scheme 1).<sup>[4]</sup> In this reaction, initial formation of a spirocyclic intermediate sets up the selective migration of one of the alkyl groups originally adjacent to the ketone carbonyl. Migration of a C–C bond antiperiplanar to the  $N_2^+$  leaving group (only possible when the latter is in an axial position as shown) affords an iminium ether that is converted into lactam by workup with aqueous base. For 1- or 3-substituted azidopropanols (not shown), 10:1 selectivities are obtained, corresponding to preferential reaction through the most stable chairlike heterocyclic ring (**A** or **B**) resulting from equatorial addition of azide relative to the *tert*-butyl group. Intermediates **A** and **B** can interconvert through conformational reorganization or by reversion to the initially formed oxonium ion followed by reclosure. In this scenario, selectivity is attained by stabilization of **A** over **B** due to traditional minimization of 1,3-diaxial interactions by placement of the  $R^1$  or  $R^3$  into equatorial positions in the former.

2-Substituted 1,3-azidopropanols present a special case that is unusually susceptible to stereoelectronic control due to three factors: (1) the methylene groups near the spiro linkage are locally isoelectronic, so the reaction cannot be controlled by “migratory aptitude”, (2) the presence of either an oxygen ether or an N– $N_2^+$  group in a 1,3 relationship to the R group means that 1,3-diaxial steric interactions will be minimized, and (3) the 1,3 relationship between axial R and  $N_2^+$  groups provides a strong opportunity for attractive electrostatic interactions to occur between these groups in intermediate **B**. In previous work, it was demonstrated that unusually low selectivities obtained in this system when R = aryl could be ascribed to preferential stabilization of intermediate **B** by attractive, non-bonded cation– $\pi$  interactions between the aromatic group and the  $N_2^+$  leaving group (Table 1).<sup>[5]</sup> Although such interactions are commonly proposed in biological systems,<sup>[6]</sup> they are rarely invoked as stereocontrolling features of small-molecule stereoselective reactions.<sup>[7]</sup>

A computational study<sup>[8]</sup> and analogy to the well-known ability of ether groups to bind cations suggested that intermediates like **B** should be even more enhanced in compounds where R = alkoxy. As shown in Figure 1, isomer **B** containing a diaxial relationship between methoxy group and leaving group was calculated to be ca. 3.8 kcal/mol more stable than the equatorial isomer for which no interaction between methoxy and  $N_2^+$  groups are possible. To test this, 1-azido-2-methoxypropanol **3** was prepared and reacted with 4-*tert*-butylcyclohexanone using  $BF_3 \cdot OEt_2$  as Lewis acid promoter. A striking 24:1 selectivity *in favor of the isomer emanating from an axially disposed methoxy group was obtained in high yield* (Table 1, entry 3).

This result suggests that the methoxy cation–*n* interaction is considerably stronger than the previously reported cation– $\pi$  effect, due to the fact that the highest **3:2** ratio observed to date was 57:43 for the electron-rich 3,4,5-trimethoxyphenyl group (not shown).<sup>[5]</sup> The fact

that the small MeO group (A value = 0.6<sup>[9]</sup>) pays a relatively small steric penalty in the axial orientation is a likely contributor to the high selectivity of this reaction as well. However, the much higher selectivity and opposite direction of stereocontrol obtained for the smaller MeO group as compared to alkyl or aryl substituents (Table 1, entries 1 and 2) is strong evidence for the proposed role of electrostatics in this reaction.

We proposed that a similar effect might be observed with a more polarizable heteroatom.<sup>[3m]</sup> Accordingly, **1d** where R = SMe, was prepared and submitted to the asymmetric Schmidt reaction protocol. Remarkably, a >98:2 dr was obtained for this system, favoring **3d**. *The selectivities obtained with both methoxy and methylthio, which depend mainly on electrostatics and feature axially disposed substituents, are higher than any previously reported, sterically-based, example of this ring-expansion reaction.*<sup>[4,5]</sup>

Although the first example of an asymmetric azido-Schmidt reaction reported utilized an azidoethanol reagent, that series has typically provided lower selectivities relative to the three-carbon-containing reagents like **1** and has more recently been shown to occur via predominant steric control, even when a phenyl group is in a position to participate in a cation- $\pi$  interaction.<sup>[5]</sup> In sharp contrast to these previous results, the reaction using reagent **4** afforded a high 97:3 ratio of **5** over **6**, in which the major product goes through an intermediate in which a *syn* relationship between the methoxy group and the leaving N<sub>2</sub><sup>+</sup> substituent is possible (Scheme 2). A computational investigation showed that the cation-*n* intermediate **C** is stabilized by 3.9 kcal/mol. Notably, the O-N<sub>2</sub><sup>+</sup> distances, energy differences, and ratios are similar between systems **B** and **C**. Previous work in the reactions of substituted 1,2-azidoethanols has shown the predominant steric feature affecting stereochemistry to exist between the migrating carbon and substituents on the five-membered heterocyclic ring.<sup>[4,5b]</sup> In cases where the alkyl group is adjacent to oxygen (i.e., across the ring from the migrating methylene group and the N<sub>2</sub><sup>+</sup> leaving group), steric effects do not play an important role in determining reaction stereochemistry, as clearly demonstrated by the non-selective cyclohexyl case shown in Scheme 2b.

The opposite situation occurs when the methoxymethyl group is placed adjacent to the azido group. In this case, there is no substantial difference in distance between the methoxy group and either isomeric intermediate, so electrostatic considerations cannot play a role and the preference for *syn* **E** over *anti* **F** drops to 0.6 kcal/mol computationally. Instead, the usual steric course of the reaction leads to the same product observed for the analogous example to **14** (Scheme 3).

The most interesting elements of this approach are that: (1) intermediates are subject to non-bonded, attractive interactions that are able to strongly favor one stereoisomeric form over the other, (2) these intermediates lead to the corresponding products in a process entirely controlled by stereoelectronic considerations, and (3) the overall stereoselectivity ultimately depends on the control of leaving group stereoselectivity at an epimerizable nitrogen atom. The high yields of these reactions combined with the utility of the lactam products suggests a high level of utility of the present reaction. Of perhaps greater long-term interest will be the attempted utilization of cation nonbonding electron stabilization in other stereoselective processes.<sup>[10]</sup>

## Supplementary Material


Refer to Web version on PubMed Central for supplementary material.

## Acknowledgments


Financial support of this work by the Institute of General Medical Sciences is gratefully acknowledged (GM 49093, JA). Acknowledgment is made to the Donors of the American Chemical Society Petroleum Research Fund for partial support of this research (to JLP). We also thank Paul Hanson and his group for their gas chromatograph.

## References


1. Deslongchamps, P. *Stereoelectronic Effects in Organic Chemistry*. Pergamon; Oxford: 1983.
2. For reviews, see: Juaristi E, Cuevas G. *Tetrahedron*. 1992; 48:5019–5087. Graczyk PP, Mikolajczyk M. *Top Stereochem*. 1994; 21:159–349.
3. For a review on electrostatic interactions between ethers and oxocarbenium ions, see: Smith DM, Woerpel KA. *Org Biomol Chem*. 2006; 4:1195–1201. [PubMed: 16557303] For selected examples from the recent literature, see: Dunkin IR, Paul AJ, Suckling CJ, Valente E, Wood HC. *J Chem Soc, Perkin Trans 1*. 1985:1323–1326. Mootoo DR, Fraser-Reid B. *J Chem Soc, Chem Commun*. 1986:1570–1571. Elvey SP, Mootoo DR. *J Am Chem Soc*. 1992; 114:9685–9686. Molander GA, Cameron KO. *J Org Chem*. 1993; 58:5931–5943. Shan W, Wilson P, Liang W, Mootoo DR. *J Org Chem*. 1994; 59:7986–7993. Andersson PG. *J Org Chem*. 1996; 61:4154–4156. [PubMed: 11667299] Suga S, Suzuki S, Yoshida J-i. *Org Lett*. 2005; 7:4717–4720. [PubMed: 16209518] Fujita M, Lee HJ, Okuyama T. *Org Lett*. 2006; 8:1399–1401. [PubMed: 16562901] Judd WR, Ban S, Aubé J. *J Am Chem Soc*. 2006; 128:13736–13741. [PubMed: 17044701] Baghdasarian G, Woerpel KA. *J Org Chem*. 2006; 71:6851–6858. [PubMed: 16930037] Shenoy SR, Smith DM, Woerpel KA. *J Am Chem Soc*. 2006; 128:8671–8677. [PubMed: 16802834] Beaver MG, Billings SB, Woerpel KA. *J Am Chem Soc*. 2008; 130:2082–2086. [PubMed: 18215038]
4. a) Gracias V, Frank KE, Milligan GL, Aubé J. *Tetrahedron*. 1997; 53:16241–16252. b) Sahasrabudhe K, Gracias V, Furness K, Smith BT, Katz CE, Reddy DS, Aube J. *J Am Chem Soc*. 2003; 125:7914–7922. [PubMed: 12823012]
5. a) Katz CE, Aubé J. *J Am Chem Soc*. 2003; 125:13948–13949. [PubMed: 14611217] b) Katz CE, Ribelin T, Withrow D, Basseri Y, Manukyan AK, Bermudez A, Nuera CG, Day VW, Powell DR, Poutsma JL, Aubé J. *J Org Chem*. 2008 in press.
6. a) Dougherty DA. *Science*. 1996; 271:163–168. [PubMed: 8539615] b) Ma JC, Dougherty DA. *Chem Rev*. 1997; 97:1303–1324. [PubMed: 11851453]
7. For a review, see: Yamada S. *Org Biomol Chem*. 2007; 5:2903–2912. [PubMed: 17728854] For some leading references, see: Lakshminarasimhan P, Sunoj RB, Chandrasekhar J, Ramamurthy V. *J Am Chem Soc*. 2000; 122:4815–4816. Stratakis M, Froudakis G. *Org Lett*. 2000; 2:1369–1372. [PubMed: 10814449] Yamada S, Misono T, Ichikawa M, Morita C. *Tetrahedron*. 2001; 57:8939–8949. Yamada S, Morita C. *J Am Chem Soc*. 2002; 124:8184–8185. [PubMed: 12105887] Ramamurthy V, Shailaja J, Kaanumalle LS, Sunoj RB, Chandrasekhar J. *Chem Commun*. 2003:1987–1999. Chiavarino B, Crestoni ME, Fornarini S. *Int J Mass Spectrom*. 2004; 235:145–154.
8. All calculations reported utilize MP2/6-311+G(d,p)//MP2/6-31G(d) free energies with CPCM dichloromethane solvent-optimized geometries.
9. Eliel, EL.; Wilen, SH.; Mander, LN. *Stereochemistry of Organic Compounds*. John Wiley & Sons; New York: 1994. p. 696-697. references cited
10. All experimental details and stereochemical assignments are described in the Supporting Information. CCDC entries 647652, 647653, 647654, and 647652 contain the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via [www.ccdc.cam.ac.uk/data\\_request/cif](http://www.ccdc.cam.ac.uk/data_request/cif).



**Figure 1.** Calculations for proposed intermediates **A** and **B** performed at the MP2/6-311+G\*\*//MP2/6-31G\* level of theory.<sup>[8]</sup>




**Scheme 1.**  
Origin of selectivity in asymmetric Schmidt reactions.



**Scheme 2.**

(a) Electrostatically controlled reaction of 1-azidoethanol derivative **4** with 4-*tert*-butylcyclohexanone (including calculated energies of proposed, minimized intermediates **C** and **D**<sup>[8]</sup>) and (b) a cyclohexyl-containing control.<sup>[5b]</sup> The model systems used for the calculations are given in the Supporting Information (Figure S1).

**Scheme 3.**

Sterically controlled reactions of (a) **10** and (b) a previously reported cyclohexanyl example.<sup>[5]</sup> The model systems used for the calculations are given in the Supporting Information (Figure S1).



**Table 1**

Selectivity of reactions of substituted 1,3-hydroxyalkyl azides.

entry	series	R <sup>2</sup>	2:3 ratio	yield (%)
1	<b>a</b>	Me <sup>[a]</sup>	74:26	98
2	<b>b</b>	Ph <sup>[b]</sup>	60:40	98
3	<b>c</b>	OMe	4:96	98
4	<b>d</b>	SMe	1.8:98.2	90

<sup>[a]</sup>Reference 4.<sup>[b]</sup>References 4 and 5.