

East Tennessee State University

Digital Commons @ East Tennessee State University

ETSU Faculty Works

Faculty Works

12-1-2020

High Glucose Distinctively Regulates Ca²⁺ Influx in Cytotoxic T Lymphocytes Upon Target Recognition and Thapsigargin Stimulation

Huajiao Zou
Central South University

Wenjuan Yang
Universitätsklinikum des Saarlandes Medizinische Fakultät der Universität des Saarlandes

Gertrud Schwär
Universitätsklinikum des Saarlandes Medizinische Fakultät der Universität des Saarlandes

Renping Zhao
Universitätsklinikum des Saarlandes Medizinische Fakultät der Universität des Saarlandes

Dalia Alansary
Universitätsklinikum des Saarlandes Medizinische Fakultät der Universität des Saarlandes

See next page for additional authors

Follow this and additional works at: <https://dc.etsu.edu/etsu-works>

Citation Information

Zou, Huajiao; Yang, Wenjuan; Schwär, Gertrud; Zhao, Renping; Alansary, Dalia; Yin, Deling; Schwarz, Eva C.; Niemeyer, Barbara A.; and Qu, Bin. 2020. High Glucose Distinctively Regulates Ca²⁺ Influx in Cytotoxic T Lymphocytes Upon Target Recognition and Thapsigargin Stimulation. *European Journal of Immunology*. Vol.50(12). 2095-2098. <https://doi.org/10.1002/eji.202048577> PMID: 32697355 ISSN: 0014-2980

This Letter is brought to you for free and open access by the Faculty Works at Digital Commons @ East Tennessee State University. It has been accepted for inclusion in ETSU Faculty Works by an authorized administrator of Digital Commons @ East Tennessee State University. For more information, please contact digilib@etsu.edu.

High Glucose Distinctively Regulates Ca²⁺ Influx in Cytotoxic T Lymphocytes Upon Target Recognition and Thapsigargin Stimulation

Copyright Statement

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

Creative Commons License







This work is licensed under a [Creative Commons Attribution-NonCommercial 4.0 International License](https://creativecommons.org/licenses/by-nc/4.0/)

Creator(s)

Huajiao Zou, Wenjuan Yang, Gertrud Schwär, Renping Zhao, Dalia Alansary, Deling Yin, Eva C. Schwarz, Barbara A. Niemeyer, and Bin Qu

invitrogen

Two data sets. One step. Zero doubt.



Confidently confirm your cell profiles with a new flow cytometer that delivers flow cytometry and imaging data simultaneously. Now, you can acquire dual data quickly and easily. The new Invitrogen™ Attune™ CytPix™ Flow Cytometer delivers both brightfield images and flow cytometry data sets simultaneously, so you can confirm cellular characteristics and sample quality confidently, without changing your protocols.



Enhance analysis and confidence at [thermofisher.com/cytpix](https://www.thermofisher.com/cytpix)

ThermoFisher
SCIENTIFIC

For Research Use Only. Not for use in diagnostic procedures. © 2021 Thermo Fisher Scientific Inc. All rights reserved. All trademarks are the property of Thermo Fisher Scientific and its subsidiaries unless otherwise specified. COL25211 0621

Letter to the Editor

[DOI: 10.1002/eji.202048577]

High glucose distinctively regulates Ca^{2+} influx in cytotoxic T lymphocytes upon target recognition and thapsigargin stimulation

The killing processes of CD8^+ cytotoxic T lymphocytes (CTLs) are highly dependent on calcium, both extracellular and cytosolic [1]. Engagement of TCRs triggers store-operated calcium entry (SOCE) mediated by stromal interaction molecules (STIM) to trap and activate ORAI channels [2]. Glucose is essential for effector functions of CTLs [3]. Elevated level of blood glucose is one of the leading symptoms of diabetes. Yet, the impact of high glucose (HG) on CTL function is not well understood.

To investigate the impact of HG on Ca^{2+} influx in CTLs, we used primary human CD8^+ T cells, stimulated with anti-CD3/anti-CD28 antibody-coated beads (CD3/CD28 beads) and cultured in normal glucose medium (NG, 5.6 mM) or HG medium (25 mM). To elicit SOCE, we used thapsigargin (Tg) to deplete Ca^{2+} from the ER [4]. For Tg-induced SOCE, we found no difference between NG- and HG-condition for unstimulated CD8^+ T cells (Fig. 1A–D) or at day 1.5 poststimulation (Fig. 1E–H). Remarkably, at days 3 and 7 after bead stimulation, Tg-induced SOCE

was higher in HG-cultured CTLs compared to their NG counterparts (Fig. 1I–P). Upon bead stimulation, CTLs exhibited a significantly enhanced SOCE, which started to decline in NG condition but not in HG condition from day 3 (Fig. 1Q, R). These results suggest that Tg-elicited SOCE in CTLs is elevated by prolonged treatment of HG.

Target cell recognition, a more physiological stimulus than Tg, also elicits SOCE in CTLs. We used staphylococcal enterotoxin A and B (SEA/SEB)-pulsed Raji cells as target cells, which induce CTL-mediated cytotoxicity [1, 5]. We observed that at day 3 poststimulation, in HG-cultured CTLs, target recognition-induced Ca^{2+} influx was significantly reduced relative to their NG counterparts, despite of the same basal level of intracellular Ca^{2+} (Fig. 2A–D). At day 7, upon target cell engagement, only a moderate difference in the Ca^{2+} influx rate and the Ca^{2+} peak was identified between NG- and HG-cultured CTLs, whereas no difference in Ca^{2+} plateau (Fig. 2E–H). Furthermore, the Raji cells without SEA/SEB-pulsing did not elicit Ca^{2+} influx in CTLs (Fig. 2I). Together, we conclude that in CTLs, target recognition-induced Ca^{2+} influx is reduced by HG-culture.

Next, we investigated how HG could differently regulate SOCE induced by Tg or target recognition in CTLs. We first examined glucose uptake using a fluorescent analog of glucose 2-NBDG, and found no difference between HG- and NG-cultured CTLs (Fig. 2J, K). Subsequently, we analyzed the expression of STIMs and ORAIs. At mRNA level, in HG-cultured CTLs, *ORAI1* stayed unchanged, while *ORAI2* and *ORAI3* were slightly upregulated; both *STIM1* and *STIM2* were downregulated

(Fig. 2L, M). Interestingly, at protein level, *STIM1*, *STIM2*, and *ORAI1* in HG-CTLs were upregulated in Donor 1 and 4, but remained unchanged in Donor 2 and 3 (Fig. 2N, O). Due to lack of reliable antibody against *ORAI2* and *ORAI3*, their protein levels could not be determined. Our findings suggest that the increase in Tg-induced SOCE in CTLs by HG could be mediated by upregulation of *STIM1* and *STIM2*. Since *ORAI2* and *ORAI3* mainly function as negative regulators of Ca^{2+} entry in T cells [6, 7], upregulation of *ORAI2* and *ORAI3* by HG could contribute to the reduction of target recognition-induced Ca^{2+} influx. Inhibitory splice variants of STIMs, for example, *STIM2.1* or phosphorylation of STIMs could also play a role.

After glucose loading, proportion of CTLs is reduced in humans [8]. Interestingly, we found that after killing started, the fraction of necrotic CTLs was higher in HG-condition after 16 h but not after 2 h (Fig. 2P, Q), suggesting that in a time scale of several days in HG, CTLs are more prone toward necrosis after execution of their killing function.

One main difference between Tg and target-recognition is that engagement of TCRs upon target-recognition activates many signaling cascades, whereas Tg only blocks the sarco/endoplasmic reticulum Ca^{2+} -ATPase, leading to passive depletion. In addition, CTLs are stimulated globally with Tg and locally with target cells at the immunological synapse (IS). Along this line, other factors could also be involved, for example, mitochondria or PMCA (plasma membrane Ca^{2+} ATPase). Both are enriched at the IS [9, 10] but scarcely distributed in Tg-stimulation. Considering that decreased intracellular




Figure 1. SOCE in CTLs is enhanced by HG-culture. Ca²⁺ imaging was conducted (20 × objective). Primary human CD8⁺ T cells were either unstimulated or stimulated with CD3/CD28 beads in NG medium (5.6 mM) or HG (25 mM) medium and then loaded with Fura-2-AM. Exemplary cells are shown in A, E, I, and M. Calcium influx curves are shown in B, F, J, N (n [independent experiments/donors] = 4/6 [Day 0/Day 1.5], 5/7 [Day 3], or 5/8 [Day 7]). Paired t-test was used except for Q and R (unpaired t-test). Results are presented as mean ± SEM. Scale bars are 5 μm.

Ca²⁺ concentration in CTLs is correlated with enhanced CTL-mediated cytotoxicity [1], our findings suggest that increase the environmental glucose for a short period of

time might be beneficial for CTLs to fight aberrant cells more efficiently. However, considering our result that HG-treated CTLs tend to die more after killing, in dia-

betic patients, it might lead to a reduction in CTL numbers, resulting in a comprised immune response against pathogens in a long run.





Figure 2. HG-culture reduces Ca²⁺ influx in CTLs upon target recognition and impairs CTL viability after killing. (A-I) Ca²⁺ imaging was conducted. CD3/CD28 bead-stimulated CTLs were loaded with Fura-2-AM (20 × objective, Scale bars: 5 μm). Exemplary cells are shown in A and E. Raji cells pulsed with (A-H) or without (I) SEA/SEB. n (independent experiments/donors) = 5/7 (A-D), 3/6 (E-H), 3/5 (I, Day3), or 3/6 (I, Day7) (J, K) Glucose uptake in CTLs. Day 3 CD3/CD28 bead-stimulated CTLs were loaded with 2-NBDG without target cells. Light gray: unstained cells. n = 5 independent experiments/3 donors. (L, M) Expression of ORAIs and STIMs in CTLs. The mRNA level was determined using qRT-PCR and either normalized to ORA11 in NG-cultured CTLs (L) or normalized to NG condition as fold change (M). n = 2 independent experiments/6 donors. (N, O) Protein levels of STIMs and ORA11. Quantification by densitometry analysis following Western blotting (N) and normalization to β-actin (O). n = 4 donors. (P, Q) Necrosis of CTLs after killing. Bead-stimulated CTLs were co-incubated with indicated target cells for 2 h (P, Raji n = 2 independent experiments/3 donors, Raji-SEA/SEB n = 4 independent experiments/5 donors.) or 16 h (Q, Raji n = 3 independent experiments/4 donors, Raji-SEA/SEB n = 5 independent experiments/6 donors). Propidium iodide (PI) was used to determine necrotic CTLs death. Gating strategy shown in Supporting information Fig. 1. Gray: no PI-staining. Paired t-test was used for all statistical analysis. Data shown as mean ± SEM.

Huajiao Zou^{1,2}, Wenjuan Yang²,
Gertrud Schwärz², Renping Zhao²,
Dalia Alansary³, Deling Yin^{1,4},
Eva C. Schwarz², Barbara A. Niemeyer³
and Bin Qu^{2,5} 

¹ Xiangya School of Pharmaceutical Sciences,
Central South University, Changsha, China

² Biophysics, Center for Integrative Physiology and
Molecular Medicine (CIPMM), School of
Medicine, Saarland University, Homburg,
Germany

³ Molecular Biophysics, CIPMM, School of
Medicine, Saarland University, Homburg,
Germany

⁴ Department of Internal Medicine, College of
Medicine, East Tennessee State University,
Johnson City, TN, USA

⁵ INM-Leibniz Institute for New Materials,
Saarbrücken, Germany

Acknowledgments: We thank the Institute for Clinical Hemostaseology and Transfusion Medicine for providing donor materials; Carmen Hässig and Susanne Renno for excellent technical help; Girish Ramesh for Western blotting; Sandra Janku for her careful and critical reading. We are grateful to Markus Hoth for constant support, inspiring discussions, and advice regarding writing of the manuscript. This project was funded by the Deutsche Forschungsgemeinschaft (SFB

1027, A2 to B.Q., C4 to B.N.; Forschungs-
großgeräte (GZ: INST 256/423-1 FUGG)
for the flow cytometer), and by the Leibniz-
Gemeinschaft (INM Fellow Program, to
B.Q.).

Open access funding enabled and orga-
nized by Projekt DEAL.

Ethical Considerations: Research car-
ried out for this study with human mate-
rial (leukocyte reduction system chambers
from human blood donors) is authorized
by the local ethic committee (declaration
from 16.4.2015 [84/15; Prof. Dr. Rettig-
Stürmer]).

Conflict of Interest: The authors declare
no commercial or financial conflict of inter-
est.

References

- Zhou, X. et al., *J. Physiol.* 2018. **596**: 2681–2698.
- Amcheslavsky, A. et al., *Biophys. J.* 2015. **108**: 237–246.
- Geltink, R. I. K. et al., *Annu. Rev. Immunol.* 2018. **36**: 461–488.
- Parekh, A. B. and Putney, J. W., Jr., *Physiol. Rev.* 2005. **85**: 757–810.
- Qu, B. et al., *J. Immunol.* 2011. **186**: 6894–6904.
- Vaeth, M. et al., *Nat. Commun.* 2017. **8**: 14714.
- Alansary, D. et al., *Biochim. Biophys. Acta* 2015. **1853**: 1541–1550.
- Miya, A. et al., *Front. Endocrinol.* 2018. **9**.
- Quintana, A. et al., *EMBO J.* 2011. **30**: 3895–3912.
- Korthals, M. et al., *Sci. Rep.* 2017. **7**: 8358.

Keywords: cytotoxic T cells • glucose • metabolism • calcium • SOCE

Full correspondence: Dr. Bin Qu, Biophysics,
CIPMM, School of Medicine, Saarland
University, 66421 Homburg, Germany.
INM-Leibniz Institute for New Materials
66123 Saarbrücken, Germany.
e-mail: bin.qu@uks.eu

The peer review history for this article is
available at <https://publons.com/publon/10.1002/eji.202048577>

Received: 6/2/2020

Revised: 11/6/2020

Accepted: 21/7/2020

Accepted article online: 22/7/2020



The detailed *Materials and methods*
for Letters to the Editor are available
online in the Supporting
information