

East Tennessee State University

Digital Commons @ East Tennessee State University

[ETSU Faculty Works](#)

[Faculty Works](#)

5-7-2017

Chrotopterus auritus (Peters, 1856) (Chiroptera, Phyllostomidae): First Record for the State of Rio Grande Do Norte, Northeastern Brazil

Gustavo Henrique Nunes Basílio
Centro

Jan Pierre Martins de Araujo
Centro

Juan Carlos Vargas Mena
Universidade Federal do Rio Grande do Norte

Patrício A. da Rocha
Universidade Federal da Paraíba

Marcelo Augusto Freitas Kramer
Centro

Follow this and additional works at: <https://dc.etsu.edu/etsu-works>

Citation Information

Basílio, Gustavo Henrique Nunes; de Araujo, Jan Pierre Martins; Mena, Juan Carlos Vargas; da Rocha, Patrício A.; and Kramer, Marcelo Augusto Freitas. 2017. Chrotopterus auritus (Peters, 1856) (Chiroptera, Phyllostomidae): First Record for the State of Rio Grande Do Norte, Northeastern Brazil. *Check List*. Vol.13(3). <https://doi.org/10.15560/13.3.2110>

This Article is brought to you for free and open access by the Faculty Works at Digital Commons @ East Tennessee State University. It has been accepted for inclusion in ETSU Faculty Works by an authorized administrator of Digital Commons @ East Tennessee State University. For more information, please contact digilib@etsu.edu.

Chrotopterus auritus (Peters, 1856) (Chiroptera, Phyllostomidae): First Record for the State of Rio Grande Do Norte, Northeastern Brazil

Copyright Statement

This is an open access article distributed under the terms of the [Creative Commons Attribution License \(CC BY 4.0\)](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Creative Commons License



This work is licensed under a [Creative Commons Attribution 4.0 International License](#).



Chrotopterus auritus (Peters, 1856) (Chiroptera, Phyllostomidae): first record for the state of Rio Grande do Norte, northeastern Brazil

Gustavo Henrique Nunes Basílio^{1,5}, Jan Pierre Martins de Araujo¹, Juan Carlos Vargas Mena²,
Patrício A. da Rocha³ & Marcelo Augusto Freitas Kramer^{1,4}

¹ Sociedade Espeleológica Potiguar. CEP 59375-000, Rua da Matriz, Centro, Acari, Rio Grande do Norte, Brazil

² Departamento de Ecologia, Programa de Pós-Graduação em Ecologia, Universidade Federal do Rio Grande do Norte, Centro de Biociências Campus Lagoa Nova, 59072-970, Natal, Rio Grande do Norte, Brazil

³ Programa de Pós-Graduação em Ciências Biológicas (Zoologia), CCEN, Universidade Federal da Paraíba, Campus I, 58051-900, João Pessoa, Paraíba, Brazil

⁴ Department of Geosciences and Don Sundquist Center of Excellence in Paleontology, East Tennessee State University, Johnson City, TN 37614, USA

⁵ Corresponding author. E-mail: gustavohenriquebiologia@gmail.com

Abstract. *Chrotopterus auritus* is a phyllostomid bat with a wide distribution in the Neotropics. It has been recorded in Brazil's 6 biomes but with few records in the Caatinga. We provide the first record of *C. auritus* for Rio Grande do Norte state, northeastern Brazil, based on records from 2 caves, Três Inchu and Gruta da Carrapateira. The nearest records are ca. 400 km southeast in Ceará state and ca. 350 km northwest in Pernambuco state. Our new records fill the northeastern distributional gap of *C. auritus* in Brazil and South America.

Key words. Bats; *Chrotopterus*; geographic distribution; Caatinga; caves

Chrotopterus auritus (Peters, 1856) is one the largest phyllostomid bats and the single representative of the genus *Chrotopterus* Peters, 1865 (SIMMONS 2005). As well as *Vampyrum spectrum* (Linnaeus, 1758) and *Trachops cirrhosus* (Spix, 1823), *C. auritus* is considered a predominantly carnivorous species (FERRAREZZI & GIMENEZ 1996) but occasionally including insects (MEDELLÍN 1988) and fruits (BERNARD 2002) in its diet. It is known to occur in southern Mexico, Central America, Colombia, Venezuela, Guyana, Surinam, French Guiana, Brazil, Ecuador, Peru, Bolivia, Paraguay, and northern Argentina (WILLIAMS & GENOWAYS 2008). In Brazil, this species has been recorded in all biomes (PAGLIA et al. 2012) but there are few records for the Caatinga (Fig. 1; Table 1). The Caatinga is restricted to Brazil and is a mosaic of tropical dry forest with xeric shrublands (LEAL et al. 2003). With a predominant semi-arid climate (MMA 2017) and located in the Northeast Region, the Caatinga has an area of over 800,000 km², which represents about 11% of Brazil's territory, and covering 9 states and the northern part Minas Gerais. Herein, we provide the first occurrence record of *Chrotopterus auritus* for the state of Rio Grande do Norte, northeastern Brazil, based

on records from 2 separate caves in the Caatinga on the west side of the state.

On 28 August 2015 at 23:40 h an individual of *C. auritus* was observed and photographed in the Gruta da Carrapateira. Capture attempts were unsuccessful but the individual was identified through direct observation using binoculars and photographic records. The calcareous cave is located in the municipality of Felipe Guerra (05°33'38.2"S, 037°39'50.3"W, altitude ca. 32 m) about 348 km west of Natal and has 5 entrances and a linear development of 242.2 m.

The second record is based on a colony of 8 individuals and an adult male captured with a hand net (Fig. 3) inside the Três Inchu cave (06°03'07"S, 037°54'03"W) on 18 December 2015. The Três Inchu is a marble cave located in the municipality of Martins, ca. 362 km from Natal, with a linear development of 146.0 m and 5 recognized entrances, some of which are skylights, providing for photic zones inside.

The Três Inchu specimen was handled in accordance with SIKES et al. (2011), euthanized and fixed in 10% formaldehyde and preserved in 70% ethanol with subsequent extraction of the skull (Fig. 2). Cranial and external measurements were taken following VIZOTTO & TADDEI (1973) using a digital caliper (0.2 mm). The specimen was collected under permit SISBIO/IBAMA 52492-1 and is deposited in the mammal collection of the Federal University of Sergipe (CMUFS) under voucher number CMUFS 259.

Our voucher of *Chrotopterus auritus* has the set of characters that distinguish this species from other members of the Phyllostominae, such as its large size (forearm 74–83 mm), presence of 1 pair of lower incisors and 3 lower premolars (Fig. 2), rudimentary tail, calcar longer than the foot, large rounded ears not connected by a band, horseshoe of noseleaf cup-shaped and continuous with spear, and long thumb with a large strongly curved claw (Fig. 3) (MEDELLÍN 1989; WILLIAMS




Figure 1. Records of *Chrotopterus auritus* in Brazil. The numbers corresponding to the records are indicated in Table 1. Blue stars represent the new records from the state of Rio Grande do Norte, Brazil.

Table 1. Locality records for *Chrotopterus auritus* in Brazil. The code numbers refer to the points shown in Figure 1.

Code	Locality	State	Latitude (S)	Longitude (W)	References
1	Parque Nacional da Serra do Divisor	AC	08°03'	073°55'	NOGUEIRA (1999)
2	Santa Isabel do Rio Negro	AM	00°04'	065°01'	MORATELLI et al. (2010)
3	Barcelos	AM	00°96'	062°91'	MORATELLI et al. (2010)
4	Parque Nacional Montanha do Tumucumaque	AP	01°06'	052°48'	MARTINS et al. (2006)
5	Floresta Nacional do Amapá	AP	00°66'	051°16'	MARTINS et al. (2006)
6	Melgaço	PA	01°08'	050°07'	MARQUES-AGUIAR et al. (2003)
7	Uruará	PA	03°28'	053°63'	CAJABA (2014)
8	São Luiz	MA	02°51'	044°03'	CRUZ et al. (2007)
9	Ubajara	CE	03°08'	040°83'	UIEDA et al. (1980)
10	Gruta da Carrapateira cave, Felipe Guerra	RN	05°55'	037°65'	This study
11	Três Inchu cave, Martins	RN	06°05'	037°09'	This study
12	Ecológico Municipal Professor João Vasconcelos Sobrinho, Caruaru	PE	08°28'	035°96'	SOUSA et al. (2004)
13	Una	BA	15°16'	039°05'	FARIA et al. (2006)
14	Chapada Diamantina	BA	13°56'	041°18'	GREGORIN & MENDES (1999)
15	Chapada Diamantina	BA	11°23'	041°71'	SBRAGA & CARDOSO (2008)
16	Médio São Francisco, Paratinga	BA	12°73'	043°09'	SÁ-NETO & MARINHO-FILHO (2013)
17	Linhares	ES	19°41'	040°05'	PERACCHI & ALBUQUERQUE (1993)

Continued

Table 1. *Continued.*

Code	Locality	State	Latitude (S)	Longitude (W)	References
18	Jaíba	MG	15°13'	043°54'	NOGUEIRA et al. (2015)
19	Piarapora	MG	17°35'	044°78'	TAVARES et al. (2010)
20	Uberlândia	MG	19°00'	048°66'	PEDRO & TADDEI (2014)
21	São Tomé das Letras	MG	21°71'	044°98'	TAVARES et al. (2010)
22	Serra Negra	MG	21°96'	043°88'	NOBRE et al. (2009)
23	Sete Lagoas	MG	19°84'	044°18'	TAVARES et al. (2010)
24	Santa Barbara	MG	19°95'	043°04'	TAVARES et al. (2010)
25	Serra da Mantiqueira	MG	19°58'	042°05'	STALLINGS et al. (1990)
26	Paraíso do Tobias, Miracema	RJ	21°04'	042°06'	ESBÉRARD et al. (2010)
27	Parque Estadual Desengano	RJ	21°83'	041°83'	MODESTO et al. (2008)
28	Silva Jardim	RJ	22°66'	042°39'	BAPTISTA & MELLO (2001)
29	Pedra Branca	RJ	22°09'	043°02'	DIAS et al. (2002)
30	Engenheiro Paulo de Frontin	RJ	22°55'	043°66'	DIAS et al. (2010)
31	Vassouras	RJ	22°04'	043°65'	DIAS et al. (2002)
32	Mangaratiba	RJ	22°98'	044°01'	LUZ et al. (2011)
33	Ilha Grande - Angra dos Reis	RJ	23°15'	044°22'	ESBÉRARD et al. (2006)
34	Itaperina	SP	22°23'	047°83'	UIEDA et al. (2007)
35	Estação Experimental de Itirapina	SP	22°32'	047°84'	SATO et al. (2015)
36	Ribeirão Preto	SP	21°22'	047°84'	PASSOS et al. (2003)
37	Estação Ecológica dos Caetetus, Gália	SP	22°27'	049°61'	PEDRO et al. (2014)
38	Parque Nacional das Emas	GO	18°25'	052°88'	RODRIGUES et al. (2002)
39	Indiara	GO	17°00'	050°00'	SILVA et al. (2011)
40	Posse	GO	14°31'	046°23'	ESBÉRARD et al. (2001)
41	APA Nascente Rio Vermelho	GO	14°28'	046°25'	ESBÉRARD et al. (2005)
42	Alvorada do Norte	GO	14°05'	046°71'	BEZERRA & MARINHO-FILHO (2010)
43	Planaltina	DF	15°55'	047°75'	BREDT & MAGALHAES (1999)
44	Paranoá	DF	15°86'	047°08'	BREDT & MAGALHAES (1999)
45	Brazlândia	DF	15°55'	048°11'	BREDT & MAGALHAES (1999)
46	Near to Nova Xavantina	MT	15°01'	052°15'	OLIVEIRA et al. (2015)
47	Cedro Trail, lower Juruena River	MT	08°83'	058°46'	DALPONTE et al. (2016)
48	Estação Ecologica Serra das Araras	MT	15°45'	057°05'	GONÇALVES & GREGORIN (2004)
49	Corumbá	MS	18°96'	057°65'	BORDIGNON & FRANÇA (2004)
50	Nhecolândia	MS	19°00'	056°00'	OLIVEIRA et al. (2011)
51	Abobral	MS	19°05'	056°5'	OLIVEIRA et al. (2011)
52	Miranda	MS	20°00'	056°05'	OLIVEIRA et al. (2011)
53	Parque Nacional Serra da Bodoquena	MS	20°99'	056°83'	CAMARGO et al. (2009)
54	Campo Grande	MS	20°43'	054°63'	BORDIGNON (2005)
55	Chapadão do Sul	MS	19°18'	052°76'	BORDIGNON et al. (2006)
56	Paraíso das Águas	MS	19°02'	053°01'	DALPONTE et al. (2016)
57	Campinhos	PR	25°03'	049°08'	ARNONE & PASSOS (2003)
58	Telêmaco Borba	PR	24°02'	050°55'	REIS et al. (1999)
59	Londrina	PR	23°38'	051°18'	REIS et al. (2003)
60	São Pedro do Ivaí	PR	23°81'	051°93'	CARNEIRO (2008)
61	Fênix	PR	23°91'	051°95'	BIANCONI et al. (2004)
62	Cianorte	PR	23°66'	052°63'	ORTÉNCIO-FILHO & REIS (2009)
63	Foz do Iguaçú	PR	25°51'	054°58'	GRACIOLLI & CARVALHO (2001)
64	Blumenau	SC	27°08'	049°08'	ALTHOFF (2007)
65	Pedras Grandes	SC	28°48'	049°25'	CARVALHO et al. (2013)
66	Vale do Taquari	RS	29°66'	050°04'	KASPER et al. (2007)
67	São Francisco de Paula	RS	29°41'	050°04'	MARQUES et al. (2011)
68	Barracão	RS	27°56'	051°53W	WITT et al. (2010)
69	Machadinho	RS	27°58'	051°06'	WITT et al. (2010)
70	Rio Grande do Sul	RS	27°66'	053°11'	PETERS et al. (2010)
71	Frederico Westphalen	RS	27°33'	053°38'	BERNARDI et al. (2009)
72	Derrubados	RS	27°33'	054°05'	KASPER et al. (2007)
73	Rio Grande	RS	31°49'	052°49' W	QUINTEL A et al. (2011)




Figure 2.Dorsal, ventral and lateral views of the skull and lateral view of the mandible (scale bar=10 mm) of *Chrotopterus auritus* (CMUFS 259) from Três Inchu cave, Martins municipality, Rio Grande do Norte state, northeast Brazil. In detail, front view of the mandible showing the presence of 2 incisors (scale bar = 5 mm).

& GENOWAYS 2008).

Cranial and external measurements of the voucher, in millimeters, are: forearm length (84.6), hind foot length (21.1), calcar (22.3), ear (36.2), greatest length of skull (37.12), condylobasal length (32.41), breadth of braincase (14.28), post-orbital constriction (6.06), breadth across upper molars (12.18), breadth across upper canines (7.92), length of mandible (24.78) and length of maxillary tooth row (15.18). These measurements are within the known range for the species (see MEDELLÍN 1988). In the case of the individual from Gruta da Carrapateira, not captured, it was possible to see external diagnostic characters of *C. auritus*, such as the cup-shaped horseshoe of the noseleaf and the long thumbs with large claw.

Our records of *Chrotopterus auritus* from Rio Grande do Norte fill the distributional gap of this species in northeastern

Brazil. They extend the distribution of *C. auritus* in ca. 400 km southeast and 350 km northwest from the Ceará (UIEDA et al. 1980) and Pernambuco (SOUSA et al. 2004) states, respectively.

The presence of *C. auritus* within the Caatinga may be unusual, since there are only 6 previous records from this biome (Fig. 1; Table 1). The records are: in xeric-woodlands of Caatinga (SÁ-NETO & MARINHO-FILHO 2013); in Chapada Diamantina National Park in Bahia state (GREGORIN & MENDES 1999; SBRAGIA & CARDOSO 2008); in transition areas of Caatinga and the Amazon (UIEDA et al. 1980); in humid forest enclaves within the Caatinga known as “Brejos de Altitude” (SOUSA et al. 2004); and in the southern limit of the Caatinga in Jaíba, northern Minas Gerais state (NOGUEIRA et al. 2015). Moreover, 4 of these records were obtained from underground cavities (GREGORIN & MENDES 1999; SBRAGIA &



Figure 3. *Chrotopterus auritus* photographed at the Gruta Três Inchu, Martins municipality, Rio Grande do Norte state, northeast Brazil. Note the long thumb (arrow a) with a large strongly curved claw and the cup-shaped horseshoe (arrow b).

CARDOSO 2008; UIEDA et al. 1980; NOGUEIRA et al. 2015).

Chrotopterus auritus is known to roost in caves, abandoned mines, buildings, and hollow trees (REID 2009). It inhabits mature evergreen forest and deciduous forests, but records in this latter formation are less common (MEDELLÍN 1989). Because *C. auritus* is a top predator, more regularly found in undisturbed habitats, its presence is considered an indicator of healthy forests (MEDELLÍN 1989; GORRENSEN & WILLIG 2004). Individuals generally form small groups from 3 to 6 individuals in their roost (ARITA & VARGAS 1995; BREDT et al. 1999). When roosting in caves, this species is commonly found in cohabitation with other bat species (COIMBRA et al. 1982; TRAJANO 1985; BREDT et al. 1999). We observed *C. auritus* cohabitating with *Desmodus rotundus* (E. Geoffroy, 1810) and *Peropteryx macrotis* (Wagner, 1843) in Três Inchu cave and with *Artibeus planirostris* (Spix, 1823), *Furipterus horrens* (F. Cuvier, 1828), *Desmodus rotundus*, *Diphylla ecaudata* Spix, 1823, *Glossophaga soricina* (Pallas, 1766) and *P. macrotis* in Gruta da Carrapateira.

Caves in the Caatinga, including those in the state of Rio Grande do Norte, may be important roosting sites for *C. auritus*, and the presence of a high density of caves in the landscape may be a good predictor of a potential occurrence of this species. Furthermore, roosting in caves may provide benefits to *C. auritus* besides providing stable roost conditions. The cohabitation with others bats inside caves may allow for the predation of other smaller bats, a behavior already reported for *C. auritus* (BORDIGNON 2005; WITT & FABIAN 2010).

Rio Grande do Norte is one of the least known states in relation to its chiropterofauna, with only 11 species currently recorded (*Desmodus rotundus*, *Diphylla ecaudata*, *Lonchorhina aurita* Tomes, 1863, *Artibeus planirostris*, *Trachops cirrhosus*, *Natalus macrourus* (Gervais, 1856), *Peropteryx macrotis*, *Dermanura cinerea* (Gervais, 1856), *Molossus molossus* (Pallas, 1766), *Myotis nigricans* (Schinz, 1821), *Glossophaga soricina*; OLIVEIRA et al. 2003; FEIJÓ & NUNES 2010; FERREIRA et al. 2010; BARROS 2014). With 949 cavities currently cataloged, the state is fourth in Brazil for the number of natural underground cavities (CECAV/ICMBio 2016), which constitute potential important roost for the bat communities in the Caatinga of the state.

An increase of bat inventories in Rio Grande do Norte, both in caves as well as in other habitats, should provide a better understanding of the distribution of *C. auritus* and more generally the species richness in this state and in the Caatinga biome.

ACKNOWLEDGEMENTS

We thank the Sociedade Espeleológica Potiguar (SEP) for providing our main source of funding, the Universidade Federal de Sergipe (UFS) and the Universidade Federal do Rio Grande do Norte (UFRN) for supporting our study. We also thank Solon Rodrigues de Almeida Neto, Aldo Guimarães, Eugenia Cordero Schmidt, and Valtencí Santana for field assistance and Blaine W. Schubert for helping improve the manuscript. JCVM thanks the National Council for Scientific and Technological Development (CNPq) for a research grant (Pes-

quisador Visitante Especial-PVE “Ecologia e Conservação de Morcegos na Caatinga Potiguar” Project: 401467/2014-7); CAPES for the post-graduation scholarship, and the Wildlife Conservation Society (WCS) of Brazil for the help with field logistics. PAR thanks CNPq for a research grant (processes 501701/2013-3 and 150407/2015-7).

LITERATURE CITED

- ARITA, H.T. & J.A. VARGAS. 1995. Natural history, interspecific association, and incidence of the cave bats of Yucatan, Mexico. The Southwestern Naturalist 40(1): 29–37. <http://www.jstor.org/stable/30054390>
- ARNONE, I.S. & F.C. PASSOS. 2007. Estrutura de comunidade da quiropterofauna (Mammalia, Chiroptera) do Parque Estadual de Campinhos, Paraná, Brasil. Revista Brasileira de Zoologia 24(3): 573–581. <https://doi.org/10.1590/0031-1049.2015.55.01>
- BAPTISTA, M. & M. MELLO. 2001. Preliminary inventory of the bat species of the Poço das Antas Reserve, RJ. Chiroptera Neotropical 7(1–2):133–135. <http://chiropteraneotropical.net/index.php/cn/article/view/123>
- BARROS, M.A. 2014. First record of *Molossusmolossus* (Pallas, 1766) (Mammalia: Chiroptera) in the state of Rio Grande do Norte, northeastern Brazil. Check List 10(6): 1520–1524. <https://doi.org/10.15560/10.6.1520>
- Bernard, E. 2002. Diet, activity and reproduction of bats species (Mammalia, Chiroptera) in Central Amazonia, Brazil. Revista Brasileira de Zoologia 19(1):173–188. <https://doi.org/10.1590/S11-81752002000100016>
- BERNARDI, I.P., J.M. D. MIRANDA, J. SPOONCHIADO, E. GROTTI, F.F. JACOMASSA, E.M. TEIXEIRA, S.H. ROANI & F.C. PASSOS. 2009. Morcegos de Frederico Westphalen, Rio Grande do Sul, Brasil (Mammalia: Chiroptera): riqueza e utilização de abrigos. Biota Neotropica 9(3): 349–354. <http://www.biota-neotropica.org.br/v9n3/pt/abstract?article+bn03009032009>
- Bezerra, A.M. and J. Marinho-Filho. 2010. Bats of the Paraná river valley, Tocantins and Goiás states, central Brazil. Zootaxa 2725: 41–56.
- BIANCONI, G.V., S.B. MIKICH & W.A. PEDRO. 2004. Diversidade de morcegos (Mammalia, Chiroptera) em remanescentes florestais do município de Fênix, noroeste do Paraná, Brasil. Revista Brasileira de Zoologia 21(4): 943–954. <https://doi.org/10.1590/S0101-81752004000400032>
- BORDIGNON, M. O. 2005. Predação de morcegos por *Chrotopterus auritus* (Peters, 1856) (Mammalia, Chiroptera) no Pantanal de Mato Grosso do Sul, Brasil. Revista Brasileira de Zoologia 22(4): 1207–1208. <https://doi.org/10.1590/S0101-81752005000400058>
- BORDIGNON, M.O. 2006. Diversidade de morcegos (Mammalia, Chiroptera) do Complexo Aporé-Sucuriú, Mato Grosso do Sul, Brasil. Revista Brasileira de Zoologia 23(4): 1002–1009. <https://doi.org/10.1590/S0101-81752006000400004>
- BORDIGNON, M.O. & A.O. FRANÇA. 2004. Análise preliminar sobre a diversidade de morcegos no Maciço do Urucum, Mato Grosso do Sul, Brasil. IV Simpósio sobre os recursos naturais e sócio-econômicos do Pantanal. Corumbá: Embrapa. <http://www.cgap.embrapa.br/agencia/simpan/sumario/artigos/aspectos/pdf/bioticos/C%2A2pia%20de%2020604RBBordignon%20OKVisto.pdf>
- BORDIGNON, M.O., N.C. CÁCERES, A.O. FRANÇA, J. CASELLA & C.F. VARGAS. 2006. Inventário da Mastofauna no Complexo Aporé-Sucuriú; pp. 129–142, in: T.C.S. PAGOTTO & P.R. SOUZA (eds.). Biodiversidade do complexo Aporé-Sucuriú: subsídios à conservação e manejo do bioma Cerrado. Campo Grande: Editora da UFMS.
- BREDT, A., W. UIEDA & E.D. MAGALHÃES. 1999. Morcegos cavernícolas da região do Distrito Federal, centro-oeste do Brasil. Revista

- Brasileira de Zoologia 16(3): 731–770. <https://doi.org/10.1590/S0101-81751999000300012>
- Cajaiba, R.L. 2014. Morcegos (Mammalia, Chiroptera) em cavernas no município de Urucá, Pará, norte do Brasil. Biota Amazônica 4(1): 81–86. <http://dx.doi.org/10.18561/2179-5746/biotaamazonia.v4n1p81-86>
- CAMARGO, G., E. FISCHER, F. GONÇALVES, G. FERNANDES & S. FERREIRA. 2009. Morcegos do Parque Nacional da Serra da Bodocuema, Mato Grosso do Sul, Brasil. Chiroptera Neotropical 15(1): 417–424. <https://doi.org/10.1111/j.0030-1299.2004.12884.x>
- CARVALHO, F., M.E. FABIAN & J.O. MENEGHETI. 2013. Vertical structure of an assemblage of bats (Mammalia: Chiroptera) in a fragment of Atlantic Forest in Southern Brazil. Zoologia 30(5): 491–498. <https://doi.org/10.1590/S1984-46702013000500004>
- CECAV/ICMBIO. 2016. Centro Nacional de Pesquisa e Conservação de Cavernas. Base de Dados Geoespecializados de Cavidades Naturais Subterrâneas do CECAV. Accessed at http://www.ibama.gov.br/cecav/%20index.php?Id_menu=228, 28 October 2016.
- COIMBRA JR, C. E. A., M. M BORGES, D.Q. GUERRA, & D.A. MELLO. 1982. Contribuição à zoogeografia e ecologia de morcegos em regiões de cerrado do Brasil Central. Boletim Técnico Revista Brasil Florestal 7: 34–38.
- CRUZ, L.D., C. MARTÍNEZ & F.R. FERNANDES. 2007. Comunidades de morcegos em habitats de uma Mata Amazônica remanescente na Ilha de São Luís, Maranhão. Acta Amazonica 37(4): 613–619. <https://doi.org/10.1590/S0044-59672007000400017>
- DALPONTE, J.C., R. GREGORIN, V.A. ESTEVEZ-COSTA, E.C. ROCHA & R. MARCELINHO. 2016. Bat survey of the lower Juruena River and five new records for the state of Mato Grosso, Brazil. Acta Amazonica 46(2): 227–232. <https://doi.org/10.1590/1809-4392201500888>
- DIAS, D., A.L. PERACCHI, & S.S.P DA SILVA. 2002. Quirópteros do Parque Estadual da Pedra Branca, Rio de Janeiro, Brasil (Mammalia, Chiroptera). Revista Brasileira de Zoologia 19(2): 113–140. <https://doi.org/10.1590/S0101-81752002000600012>
- DIAS, D., S.N. PEREIRA, A.C. MAAS, M.A. MARTINS, M.A., D.P. BOLZAN & A.L. PERACCHI. 2010. Quirópteros das regiões Centro-Sul e Médio Paraíba do estado do Rio de Janeiro (Mammalia, Chiroptera). Chiroptera Neotropical 16(1): 579–585. <http://chiropteraneotropical.net/index.php/cn/article/view/231>
- ESBÉRARD, C. E.L., J. A. MOTTA, E. M. CALVO, V. M. FERREIRA, J. C., CARVALHO, C.C. CARVALHO, C.R.P. SOUZA, E.A. PIRES, G.M.V. ROSA, J.S. REIS, J.N. ARAÚJO, K.E. QUEGE. 2001. Morcegos Cavernícolas de Mambai e Arredores, Goiás, Brasil. Brasília:26vo Congresso Brasileiro de espeleologia. http://www.sbe.com.br/anais26cbe/26CBE_361-364.pdf
- ESBÉRARD, C.E.L., J.A. MOTTA & C. PERIGO. 2005. Morcegos cavernícolas da Área de Proteção Ambiental (APA) Nascentes do Rio vermelho, Goiás. Revista Brasileira de Zoociências 7(2): 311–325. <https://zoociencias.ufjf.emnuvens.com.br/zoociencias/article/view/165>
- ESBÉRARD, C.E. L., T. JORDÃO-NOGUEIRA, J.L. LUZ, G.G.S. MELO, R. MANGOLIN, N. JUCÁ, D.S.L. RAÍCES, M.C. ENRICI & H.G. BERGALLO. 2006. Morcegos da Ilha Grande, Angra dos Reis, RJ, Sudeste do Brasil. Revista Brasileira de Zoociências 8(2): 147–153. <https://zoociencias.ufjf.emnuvens.com.br/zoociencias/article/view/110>
- ESBÉRARD, C.E.L., M. BAPTISTA, L.D. M. COSTA, J.L. LUZ & E.C. LOURENÇO. 2010. Bats from Paraíso do Tobias, Northwest of Rio de Janeiro, Brazil. Biota Neotropica 10(4): 249–255. <https://doi.org/10.1590/S1676-06032010000400030>
- FARIA, D., B. SOARES-SANTOS & E. SAMPAIO. 2006. Bats from the Atlantic rainforest of southern Bahia, Brazil. Biota Neotropica 6(2): 1–13. <https://doi.org/10.1590/S1676-06032006000200022>
- FEIJÓ, J.A. & H.L. NUNES. 2010. Primeiro registro de *Myotis nigricans* (Schinz, 1821) para o estado do Rio Grande do Norte, nordeste do Brasil. Chiroptera Neotropical 16(1): 531–534. <https://doi.org/10.1111/j.0030-1299.2004.12884.x>
- chiroptera.unb.br/index.php/cn/article/view/40
- FERRAREZZI, H. & E.A. GIMENEZ. 1996. Systematic patterns and the evolution of feeding habits in Chiroptera (Archonta: Mammalia). Journal of Comparative Biology 1(3):75–94.
- FERREIRA, R.L., X. PROUS, B.L. F. DE OLIVEIRA & M. SOUZA-SILVA. 2010. Fauna subterrânea do Estado do Rio Grande do Norte: Caracterização e impactos. Revista Brasileira de Espeleologia 1(1): 25–51. <http://www.icmbio.gov.br/revistaelectronica/index.php/RBEsp/article/view/70>
- GRACIOLLI, G. & C.J.B.D. CARVALHO. 2001. Moscas ectoparasitas (Diptera, Hippoboscidae, Nycteriidae) de morcegos (Mammalia, Chiroptera) do Estado do Paraná, Brasil. I. Basilia, taxonomia e chave pictórica para as espécies. Revista Brasileira de Zoologia 18(1): 33–49. <http://www.scielo.br/pdf/rbzoool/vl8sl/vl8suplla02.pdf>
- GREGORÍN, R. & L.D.F. MENDES. 1999. Sobre quirópteros (Emballonuridae, Phyllostomidae, Natalidae) de duas cavernas da Chapada Diamantina, Bahia, Brasil. Iheringia, Série Zoologia 86: 121–124.
- GONÇALVES, E. & R. GREGORIN. 2004. Quirópteros da Estação Ecológica da Serra das Araras, Mato Grosso, Brasil, com o primeiro registro de *Artibeus gnomus* e *A. anderseni* para o cerrado. Lundiana 5(2): 143–149.
- GORRENSEN P.A. & M.R. WILLIG. 2004. Landscape responses of bats to habitat fragmentation in Atlantic forest of Paraguay. Journal of Mammalogy 85(4): 688–697. <https://doi.org/10.1644/BWG-125>
- KASPER, C.B., M.J. FELDEN, F.D. MAZIM, A. SCHNEIDER, C.V. CADEMARTORI & H.C.Z. GRILLO. 2007. Mamíferos do Vale do Taquari, região central do Rio Grande do Sul. Biociências 15(1): 53–62. <http://revistaselectronicas.pucrs.br/ojs/index.php/fabio/article/view/257/1742>
- LEAL, I.R. & J.M.C. DA SILVA. 2003. Ecologia e conservação da Caatinga. Editora Universitária da Universidade Federal de Pernambuco, 3rd edition. Recife, Pernambuco, Brazil. 797 pp.
- LUZ, J.L., L.D.M. COSTA, E.C. LOURENÇO & C.E.L. ESBÉRARD. 2011. Bats (Mammalia, Chiroptera) from Reserva Rio das Pedras, Rio de Janeiro, Southeastern Brazil. Biota Neotropica 11(1): 95–101. <https://doi.org/10.1590/S1676-06032011000100009>
- MARQUES-AGUIAR, S.A., M.V. DEL AGUILA, G.F. AGUIAR, N. SALDANHA, J.D.S. SILVA-JUNIOR & M.M. ROCHA. 2003. Caracterização e perspectivas de estudo dos quirópteros da Estação Científica Ferreira Penna-município de Melgaço-PA. Idéias e Debates, Belém 6: 1–3. & MARQUES, R.V., V.C. CRISTINA & M.P. SUSI. 2011. Mastofauna no Planalto das Araucárias, Rio Grande do Sul, Brasil. Revista Brasileira de Biociências 9(3): 278–288. <http://www.ufrgs.br/seerbio/ojs/index.php/rbb/article/view/1414>
- MARTINS, A.C., E. BERNARD & R. GREGORIN. 2006. Inventários biológicos rápidos de morcegos (Mammalia, Chiroptera) em três unidades de conservação do Amapá, Brasil. Revista Brasileira de Zoologia 23(4): 1175–1184. <https://doi.org/10.1590/S0101-81752006000400026>
- MEDELLÍN, R.A. 1988. Prey of *Chrotopterus auritus*, with notes on feeding behavior. Journal of Mammalogy 69(4): 841–844. <http://www.jstor.org/stable/1381644>
- MEDELLÍN, R.A. 1989. *Chrotopterus auritus*. Mammalian Species 343: 1–5. <https://doi.org/10.2307/3504232>
- MMA. 2017. Ministério do Meio Ambiente. Caatinga. Accessed at <http://www.mma.gov.br/biomass/caatinga>, 24 February 2017.
- MODESTO, T.C., F.S. PESSÔA, M.C. ENRICI, N. ATTIAS, T. JORDÃO-NOGUEIRA, L.D.M. COSTA, H.G. ALBUQUERQUE & H.D.G. BERGALLO. 2008. Mamíferos do Parque Estadual do Desengano, Rio de Janeiro, Brasil. Biota Neotropica 8(4): 153–159. <http://www.biotaneotropica.org.br/v8n4/pt/abstract?article+bn01408042008>
- MORATELLI, R., D. DIAS & C.R. BONVICINO. 2010. Estrutura e análise zoogeográfica de uma taxocenose de morcegos no norte do Estado do Amazonas, Brasil. Chiroptera Neotropical 16(1): 661–671. <http://www.chiropteraneotropical.net/index.php/cn/article/view/254>
- NOBRE, P.H., A.D.S. RODRIGUES, I.D.A. COSTA, A.E.D.S. MOREIRA

- & H.H. MOREIRA. 2009. Similaridade da fauna de Chiroptera (Mammalia), da Serra Negra, municípios de Rio Preto e Santa Bárbara do Monte Verde, Minas Gerais, com outras localidades da Mata Atlântica. Biota Neotropica 9(3): 151–156. <http://www.biota-neotropica.org.br/v9n3/pt/abstract?article+bn03309032009>
- NOGUEIRA, M.N., A. POL & A.L. PERACCHI. 1999. New records of bats from Brazil with a list of additional species for the chiropteran fauna of the state of Acre, western Amazon basin. Mammalia 63(3): 363–367. <https://doi.org/10.1515/mamm.1999.63.3.363>
- NOGUEIRA, M.R., POL, A., PESSÔA, L.M., J.A.D., OLIVEIRA & A.L. PERACCHI. 2015. Small mammals (Chiroptera, Didelphimorphia, and Rodentia) from Jaíba, middle Rio São Francisco, northern Minas Gerais State, Brazil. Biota Neotropica 15(2): 1–18. <https://doi.org/10.1590/1676-06032015012614>
- OLIVEIRA, J.A., P.R. GONÇALVES & C.R. BONVICINO. 2003. Mamíferos da Caatinga; pp. 275–303, in: I.R. LEAL, M. TABARELLI & J.M.C. SILVA (eds.) Ecologia e Conservação da Caatinga. Recife: Universidade Federal de Pernambuco.
- OLIVEIRA, A.K.M., M.D. OLIVEIRA, S. FAVERO, L.F. DE OLIVEIRA. 2011. Diversity, similarity and trophic guild of chiropterofauna in 3 southern Pantanal sub-regions, state of Mato Grosso do Sul, Brazil. Biological Sciences 34(1): 33–39. <https://doi.org/10.4025/actascibiolsci.v34i1.7596>
- OLIVEIRA, S.L., L.A.S. SOUZA, H.K. SILVA & K.C. FARIA. 2015. Spatial configuration of the occurrence of bat species (Mammalia: Chiroptera) in eastern Mato Grosso, Brazil. Biota Neotropica 15(1): e20140122. <https://doi.org/10.1590/1676-06032014012214>
- ORTÊNCIO-FILHO, H. & N.R. REIS. 2009. Species richness and abundance of bats in fragments of the stationalsemideciduous forest, Upper Paraná River, southern Brazil. Brazilian Journal of Biology 69(2): 727–734. <https://doi.org/10.1590/S1519-69842009000300026>
- PAGLIA, A.P., G.A. DA FONSECA, A.B. RYLANDS, G. HERRMANN, L.M. AGUIAR, A.G. CHIARELLO, Y.R.L. LEITE, L.P. COSTA, S. SICILIANO, M.C.M. KIERULFF, S.L. MENDES, V.C. TAVARES, R.A. MITTERMEIER & J.M. PATTON. 2012. Lista anotada dos mamíferos do Brasil, 2^a Edição. Annotated checklist of Brazilian mammals. Occasional papers in Conservation Biology 6: 76 pp.
- PASSOS, F.C., W.R. SILVA, W.A. PEDRO & M.R. BONIN. 2003. Frugivory em morcegos (Mammalia, Chiroptera) no Parque Estadual Intervales, sudeste do Brasil. Revista Brasileira de Zoologia 20(3): 511–517. <https://doi.org/10.1590/S0101-81752003000300024>
- PEDRO, W.A., F.C. PASSOS & B.K. LIM. 2014. Morcegos (Chiroptera: Mammalia) da Estação Ecológica dos Caetetus, estado de São Paulo. Chiroptera Neotropical 7(1–2): 136–140. <http://www.chiropteraneotropical.net/index.php/cn/article/view/124>
- PEDRO, W.A. & V.A. TADDEI. 2014. Bats from southwestern Minas Gerais, Brazil (Mammalia: Chiroptera). Chiroptera Neotropical 4(1): 85–88. <http://www.chiropteraneotropical.net/index.php/cn/article/view/179>
- PERACCHI, A.L. & S.T. ALBUQUERQUE. 1993. Quirópteros do município de Linhares, Estado do Espírito Santo, Brasil (Mammalia, Chiroptera). Revista Brasileira de Biologia 53(4): 575–581. <http://rl.ufrj.br/labmasto/publicacoes/32.pdf>
- PETERS, F.B., R.P.R. DE O. ROTH, L.F. MACHADO, E. DE L. COELHO, D.M.H. JUNG & A.U. CHRISTOFF. 2010. Assembléia de mamíferos dos agroecossistemas constituintes da bacia hidrográfica do rio da Várzea, Rio Grande do Sul. Biotemas 23(4): 91–107. <https://doi.org/10.5007/2175-7925.2010v23n4p91>
- QUINTELA, F.M., C. IBARRA, S.V.D. OLIVEIRA, I.G. MEDVEDOVISKY, F. CORREA, D. GIANUCA, A. GAVA & S.M. PACHECO. 2011. Mammalia, Chiroptera, Rio Grande, state of Rio Grande do Sul, Brazil. Check List 7(4): 443–447. <https://doi.org/10.15560/7.4.443>
- REID, F. 2009. A field guide to the mammals of Central America and southeast Mexico. 2nd Edition. New York: Oxford University Press. 346 pp.
- REIS, N.R. dos., A.L. PERACCHI & M.L. SEKIAMA. 1999. Morcegos da Fazenda Monte Alegre, Telêmaco Borba, Paraná (Mammalia, Chiroptera). Revista Brasileira de Zoologia 16 (2): 501–505. <http://www.scielo.br/pdf/rbzool/v16n2/v16n2a15>
- REIS, N.R., M.L.S. BARBIERI, I.P. LIMA & A.L. PERACCHI. 2003. O que é melhor para manter a riqueza de espécies de morcegos (Mammalia, Chiroptera): um fragmento florestal grande ou vários fragmentos de pequeno tamanho. Revista Brasileira de Zoologia 20(2): 225–230. <https://doi.org/10.1590/S0101-81752003000200009>
- RODRIGUES, F.H., L. SILVEIRA, A.P. CARMIGNOTTO, A.M. BEZERRA, D.C. COELHO, H. GARBONINI, J. PAGNOZZI, & A. HASS. 2002. Composição e caracterização da fauna de mamíferos do Parque Nacional das Emas, Goiás, Brasil. Revista Brasileira de Zoologia 19(2): 589–600. <https://doi.org/10.1590/S0101-81752002000200015>
- SATO, T.M., M.C. CARVALHO-RICARDO, W. UIEDA & F.C. PASSOS. 2015. Estrutura da comunidade de morcegos (Mammalia, Chiroptera) da Estação Experimental de Itirapina, estado de São Paulo, Brasil. Papeis Avulsos Zoologia. 55(1): 1–11. <https://doi.org/10.1590/0031-1049.2015.55.01>
- SÁ-NETO, R.J. & J. MARINHO-FILHO. 2013. Bats in fragments of xeric woodland caatinga in Brazilian semiarid. Journal of Arid Environments 90: 88–94. <https://doi.org/10.1016/j.jaridenv.2012.10.007>
- SBRAGIA, I.A. & A. CARDOSO. 2008. Quiropterofauna (Mammalia: Chiroptera) cavernícola da Chapada Diamantina, Bahia, Brasil. Chiroptera Neotropical 14(1): 360–365. <https://chiroptera.unb.br/index.php/cn/article/view/107>
- SIKES, R.S. & W.L. GANNON. 2011. Animal Care and Use Committee of the American Society of Mammalogists. Guidelines of the American Society of Mammalogists for the use of wild mammals in research. Journal of Mammalogy 92(1): 235–253. <https://doi.org/10.1644/10-mamm-f-355.1>
- SILVA, J.P.A., A.R. CARVALHO & J.A. DE O. MOTTA. 2009. Fauna de morcegos (Mammalia, Chiroptera) em cavernas do bioma Cerrado na região de Indiara (Goiás). Revista Brasileira de Zootecnias 11(3): 209–217. <https://zoociencias.ufjf.emnuvens.com.br/zoociencias/article/view/1259/995>
- SIMMONS, N.B. 2005. Order Chiroptera; pp. 312–529, in: D.E. WILSON & D.M. REEDER (eds.). Mammal species of the world: a taxonomic and geographic reference. Baltimore: Johns Hopkins University Press.
- SOUZA, M.A.N., A. LANGGUTH & G.E. DO AMARAL. 2004. Mamíferos dos brejos de altitude Paraíba e Pernambuco; pp. 229–254, in: K.C. PÓRTO, J.J.P. CABRAL & M. TABARELLI (eds.). Brejos de altitude em Pernambuco e Paraíba: história natural, ecologia e conservação. Brasília: Ministério do Meio Ambiente. http://www.mma.gov.br/estruturas/chm/_arquivos/parte7_brejos.pdf
- STALLINGS, J.R., G.A. DA FONSECA, L.P.D.S. PINTO, L.M.D.S., AGUIAR, E.L. SÁBATO. 1990. Mamíferos do Parque Florestal Estadual do Rio Doce, Minas Gerais, Brasil. Revista Brasileira de Zoologia 7(4): 663–677. <https://doi.org/10.1590/S0101-81751990000400022>
- TAVARES, V.C., L.M. AGUIAR, F.A. PERINI, F.C. FALCÃO & R. GREGORÍN. 2010. Bats of the state of Minas Gerais, southeastern Brazil. Chiroptera Neotropical 16(1): 675–705. <http://chiroptera-neotropical.net/index.php/cn/article/view/257>
- TRAJANO, E. 1985. Ecologia de populações de morcegos cavernícolas em uma região cárstica do sudeste do Brasil. Revista Brasileira de Zoologia 2(5): 255–320. <https://doi.org/10.1590/S0101-8175198000100001>
- UIEDA, W., T.M. SATO, M.C. DE CARVALHO & V. BONATO. 2007. Fruits as unusual food items of the carnivorous bat *Chrotopterus auritus* (Mammalia, Phyllostomidae) from southeastern Brazil. Revista Brasileira de Zoologia 24(3): 844–847. <https://doi.org/10.1590/S0101-81752007000300035>
- UIEDA, W., I. SAZIMA & A. STORTI FILHO. 1980. Aspectos da biologia do morcego *Furipterus horrens* (Mammalia, Chiroptera, Furipteridae). Revista Brasileira de Biologia 40(1): 59–66. <http://>

- repositorio.inpa.gov.br/handle/123/5284
- VIZOTTO, L.D. & V.A. TADDEI. 1973. Chave para determinação de quirópteros brasileiros. São José do Rio Preto: Editora da UNESP. 72 pp.
- WILLIAMS, S.L. & H.H. GENOWAYS. 2008. Subfamily Phyllostominae Gray, 1825; pp. 255–300, in: A.L. GARDNER (ed.). Mammals of South America. Volume 1: marsupials, xenarthrans, shrews, and bats. Chicago and London: The University of Chicago Press, Chicago, IL.
- WITT, A.A. & M.E. FABIAN. 2010. Hábitos alimentares e uso de abrigos por *Chrotopterus auritus* (Chiroptera, Phyllostomidae). *Mastozoología Neotropical* 17(2): 353–360. <http://ref.scielo.org/nsfdyv>

Authors' contributions. GHNB contributed in collection of data and wrote manuscript; JPMA and JCVM contributed in collecting data in the field, revised and wrote part of the manuscript; PAR did cranial analysis, map and manuscript revision; MAFK revised the manuscript.

Received: 30 October 2016

Accepted: 31 March 2017

Academic editor: Annia Rodriguez-San Pedro