



LJMU Research Online

Burgess, LC, Venugopalan, L, Badger, J, Street, T, Alon, G, Jarvis, JC, Wainwright, TW, Everington, T, Taylor, P and Swain, ID

Effect of neuromuscular electrical stimulation on the recovery of people with COVID-19 admitted to the intensive care unit: A narrative review

<http://researchonline.ljmu.ac.uk/id/eprint/17042/>

Article

Citation (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

Burgess, LC, Venugopalan, L, Badger, J, Street, T, Alon, G, Jarvis, JC, Wainwright, TW, Everington, T, Taylor, P and Swain, ID (2021) Effect of neuromuscular electrical stimulation on the recovery of people with COVID-19 admitted to the intensive care unit: A narrative review. Journal of

LJMU has developed [LJMU Research Online](#) for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact researchonline@ljmu.ac.uk

<http://researchonline.ljmu.ac.uk/>



EFFECT OF NEUROMUSCULAR ELECTRICAL STIMULATION ON THE RECOVERY OF PEOPLE WITH COVID-19 ADMITTED TO THE INTENSIVE CARE UNIT: A NARRATIVE REVIEW

Louise C. BURGESS, BSc¹, Lalitha VENUGOPALAN, PhD², James BADGER, MMedSc³, Tamsyn STREET, PhD^{4,5}, Gad ALON, PhD⁶, Jonathan C. JARVIS, PhD⁷, Thomas W. WAINWRIGHT, PgDip PgCert^{1,8}, Tamara EVERINGTON, PhD⁹, Paul TAYLOR, PhD^{4,5} and Ian D. SWAIN, PhD¹

From the ¹Orthopaedic Research Institute, Bournemouth University, Bournemouth, UK, ²The Department of Biomedical Engineering, Saveetha Engineering College, Thandalam, Chennai, India, ³Department of Anaesthetics, University Hospital Southampton NHS Foundation Trust, Southampton, ⁴National Clinical FES Centre, Salisbury NHS Foundation Trust, Salisbury, ⁵Faculty of Health and Social Science, Bournemouth University, Bournemouth, UK, ⁶University of Maryland, School of Medicine, Department of Physical Therapy and Rehabilitation Science, Baltimore, Maryland, MD, USA, ⁷School of Sport and Exercise Science, Liverpool John Moores University, Liverpool, ⁸Physiotherapy Department, University Hospitals Dorset NHS Foundation Trust, Bournemouth, and ⁹Department of Haematology, Hampshire Hospitals NHS Foundation Trust, Basingstoke, UK.

The rehabilitation of patients with COVID-19 after prolonged treatment in the intensive care unit is often complex and challenging. Patients may develop a myriad of long-term multi-organ impairments, affecting the respiratory, cardiac, neurological, digestive and musculoskeletal systems. Skeletal muscle dysfunction of respiratory and limb muscles, commonly referred to as intensive care unit acquired weakness, occurs in approximately 40% of all patients admitted to intensive care. The impact on mobility and return to activities of daily living is severe. Furthermore, many patients experience ongoing symptoms of fatigue, weakness and shortness of breath, in what is being described as “long COVID”. Neuromuscular electrical stimulation is a technique in which small electrical impulses are applied to skeletal muscle to cause contractions when voluntary muscle contraction is difficult or impossible. Neuromuscular electrical stimulation can prevent muscle atrophy, improve muscle strength and function, maintain blood flow and reduce oedema. This review examines the evidence, current guidelines, and proposed benefits of using neuromuscular electrical stimulation with patients admitted to the intensive care unit. Practical recommendations for using electrical muscle stimulation in patients with COVID-19 are provided, and suggestions for further research are proposed. Evidence suggests NMES may play a role in the weaning of patients from ventilators and can be continued in the post-acute and longer-term phases of recovery. As such, NMES may be a suitable treatment modality to implement within rehabilitation pathways for COVID-19, with consideration of the practical and safety issues highlighted within this review.

Key words: critical care; rehabilitation; neuromuscular electrical stimulation; muscular atrophy; coronavirus infection; COVID-19.

Accepted Jan 29, 2021; Epub ahead of print Feb 26, 2021

J Rehabil Med 2021; 53: jrm00164

Correspondence address: Ian D. Swain, Orthopaedic Research Institute, Bournemouth University, Bournemouth, UK. E-mail: iswain@bournemouth.ac.uk

LAY ABSTRACT

Many patients with COVID-19 are admitted to the intensive care unit with ongoing symptoms of fatigue, weakness and shortness of breath. Neuromuscular electrical stimulation is a technique in which small electrical impulses are applied to skeletal muscle to cause contractions when voluntary muscle contraction is difficult or impossible. It can prevent muscle atrophy, improve muscle strength and function, maintain blood flow and reduce oedema. This review examines the evidence, current guidelines, and proposed benefits of using neuromuscular electrical stimulation with patients admitted to the intensive care unit. Practical recommendations for using electrical muscle stimulation with COVID-19 patients are provided and suggestions for further research are proposed. Evidence suggests NMES may play a role in the weaning of patients from ventilators and can be continued in the post-acute and longer-term phases of recovery. As such, NMES may be a suitable treatment modality to implement within rehabilitation pathways for COVID-19, with consideration of the practical and safety issues highlighted within this review.

The COVID-19 pandemic has seen unprecedented numbers of people being treated in intensive care units (ICUs) worldwide. Many patients have received artificial ventilation, and some have been ventilated for many weeks. Those that survive are often left with long-term disabilities as a result of the effects of both the disease and of the treatments necessary to keep them alive. A myriad of multi-organ impairments is associated with COVID-19 including respiratory, cardiac, neurological, bowel and kidney dysfunction (1). The unexpectedly large number of COVID-19 patients requiring a prolonged stay in ICU additionally increases the risk of dysfunction of both respiratory and skeletal muscle, commonly referred to as ICU-acquired weakness (ICUAW). A conspicuous feature of COVID-19 is the persistence of symptoms, which may appear to resolve, but then recur. As a result, many survivors are left needing significant rehabilitation at a time when such services are under great

stress. This has led to the blanket term “long COVID”, which describes ongoing symptoms, which may include fatigue, weakness and delayed recovery (2).

Strikingly, in the first 7 months of 2020, there were more than 10,000 COVID-19 admissions to critical care in the UK National Health Service (NHS), which is 4 times greater than historic annual cases of viral pneumonia (3). Our experience of COVID-19 in the UK is that critically unwell patients generally require a longer course of respiratory support, exacerbating other risk factors for ICUAW (Table I) (3). At present, ICUAW is seen in approximately 20–50% of patients with COVID-19 admitted to the ICU (4). General deconditioning, muscle atrophy, inflammation, and functional disability often necessitate transfer from the ICU to a long-term care facility. Exacerbations of chronic comorbidities and the cycle of prolonged bed rest, ongoing inflammation and malnutrition can lead to continued functional disability, immobility and continued ventilation support. Data from the UK Intensive Care National Audit and Research Centre (ICNARC) database indicates that older age, obesity, multiple deprivation, and the requirement for assistance in activities of daily living (ADL) are predictors for severe disease requiring admission to critical care (3). These risk factors are associated with a reduced level of background fitness, malnutrition and neuropathy. Infection with COVID-19 characteristically causes myalgia, lethargy and a loss of appetite, which are likely to exacerbate this pre-morbid condition. Further deconditioning may result from constrained normal daily activities. This may be due to the disease itself, causing shortness of breath on exertion or delirium (5), or may be the result of supportive interventions and infection control measures. It is also noteworthy that proximal myopathy is associated with the use of therapeutic dexamethasone, a drug that has been shown to reduce 28-day mortality in COVID-19 (6).

After leaving hospital, almost 90% of survivors experience ongoing symptoms for more than 2 months, such as fatigue and shortness of breath, which are likely to limit rehabilitation and potentiate deconditioning (7). ICUAW is associated with worse outcomes,

including a nearly 2-fold increase in 1-year mortality, and decreased quality of life (QoL) (8, 9). A major challenge within current practice is how to ameliorate profound physical and functional deficits in COVID-19 survivors at a time when traditional services are stretched. Innovations that reduce the duration and improve the outcome of rehabilitation will alleviate the burden of suffering and economic damage caused by COVID-19.

Neuromuscular electrical stimulation

Neuromuscular electrical stimulation (NMES) is the application of small electrical impulses to nerves supplying muscles, using electrodes applied to the skin. NMES has long been used as a treatment for muscle weakness (10). NMES can be used to induce a muscle contraction when it is difficult or impossible for the person to achieve this voluntarily, thereby allowing effective exercise and the strengthening of muscles. NMES has been proposed as an intervention to address immobilization and ICUAW in patients with severe COVID-19 (11), however details on when and how to utilize NMES are lacking. As post-acute rehabilitation services respond to the increasing demand on services, recommendations are required to guide the delivery of rehabilitation models.

Aim

This narrative review critically examines the evidence for using NMES in the ICU and offers suggestions for clinical practice among patients with COVID-19. This article provides practical recommendations using a continuum of care model for clinicians interested in using electrical stimulation for patients during and after prolonged ICU treatment.

METHODS

This narrative review was informed by the findings of a web-based literature search, completed in October 2020. The search aimed to identify studies that have investigated the role of electrical stimulation in the recovery of patients admitted to the ICU, published in the last 10 years (January 2010–October 2020). A search strategy (Table S1¹) was developed to capture randomized controlled trials (RCTs) or non-randomized clinical trials that have evaluated an intervention of electrical stimulation (functional electrical stimulation (FES) or NMES) in patients admitted to the ICU. Specifically, we sought studies of adults (aged over 18 years), admitted to the ICU due to chronic illness or following non-elective surgery, who received an intervention of electrical stimulation, (i) during their stay in the ICU, (ii) during the acute recovery phase in hospital, or (iii) following discharge from hospital. The databases searched included: PubMed, EMBASE, Medline, CINAHL Complete, and The Cochrane Library. Articles were systematically reviewed by the research team to ensure they met the eligibility criteria

Table I. Risk factors for deconditioning and intensive care unit associated weakness (ICUAW) in patients with COVID-19 in comparison with those with viral pneumonia (3)

Risk factor for deconditioning/ICUAW	COVID-19 (n = 10,557)	Viral pneumonia, 2017 to 2019 (n = 5,782)
Duration of advanced respiratory support, median days (IQR)	13 (7–23)	9 (4–17)
Multi-organ failure, %	40.8	26.3
Age, mean (SD)	58.8 (12.7)	58 (17.4)
Very severe comorbidities, %	13.6	24
Dependency prior to hospital admission, %	10.3	26.4

ICU: intensive care unit; IQR: interquartile range; SD: standard deviation.

(Table SII¹) and were subsequently used to inform this critical analysis and recommendations for future practice. Studies were only included if they reported a replicable NMES protocol. In addition, recently published guidelines recommending the use of home-based NMES for chronic respiratory conditions, such as chronic obstructive pulmonary disease (COPD) from the National Institute of Clinical Excellence (NICE) were used to inform recommendations (12). A narrative review was considered the most appropriate methodology so that the research team could use a broad survey of the literature, in combination with expert opinion, to inform clinical recommendations. The research team is a multinational, multidisciplinary group of experts with many years clinical experience of NMES. The group includes biomedical engineers, physiotherapists, intensive care clinicians, physiologists, and haematologists.

RECOMMENDATIONS

Physiological considerations

Fundamental to treatment with NMES is an understanding of the electrophysiological mechanisms associated with skeletal muscle function. Skeletal muscles, including diaphragm and accessory respiratory muscles, are made up of long, multinucleate, approximately cylindrical cells containing sarcomeres, in which the contractile proteins actin and myosin interact to generate force and shortening. Skeletal muscle powers voluntary movement, including speech and breathing, buffers circulating glucose, and is surprisingly labile. Disuse during bedrest causes loss of muscle mass by active cellular mechanisms. This presents a severe problem in ventilated patients. The domed diaphragm muscle normally flattens by shortening to generate a lower than atmospheric pressure in the pleural space, so the lungs inflate. During mechanical ventilation the diaphragm muscle quickly loses mass, so that after ventilatory assistance, diaphragm function is reduced (13). The extreme reduction in activity from contraction during every breath, to zero, may explain why the diaphragm loses mass more quickly than, say, the pectoral muscles. In healthy persons, growth of muscle is often considered to be slower than the loss of muscle with disuse; to gain 1 kg of leg muscle might take 12 weeks of resistance training, whereas 1 kg of mass is lost in 1 week with complete disuse (14). The magnitude of the difference in activity before and after is very different in these scenarios; hence, the prevention of atrophy using early activity-based methods may reduce the human and financial cost of rehabilitation after critical illness.

To activate muscle contractions from outside the body, action potentials must be generated in the muscle membrane. Stimulation is usually applied where the

nerve that contains the target motor neurones is most accessible. Muscles respond to single action potentials with a brief period of activation then relaxation. The force response to a single stimulus is a very brief twitch with a low force. To produce stronger contractions, successive activations must be applied before the relaxation of the prior stimulus, and so frequencies in humans of 20–50 impulses per second are used (20–50 Hz). Muscles require a continuous supply of oxygen and glucose to generate sustained work, and therefore contractions must be intermittent, because blood flow is excluded during strong contractions. The activity/rest cycle and the number of contractions in a session provides a huge number of possible combinations. Exercise is often prescribed in terms of a number of sets of repetitions (single contractions), with a rest period between sets. As a result, unless otherwise stated, cyclic electrical stimulation was used in the articles considered in this review, rather than any other NMES (for example, electromyography (EMG)-triggered stimulation).

Neuromuscular electrical stimulation and intensive care unit acquired weakness

The application of NMES to treat ICUAW is well documented within the evidence-base (15–18). The primary objective of interventions has been to induce intermittent muscle contractions with electrical stimulation to minimize the loss of muscle mass and excitability, to strengthen these muscles and to enhance the recovery of mobility during and after discharge from the ICU (19). The findings from pre-clinical work on underlying electrophysiological mechanisms from healthy participants and data from critical care patients suggest that, to prevent ICUAW, an NMES programme should begin in the ICU as soon as medically feasible. This is particularly relevant to people with COVID-19, as early intervention is advised due to the often-prolonged



Fig. 1. Electrode positioning for electrical stimulation of the quadriceps (posed with a mannequin).

¹<http://www.medicaljournals.se/jrm/content/?doi=10.2340/16501977-2805>

stay and risk of subsequent long-term ICUAW. Reducing initial muscle atrophy is preferable to extending rehabilitation, due to the extended amount of time it takes to recover pre-ICU muscle strength (20). Those with risk factors for ICUAW should be prioritized because there is a small amount of evidence that NMES can reduce the prevalence of ICUAW (21).

Many studies have activated the quadriceps (Fig. 1) along with another muscle group such as the hamstrings, whereas others have targeted the abdominal musculature. Stimulation parameters commonly used are a frequency between 30–50Hz, pulse duration of 250–400 μ s and an intensity adjusted up to maximum sensory tolerance, so that contractions are easily visible and palpable. Most studies have included one 60-min session or 2 30-min sessions per day. There has been enough commonality to conduct systematic reviews and a meta-analysis using the Medical Research Council's (MRC) score for muscle strength as an outcome measure. Liu et al. (15) found a significant improvement in muscle strength for NMES over control (mean difference (MD)=1.78, 95% CI 0.44, 3.12 ($p=0.009$)). All studies included in the review used the MRC scale to evaluate the strength of the surrounding muscles, with a score of <48 to diagnose ICUAW (22–26). The results of several previous systematic reviews in this area are largely consistent with these findings (16–18).

The current most common protocols used in the ICU suggest that NMES at this stage for a limited amount of time might be sufficient to maintain muscle volume, but does not increase it. In one of the larger studies, Dall'Acqua et al. (27) did not find a significant improvement in abdominal muscle thickness with NMES, but, interestingly, found a significant decline in the control. Further support for this hypothesis is suggested in a recent study from Nakamura et al, (28) who examined the effects of a 20-min daily dose of NMES (171 contractions per day) on femoral muscle volume. Researchers found a significant decrease in muscle volume for both the control and intervention group; however, the mean rate of muscle volume reduction was significantly less for the NMES group (NMES (standard deviation; SD)=10.4% (SD 10.1%), control=17.7% (SD 10.8%) ($p=0.04$)). The data from these studies and longer-term treatment, for example, up to 9 weeks (29) suggest that NMES can be used in the ICU to slow down muscle wasting, but it is necessary for participants to then use home-based NMES to maintain and strengthen muscles post-ICU. Interestingly, recent research by Nakashini et al. (30) suggests that identifying the motor-point to elicit the strongest contraction, as well as increasing the number of contractions in a session, may maintain muscle strength

more effectively. Researchers included a 30-min daily session (180 contractions) for 5 days to the NMES group, while the control had usual care. A significant difference in muscle volume and strength was found, but no difference in ICUAW was found. This suggests that further research should be conducted into optimal dosing for ICU patients and is supportive of a period of post-ICU NMES treatment for maintenance and recovery of strength.

Neuromuscular electrical stimulation in chronic obstructive pulmonary disease

As COVID-19 is a chronic respiratory condition, patients may share some similarities with patients with COPD in terms of symptoms and complications (e.g. shortness of breath, respiratory infection, heart problems and peripheral muscle weakness), and thus it is beneficial to review the evidence for NMES within COPD patient groups. Recently published NICE guidelines for the use of NMES to strengthen muscles in patients with chronic respiratory disease recommend that, for those who are unable to exercise, evidence supports the use of electrical muscle stimulation. However, standard arrangements must be in place for clinical governance, consent and audit (12). A meta-analysis of 9 studies, including 276 patients with moderate-to-severe COPD, found improvement in quadriceps muscle strength (standardized mean difference (SMD)=1.12, 95% CI 0.64–1.59 ($p<0.001$); 6 studies of 207 patients) with NMES (31). In a recent Cochrane review, improvements were found for peripheral muscle endurance (SMD=1.36, 95% CI 0.59–2.12, ($p<0.001$); 2 studies of 35 patients) and these improvements translated into improved 6-min walking distance (MD=39.26 m, 95% CI 16.31–62.22, ($p<0.001$); 2 studies of 72 patients) (32). An improvement in exercise endurance was also found (MD=3.62 min, 95% CI 2.33–4.91, ($p<0.001$); 3 studies of 55 patients) and days to first transfer out of bed was decreased for the NMES group (MD=-4.98 days, 95% CI -8.55 to -1.41, ($p=0.006$); 2 studies of 44 patients) (32). However, NMES was not associated with improvements in health-related quality of life (HRQoL) (32), and thus the actual value of NMES for improved QoL remains uncertain (31).

NMES stimulation parameters for COPD vary considerably among studies, with stimulation frequency set to a median value of 50 Hz (range 15–75 Hz), pulse duration 400 μ s (200–700), target duty cycle 33% (13–75), session length 30 min (18–240), session frequency 5 times (2–7) each week, and programme duration 6 weeks (4–11) (31). All studies set stimulation amplitude to elicit a visible muscle contraction within the participant's tolerance and most found that

the amplitude could be increased over the course of the programme. However, the high variability in length of time, parameters and different type of outcome measures used in the studies made comparisons difficult.

Neuromuscular electrical stimulation to wean critically ill patients off ventilators

Neuromuscular electrical stimulation may be considered to help wean critically ill patients off ventilators, and is advantageous when the patient cannot participate in voluntary exercise. Preliminary work supporting the added value of an NMES programme to wean patients from dependence on ventilators is supportive of further research in this area. McCaughey et al. (33) provided the most credible, albeit preliminary data, that earlier weaning is possible. They applied NMES over the posterior-lateral abdominal wall to activate the transversus abdominis and internal and external oblique muscles during exhalation, automatically synchronized with the participant's breathing pattern. Stimulation was applied for 30 min, twice per day, 5 days per week, until discharge from the ICU. The study compared an active group receiving stimulation that caused a strong visible muscle contraction (30 Hz frequency and a pulse-width of 350 μ s) with a control group that received sensory level stimulation (10 Hz frequency and 350 μ s pulse-width, but with an amplitude sufficient to be felt on the skin, but not to cause muscle contraction). A survival analysis found ICU length of stay (median 11 vs not estimable days, ($p=0.011$)) and ventilation duration (median 6.5 vs 34 days, ($p=0.039$)) were shorter in the intervention compared with the control group. Dall'Acqua and colleagues (27) stimulated the pectoral and rectus abdominis muscles bilaterally for 30 min daily, using 300 μ s phase duration, 50 Hz pulse rate to induce a 3-s contraction followed by 10 s of relaxation and compared it with a sensory threshold stimulation group. Time to weaning off the ventilator was not recorded, but the length of ICU stay was shorter in the NMES group (mean: 10 \pm 4 days) compared with the control group (mean: 16 \pm 9) ($p=0.045$). Other investigators used NMES to activate the deltoid and quadriceps muscles bilaterally, applied concurrently with active exercises or without exercises or exercise only, and found no difference between groups in terms of time to discharge from the ICU (34). None of these 3 groups of investigators reported any adverse response or interference with the recovery of and discharge from the ICU.

Neuromuscular electrical stimulation and prevention of venous thromboembolism

Venous thromboembolism (VTE), encompassing pulmonary embolism (PE) and deep vein throm-

bolism (DVT), is a common and severe complication of critical illness (35, 36). Many critically ill patients have multiple risk factors that predate ICU admission; including, recent surgery or trauma, sepsis, malignancy, immobilization, increased age, and cardiac or respiratory failure (37). Once admitted, patients who need treatment on the ICU are exposed to additional VTE risk factors, including prolonged immobilization, pharmacological paralysis, central venous catheterization, haemodialysis and treatment with vasopressors (37–39). In 4 recent meta-analyses of hospitalized COVID-19 patients, incidence of thrombotic complications was reported as between 22.7% and 31%, and risk persisted even in those receiving anticoagulation (40–43).

Prophylaxis aims to combat the 3 predisposing factors to VTE; venous stasis, hypercoagulability, and endothelial injury (44). Traditional prevention strategies include pharmacological agents, such as unfractionated heparin, low molecular weight heparin (LMWH), direct oral anticoagulants, and mechanical devices, such as graduated compression stockings or intermittent pneumatic compression of the limbs (45). Interim guidance for COVID-19 recommends treatment with LMWH administered at prophylaxis doses pending the emergence of additional data and guidance (46). Despite receiving anticoagulation for thromboprophylaxis, a high rate of VTE has been observed among patients with COVID-19 admitted to the ICU (43).

NMES has been approved by NICE as an alternative prophylaxis when other mechanical and pharmacological methods of prophylaxis are impractical or contraindicated (47, 48). The transcutaneous application of electrical impulses stimulates the common peroneal nerve to generate dorsiflexion in the lower limb, which, in turn, activates the calf muscle pump, emulating the normal physiological response achieved by walking, without the patient having to mobilize. Electrode positioning is demonstrated in Fig. 2. NMES has been shown to



Fig. 2. Electrode position for electrical stimulation of the peroneal nerve for increased blood flow to the lower limb (posed with a mannequin).

ICU: Mechanically ventilated >48 hours	ICU: Weaned off ventilator	ICU: Recovery	Acute rehabilitation facility	At home: post-discharge
<ul style="list-style-type: none"> • Immobilised • Risk of muscle atrophy • Risk of DVT • Inability to respond to verbal commands • Unable to engage in rehab 	<ul style="list-style-type: none"> • Hypoactive delirium • Unable to engage in rehab • Disorientation • Risk of muscle atrophy • Risk of DVT 	<ul style="list-style-type: none"> • Alert and cooperative • Limited mobility • ICUAW • Risk of DVT • Respiratory distress 	<ul style="list-style-type: none"> • ICUAW • Functionally limited • Persistent fatigue • Respiratory impairment • Neurological symptoms 	<ul style="list-style-type: none"> • ICUAW • Fatigue on exertion • Impaired lung function • Dyspnoea • Dizziness and headaches

Fig. 3. Characteristics of a patient admitted to the intensive care unit (ICU) with COVID-19. ICUAW: intensive care unit acquired weakness; DVT: deep vein thrombosis.

be effective in reducing fibrinogen, D-dimer and tissue plasminogen activator (tPA) levels, and increasing venous, arterial and microcirculatory flow, thus preventing venous stasis and oedema (49–58). Moreover, clinical evidence has shown effectiveness of NMES for reducing the incidence of DVT in hospitalized patients (59–66).

In line with recommendations from NICE, NMES should be considered as an alternative or adjunct prophylaxis in patients with COVID-19 where other mechanical and pharmacological prophylaxis are impractical (47). It may be most effective when used prior to the formation of oedema, to prevent venous stasis and reduce risk of VTE. Devices should be used in accordance with guidance (47) and individual instructions for use of specific devices. If NMES is used for other treatment aims (such as muscle strengthening), it should be acknowledged that a circulatory effect will be delivered simultaneously, and so competing treatment aims may be balanced by preferentially aiming NMES settings for muscle strengthening parameters. Furthermore, NMES may provide the most benefit to patients who are immobilized or positioned where the leg is lower than the body.

Neuromuscular electrical stimulation and the continuum of care model

NMES may be advantageous in COVID-19, as it can be used throughout the patient’s recovery to address a number of physiological and clinical deficits (Fig. 3) in a continuum of care model. Example applications of NMES for patients admitted to the ICU with COVID-19 are illustrated in Fig. 4.

While minimizing the amount of stimulation is pragmatic on the ICU unit, following discharge, similarly to any exercise programme, NMES can be increased progressively subject to patient tolerance and measurable benefits. As the patients begin to mobilize out of bed, a structured mobility programme has been recommended (67). Adding the NMES to a structured physical exercise programme appears advisable compared with applying the NMES in isolation (68). From a practical perspective, as long as the patient is non-responsive to verbal commands, the NMES can be combined with passive range of motion (PROM)

exercises. Once responsive, the patient should be encouraged to add volitional contraction and active range of motion (AROM) combined with the NMES. In studies with neurological patients, volitional contraction has been found to be more effective at inducing useful therapeutic improvements (69).

Following discharge from hospital, ongoing use of NMES may also be considered, to address persistent symptoms and functional limitations. NMES can be applied independently in the home environment and is considered an attractive adjunct to enhance the hypertrophic effect of traditional exercise (10). Likewise, following discharge it may be appropriate to consider the ongoing use of NMES to increase blood flow and prevent oedema or DVT.

Nonetheless, one of the main shortcomings of current research on NMES in the ICU is the lack of long-term follow-up, because most studies only use NMES for the duration of hospital stay (5–14 days). This may be reflective of the lack of long-term rehabilitation and follow-up for these patients once they leave the ICU and hospital. Further research, including long term follow-up, should be conducted, as currently it is unknown whether patients who appear to benefit during their stay in ICU continue to benefit after a relatively short period of treatment. Further research should also examine the potential benefits of home-based NMES post-ICU as part of a continuum of care. Using NMES for a period of 9 weeks, as in previous investigations (29), or for a minimum of 6 weeks, as in many of the COPD studies, may lead to sustained longer-term benefits (31).

PRACTICAL CONSIDERATIONS

Early rehabilitation has generally been accepted as a safe and effective intervention in critical care (70–74). However, there are several practical issues that make the implementation of these interventions challenging, especially in those with COVID-19. Such issues include deep sedation for facilitating mechanical ventilation; delirium; prone positioning; access to appropriate number or type of personnel; physiological stability; and obesity. An observation study in France demonstrated




Fig. 4. Examples of neuromuscular electrical stimulation (NMES) application for patients admitted to the intensive care unit (ICU) with COVID-19, by indication. ICUAW: intensive care unit acquired weakness; DVT: deep vein thrombosis; FES: functional electrical stimulation; VTE: Venous thromboembolism.

that 65% of those with COVID-19 admitted to critical care experienced delirium, and therefore a significant number of patients presumably would have been unable to safely/successfully participate in active physiotherapy regimes while affected (75). Furthermore, even passive interventions, such as in-bed cycle ergometry, are restricted to those in the supine position, rendering them unsuitable for patients with COVID-19, for whom prone positioning for more than 12 h per day is a widely accepted strategy for improving oxygenation (76). In addition, in-bed cycling is purely passive and, although it will help maintain range of movement, it will not increase muscle bulk or strength. Another consideration is weight restrictions on rehabilitation equipment, which may preclude the 7.9% of morbidly obese patients (3) admitted to critical care with COVID-19 from receiving a number of interventions. Finally, accepting that COVID-19 has resulted in an increase in intensive care admissions and physiotherapy demand, more efficient rehabilitation interventions and use of staff is required.

SAFETY CONSIDERATIONS

Common equipment in an ICU includes mechanical ventilators to assist breathing through an endotracheal tube or a tracheostomy tube; monitors of cardiac functions; equipment for the constant monitoring of bodily functions; a web of intravenous lines, feeding tubes, nasogastric tubes, suction pumps, drains, and catheters; syringe pumps; and a wide array of drugs to treat the primary condition(s) of hospitalization. Accordingly, the clinical team must verify the compatibility of the stimulation system to ensure there is no interference with the electronic systems, such as electrocardiography (ECG) and electroencephalography (EEG) monitors, pacemakers, defibrillators, or other implanted stimulators. Iwatsu et al. (77) provided evidence assuring the safety of stimulation in the ICU. Furthermore, none of the other published clinical trials that used non-invasive electrical stimulation in the ICU reported interference with the ICU equipment (27, 28, 30, 33, 34, 78). Interference with pacemakers and implantable cardioverter defibrillators

appears to depend on the proximity of the electrodes to the implanted device; lower limb stimulation, in particular, appears safe in this group, but clinicians must be aware of, and monitor for, such an interaction (79), especially if stimulation of respiratory muscles is indicated. In addition, the stimulation system must meet all hygiene, disinfection and sterilization standards required by the hospital. When applying the electrical stimulation, clinicians must not apply the electrode over open wounds and should avoid any contact of the electrodes with external fixation hardware. In contrast, applying NMES over internal hardware appears safe (80, 81). Electrical stimulation is known to increase muscle perfusion and oxygen consumption in a similar way to light intensity exercise. Given that changes are small and reversible, it is likely to be safe in those receiving cardiovascular support, and studies in this cohort have not reported any adverse effects (78). Finally, when applying electrical stimulation to those with reduced consciousness, special care must be taken regarding skin integrity, as the patient will not be able to report pain.

SUMMARY

Innovations that save time and improve the outcome of rehabilitation will alleviate the burden of suffering and economic damage caused by COVID-19. Current evidence suggests that NMES can reduce the rate of muscle atrophy for patients admitted to the ICU. Whilst the evidence for increasing muscle mass is less clear, reduction in atrophy is a worthwhile goal in the pursuit of expedited recovery and return to independence. For the immobilized patient, NMES increases blood flow, reduces oedema, and can be used as an alternative prophylaxis in cases where traditional methods are contraindicated. Evidence suggests NMES may play a role in the weaning of patients from ventilators and should be continued in the post-acute and longer-term phases of recovery. As such, NMES may be a suitable treatment modality to implement within rehabilitation pathways for COVID-19, with consideration of the practical and safety issues highlighted within this review.

Future research endeavours should aim to evaluate the specific application of NMES to patients with COVID-19, the longer-term effects of NMES, and the most effective parameters to influence underlying electrophysiological mechanisms.

ACKNOWLEDGEMENTS

The authors would like to thank Salisbury NHS Foundation Trust Simulation Suite for supporting the images used in Figs 1 and 2.

The authors have no conflicts of interest to declare.

REFERENCES

1. Zaim S, Chong JH, Sankaranarayanan V, Harky A. COVID-19 and multiorgan response. *Curr Probl Cardiol* 2020; 45: 100618.
2. Mahase E. Covid-19: What do we know about "long covid"? *BMJ* 2020; 370: m2815.
3. Intensive Care National Audit and Research Centre. IC-NARC report on COVID-19 in critical care. 2020 Jul 17. Available from: <https://www.icnarc.org/Our-Audit/Audits/Cmp/Reports>.
4. The Faculty of Intensive Care Medicine. Clinical guide for the management of critical care for adults with COVID-19 during the coronavirus pandemic; 2020 Sep 11. Available from: <https://static1.squarespace.com/static/5e6613a1dc75b87df82b78e1/t/5f999cd5b3df86542e85d0ab/1603902680560/AdultCriticalCare-COVID-19-October2020.pdf>.
5. World Health Organisation (2020). Clinical Management of COVID-19: interim guidance, 2020 May 27. Available from: <https://apps.who.int/iris/handle/10665/332196>.
6. The RECOVERY Collaborative Group, Horby P, Lim WS, Emberson JR, Mafham M, Bell JL, et al. Dexamethasone in Hospitalized patients with covid-19 – preliminary report. *N Engl J Med* 2020; 10.1056/NEJMoa2021436.
7. Carfi A, Bernabei R, Landi F, Gemelli Against C-P-ACSG. Persistent symptoms in patients after acute COVID-19. *JAMA* 2020; 10.1001/jama.2020.12603.
8. Hermans G, Van Mechelen H, Clerckx B, Vanhullebusch T, Mesotten D, Wilmer A, et al. Acute outcomes and 1-year mortality of intensive care unit-acquired weakness. A cohort study and propensity-matched analysis. *Am J Respir Crit Care Med* 2014; 190: 410–420.
9. Fan E, Dowdy DW, Colantuoni E, Mendez-Tellez PA, Sevransky JE, Shanholtz C, et al. Physical complications in acute lung injury survivors: a two-year longitudinal prospective study. *Crit Care Med* 2014; 42: 849–859.
10. Nussbaum EL, Houghton P, Anthony J, Rennie S, Shay B, Hoens AM. Neuromuscular electrical stimulation for treatment of muscle impairment: critical review and recommendations for clinical practice. *Physiother Can* 2017; 69: 1–76.
11. Nakamura K, Nakano H, Naraba H, Mochizuki M, Hashimoto H. Early rehabilitation with dedicated use of belt-type electrical muscle stimulation for severe COVID-19 patients. *Crit Care* 2020; 24: 342.
12. National Institute of Health and Care Excellence. Electrical stimulation to improve muscle strength in chronic respiratory conditions, chronic heart failure and chronic kidney disease. *Interventional procedures guidance [IPG677]*. 2020. Available from: <https://www.nice.org.uk/guidance/ipg677>.
13. Schepens T, Verbrugge W, Dams K, Corthouts B, Parizel PM, Jorens PG. The course of diaphragm atrophy in ventilated patients assessed with ultrasound: a longitudinal cohort study. *Critical Care (London, England)* 2015; 19: 422.
14. Galvan E, Arentson-Lantz E, Lamon S, Paddon-Jones D. Protecting skeletal muscle with protein and amino acid during periods of disuse. *Nutrients* 2016; 8: 404.
15. Liu M, Luo J, Zhou J, Zhu X. Intervention effect of neuromuscular electrical stimulation on ICU acquired weakness: a meta-analysis. *Int J Nurs Sci* 2020; 7: 228–237.
16. Maffioletti NA, Roig M, Karatzanos E, Nanas S. Neuromuscular electrical stimulation for preventing skeletal-muscle weakness and wasting in critically ill patients: a systematic review. *BMC Med* 2013; 11: 137.
17. Wageck B, Nunes GS, Silva FL, Damasceno MC, de Noronha M. Application and effects of neuromuscular electrical stimulation in critically ill patients: systematic review. *Med Intensiva* 2014; 38: 444–454.
18. Burke D, Gorman E, Stokes E, Lennon O. An evaluation of neuromuscular electrical stimulation in critical care

using the ICF framework: a systematic review and meta-analysis. *Clin Respir J* 2016; 10: 407–420.

19. Piva S, Fagoni N, Latronico N. Intensive care unit-acquired weakness: unanswered questions and targets for future research. *F1000Res* 2019; 8: F1000 Faculty Rev-508
20. Parry SM, Puthuchery ZA. The impact of extended bed rest on the musculoskeletal system in the critical care environment. *Extrem Physiol Med* 2015; 4: 16.
21. Routsis C, Gerovasili V, Vasileiadis I, Karatzanos E, Pitsolis T, Tripodaki, E, Markaki V, et al. Electrical muscle stimulation prevents critical illness polyneuropathy: a randomized parallel intervention trial. *Critical Care (London, England)* 2010; 14: R74.
22. Kho ME, Truong AD, Zanni JM, Ciesla ND, Brower RG, Palmer JB, Needham D M. Neuromuscular electrical stimulation in mechanically ventilated patients: a randomized, sham-controlled pilot trial with blinded outcome assessment. *J Crit Care* 2015; 30: 32–39.
23. Koutsioumpa E, Makris D, Theochari A, Bagka D, Stathakis S, Manoulakas E, Sgantzios, M et al. Effect of transcutaneous electrical neuromuscular stimulation on myopathy in intensive care patients. *Am J Crit Care* 2018; 27: 495–503.
24. Leite MA, Osaku EF, Albert J, Costa CRLM, Garcia AM, Czapiessvski FDN et al. Effects of Neuromuscular Electrical Stimulation of the Quadriceps and Diaphragm in Critically Ill Patients: A Pilot Study. *Crit Care Res Pract.* 2018 Jul 8;2018:4298583. doi: 10.1155/2018/4298583.
25. Patsaki I, Gerovasili V, Sidiras G, Karatzanos E, Mitsiou G, Papadopoulos E, Christakou A, Routsis C, Kotanidou A, Nanas S. Effect of neuromuscular stimulation and individualized rehabilitation on muscle strength in Intensive Care Unit survivors: A randomized trial. *J Crit Care* 2017; 40: 76–82.
26. Chen YH, Hsiao HF, Li LF, Chen NH, Huang CC. Effects of Electrical Muscle Stimulation in Subjects Undergoing Prolonged Mechanical Ventilation. *Respir Care* 2019; 64: 262–271.
27. Dall'Acqua AM, Sachetti A, Santos LJ, Lemos FA, Bianchi T, Naue WS, Dias AS, Sbruzzi G, Vieira SR; MoVe- ICU Group. Use of neuromuscular electrical stimulation to preserve the thickness of abdominal and chest muscles of critically ill patients: a randomized clinical trial. *J Rehab Med* 2017; 49: 40–48.
28. Nakamura K, Kihata A, Naraba H, Kanda N, Takahashi Y, Sonoo T, Hashimoto H, Morimura N. Efficacy of belt electrode skeletal muscle electrical stimulation on reducing the rate of muscle volume loss in critically ill patients: a randomized controlled trial. *J Rehab Med* 2019; 51: 705–711.
29. Zampieri S, Mosole S, Lofler S, Fruhmman H, Burggraf S, Cvecka J, et al. Physical exercise in aging: nine weeks of leg press or electrical stimulation training in 70 years old sedentary elderly people. *Eur J Transl Myol* 2015; 25: 237–242.
30. Nakanishi N, Oto J, Tsutsumi R, Yamamoto T, Ueno Y, Nakataki E, et al. Effect of electrical muscle stimulation on upper and lower limb muscles in critically ill patients: a two-center randomized controlled trial. *Crit Care Med* 2020; 10.1097/CCM.0000000000004522.
31. Chen RC, Li XY, Guan LL, Guo BP, Wu WL, Zhou ZQ, Huo YT, Chen X, Zhou LQ. Effectiveness of neuromuscular electrical stimulation for the rehabilitation of moderate-to-severe COPD: a meta-analysis. *Int J Chronic Obstr Pulm Dis* 2016; 11: 2965–2975.
32. Hill K, Cavalheri V, Mathur S, Roig M, Janaudis-Ferreira T, Robles P, Dolmage TE, Goldstein R. Neuromuscular electrostimulation for adults with chronic obstructive pulmonary disease. *Cochrane Database Syst Rev* 2018; 5: CD010821–CD010821.
33. McCaughey EJ, Jonkman AH, Boswell-Ruys CL, McBain RA, Bye EA, Hudson AL, Collins DW, Heunks LMA, McLachlan AJ, Gandevia SC, Butler JE. Abdominal functional electrical stimulation to assist ventilator weaning in critical illness: a double-blinded, randomised, sham-controlled pilot study. *Critical Care (London, England)* 2019; 23: 261.
34. Akar O, Günay E, Sarinc Ulasli S, Ulasli AM, Kacar E, Sariaydin M, Solak Ö, Celik S, Ünlü M. Efficacy of neuromuscular electrical stimulation in patients with COPD followed in intensive care unit. *Clin Respir J* 2017; 11: 743–750.
35. Attia J, Ray JG, Cook DJ, Douketis J, Ginsberg JS, Geerts WH. Deep vein thrombosis and its prevention in critically ill adults. *Arch Intern Med* 2001; 161: 1268–1279.
36. Cook D, Crowther M, Meade M, Rabbat C, Griffith L, Schiff D, et al. Deep venous thrombosis in medical-surgical critically ill patients: prevalence, incidence, and risk factors. *Crit Care Med* 2005; 33: 1565–1571.
37. Cook D, Attia J, Weaver B, McDonald E, Meade M, Crowther M. Venous thromboembolic disease: an observational study in medical-surgical intensive care unit patients. *J Crit Care* 2000; 15: 127–132.
38. Minet C, Potton L, Bonadona A, Hamidfar-Roy R, Somohano CA, Lugosi M, et al. Venous thromboembolism in the ICU: main characteristics, diagnosis and thromboprophylaxis. *Crit Care* 2015; 19: 287.
39. Anderson FA, Jr, Spencer FA. Risk factors for venous thromboembolism. *Circulation* 2003; 107: 19–16.
40. Chi G, Lee JJ, Jamil A, Gunnam V, Najafi H, Memar Montazerin S, et al. Venous thromboembolism among hospitalized patients with COVID-19 undergoing thromboprophylaxis: a systematic review and meta-analysis. *J Clin Med* 2020; 9.
41. Nopp S, Moik F, Jilma B, Pabinger I, Ay C. Risk of venous thromboembolism in patients with COVID-19: a systematic review and meta-analysis. *Res Pract Thromb Haemost* 2020; 10.1002/rth2.12439.
42. Porfidia A, Valeriani E, Pola R, Porreca E, Rutjes AWS, Di Nisio M. Venous thromboembolism in patients with COVID-19: systematic review and meta-analysis. *Thromb Res* 2020; 196: 67–74.
43. Hasan SS, Radford S, Kow CS, Zaidi STR. Venous thromboembolism in critically ill COVID-19 patients receiving prophylactic or therapeutic anticoagulation: a systematic review and meta-analysis. *J Thromb Thrombolysis* 2020; 50: 814–821.
44. Kushner A, West WP, Pillarisetty LS. Virchow triad. StatPearls. Treasure Island, Florida, 2020. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK539697/#:~:text=The%20three%20factors%20of%20Virchow's,presence%20of%20a%20hypercoagulable%20state>.
45. Kakkos SK, Warwick D, Nicolaidis AN, Stansby GP, Tsolakis IA. Combined (mechanical and pharmacological) modalities for the prevention of venous thromboembolism in joint replacement surgery. *J Bone Joint Surg Br* 2012; 94: 729–734.
46. Thachil J, Tang N, Gando S, Falanga A, Cattaneo M, Levi M, et al. ISTH interim guidance on recognition and management of coagulopathy in COVID-19. *J Thromb Haemost* 2020; 18: 1023–1026.
47. National Institute of Health and Care Excellence. The geko device for reducing the risk of venous thromboembolism. 2014 [cited 2020 Jan 27 Jan]; Available from: <https://www.nice.org.uk/guidance/mtg19/chapter/1-Recommendations>.
48. Summers JA, Clinch J, Radhakrishnan M, Healy A, McMillan V, Morris E, Rua T, Ofuya M, Wang Y, Dimmock PW, Lewis C, Peacock JL, Keevil SF. The geko™ electro-stimulation device for venous thromboembolism prophylaxis: a NICE medical technology guidance. *Appl Health Econ Health Policy* 2015; 13: 135–147.
49. Barnes R, Madden LA, Chetter IC. Fibrinolytic effects of peroneal nerve stimulation in patients with lower limb vascular disease. *Blood Coagul Fibrinolysis* 2016; 27: 275–280.
50. Jingwei L ZZ, Xuesong L, Hui J, Ning L, Wenbo W Clinical observation of neuromuscular electrical stimulation in prevention of deep venous thrombosis after total hip replacement. *Chin J Bone Joint Injury* 2017; 32: 615–616.
51. Broderick BJ, Breathnach O, Masterson E, Breen PP, G OL. Hemodynamic performance of NMES in the early post operative period following orthopaedic surgery. *Conf Proc IEEE Eng Med Biol Soc* 2011; 2011: 7630–7633.
52. Broderick BJ, O'Briain DE, Breen PP, Kearns SR, O'laighin G. A pilot evaluation of a neuromuscular electrical stimulation

- (NMES) based methodology for the prevention of venous stasis during bed rest. *Med Eng Phys* 2010; 32: 349–355.
53. Tucker A, Maass A, Bain D, Chen LH, Azzam M, Dawson H, et al. Augmentation of venous, arterial and microvascular blood supply in the leg by isometric neuromuscular stimulation via the peroneal nerve. *Int J Angiol* 2010; 19: e31–e37.
 54. Burgess LC, Immins T, Swain I, Wainwright, TW. Effectiveness of neuromuscular electrical stimulation for reducing oedema: a systematic review. *J Rehab Med* 2019; 51: 237–243.
 55. Wainwright TW, Burgess LC, Middleton RG. A feasibility randomised controlled trial to evaluate the effectiveness of a novel neuromuscular electro-stimulation device in preventing the formation of oedema following total hip replacement surgery. *Heliyon* 2018; 4: e00697.
 56. Badger J, Taylor P, Papworth N, Swain I. Electrical stimulation devices for the prevention of venous thromboembolism: preliminary studies of physiological efficacy and user satisfaction. *J Rehabil Assist Technol Eng* 2018; 5: 2055668318800218.
 57. Bahadori S, Immins T, Wainwright TW. The effect of calf neuromuscular electrical stimulation and intermittent pneumatic compression on thigh microcirculation. *Microvasc Res* 2017; 111: 37–41.
 58. Ojima M, Takegawa R, Hirose T, Ohnishi M, Shiozaki T, Shimazu T. Hemodynamic effects of electrical muscle stimulation in the prophylaxis of deep vein thrombosis for intensive care unit patients: a randomized trial. *J Intens Care* 2017; 5: 9–9.
 59. Lindstrom B, Holmdahl C, Jonsson O, Korsan-Bengtson K, Lindberg S, Petrusson B, et al. Prediction and prophylaxis of postoperative thromboembolism – a comparison between peroperative calf muscle stimulation with groups of impulses and dextran 40. *Br J Surg* 1982; 69: 633–637.
 60. Browse NL, Negus D. Prevention of postoperative leg vein thrombosis by electrical muscle stimulation. An evaluation with 125I-labelled fibrinogen. *Br Med J* 1970; 3: 615–618.
 61. Wainwright TW, Burgess LC, Middleton RG. A single-centre feasibility randomised controlled trial comparing the incidence of asymptomatic and symptomatic deep vein thrombosis between a neuromuscular electrostimulation device and thromboembolism deterrent stockings in post-operative patients recovering from elective total hip replacement surgery. *Surg Technol Int* 2020; 36: 289–298.
 62. Doran FS, White HM. A demonstration that the risk of postoperative deep venous thrombosis is reduced by stimulating the calf muscles electrically during the operation. *Br J Surg* 1967; 54: 686–689.
 63. Dejode LR, Khurshid M, Walther WW. The influence of electrical stimulation of the leg during surgical operations on the subsequent development of deep-vein thrombosis. *Br J Surg* 1973; 60: 31–32.
 64. Hajibandeh S, Hajibandeh S, Antoniou GA, Scurr JR, Torella F. Neuromuscular electrical stimulation for the prevention of venous thromboembolism. *Cochrane Database Syst Rev* 2017; 11: CD011764.
 65. Hajibandeh S, Hajibandeh S, Antoniou GA, Scurr JR, Torella F. Neuromuscular electrical stimulation for thromboprophylaxis: A systematic review. *Phlebology* 2015; 30: 589–602.
 66. Ravikumar R, Williams KJ, Babber A, Moore, HM, Lane TRA, Shalhoub J et al. Neuromuscular electrical stimulation for the prevention of venous thromboembolism. *Phlebology* 2018 ; 33: 367–378.
 67. Sosnowski K, Mitchell ML, White H, Morrison L, Sutton J, Sharratt J, Lin F. A feasibility study of a randomised controlled trial to examine the impact of the ABCDE bundle on quality of life in ICU survivors. *Pilot Feasibil Studies* 2018; 4: 32.
 68. Dos Santos FV, Cipriano G, Jr., Vieira L, Guntzel Chiappa AM, Cipriano GBF, Vieira P, et al. Neuromuscular electrical stimulation combined with exercise decreases duration of mechanical ventilation in ICU patients: a randomized controlled trial. *Physiother Theory Pract* 2018; 10.1080/09593985.2018.1490363: 1–9.
 69. Duffell LD, Donaldson NN. A Comparison of FES and SCS for neuroplastic recovery after SCI: historical perspectives and future directions. *Front Neurol* 2020; 11: 607.
 70. Latronico N, Bolton CF. Critical illness polyneuropathy and myopathy: a major cause of muscle weakness and paralysis. *Lancet Neurol* 2011; 10: 931–941.
 71. Schweickert WD, Pohlman MC, Pohlman AS, Nigos C, Pawlik AJ, Esbrook CL, et al. Early physical and occupational therapy in mechanically ventilated, critically ill patients: a randomised controlled trial. *Lancet* 2009; 373: 1874–1882.
 72. Kasotakis G, Schmidt U, Perry D, Grosse-Sundrup M, Benjamin J, Ryan C, et al. The surgical intensive care unit optimal mobility score predicts mortality and length of stay. *Crit Care Med* 2012; 40: 1122–1128.
 73. Calvo-Ayala E, Khan BA, Farber MO, Ely EW, Boustani MA. Interventions to improve the physical function of ICU survivors: a systematic review. *Chest* 2013; 144: 1469–1480.
 74. Kho ME, Martin RA, Toonstra AL, Zanni JM, Manthey EC, Nelliott A, et al. Feasibility and safety of in-bed cycling for physical rehabilitation in the intensive care unit. *J Crit Care* 2015; 30: 1419 e1411–1415.
 75. Helms J, Kremer S, Merdji H, Clere-Jehl R, Schenck M, Kumarlen C, et al. Neurologic features in severe SARS-CoV-2 infection. *N Engl J Med* 2020; 382: 2268–2270.
 76. Munshi L, Del Sorbo L, Adhikari NKJ, Hodgson CL, Wunsch H, Meade MO, et al. Prone position for acute respiratory distress syndrome: a systematic review and meta-analysis. *Ann Am Thorac Soc* 2017; 14: S280–S288.
 77. Iwatsu K, Yamada S, Iida Y, Sampei H, Kobayashi K, Kainuma M, Usui A. Feasibility of neuromuscular electrical stimulation immediately after cardiovascular surgery. *Arch Phys Med Rehabil* 2015; 96: 63–68.
 78. Segers J, Hermans G, Bruyninckx F, Meyfroidt G, Langer D, Gosselink R. Feasibility of neuromuscular electrical stimulation in critically ill patients. *J Crit Care* 2014; 29: 1082–1088.
 79. Badger J, Taylor P, Swain I. The safety of electrical stimulation in patients with pacemakers and implantable cardioverter defibrillators: a systematic review. *J Rehabil Assist Technol Eng* 2017; 4: 2055668317745498.
 80. Pal S, Chughtai M, Sultan AA, Khlopas A, Sodhi N, George NE et al. Impact of neuromuscular electrical stimulation (NMES) on 90-day episode costs and post-acute care utilization in total knee replacement patients with disuse atrophy. *Surg Technol Int* 2017; 31: 384–388.
 81. Stevens-Lapsley JE, Balter JE, Wolfe P, Eckhoff DG, Schwartz RS, Schenkman M, et al. Relationship between intensity of quadriceps muscle neuromuscular electrical stimulation and strength recovery after total knee arthroplasty. *Phys Ther* 2012; 92: 1187–1196.