



## LJMU Research Online

Zhao, W, Peng, Y, Hu, Y, Guo, XE, Li, J, Cao, J, Pan, J, Feng, JQ, Cardozo, C, Jarvis, JC, Bauman, WA and Qin, W

**Electrical stimulation of hindlimb skeletal muscle has beneficial effects on sublesional bone in a rat model of spinal cord injury**

<http://researchonline.ljmu.ac.uk/id/eprint/17085/>

### Article

**Citation** (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

**Zhao, W, Peng, Y, Hu, Y, Guo, XE, Li, J, Cao, J, Pan, J, Feng, JQ, Cardozo, C, Jarvis, JC, Bauman, WA and Qin, W (2021) Electrical stimulation of hindlimb skeletal muscle has beneficial effects on sublesional bone in a rat model of spinal cord injury. Bone. 144. ISSN 8756-3282**

LJMU has developed [LJMU Research Online](http://researchonline.ljmu.ac.uk/) for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact [researchonline@ljmu.ac.uk](mailto:researchonline@ljmu.ac.uk)

<http://researchonline.ljmu.ac.uk/>

1 **Electrical Stimulation of Hindlimb Skeletal Muscle has Beneficial Effects on Sublesional**  
2 **Bone in a Rat Model of Spinal Cord Injury**

3  
4 Wei Zhao<sup>1,2</sup>, Yuanzhen Peng<sup>1</sup>, Yizhong Hu<sup>4</sup>, X. Edward Guo<sup>4</sup>, Jiliang Li<sup>5</sup>, Jay Cao<sup>6</sup>, Jiangping Pan<sup>1</sup>,  
5 Jian Q. Feng<sup>7</sup>, Christopher Cardozo<sup>1,2,3</sup>, Jonathan Jarvis<sup>8</sup>, William A. Bauman<sup>1,2,3</sup>, and Weiping  
6 Qin<sup>1,2\*</sup>

7 <sup>1</sup> National Center for the Medical Consequences of SCI, James J. Peters VA Medical Center,  
8 Bronx, NY; <sup>2</sup> Departments of Medicine, and <sup>3</sup> Rehabilitation Medicine, Icahn School of Medicine  
9 at Mount Sinai, New York, NY; <sup>4</sup> Department of Biomedical Engineering, Columbia University,  
10 New York, NY; <sup>5</sup> Indiana University Purdue University Indianapolis, Indianapolis, IN, USA;  
11 <sup>6</sup> United States Department of Agriculture Agricultural Research Service Human Nutrition  
12 Research Center, Grand Forks, North Dakota; <sup>7</sup> Baylor College of Dentistry, TX A&M, Dallas, TX,  
13 USA; <sup>8</sup> School of Sport and Exercise Sciences, Liverpool John Moores University, Liverpool L3  
14 3AF, United Kingdom.

15  
16 \* Corresponding Author:

17 Weiping Qin, MD, Ph.D.

18 James J. Peters Veteran Affairs Medical Center  
19 130 West Kingsbridge Road, Bronx, NY 10468, USA

20 Tel: 718-584-9000 X1831

21 Email: [weiping.qin@mssm.edu](mailto:weiping.qin@mssm.edu) or [weiping.qin@va.gov](mailto:weiping.qin@va.gov)

22

23 **Running title:** Benefits of ES on bone after SCI.

24

1 **Abstract**

2 Spinal cord injury (SCI) results in marked atrophy of sublesional skeletal muscle and substantial  
3 loss of bone. In this study, the effects of prolonged electrical stimulation (ES) and /or  
4 testosterone enanthate (TE) on muscle mass and bone formation in a rat model of SCI were  
5 tested. Compared to sham-transected animals, a significant reduction of the mass of soleus,  
6 plantaris and extensor digitorum longus (EDL) muscles was observed in animals 6 weeks post-  
7 SCI. Notably, ES or ES+TE resulted in the increased mass of the EDL muscles. ES or ES+TE  
8 significantly decreased mRNA levels of muscle atrophy markers (e.g., MAFbx and MurF1) in the  
9 EDL. Significant decreases in bone mineral density (BMD) (-27%) and trabecular bone volume  
10 (-49.3%) at the distal femur were observed in animals 6 weeks post injury. TE, ES and ES+TE  
11 treatment significantly increased BMD by +6.4%, +5.4%, +8.5% and bone volume by +22.2%,  
12 and +56.2% and+ 60.2%, respectively. Notably, ES alone or ES+TE resulted in almost  
13 complete restoration of cortical stiffness estimated by finite element analysis in SCI animals.  
14 Osteoblastogenesis was evaluated by colony-forming unit-fibroblastic (CFU-F) staining using  
15 bone marrow mesenchymal stem cells obtained from the femur. SCI decreased the CFU-F+  
16 cells by -56.8% compared to sham animals. TE or ES+TE treatment after SCI increased  
17 osteoblastogenesis by +74.6% and +67.2%, respectively. An osteoclastogenesis assay  
18 revealed significantly increased TRAP+ multinucleated cells (+34.8%) in SCI animals compared  
19 to sham animals. TE, ES and TE+ES treatment following SCI markedly decreased TRAP+ cells  
20 by -51.3%, -40.3% and -46.9%, respectively. Each intervention greatly reduced the ratio of  
21 RANKL to OPG mRNA of sublesional long bone. Collectively, our findings demonstrate that  
22 after neurologically complete paralysis, dynamic muscle resistance exercise by ES reduced  
23 muscle atrophy, downregulatd genes involved in muscle wasting, and restored mechanical  
24 loading to sublesional bone to a degree that allowed for the preservation of bone by inhibition of  
25 bone resorption and/or by facilitating bone formation.

26 **Keywords:** spinal cord injury; electrical stimulation; muscle; bone

## 1 **Introduction**

2           Immobilization results in muscle atrophy and decreased bone mass and structural  
3 integrity. Spinal cord injury (SCI) results in paresis or paralysis and protracted immobilization  
4 associated with extensive sublesional muscle atrophy and loss of bone mass and structural  
5 integrity [1-3]. In rats, bone loss following SCI develops rapidly, being evident within 2 days after  
6 complete spinal cord transection [4] and 2 weeks after a severe spinal cord contusion, and  
7 sublesional skeletal deterioration is most predominant for trabecular bone at the metaphysis  
8 and epiphysis of the femur and tibia where bone volume/total volume may be reduced by more  
9 than 50 percent [4-7]. Mechanisms implicated in bone loss in the period shortly after SCI are  
10 thought to primarily involve increased osteoclastic bone resorption with unchanged or slightly  
11 increased bone formation, apparently attributable to elevated cellular expression of receptor  
12 activator of NF- $\kappa$ B ligand (RANKL) and SOST, which encodes the protein sclerostin, a potent  
13 inhibitor of bone formation [5, 6, 8]. Loss of cortical bone has been reported in patients with SCI  
14 [2, 9], mice [10] and rats [7], although such loss occurs more slowly and to a lesser extent than  
15 that of trabecular bone, perhaps due to the smaller surface area accessible to osteoclasts for  
16 bone resorption.

17           Muscle atrophy following SCI is rapid, with muscle weights being reduced by 40 to 60%  
18 at two weeks after spinal cord transection when compared to sham-operated controls [11].  
19 Mechanisms responsible are thought to involve primarily a marked upregulation of proteolytic  
20 degradation of muscle proteins likely attributable to the muscle-restricted E3 ubiquitin ligases  
21 muscle atrophy F-box (MAFBx, atrogin-1) and muscle ring finger-1 (MuRF1, Trim63), the  
22 expression of which is increased within days after SCI [1, 3, 11]. Both muscle atrophy and  
23 sublesional bone loss are mitigated in a dose-dependent manner by testosterone [12].  
24 Beneficial effects of testosterone on muscle atrophy have been reported to have numerous  
25 potential mechanisms of action that include suppression of MAFbx expression, upregulation of

1 PGC-1 $\alpha$  expression, activation of mTOR and suppression of FOXO activation [13-16]. In mice,  
2 androgen receptor knockouts reduce trabecular bone, apparently due to increased bone  
3 resorption [17]. Additionally, testosterone can undergo aromatization to estrogen which appears  
4 to be beneficial to preserving bone as demonstrated by findings deletion of the estrogen  
5 receptor alpha resulted in the reduction of trabecular bone [18].

6 An increase in physical activity, whether voluntary in the able-bodied or through use of  
7 surface or implantable electrical stimulation (ES) of nerves to elicit muscle contraction, has been  
8 shown to preserve muscle mass when initiated acutely after SCI, and restore, at least partially,  
9 muscle size and function once atrophy has occurred [19-21]. In rats with SCI, ES has been used  
10 to study acute responses of paralyzed muscle and bone to reloading [5, 22]. In those studies,  
11 ES was initiated approximately 4 months after SCI and continued for 7-days. While soleus  
12 muscles increased in weight in the SCI animals provided ES, expected changes in  
13 mitochondrial metabolism of carbohydrates and fats, and oxidative phosphorylation were not  
14 observed in contrast to controls in which muscle overloading by gastrocnemius ablation was  
15 conducted for the same period [22]. Cultured bone marrow progenitors were studied to  
16 understand how ES for 7 days may have influenced bone cells post-SCI [5]. Among the  
17 changes noted, numbers of osteoclast colonies, which increased in marrow from SCI animals,  
18 were decreased by ES, and, of interest, in cultures of marrow stromal cells under conditions  
19 favoring differentiation to osteoblasts, ES lowered SOST mRNA and raised mRNA for  
20 osteoprotegrin, a decoy receptor to RANKL, thus inhibiting the ability of RANKL to stimulate  
21 osteoclastogenesis and osteoclast activity [5]. However, no change in bone parameters was  
22 observed after 7 days of ES when assessed by microCT, most probably due to the short  
23 duration of ES. The purpose of this study was to determine the effects of an extended period of  
24 ES on muscle and bone in a rat model of severe SCI, and to test for synergist effects of ES  
25 when combined with supraphysiologic doses of testosterone.

## 1 **Materials and Methods**

### 2 Animals:

3 Animals: All animals were maintained on a 12:12-h light/dark cycle with lights on at 07:00 h in a  
4 temperature-controlled ( $20 \pm 2^\circ\text{C}$ ) vivarium. All procedures were approved by the JJP VA  
5 Medical Center Institutional Animal Care and Use Committee and were performed in  
6 accordance with applicable requirements of NIH, PHS and VA.

7  
8 *Experimental Design:* Animals were randomly assigned to either Sham SCI (laminectomy only)  
9 or SCI (spinal cord transection at T4). The SCI groups were randomly assigned to receive ES,  
10 testosterone enanthate (TE) or the combination. Interventions were initiated 2 weeks post  
11 surgery and continued for 4 weeks at which time tissue samples were collected for processing  
12 and animals were euthanized. The electrical stimulation for 4 weeks in this rat study is  
13 equivalent of approximately 2.5 years in humans (Ruth 1935) [23], representing a sufficient  
14 treatment duration when compared to months of intervention in most clinical studies with  
15 electrical stimulation [24]. TE was administered intramuscularly to the quadriceps of the  
16 unstimulated right leg once a week at a dose of 4.0 mg; vehicle (sesame oil) was used in SCI or  
17 SCI+ES animals. Group sizes are as follows: Sham, n=13; SCI, n=13; SCI+TE, n=13; SCI+ES,  
18 n=10; and SCI+ES+TE, n=11.

19  
20 *ES Stimulators:* ES was provided by implantable microstimulators with a single channel.  
21 Microstimulators were custom-designed and fabricated and encapsulated in silastic and gas  
22 sterilized. Stimulators were designed to be activated using a small magnet and had three modes  
23 of operation: off, testing and active. In the active mode, stimulators delivered 1.5 V at 40 Hz for  
24 2 seconds followed by 18 seconds rest, as we described previously [5, 25]. Stimulation after  
25 severing the insertion of the gastrocnemius muscle into the Achilles tendon (see below for

1 details) elicited some movement of the left foot and ankle with the ankle in a neutral position at  
2 approximately 90 degrees of dorsiflexion. Stimulation was provided for 60 min on each training  
3 day, five days a week.

4  
5 *Surgeries:* Male Wistar rats (13 months old) underwent a complete spinal cord transection (T3-  
6 4) as previously described [4, 5, 22, 26-28]. Sham controls received laminectomy only at the  
7 same site, as previously described [5, 22]. Implantation of the stimulators was performed  
8 immediately after spinal transection, as previously described [5, 22], with some modifications. In  
9 brief, a small incision (~3 cm) was made over the left hip parallel to the femur. A pocket was  
10 created under the skin of the lower back for the stimulator. The pocket began at the point of  
11 origin of the incision with a size of approximately 5 by 5 cm. The bifurcation of the left sciatic  
12 nerve was exposed by blunt dissection. The stimulator was then inserted into the pocket and  
13 sutured in place with a single loop through a webbing tab; the two electrodes were tunneled  
14 subcutaneously from the back to the incision site over the bifurcation of the sciatic nerve. The  
15 first electrode was placed such that it nearly touched the anterior tibial nerve just cephalad to  
16 the nerve and about 0.8 cm distal to the bifurcation of the sciatic nerve. The second electrode  
17 was placed about 0.2 cm caudal from the common peroneal nerve at about 1 cm distal to the  
18 bifurcation. Tenotomy of the left gastrocnemius muscle was performed in all experimental  
19 groups by separating the tendons of the gastrocnemius, soleus and plantaris muscles and  
20 transecting the gastrocnemius muscle tendon. As demonstrated in our previous studies [5, 25],  
21 cutting the distal insertion of the gastrocnemius prevents concurrent contraction of this muscle  
22 from overwhelming the opposing force of the contracting tibialis anterior. SCI and Sham animals  
23 underwent an identical procedure except that no stimulator was implanted. We note that  
24 compared to our initial work with implanting the stimulators [5, 25], the size of the incision  
25 through which the electrodes are placed has been greatly reduced that was observed to  
26 considerably reduce excessive grooming and potential damage to the electrodes.

1 *Fluorometric labeling of bone:* Bones were labeled with fluorochromes by subcutaneous  
2 injection of calcein (10 mg/kg body weight) and xylenol orange (90 mg/kg body weight) on day  
3 -6 and day -2 before euthanasia, respectively [5, 26, 28].

#### 4 5 ELISA assays

6 Serum C-terminal telopeptide of type I collagen (CTX) levels were measured using a  
7 RatLaps™ enzyme-immunoassay kit from Immunodiagnostic Systems (Fountain Hills, AZ).  
8 Serum concentrations of osteocalcin were measured using a rat osteocalcin immunoassay kit  
9 (Alfa Aesar). Serum levels of irisin were determined using a rat irisin competitive ELISA kit  
10 (Adipogen). All samples were assayed in duplicate, following the manufacturer's protocols.

#### 11 12 Bone Density, Structure, and Strength

13 Areal bone mineral density (BMD) measurement was performed with a small animal dual energy  
14 X-ray absorptiometer (DXA) (Lunar Piximus, Fitchburg, WI, USA) as previously described [4, 6,  
15 26, 27, 29, 30]. Volumetric BMD and bone architecture of the left distal femur and midshaft were  
16 assessed by a Scanco  $\mu$ CT scanner ( $\mu$ CT-40; Scanco Medical AG, Bassersdorf, Switzerland) at  
17 21  $\mu$ m isotropic voxel size as previously described [31, 32]. The  $\mu$ CT images were evaluated  
18 using standard software of the manufacturer to evaluate relative bone volume (BV/TV),  
19 trabecular number (Tb.N), trabecular thickness (Tb.Th), trabecular separation (Tb.Sp), structure  
20 model index (SMI), connectivity density (Conn.D), total cortical area (Ct.Ar), cortical thickness  
21 (Ct.Th), periosteal perimeter, and endosteal perimeter. Bone stiffness was estimated from micro  
22 finite element analysis ( $\mu$ FEA), following the manufacturer's recommended procedures, as  
23 previously described [4, 28, 33-37]. Briefly,  $\mu$ FEA models were produced by converting each  
24 bone voxel to an 8-node brick element. Bone tissue was computationally subjected to applied  
25 uniaxial compression, with an elastic modulus of 15 GPa and Poisson's ratio of 0.3 for each



1 element. A linear elastic analysis was used to estimate the bone stiffness. Please see the  
2 Supplemental Materials and Methods for additional details.

### 3 Bone Histomorphometric Studies

4 For fluorochrome-based determination of rates of bone formation at the left distal femur by  
5 dynamic histomorphometry, 6 mm frozen sections embedded in methyl methacrylate plastic  
6 were cut using a Reichert-Jung sledge microtome. Xylenol orange and calcein were visualized  
7 by fluorescent microscopy and the distance between labeled layers was used as a measure of  
8 the rate of bone formation as determined by morphometry software. To quantify the osteoclast  
9 number and activity, tartrate-resistant acid phosphatase (TRAP) staining was used to  
10 specifically label osteoclasts in deplasticized distal femur sections. Slides were counterstained  
11 with hematoxylin and eosin (H&E). Osteoclasts were measured under bright field microscopy  
12 using an Olympus microscope with an OsteoMeasure system. Additional details regarding the  
13 procedures performed are provided in Supplemental Materials.

14

### 15 Extraction of total RNA from bone

16 Total bone RNA was extracted as previously described with some modifications [4, 38]. Briefly,  
17 long bones were dissected free of soft tissues, and bone marrow were flushed away with PBS  
18 using a 27G<sup>1/2</sup> needle-syringe. The bone samples (~1g) were longitudinally cut into small piece  
19 and then digested 3 times with 2 mg/ml collagenase type I (Gibco, >150 Units; 20ml), one time  
20 with 5mM EDTA (Sigma-Aldrich, 10 ml)), and one more with the collagenase and EDTA, each  
21 for 25 min on a Shaker with rotation at 150 rpm at 37°C. Following the digestions, the bone  
22 samples were crushed using a mortar and pestle in liquid nitrogen. RNA was extracted from the  
23 lysate using the TRizol reagent (Sigma Aldrich) according to the manufacturer's instructions.

24

25

1 *Ex vivo Osteoblastogenesis and Osteoclastogenesis Assay*

2 Procedures for osteoblast and osteoclast formation from bone marrow stem cells were  
3 performed, as previously described [5, 27, 29-31, 39] and are described in greater detail in the  
4 Supplemental Materials.

5

6 *RNA Extraction from Bone Marrow Cultures and Quantitative PCR*

7 Total RNAs were extracted from bone marrow cell cultures using the TRI reagent (Sigma-  
8 Aldrich). One  $\mu\text{g}$  of total RNA was used to synthesize the first strand cDNA by the High Capacity  
9 cDNA Reverse Transcription Kit (Applied Biosystems). Real-time PCR determination of mRNA  
10 levels for specific mRNAs was determined with commercial primers from Taqman Assay On  
11 Demand probesets (Applied Biosystems) using the ViiA7 system (Applied Biosystems) as  
12 described previously [5, 27, 29-31, 39]. Relative expression levels were calculated using the 2-  
13  $\Delta\Delta\text{Ct}$  method with 18S RNA as an internal control [40]. Additional details regarding procedures  
14 performed are provided in Supplemental Materials.

15

16 *Statistics*

17 The mean values for study endpoints of stimulated (ipsilateral) left hindlimb among the  
18 experimental groups were compared. Data are expressed as mean  $\pm$  standard error of the  
19 mean (SEM). The number of independent samples (n) is provided in the legend of each figure.  
20 The statistical significance of differences among means was tested using one-way analysis of  
21 variance and a Newman-Keuls post hoc test to determine the significance of differences  
22 between individual pairs of means using a p value of  $<0.05$  to attain significance. Statistical  
23 calculations were performed using Prism 4.0c (GraphPad Software, La Jolla, CA, USA).

24

25

1 **Results**

2 Spinal cord transected animals lost approximately  $10\pm 3\%$  percent of their preoperative body  
3 weight (Supplemental Fig. 1), a finding which is consistent with prior studies. Animals began  
4 gaining weight by day 10 post-injury and continued to gain weight until the end of the study at  
5 day 42 post injury (Supplemental Fig. 1). Implantation of a stimulator and / or administration of  
6 TE in SCI animals did not result in significant body weight change.

7  
8 *Beneficial Effects of ES and/or TE on Muscle:*

9  
10 To verify the effectiveness of ES in stimulating muscle overloading, weights of soleus, plantaris  
11 and extensor digitorum longus (EDL) muscles from hindlimbs were removed by careful  
12 dissection and weighed. The left (stimulated) muscles were evaluated and all muscle results are  
13 relative mass that has been corrected for body mass. As compared to sham-transected animals,  
14 significant reduction of the left soleus, plantaris and EDL muscle mass was found in animals  
15 that received complete spinal cord transection (Fig. 1A-C). Although it did not reach statistical  
16 significance, TE, ES or the combination of these two treatments was linked to a numerical  
17 increase of the left (e.g., stimulated) plantaris muscle mass (+6.8%, +7.2% and +12.7%) and  
18 soleus muscle (+4.4%, +8.7% and +9.9%), respectively (Fig. 1A-B). Notably, ES alone or  
19 ES+TE treatment resulted in a clear and significant ( $\sim 13\%$ ,  $p < 0.05$ ) increase of the left EDL  
20 muscle mass (Fig. 1C). Since SCI results in muscle atrophy and a shift of slow oxidative (e.g.,  
21 soleus muscle) to fast glycolytic fibers (e.g., EDL muscle) [5, 41] and the observed changes on  
22 EDL muscle mass in response to ES alone or ES+TE treatment in this study (Fig. 1C),  
23 subsequent analyses will focus on the EDL muscle.

24  
25 To further examine the effects of ES on muscle reloading, mRNA levels of two muscle atrophy  
26 related muscle-specific E3 ubiquitin ligases, MAFbx/atrogen-1 and Muscle RING Finger-1  
27 (MuRF1) in the left EDL were examined. SCI resulted in a significant increase of MAFbx mRNA

1 expression (Fig. 1D,  $p < 0.05$ ). TE injection did not change the MAFbx expression level, whereas  
2 a significant decrease of MAFbx mRNA was found upon ES alone or ES +TE treatment (Fig.  
3 1D,  $p < 0.001$  and  $p < 0.05$ , respectively). Of note, although no change of MurF1 mRNA  
4 expression was detected after SCI, each intervention, TE, ES or ES+TE, significantly reduced  
5 MurF1 mRNA levels (Fig. 1E,  $p < 0.05$ ,  $p < 0.01$  and  $p < 0.001$ , respectively). The mRNA levels of  
6 the Wnt antagonists secreted frizzled-related protein (sFRP) 1 in the EDL muscle were  
7 increased after SCI, and were drastically reduced by TE injection, ES or ES+TE treatment (Fig.  
8 1F,  $p < 0.001$ ). Compared to SCI animals, there was a non-significant trend of a decrease of  
9 sFRP2 after TE injection, but no change after ES or ES+TE (Fig. 1G).

10  
11 Collectively, these data from sublesional skeletal muscle indicate that dynamic muscle  
12 resistance exercise by ES reduced muscle atrophy, downregulating key genes associated with  
13 muscle wasting. The alterations in muscle mass and total gene expression reflect biological  
14 changes as a direct consequence of ES intervention. The potential impact of mechanical  
15 reloading of muscle by ES on bone will be examined in subsequent studies.

16

#### 17 *Beneficial Effects of ES and/or TE on Bone Mass, Bone Structure and Bone Strength:*

18 Dual energy x-ray absorptiometry (DXA) was performed using a small animal densitometer to  
19 determine bone mineral density (BMD) in rats after a complete spinal cord transection and TE  
20 injection, ES or TE+ES. At the left distal femur (Fig. 2A), BMD were decreased by -27%  
21 ( $p < 0.001$ ) after SCI, and significantly increased by +5.4% ( $p = 0.07$ ) and +8.5% ( $p < 0.01$ ) after ES  
22 and ES+TE treatments, respectively. An almost identical pattern of BMD change was also  
23 detected at the left proximal tibia (Fig. 2B) where BMD was reduced by -23.5% ( $p < 0.001$ ) after  
24 SCI. Compared to SCI, BMD at the proximal tibia increased +5.9% ( $p < 0.05$ ) and +11%  
25 ( $p < 0.001$ ) following ES and ES+TE treatment, respectively. The larger magnitude of changes on  
26 BMD in distal femur and proximal tibia in ES+TE animals as compared to TE or ES animals

1 suggests a potential synergistic effect of TE and ES on bone mass. SCI also resulted in a -  
2 13.9% decrease of BMD at L3-5 (Fig. 2C). No effect of TE, ES or the combination was observed  
3 on BMD at L3-L5.

4 Bone architecture was examined by high-resolution  $\mu$ CT to access changes in trabecular bone  
5 at the distal femoral metaphysis (Fig. 3A). After SCI, trabecular bone volume (BV/TV) at this site  
6 was significantly reduced ( $p < 0.001$ , Fig. 3B), with decreased trabecular number (Tb. N) (Fig.  
7 3C), increased trabecular separation (Tb. Sp) (Fig. 3E) and no change in trabecular thickness  
8 (Tb. Th) (Fig. 3D). Trabecular connectivity (Conn. D) was greatly reduced (Fig. 3F), associated  
9 with transformation from plate-like to rod-like structures (Fig. 3G). Electrical stimulation alone or  
10 combined with TE significantly restored trabecular bone volume (Fig. 3B) by increasing  
11 trabecular number (Tb.N), which is associated with the increased connectivity (Fig. 3F) and the  
12 decreased value of the structure model index (SMI, Fig. 3G).

13  
14 Cortical bone structure at the femoral midshaft was also examined by high-resolution  $\mu$ CT (Fig.  
15 4A). SCI decreased cortical thickness (-9.0%,  $p < 0.05$ , Fig. 4B). The decreased cortical  
16 thickness could be explained by a decreased periosteal perimeter (-7.0%,  $p < 0.05$ , Fig. 4E) with  
17 no change on endosteal perimeter (Fig. 4F). Electrical stimulation in SCI animals partially  
18 restored periosteal perimeter (+4.9%,  $p < 0.05$ , Fig. 4E). ES+TE administration completely  
19 restored cortical thickness (+9.5%,  $p < 0.05$ , Fig. 4G), suggesting a synergistic effect of ES+TE.

20  
21 Finite element analysis (FEA) based on  $\mu$ CT images was performed to predict the effect of ES  
22 and / or TE on bone mechanical property and strength. After SCI, midshaft bone stiffness  
23 estimated by FEA was reduced by -12.8% ( $p < 0.05$ , Fig. 4F). ES alone or ES+TE led to a nearly  
24 complete restoration of cortical stiffness [+11.4% vs. SCI ( $p < 0.05$ ) and +12.8% vs. SCI ( $p < 0.01$ ),  
25 respectively, Fig. 4E].

1

2 Effects of ES and/or TE on Serum Levels of Bone Biomarkers and Bone Gene Expressions

3 ELISA assays were performed to examine the serum levels of bone biomarkers for formation,  
4 osteocalcin and resorption, CTX. Although no change of osteocalcin or CTX were detected  
5 following SCI, TE treatment alone or ES alone significantly increased the serum osteocalcin level  
6 ( $p<0.05$  or  $p<0.01$ , respectively, Fig. 5A), while ES+TE treatment marginally decreased serum  
7 CTX level ( $p=0.08$ , Fig. 5B). Recent evidence showed that irisin, a myokine released after  
8 physical activity, increased cortical bone mass [42]. However, our ELISA assay showed that SCI  
9 did not change the serum irisin level, but ES alone, TE alone or ES+TE combined treatment after  
10 SCI all resulted in significantly reduced Irisin level compared to sham animals (Fig. 5C).

11

12 Total RNA from whole bone of hindlimb was extracted, and RANKL and OPG gene expression  
13 that are responsible for bone resorption and formation, respectively, was analyzed by quantitative  
14 PCR analysis. Following SCI, increases of the RANKL/OPG ratio ( $p<0.01$ , Fig. 5D) were observed  
15 in samples from SCI group compared with Sham-transected animals. The RANKL/OPG ratio was  
16 greatly reduced by ES or TE. The combination of ES+TE led to almost a complete normalization  
17 of RANKL/OPG ratio, approaching the ratio observed in Sham group (Fig. 5D).

18

19 Effects of ES and/or TE on Bone Formation and Bone Resorption

20 Dynamic histomorphometric analysis was performed to examine the effect of ES and/or TE on  
21 bone formation (Fig. 6A). Significant decreases in bone formation rate (BFR/TV, Fig. 6B), bone  
22 volume (BV/TV, Fig. 6F) and bone surface (B. Pm, Fig. 6G) were detected at the distal femur in  
23 animals with SCI. TE treatment significantly increased new bone formation at the trabecular  
24 bone, as reflected by increases in the bone formation rate (BFR/TV, +110%,  $p<0.05$ , Fig. 6B),  
25 mineralizing surface (MS/BS, +22%,  $p<0.05$ , Fig. 6C), bone volume (BV/TV, +138%,  $p<0.05$ ,  
26 Fig. 6F), bone surface (B. Pm, +109%,  $p<0.05$ , Fig. 6G), and dL Pm (+163%,  $p<0.05$ ,

1 Supplemental Fig 2B). ES significantly increased bone formation rate (Fig. 6B) and stimulated a  
2 trend for increases in mineral apposition rate (MAR, Fig. 6D), bone volume (Fig. 6F), and bone  
3 surface (Fig. 6G). ES tends to increase dL Pm (+36%,  $p=0.06$ , Supplemental Fig 2B) ES+TE  
4 treatment also partially restored single-labeled surface (sl.Pm) but not double-labeled surface  
5 (dl.Pm), both of which were decreased following SCI (Supplemental Figure 2A&B).

6  
7 Sections of trabecular bone from the femoral metaphysis were immune-stained for TRAP (Fig.  
8 7A). SCI resulted in trends for increases in both osteoclast surface (Fig. 7B) and osteoclast  
9 number (Fig. 7C). TE treatment alone or ES+TE treatments significantly decreased osteoclast  
10 surface and number (Fig. 7B-C), while ES alone only showed a trend for a decrease in  
11 osteoclast surface and number.

12

### 13 *Effects of ES and/or TE on the Differentiation Potential of Bone Marrow Stem Cells*

14 How ES and/or TE alter numbers of osteoclasts and osteoblasts in cultures of bone marrow  
15 cells was determined. The effects on bone cells of TE+ ES with the effects observed for ES or  
16 TE were compared. Osteoblastogenesis was evaluated by colony-forming unit-fibroblastic  
17 (CFU-F) staining using bone marrow mesenchymal stem cells derived from femurs from the left  
18 (e.g., stimulated) hindlimb (Fig. 8A). Consistent with our previous findings [5, 6, 27, 30], SCI  
19 decreased the numbers of CFU-F<sup>+</sup> cells by -56.8% ( $p<0.05$ ) as compared to sham animals.  
20 Either TE or ES+TE treatment after SCI increased numbers of CFU-F<sup>+</sup> cells by +74.6% ( $p<0.01$ )  
21 and +67.2% ( $p<0.05$ ), respectively, indicating greater potential of marrow stromal cells for  
22 osteoblastic differentiation. After SCI, decreased expression of Runx2 (Fig. 8C) and LRP5 (Fig.  
23 8E) mRNA, as well as increased expression of SOST (Fig. 8D) and sFRP1 (Fig. 8F) mRNA  
24 were detected by quantitative PCR analysis. TE or ES+TE treatment significantly elevated the  
25 mRNA level of Runx2 and LRP5 (Fig. 8C, E); ES also increased the expression of these two  
26 osteoblastogenic markers but to a lesser extent. Interestingly, only ES+TE treatment decreased

1 the expression of SOST and sFRP1 mRNA (Fig. 8D, F), suggesting a synergistic effect of ES in  
2 combination with TE to attenuate the expression of these markers for anti-anabolic action.  
3  
4 Consistent with our previous findings [5, 6, 27, 30], an osteoclastogenesis assay using bone  
5 marrow hematopoietic stem cells derived from the unstimulated femurs of SCI animals revealed  
6 a significant increase of +34.8% for TRAP<sup>+</sup> multinucleated cells ( $p < 0.001$ ) when compared to  
7 sham animals (Fig. 9B). Importantly, both TE treatment and TE-ES markedly decreased TRAP<sup>+</sup>  
8 cells by -51.3% ( $p < 0.001$ ) and -46.9% ( $p < 0.001$ ), respectively. Following SCI, quantitative PCR  
9 analysis revealed an increased expression in cultures of bone-marrow-derived osteoclasts of  
10 osteoclast marker genes including TRAP (Fig. 9C), integrin  $\beta 3$  (Fig. 9D) and calcitonin receptor  
11 (CTR, Fig. 9E). ES or ES+TE treatment significantly decreased the mRNA level of these three  
12 biomarkers for bone resorption, whereas TE treatment only decreased TRAP and CTR.  
13



## 1 Discussion

2 The findings reported herein support several conclusions. Under the conditions of the  
3 experiment, ES and TE each improved some bone parameters (BMD at distal femur and  
4 proximal tibia, BV/TV by histomorphometry at the distal femur, BFR/TV by dynamic  
5 histomorphometry, and mechanical strength) and ES increased muscle mass, as might have  
6 been predicted from our prior studies in rats [22] and recent publications in persons with SCI  
7 [21]. While the combination of ES+TE caused numerical increases in BMD at these sites as  
8 compared to either treatment alone, there was no difference in BV/TV by microCT or  
9 histomorphometry between ES and ES+TE groups. Similarly, the combination of ES+TE did not  
10 increase muscle mass beyond that observed with ES alone. CFU-F<sup>+</sup> was reduced after SCI  
11 while numbers of TRAP<sup>+</sup> cells present in bone marrow cultures were increased after SCI. The  
12 pattern of changes in biochemical markers of osteoblast and osteoclast differentiation and  
13 activity generally paralleled the effects of ES and TE on bone mass. For example, ES and TE,  
14 alone or in combination, reduced numbers of TRAP<sup>+</sup> cells and osteoclast markers (TRAP mRNA  
15 and CTR mRNA). There was, however, some discordance between *ex-vivo* biological changes  
16 induced by ES alone or combined with TE as compared to effects of these interventions on  
17 bone parameters. For example, TE but not ES increased the number CFU-F<sup>+</sup> colonies; the  
18 pattern of effects of TE, ES or the combination on expression of Runx2 also suggested greater  
19 effects of TE as compared to ES, although criteria for statistical significance of differences  
20 between these groups were not met. When interpreting these discordant data, it should be  
21 appreciated that, while cell culture systems provide powerful tools that seem to recapitulate  
22 many important aspects of bone biology in disease, these cultures have limitations. Specifically,  
23 the process of isolation, fractionation and culture of cells may alter their biological properties  
24 when compared to their behavior *in vivo*.

25 Taken together, the data reported here do not demonstrate that restoring loading of bone by  
26 ES is synergistic with the bone-sparing activity of TE in a model of spinal cord transection

1 performed in young adult rats. However, the findings do not exclude the possibility of synergy  
2 and there are trends in the data in which SOST and sFRP1 mRNA were lower in SCI+TE+ES  
3 (Fig. 8) while cortical bone outcomes of ES+TE were group higher (Fig. 4) relative to those in  
4 TE or ES alone groups, suggesting that a weak synergy could exist that could reach  
5 significance with larger sample size. The rather modest effect of ES on muscle mass was  
6 surprising given prior studies that demonstrated that the even small amounts of ES are sufficient  
7 to slow or prevent muscle atrophy after SCI in rats [19] or man [20]. In rats, muscle atrophy  
8 progresses rapidly after spinal cord transection; in a prior report, muscles from SCI animals  
9 were 40-60% smaller at 2 weeks compared with sham-transected controls [43]. Perhaps once  
10 atrophy is established, it is more difficult to reverse such atrophy by the application of ES.  
11 Alternatively, different ES paradigms from those used in the current study are required with  
12 respect to intensity, frequency or other parameters to restore, as opposed to prevent, post-SCI  
13 muscle atrophy. The skeleton of older animals is generally assumed to respond less to  
14 mechanical loading than that of younger animals; because 13-month old male rats were  
15 employed in the present study, it may have been anticipated that younger rats may have had a  
16 more robust response to ES. At present time, it is not known why treatment with TE and/or ES  
17 induced more changes in the mass of EDL (mainly fast glycolytic muscle fibers) than in that of  
18 soleus (mainly slow oxidative muscle fibers) after SCI. Future study of muscle contractility,  
19 muscle fiber type and the associated molecular alterations can provide a better understanding  
20 of the changes observed in muscle function.

21 Androgenic steroids, including testosterone and nandrolone, have anabolic effects on bone  
22 [6]. Androgen ablation therapy for prostate cancer results in decreased bone mass [7]. In men  
23 treated chronically with glucocorticoids, testosterone increased lumbar spine BMD [8]. In women  
24 with osteoporosis, nandrolone increased BMD of the lumbar spine and femoral neck and  
25 reduced fracture risk [9]. Recently, a number of publications from several groups, including ours,  
26 suggest that the reductions in circulating levels of testosterone, such as those that frequently

1 occur in men after SCI [10, 11], may accelerate SCI-related bone loss [12] and also suggest that  
2 androgens may reduce such bone loss [13-15], which perhaps is also the case in animals that  
3 are hypogonadal secondary to SCI [14]. Therefore, it is not surprising that although these rats  
4 with SCI had intact testes in the present study, testosterone administration has provided an  
5 anabolic effect on bone. Estrogen has an important role in muscle and bone physiology. As  
6 such, it would of interest to know if DHT, a form of testosterone which is not aromatizable, would  
7 have the same effects as the TE. Recent findings have demonstrated that DHT induced  
8 differentiation of a preosteoblast cell line in culture, and this action of DHT required Wnt  
9 signaling [16]. Similarly, our recent laboratory observation indicates that DHT inhibits expression  
10 of SOST mRNA (a key factor for controlling bone remodeling) in cultured osteocytic cells (data  
11 not shown). Therefore, we would expect that DHT would have a similar effect to that of TE on  
12 bone after SCI. TE was chosen as the hormonal intervention in the present study for several  
13 reasons. Testosterone replacement therapy is currently a therapy for older hypogonadal men  
14 [17] and for men who have loss of testosterone due to surgery [7]. Testosterone has been  
15 demonstrated to prevent loss of cancellous bone following SCI in a rat model [13-15], and it has  
16 shown a better safety profile on heart than DHT, although DHT can mimic most of the beneficial  
17 effects of testosterone.

18 Effects of ES on bone were more encouraging, and extend a prior report using similar  
19 stimulators, ES parameters and surgical models of SCI, including ablation of the insertion of the  
20 gastrocnemius into the Achilles tendon [5]. In that report, ES reduced numbers of TRAP+  
21 colonies in cultures of osteoclast precursors associated with downregulation of mRNA  
22 transcripts for osteoclast differentiation and activation genes, together with many favorable  
23 changes in mRNA levels in cultures of marrow stromal cells under osteoblastogenic conditions;  
24 these alterations included lower SOST and higher Wnt3a and osteoprotegrin mRNA levels. In  
25 the study reported here, four weeks of ES resulted in increases in multiple measures of  
26 metaphyseal trabecular bone mass and higher BFR/TV while lowering osteoclast number in

1 bone tissues and almost complete restoration of cortical mechanical strength, as predicted  
2 using FE modeling in the SCI animals at 6 weeks (a subacute phase of injury), suggesting a net  
3 anabolic effect on bone. Moreover, the mRNA ratio of RANKL to OPG in sublesional long bone  
4 was greatly reduced after the administration of ES, TE or ES+TE, suggesting that each  
5 intervention, or the combination, and the consequent reduction in bone resorption, at least in  
6 part, contributes to mitigate bone loss after SCI. One rather striking, and unexpected, finding is  
7 that osseous effects of ES appeared to be more prominent than effects of ES on our muscle  
8 endpoints. Whether this reflects that the fact that bone loss progresses more slowly than muscle  
9 atrophy in this model system, such that bone is better able to respond to restoration of  
10 mechanical inputs, remains unclear.

11 As stated earlier, the reductions in circulating levels of testosterone, such as those that  
12 frequently occur in men after SCI [48, 49], may accelerate SCI-related bone loss [51] and also  
13 suggest that androgens may reduce such bone loss. In this study, we found that SCI+TE group  
14 had higher MS/BS and lower osteoclast numbers in bone sections, coincidental with more CFU-  
15 F<sup>+</sup> cells and higher Runx2 mRNA (meaning increased osteoblastogenesis) and lower TRAP<sup>+</sup>  
16 cells (meaning the reduced osteoclastogenesis) in bone marrow cell cultures, These changes  
17 are tightly associated with higher trabecular bone measures at distal femur. Our data confirm  
18 prior reports, including a recent study in which administration of TE spared bone and muscle  
19 following severe spinal cord contusion in rats [12], further suggesting the musculoskeletal  
20 benefits of TE administration. It should be noted that in the latter study, TE treatment was  
21 started immediately after SCI, rather than 2 weeks after injury as was performed in the current  
22 study. Thus, the smaller effect of TE on muscle and bone in the present study may to some  
23 degree relate to the timing of the delay in the initiation of treatment. For example, TE may be  
24 more effective in preventing muscle atrophy or bone resorption than its action to promote gains  
25 in muscle or bone mass once losses have occurred. It is interesting to note that TE suppressed

1 MuRf1 and sFRP RNA in EDL and not MAFbx RNA. Future study will determine what it might  
2 imply about TE and how these factors may influence TE-mediated effects on muscle loss.

3 The myokine irisin is released from skeletal muscle upon exercise and contributes to  
4 certain favorable effects of physical activity. The roles of irisin in bone biology remains unsettled.  
5 Recent studies have demonstrated that administration of irisin can affect skeletal remodeling.  
6 For example, very low-dose irisin injections, administered intermittently, have been shown to  
7 improve cortical bone mineral density and strength in mice (Colaianni *et al.*, 2015, 2017; [2, 3] ).  
8 These bone effects were consistent with *in vitro* studies showing that irisin could promote  
9 osteoblast differentiation (Qiao *et al.*, 2016; [4]). However, Bonewald's group and her  
10 collaborators recently reported that irisin increases sclerostin expression in osteocytes to induce  
11 bone resorption (Kim *et al.*, 2018; [5]). In the present study we observed that levels of irisin are  
12 decreased after each treatment performed, which is associated with a reduction in bone  
13 resorption and an increase in bone mass. This observation is consistent with the possibility that  
14 inhibition of irisin can reduce bone resorption and bone loss, whereas the induction of irisin can  
15 increase bone resorption and reduce bone mass, which appears to be in line with the studies by  
16 Kim *et al.* [5]. Similarly, there is lack of any consistent pattern observed with regard to changes  
17 of serum myostatin (another myokine) levels among the experimental groups (data not shown),  
18 which might be partially due to the generally appreciated cross-reactivity of myostatin with  
19 several members of the transforming growth factor beta (TGF-beta) family using commercial  
20 kits. Therefore, our present study is not able to state with any degree of confidence the role of  
21 either irisin or myostatin in the context of physical activity after SCI. Additional work is required  
22 to address the level and role of these two myokines or other soluble factors in improving our  
23 understanding of the ES-mediated beneficial effects on bone.

24 For future perspective, further study is needed to fully appreciate the potential of ES or ES-  
25 based combinatorial approaches to preserve or restore bone more chronically after SCI for an  
26 extended period of time with substantial sublesional bone loss, which represents the vast

1 majority of the SCI population. Our initial analysis suggested the unexpected benefits on  
2 accrual of muscle and bone mass of the unstimulated (contralateral) right hindlimb that was  
3 observed in the absence of muscle contraction and bone reloading by ES (data not shown),  
4 leading to speculate that soluble factors that was released from muscle and/or bone upon ES-  
5 induced muscle contraction and bone reloading may be transported to the unstimulated  
6 hindlimb and, thereby, exert systemic musculoskeletal benefits. Future studies are warranted to  
7 further examine systemic alterations in bone homeostasis and remodeling and define the  
8 potential role of soluble mediators originating in skeletal muscle and/or bone that are  
9 responsible for humoral muscle-bone interactions in response to muscle contraction by ES.  
10 Additionally, the improvement of bone parameters in long bones (even contralateral long bones)  
11 but not the axial skeleton (L3-L5) is an important observation. It is unclear why the discrepancy  
12 was observed between the axial and appendicular skeleton, which remains a provocative topic  
13 to address in future investigation.

14

## 15 **Acknowledgments**

16 Sources of Research Support: This work was supported by the Veterans Health Administration,  
17 Rehabilitation Research and Development Service (grants 5I01RX000687, 5I01RX001313 and  
18 5I01RX02089-A2 to WQ; B9212-C and B2020-C to WAB), NIH R21 NS111393-01A1 to WQ,  
19 and the U.S. Department of Agriculture, Agricultural Research Service project plan #3062-  
20 51000-053-00D. USDA is an equal opportunity provider and employer. Mention of trade names  
21 or commercial products in this publication is solely for the purpose of providing specific  
22 information and does not imply recommendation or endorsement by the U.S. Department of  
23 Agriculture. The findings and conclusions in this manuscript are those of the authors and should  
24 not be construed to represent any official USDA of U.S. Government determination or policy.

25

1 **Author Contributions**

2 WQ and WZ were responsible for study design and data analysis. WZ, YP, YH, EG, JL, JC, JP,  
3 JQF, JJ and WAB provided consultation on and performed the experimental procedures to  
4 generate the data as well as data interpretation for the study. The manuscript was written by  
5 WQ, WZ, and CC and was revised and approved by all authors. WQ takes responsibility for the  
6 integrity of the data analysis.

7

8 **Author Disclosure Statement.**

9 No competing financial interests exist.

10

11

1 **Figure Legends**

2

3 **Figure 1. ES or ES+TE combination treatment reduced muscle loss after SCI.** (A-C) ES  
4 on muscle weight: (A) soleus muscle weight, (B) plantaris muscle weight, and (C) EDL muscle  
5 weight. (D-G) Effect of ES on the expression of muscle markers by measurement of changes in  
6 mRNA levels of EDL muscle genes by real-time PCR: (D) MAFbx, (E) MurF1, (F) sFRP1, and  
7 (G) sFRP2. Data are expressed as mean  $\pm$  SE. N=10-13 per group. \*p<0.05, \*\*P<0.01,  
8 \*\*\*P<0.001 by one-way ANOVA.

9

10 **Figure 2. ES or ES+TE treatment reduced bone loss after SCI.** (A-C) Areal BMD  
11 measurements after complete spinal cord transection (T3-T4) at the: (A) distal femur, (B)  
12 proximal tibia, and (C) spine (L3-5). Sham-operated animals were used as controls. Data are  
13 expressed as mean  $\pm$  SE. N = 10-13 per group. Significance of differences was determined  
14 using one-way analysis of variance with a Newman–Keuls post hoc test. \*P < 0.05, \*\*P < 0.01,  
15 \*\*\*P<0.001 versus the indicated group.

16

17 **Figure 3. Effects of ES or ES+TE on trabecular architecture of the distal femur.** (A-F)  
18 Parameters of trabecular architecture: (A) representative  $\mu$ CT 3D images of trabecular  
19 microarchitecture; (B) trabecular bone volume per total volume (BV/TV%), (C) trabecular  
20 number (Tb.N), (D) trabecular thickness (Tb.Th), (E) trabecular space (Tb.Sp), (F) connectivity  
21 density (Conn.D), and (G) structure model index (SMI). Data are expressed as mean  $\pm$  SE. N =  
22 6 – 8 per group. Significance of differences was determined by using one-way ANOVA with a  
23 Newman–Keuls post hoc test. \*P < 0.05, \*\*P < 0.01 and \*\*\*P < 0.001 versus the indicated  
24 group.

25



1 **Figure 4. Effects of ES+TE on cortical architecture and strength of the femur midshaft.** (A-  
2 D) Parameters of cortical architecture: (A) representative  $\mu$ CT 3D images of cortical  
3 microarchitecture; (B) cortical thickness; (C) total tissue area; (D) cortical bone area; (E)  
4 periosteal perimeter, and (F) endosteal perimeter; Parameter of cortical strength: (G) stiffness.  
5 Data are expressed as mean  $\pm$  SEM. N =6– 8 per group. Significance of differences was  
6 determined by using one-way ANOVA with Newman–Keuls post hoc test. \*P < 0.05,  
7 \*\*\*P<0.001.

8  
9 **Figure 5. Effects of ES or ES+TE on levels of serum biomarkers of bone and RANKL/OPG**  
10 **mRNA ratio in long bone.** (A-C) ELISA tests for: (A) osteocalcin, (B) CTX, (C) irisin and (D)  
11 Using RNA extracted from sublesional long bones, RANKL and OPG gene expressions were  
12 determined by real-time PCR analysis and the ratio of RANKL/OPG was calculated. Data are  
13 expressed as mean  $\pm$  SE, n=10-13 per group. \*p<0.05, \*\*p<0.01, \*\*\*p<0.001 by one-way  
14 ANOVA with Newman–Keuls post hoc test; #p<0.05, ##p<0.01 by two-tailed t-test.

15  
16 **Figure 6. Effects of ES or ES+TE on formation of trabecular bone at the distal femur.** (A)  
17 Representative images of 6- $\mu$ m-thick bone specimens showing double labeling of calcein and  
18 xylene orange under fluorescence microscopy (magnification  $\times$ 20). (B-G) Parameters of bone  
19 formation, volume and perimeter: (B) bone formation rate over tissue volume (BFR/TV), (C)  
20 mineralizing surface over bone surface (MS/BS), (D) mineral apposition rate (MAR), (E) bone  
21 formation rate over bone surface (BFR/BS), (F) bone volume over tissue volume (BV/TV), and  
22 (G) bone perimeter (B.pm). Data are expressed as mean  $\pm$  SE, \*\*p < 0.01, \*\*\*p < 0.001 versus  
23 the indicated group by one-way ANOVA plus Newman–Keuls post hoc test, n = 6-7 animals per  
24 group.

25

1 **Figure 7. Effects of ES or ES+TE on bone resorption of trabecular bone at the distal**  
2 **femur.** (A) Representative sections of trabecular bone from the femoral metaphysis  
3 immunostained for TRAP (20x). The reddish areas of TRAP staining on trabecular surfaces  
4 representing osteoclasts. (B–C) Parameters of trabecular bone resorption by histomorphometric  
5 quantification: (B) osteoclast surface over bone surface (Oc.S/BS) and (C) osteoclast number  
6 over bone perimeter (N.Oc/B. Pm). Data are expressed as mean  $\pm$  SE. N= 6-7 per group.  
7 \*p<0.05 versus the indicated group by one-way ANOVA plus Newman–Keuls post hoc test.

8  
9 **Figure 8. Effects of ES or ES+TE treatment on osteoblastic differentiation.** (A-B)  
10 Representative images of bone formation: (A) alkaline phosphatase staining (CFU-F) of cultured  
11 bone marrow stromal cells, and (B) quantification of CFU-F<sup>+</sup> cells. (C-F) Changes in the gene  
12 expression of bone formation markers in cultured bone marrow stromal cells by quantitative  
13 PCR: (C) Runx2, (D) sclerostin (SOST), (E) LRP5, and (F) sFRP1. \*p<0.05, \*\*\*p<0.001 by one-  
14 way ANOVA plus Newman–Keuls post hoc test, n=6-8 per group.

15  
16 **Figure 9. ES or ES+TE inhibited the osteoclastogenic potential of bone marrow**  
17 **hematopoietic cells.** (A) TRAP staining of cultured bone marrow hematopoietic cells, and (B)  
18 quantification of TRAP<sup>+</sup> cells. (C-E) Changes in the gene expression of bone resorption markers  
19 in cultured bone marrow hematopoietic cells determined by quantitative PCR: (C) TRAP, (D)  
20 integrin  $\beta$ 3, and (E) calcitonin receptor (CTR). \*p<0.05, \*\*p<0.01, \*\*\*p<0.001 by one-way  
21 ANOVA plus Newman–Keuls post hoc test, n=6-8 per group.

22  
23

## 1 References

- 2 1. Qin, W., W.A. Bauman, and C. Cardozo (2010). Bone and muscle  
3 loss after spinal cord injury: organ interactions. *Ann N Y*  
4 *Acad Sci* 1211 (1), 66-84.
- 5 2. Bauman, W.A. and C.P. Cardozo (2014). Osteoporosis in  
6 Individuals with Spinal Cord Injury. *PM R*.
- 7 3. Gorgey, A.S., O. Witt, L. O'Brien, C. Cardozo, Q. Chen,  
8 E.J. Lesnefsky, and Z.A. Graham (2019). Mitochondrial health  
9 and muscle plasticity after spinal cord injury. *Eur J Appl*  
10 *Physiol* 119 (2), 315-331.
- 11 4. Peng, Y., W. Zhao, Y. Hu, F. Li, X.E. Guo, D. Wang, W.A.  
12 Bauman, and W. Qin (2019). Rapid bone loss occurs as early  
13 as 2 days after complete spinal cord transection in young  
14 adult rats. *Spinal Cord*.
- 15 5. Qin, W., L. Sun, J. Cao, Y. Peng, L. Collier, Y. Wu, G.  
16 Creasey, J. Li, Y. Qin, J. Jarvis, W.A. Bauman, M. Zaidi,  
17 and C. Cardozo (2013). The Central Nervous System (CNS)-  
18 independent Anti-bone-resorptive Activity of Muscle  
19 Contraction and the Underlying Molecular and Cellular  
20 Signatures. *J Biol Chem* 288 (19), 13511-21.
- 21 6. Qin, W., X. Li, Y. Peng, L.M. Harlow, Y. Ren, Y. Wu, J. Li,  
22 Y. Qin, J. Sun, S. Zheng, T. Brown, J.Q. Feng, H.Z. Ke,  
23 W.A. Bauman, and C.C. Cardozo (2015). Sclerostin antibody  
24 preserves the morphology and structure of osteocytes and  
25 blocks the severe skeletal deterioration after motor-  
26 complete spinal cord injury in rats. *J Bone Miner Res*.
- 27 7. Otzel, D.M., C.F. Conover, F. Ye, E.G. Phillips, T.  
28 Bassett, R.D. Wnek, M. Flores, A. Catter, P. Ghosh, A.  
29 Balaez, J. Petusevsky, C. Chen, Y. Gao, Y. Zhang, J.M.  
30 Jiron, P.K. Bose, S.E. Borst, T.J. Wronski, J.I. Aguirre,  
31 and J.F. Yarrow (2019). Longitudinal Examination of Bone  
32 Loss in Male Rats After Moderate-Severe Contusion Spinal  
33 Cord Injury. *Calcif Tissue Int* 104 (1), 79-91.
- 34 8. Jiang, S.D., L.S. Jiang, and L.Y. Dai (2007). Changes in  
35 bone mass, bone structure, bone biomechanical properties,  
36 and bone metabolism after spinal cord injury: a 6-month  
37 longitudinal study in growing rats. *Calcif Tissue Int* 80  
38 (3), 167-75.
- 39 9. Eser, P., A. Frotzler, Y. Zehnder, L. Wick, H. Knecht, J.  
40 Denoth, and H. Schiessl (2004). Relationship between the  
41 duration of paralysis and bone structure: a pQCT study of  
42 spinal cord injured individuals. *Bone* 34 (5), 869-80.
- 43 10. Harlow, L., K. Sahbani, J.S. Nyman, C.P. Cardozo, W.A.  
44 Bauman, and H.A. Tawfeek (2017). Daily parathyroid hormone  
45 administration enhances bone turnover and preserves bone

- 1 structure after severe immobilization-induced bone loss.  
2 *Physiol Rep* 5 (18).
- 3 11. Zeman, R.J., J. Zhao, Y. Zhang, W. Zhao, X. Wen, Y. Wu, J.  
4 Pan, W.A. Bauman, and C. Cardozo (2009). Differential  
5 skeletal muscle gene expression after upper or lower motor  
6 neuron transection. *Pflugers Arch* 458 (3), 525-35.
- 7 12. Yarrow, J.F., C.F. Conover, L.A. Beggs, D.T. Beck, D.M.  
8 Otzel, A. Balaez, S.M. Combs, J.R. Miller, F. Ye, J.I.  
9 Aguirre, K.G. Neuville, A.A. Williams, B.P. Conrad, C.M.  
10 Gregory, T.J. Wronski, P.K. Bose, and S.E. Borst  
11 (2013). Testosterone dose-dependently prevents bone and  
12 muscle loss in rodents following spinal cord injury. *J*  
13 *Neurotrauma*.
- 14 13. Zhao, W., J. Pan, X. Wang, Y. Wu, W.A. Bauman, and C.P.  
15 Cardozo (2008). Expression of the muscle atrophy factor  
16 muscle atrophy F-box is suppressed by testosterone.  
17 *Endocrinology* 149 (11), 5449-60.
- 18 14. Zhao, W., J. Pan, X. Wang, Y. Wu, W.A. Bauman, and C.P.  
19 Cardozo (2008). Expression of the muscle atrophy factor MAFbx  
20 is suppressed by testosterone. *Endocrinology* 149 (11),  
21 5449-5460.
- 22 15. Qin, W., J. Pan, Y. Wu, W.A. Bauman, and C. Cardozo  
23 (2010). Protection against dexamethasone-induced muscle  
24 atrophy is related to modulation by testosterone of FOXO1  
25 and PGC-1alpha. *Biochem Biophys Res Commun* 403 (3-4),  
26 473-8.
- 27 16. Wu, Y., W.A. Bauman, R.D. Blitzer, and C. Cardozo  
28 (2010). Testosterone-induced hypertrophy of L6 myoblasts is  
29 dependent upon Erk and mTOR. *Biochem Biophys Res Commun*  
30 400 (4), 679-83.
- 31 17. Notini, A.J., J.F. McManus, A. Moore, M. Bouxsein, M.  
32 Jimenez, W.S. Chiu, V. Glatt, B.E. Kream, D.J. Handelsman,  
33 H.A. Morris, J.D. Zajac, and R.A. Davey (2007). Osteoblast  
34 deletion of exon 3 of the androgen receptor gene results in  
35 trabecular bone loss in adult male mice. *J Bone Miner Res*  
36 22 (3), 347-56.
- 37 18. Rooney, A.M. and M.C.H. van der Meulen (2017). Mouse models  
38 to evaluate the role of estrogen receptor alpha in skeletal  
39 maintenance and adaptation. *Ann N Y Acad Sci* 1410 (1),  
40 85-92.
- 41 19. Kim, S.J., R.R. Roy, J.A. Kim, H. Zhong, F. Haddad, K.M.  
42 Baldwin, and V.R. Edgerton (2008). Gene expression during  
43 inactivity-induced muscle atrophy: effects of brief bouts  
44 of a forceful contraction countermeasure. *J Appl Physiol*  
45 (1985) 105 (4), 1246-54.
- 46 20. Baldi, J.C., R.D. Jackson, R. Moraille, and W.J. Mysiw  
47 (1998). Muscle atrophy is prevented in patients with acute

1 spinal cord injury using functional electrical stimulation.  
2 Spinal Cord 36 (7), 463-9.

3 21. Gorgey, A.S., R.E. Khalil, R. Gill, D.R. Gater, T.D. Lavis,  
4 C.P. Cardozo, and R.A. Adler (2019).Low-Dose Testosterone  
5 and Evoked Resistance Exercise after Spinal Cord Injury on  
6 Cardio-Metabolic Risk Factors: An Open-Label Randomized  
7 Clinical Trial. J Neurotrauma 36 (18), 2631-2645.

8 22. Wu, Y., L. Collier, W. Qin, G. Creasey, W.A. Bauman, J.  
9 Jarvis, and C. Cardozo (2013).Electrical stimulation  
10 modulates Wnt signaling and regulates genes for the motor  
11 endplate and calcium binding in muscle of rats with spinal  
12 cord transection. BMC Neurosci 14 (1), 81.

13 23. How old is a rat in human years?  
14 <http://www.ratbehavior.org/RatYears.htm>.

15 24. Martin, R., C. Sadowsky, K. Obst, B. Meyer, and J. McDonald  
16 (2012).Functional electrical stimulation in spinal cord  
17 injury:: from theory to practice. Top Spinal Cord Inj  
18 Rehabil 18 (1), 28-33.

19 25. Wu, Y., L. Collier, W. Qin, G. Creasey, W.A. Bauman, J.  
20 Jarvis, and C. Cardozo (2013).Electrical stimulation  
21 modulates Wnt signaling and regulates genes for the motor  
22 endplate and calcium binding in muscle of rats with spinal  
23 cord transection. BMC Neurosci 14 81.

24 26. Qin, W., X. Li, Y. Peng, L.M. Harlow, Y. Ren, Y. Wu, J. Li,  
25 Y. Qin, J. Sun, S. Zheng, T. Brown, J.Q. Feng, H.Z. Ke,  
26 W.A. Bauman, and C.P. Cardozo (2016).Sclerostin Antibody  
27 Preserves the Morphology and Structure of Osteocytes and  
28 Blocks the Severe Skeletal Deterioration After Motor-  
29 Complete Spinal Cord Injury in Rats. J Bone Miner Res 31  
30 (7), 1482.

31 27. Sun, L., J. Pan, Y. Peng, Y. Wu, J. Li, X. Liu, Y. Qin,  
32 W.A. Bauman, C. Cardozo, M. Zaidi, and W. Qin  
33 (2013).Anabolic steroids reduce spinal cord injury-related  
34 bone loss in rats associated with increased Wnt signaling.  
35 J Spinal Cord Med 36 (6), 616-22.

36 28. Zhao, W., X. Li, Y. Peng, Y. Qin, J. Pan, J. Li, A. Xu,  
37 M.S. Ominsky, C. Cardozo, J.Q. Feng, H.Z. Ke, W.A. Bauman,  
38 and W. Qin (2018).Sclerostin Antibody Reverses the Severe  
39 Sublesional Bone Loss in Rats After Chronic Spinal Cord  
40 Injury. Calcif Tissue Int 103 (4), 443-454.

41 29. Cardozo, C.P., W. Qin, Y. Peng, X. Liu, Y. Wu, J. Pan, W.A.  
42 Bauman, M. Zaidi, and L. Sun (2010).Nandrolone slows  
43 hindlimb bone loss in a rat model of bone loss due to  
44 denervation. Ann N Y Acad Sci 1192 303-6.

45 30. Bramlett, H.M., W.D. Dietrich, A. Marcillo, L.J. Mawhinney,  
46 O. Furones-Alonso, A. Bregy, Y. Peng, Y. Wu, J. Pan, J.  
47 Wang, X.E. Guo, W.A. Bauman, C. Cardozo, and W. Qin

1 (2014).Effects of low intensity vibration on bone and  
2 muscle in rats with spinal cord injury. *Osteoporos Int* 25  
3 (9), 2209-19.

4 31. Qin, W., X. Li, Y. Peng, L.M. Harlow, Y. Ren, Y. Wu, J. Li,  
5 Y. Qin, J. Sun, S. Zheng, T. Brown, J.Q. Feng, H.Z. Ke,  
6 W.A. Bauman, and C.C. Cardozo (2015).Sclerostin antibody  
7 preserves the morphology and structure of osteocytes and  
8 blocks the severe skeletal deterioration after motor-  
9 complete spinal cord injury in rats. *J Bone Miner Res* 30  
10 (11), 1994-2004.

11 32. Bouxsein, M.L., S.K. Boyd, B.A. Christiansen, R.E.  
12 Guldberg, K.J. Jepsen, and R. Muller (2010).Guidelines for  
13 assessment of bone microstructure in rodents using micro-  
14 computed tomography. *J Bone Miner Res* 25 (7), 1468-86.

15 33. Zysset, P.K., E. Dall'ara, P. Varga, and D.H. Pahr  
16 (2013).Finite element analysis for prediction of bone  
17 strength. *Bonekey Rep* 2 386.

18 34. Lin, T., W. Tong, A. Chandra, S.Y. Hsu, H. Jia, J. Zhu,  
19 W.J. Tseng, M.A. Levine, Y. Zhang, S.G. Yan, X.S. Liu, D.  
20 Sun, W. Young, and L. Qin (2015).A comprehensive study of  
21 long-term skeletal changes after spinal cord injury in  
22 adult rats. *Bone Res* 3 15028.

23 35. Pistoia, W., B. van Rietbergen, E.M. Lochmuller, C.A. Lill,  
24 F. Eckstein, and P. Ruegsegger (2002).Estimation of distal  
25 radius failure load with micro-finite element analysis  
26 models based on three-dimensional peripheral quantitative  
27 computed tomography images. *Bone* 30 (6), 842-8.

28 36. Lan, S., S. Luo, B.K. Huh, A. Chandra, A.R. Altman, L. Qin,  
29 and X.S. Liu (2013).3D image registration is critical to  
30 ensure accurate detection of longitudinal changes in  
31 trabecular bone density, microstructure, and stiffness  
32 measurements in rat tibiae by in vivo microcomputed  
33 tomography (muCT). *Bone* 56 (1), 83-90.

34 37. Liu, X.S., X.H. Zhang, K.K. Sekhon, M.F. Adams, D.J.  
35 McMahon, J.P. Bilezikian, E. Shane, and X.E. Guo  
36 (2010).High-resolution peripheral quantitative computed  
37 tomography can assess microstructural and mechanical  
38 properties of human distal tibial bone. *J Bone Miner Res*  
39 25 (4), 746-56.

40 38. Kitase, Y., J.A. Vallejo, W. Gutheil, H. Vemula, K. Jahn,  
41 J. Yi, J. Zhou, M. Brotto, and L.F. Bonewald (2018).beta-  
42 aminoisobutyric Acid, l-BAIBA, Is a Muscle-Derived  
43 Osteocyte Survival Factor. *Cell Rep* 22 (6), 1531-1544.

44 39. Qin, W., W. Zhao, X. Li, Y. Peng, L.M. Harlow, J. Li, Y.  
45 Qin, J. Pan, Y. Wu, L. Ran, H.Z. Ke, C.P. Cardozo, and W.A.  
46 Bauman (2016).Mice with sclerostin gene deletion are

- 1 resistant to the severe sublesional bone loss induced by  
2 spinal cord injury. *Osteoporos Int*.
- 3 40. Livak, K.J. and T.D. Schmittgen (2001). Analysis of relative  
4 gene expression data using real-time quantitative PCR and  
5 the 2(-Delta Delta C(T)) Method. *Methods* 25 (4), 402-8.
- 6 41. Qin, W., W.A. Bauman, and C. Cardozo (2010). Bone and muscle  
7 loss after spinal cord injury: organ interactions. *Ann N Y*  
8 *Acad Sci* 1211 66-84.
- 9 42. Colaianni, G., C. Cuscito, T. Mongelli, P. Pignataro, C.  
10 Buccoliero, P. Liu, P. Lu, L. Sartini, M. Di Comite, G.  
11 Mori, A. Di Benedetto, G. Brunetti, T. Yuen, L. Sun, J.E.  
12 Reseland, S. Colucci, M.I. New, M. Zaidi, S. Cinti, and M.  
13 Grano (2015). The myokine irisin increases cortical bone  
14 mass. *Proc Natl Acad Sci U S A* 112 (39), 12157-62.
- 15 43. Zeman, R.J., W.A. Bauman, X. Wen, N. Ouyang, J.D. Etlinger,  
16 and C.P. Cardozo (2009). Improved functional recovery with  
17 oxandrolone after spinal cord injury in rats. *Neuroreport*  
18 20 (9), 864-8.
- 19 44. Clarke, B.L. and S. Khosla (2009). Androgens and bone.  
20 *Steroids* 74 (3), 296-305.
- 21 45. Schwandt, A. and J.A. Garcia (2009). Complications of  
22 androgen deprivation therapy in prostate cancer. *Curr Opin*  
23 *Urol* 19 (3), 322-6.
- 24 46. Crawford, B.A., P.Y. Liu, M.T. Kean, J.F. Bleasel, and D.J.  
25 Handelsman (2003). Randomized placebo-controlled trial of  
26 androgen effects on muscle and bone in men requiring long-  
27 term systemic glucocorticoid treatment. *J Clin Endocrinol*  
28 *Metab* 88 (7), 3167-76.
- 29 47. Frisoli, A., Jr., P.H. Chaves, M.M. Pinheiro, and V.L.  
30 Szejnfeld (2005). The effect of nandrolone decanoate on bone  
31 mineral density, muscle mass, and hemoglobin levels in  
32 elderly women with osteoporosis: a double-blind,  
33 randomized, placebo-controlled clinical trial. *J Gerontol A*  
34 *Biol Sci Med Sci* 60 (5), 648-53.
- 35 48. Tsitouras, P.D., Y.G. Zhong, A.M. Spungen, and W.A. Bauman  
36 (1995). Serum testosterone and growth hormone/insulin-like  
37 growth factor-I in adults with spinal cord injury. *Horm*  
38 *Metab Res* 27 (6), 287-92.
- 39 49. Kostovski, E., P.O. Iversen, K. Birkeland, P.A. Torjesen,  
40 and N. Hjeltnes (2008). Decreased levels of testosterone and  
41 gonadotrophins in men with long-standing tetraplegia.  
42 *Spinal Cord* 46 (8), 559-64.
- 43 50. Bauman, W.A., M.F. La Fountaine, and A.M. Spungen  
44 (2014). Age-related prevalence of low testosterone in men  
45 with spinal cord injury. *J Spinal Cord Med* 37 (1), 32-9.

- 1 51. Qin, W., W.A. Bauman, and C.P. Cardozo (2010).Evolving  
2 concepts in neurogenic osteoporosis. *Curr Osteoporos Rep* 8  
3 (4), 212-8.
- 4 52. Beggs, L.A., F. Ye, P. Ghosh, D.T. Beck, C.F. Conover, A.  
5 Balaez, J.R. Miller, E.G. Phillips, N. Zheng, A.A.  
6 Williams, J.I. Aguirre, T.J. Wronski, P.K. Bose, S.E.  
7 Borst, and J.F. Yarrow (2015).Sclerostin inhibition  
8 prevents spinal cord injury-induced cancellous bone loss. *J*  
9 *Bone Miner Res* 30 (4), 681-9.
- 10 53. Yarrow, J.F., C.F. Conover, L.A. Beggs, D.T. Beck, D.M.  
11 Otzel, A. Balaez, S.M. Combs, J.R. Miller, F. Ye, J.I.  
12 Aguirre, K.G. Neuville, A.A. Williams, B.P. Conrad, C.M.  
13 Gregory, T.J. Wronski, P.K. Bose, and S.E. Borst  
14 (2014).Testosterone dose dependently prevents bone and  
15 muscle loss in rodents after spinal cord injury. *J*  
16 *Neurotrauma* 31 (9), 834-45.
- 17 54. Liu, X.H., A. Kirschenbaum, S. Yao, and A.C. Levine  
18 (2007).Androgens promote preosteoblast differentiation via  
19 activation of the canonical Wnt signaling pathway. *Ann N Y*  
20 *Acad Sci* 1116 423-31.
- 21 55. Kenny, A.M., A. Kleppinger, K. Annis, M. Rathier, B.  
22 Browner, J.O. Judge, and D. McGee (2010).Effects of  
23 transdermal testosterone on bone and muscle in older men  
24 with low bioavailable testosterone levels, low bone mass,  
25 and physical frailty. *J Am Geriatr Soc* 58 (6), 1134-43.
- 26 56. Rubio-Gayosso, I., I. Ramirez-Sanchez, I. Ita-Islas, P.  
27 Ortiz-Vilchis, G. Gutierrez-Salmean, A. Meaney, I. Palma,  
28 I. Olivares, R. Garcia, E. Meaney, and G. Ceballos  
29 (2013).Testosterone metabolites mediate its effects on  
30 myocardial damage induced by ischemia/reperfusion in male  
31 Wistar rats. *Steroids* 78 (3), 362-9.
- 32 57. Colaianni, G., T. Mongelli, C. Cuscito, P. Pignataro, L.  
33 Lippo, G. Spiro, A. Notarnicola, I. Severi, G. Passeri, G.  
34 Mori, G. Brunetti, B. Moretti, U. Tarantino, S.C. Colucci,  
35 J.E. Reseland, R. Vettor, S. Cinti, and M. Grano  
36 (2017).Irisin prevents and restores bone loss and muscle  
37 atrophy in hind-limb suspended mice. *Sci Rep* 7 (1), 2811.
- 38 58. Qiao, X., Y. Nie, Y. Ma, Y. Chen, R. Cheng, W. Yin, Y. Hu,  
39 W. Xu, and L. Xu (2016).Irisin promotes osteoblast  
40 proliferation and differentiation via activating the MAP  
41 kinase signaling pathways. *Sci Rep* 6 18732.
- 42 59. Kim, H., C.D. Wrann, M. Jedrychowski, S. Vidoni, Y. Kitase,  
43 K. Nagano, C. Zhou, J. Chou, V.A. Parkman, S.J. Novick,  
44 T.S. Strutzenberg, B.D. Pascal, P.T. Le, D.J. Brooks, A.M.  
45 Roche, K.K. Gerber, L. Mattheis, W. Chen, H. Tu, M.L.  
46 Bouxsein, P.R. Griffin, R. Baron, C.J. Rosen, L.F.  
47 Bonewald, and B.M. Spiegelman (2018).Irisin Mediates



1 Effects on Bone and Fat via alphaV Integrin Receptors. Cell  
2 175 (7), 1756-1768 e17.  
3