

# 神経回路再構築時における脊髄内及び脊髄神経内の 接着分子の役割：接着分子E型カドヘリンとカドヘ リン結合蛋白質カテニンに注目して

著者	長谷川 光広
著者別表示	Hasegawa Mitsuhiro
雑誌名	平成8(1996)年度 科学研究費補助金 基盤研究(C) 研究成果報告書概要
巻	1995 1996
ページ	2p.
発行年	1999-03-08
URL	<a href="http://doi.org/10.24517/00066290">http://doi.org/10.24517/00066290</a>



# 1996 Fiscal Year Final Research Report Summary

---

The role of cell adhesion molecules in reconstruction of neural network in spinal cord and spinal nerves with special attention paid to E-cadherin and its associated protein catenin

Research Project

## Project/Area Number

---

07671505

## Research Category

---

Grant-in-Aid for Scientific Research (C)

## Allocation Type

---

Single-year Grants

## Section

---

一般

## Research Field

---

Cerebral neurosurgery

## Research Institution

---

Kanazawa University

## Principal Investigator

---

**HASEGAWA Mitsuhiro** Kanazawa University Department of Neurosurgery Assistant Professor, 医学部附属病院, 講師 (70218460)

## Co-Investigator(Kenkyū-buntansha)

---

**TAKEICHI Masatoshi** Kyoto University Department of Biophysics Faculty of Science Professor, 理学部, 教授 (00025454)

## Project Period (FY)

---

1995 – 1996

## Keywords

---

## Research Abstract

An increasing number of cell adhesive proteins is being identified as involved in cell-cell and cell-extracellular matrix interactions. Extensive studies have focused on several members of molecules including L1, N-CAM, MAG during the development and regeneration of the nervous system. We have recently found that Ca<sup>2+</sup>-dependent cell adhesion molecule E-cadherin plays an important role in the selective fasciculation of a particular subset of unmyelinated sensory fibers and that E-cadherin is exclusively expressed in lamina II of Rexed in the spinal cord dorsal horn. In this study, the temporal profile of the cellular and subcellular expression of E-cadherin and cadherin-associated protein alpha N-catenin in both spinal nerves and lamina II of the spinal cord was examined after crushing, transecting or ligaturing the sciatic nerve in mice. Peripheral nerve injury results in histological and histochemical changes in neurons and glia. Schwann cells proliferated and migrated to form Schwann cell columns (Bungner's bands) as initial responses to denervation, and expressed E-cadherin at their site of contact with each other and later with sprouting axons. At the initial stage of myelin formation, slender processes of a single Schwann cell interdigitated with and enveloped axons, and expressed E-cadherin at the contact site elaborated by a single Schwann cell. The expression of E-cadherin in lamina II disappeared by day 7 after axotomy and reappeared following nerve ligation (partial axonal regeneration model) on day 63. In contrast, it remained at the reduced level after nerve clipping (permanent axonal degeneration model). The alteration of alpha N-catenin immunoreactivity showed the similar pattern. On the basis of present data, it is suggested that E-cadherin might be involved in the stabilization of peripheral glial network which provides the guidance of sprouting axons and myelination, and that E-cadherin-alpha N-catenin complex might be crucial for plasticity of the spinal cord dorsal horn after peripheral axotomy.▲  
Less

## Research Products (2 results)

All Other

All Publications (2 results)

[Publications] M. Hasegawa,; "Localization of E-cadherin in peripheral glia after nerve injury and repair" J Neuropathol Exp Neurol. 55 (4). 424-434 (1996) ▼

[Publications] M.Hasegawa, A.Seto, N.Uchiyama, S.Kida, T.Yamashima, J.Yamashita: "Localization of E-cadherin in peripheral glia after nerve injury and repair" J Neuropathol Exp Neurol. 55(4). 424-434 (1996) ▼

URL: [https://kaken.nii.ac.jp/report/KAKENHI-PROJECT-07671505/076715051996kenkyu\\_seika\\_hokoku\\_](https://kaken.nii.ac.jp/report/KAKENHI-PROJECT-07671505/076715051996kenkyu_seika_hokoku_)

Published: 1999-03-08