

脳血流再開後の二次的脳損傷に対する抗細胞間接着分子モノクローナル抗体療法

著者	池田 清延
著者別表示	Ikeda Kiyonobu
雑誌名	平成7(1995)年度 科学研究費補助金 一般研究(C) 研究成果報告書概要
巻	1994 1995
ページ	2p.
発行年	1999-03-08
URL	http://doi.org/10.24517/00066424



1995 Fiscal Year Final Research Report Summary

Study on the therapy against post-ischemic reperfusion brain injury using neutralizing monoclonal antibody

Research Project

Project/Area Number

06671379

Research Category

Grant-in-Aid for General Scientific Research (C)

Allocation Type

Single-year Grants

Research Field

Cerebral neurosurgery

Research Institution

Kanazawa University

Principal Investigator

IKEDA Kiyonobu Kanazawa University, Department of Neurosurgery, Assistant Professor, 医学部・付属病院, 講師 (80126565)

Co-Investigator(Kenkyū-buntansha)

KIDA Sin-ya Kanazawa University, Department of Neurosurgery, Assistant, 医学部・付属病院, 助手 (10214826)

Project Period (FY)

1994 - 1995

Keywords

adhesion molecule / brain ischemia / reperfusion injury / monoclonal antibody / L-selectin / E-selectin

Research Abstract

The effects of the anti-L selectin antibody on expression of E selectin were studied in a rabbit model of post-ischemic brain reperfusion. Anesthetized rabbits underwent 2.5 hours occlusion of the internal carotid, middle cerebral, and anterior cerebral artery with the trans-orbital approach, followed by 3 and 6 hours reperfusion. Just after the start of reperfusion, they were treated with either i.v.2 mg of anti-L selectin neutralizing antibody

(treated group) or normal saline (untreated group). For these 2 groups, 1)the immunohisto-pathological study was carried out for evaluating the localization of the expressed E-selectin (Endothelial Leukocyte Adhesion Molecule : ELAM-1), and 2)The northern blotting was done for evaluating the expression of ELAM-1 mRNA with the single strand DNA probe of ELAM-1 by RT-PCR cloning using the oligonucleotide primer. Immunohistopathologically, ELAM-1 expressed more strongly in the infarct area and its neighbor 3 hours after reperfusion than 6 hours. And it expressed not only in the small vessels but also in the cytoplasm of astrocytes and neurons around the infarct area. The administration of the neutralizing anti-L selectin antibody weakend the immunoreactivity to ELAM-1 in the 3 hours reperfusion model. Northern blotting showed no difference in expression of ELAM-1 mRNA between the treated group and the untreated one. In conclusion, L-selectin might play no role in upregulation of the ELAM-1 mRNA expression in the early stage of post-ischemic reperfusion brain injury. However, it might stimulate the release of ELAM-1, which is induced in the endothelial cells, astrocytes, and neurons by inflammatory cytokines and stored in them, to the extracellular space.

URL: https://kaken.nii.ac.jp/report/KAKENHI-PROJECT-06671379/066713791995kenkyu_seika_hokoku_

Published: 1999-03-08