

World Journal of *Orthopedics*

World J Orthop 2021 January 18; 12(1): 1-55



OPINION REVIEW

- 1 Anterior glenohumeral instability: Current review with technical pearls and pitfalls of arthroscopic soft-tissue stabilization
Apostolakos JM, Wright-Chisem J, Gulotta LV, Taylor SA, Dines JS

ORIGINAL ARTICLE**Observational Study**

- 14 Ceramic-on-ceramic vs ceramic-on-polyethylene, a comparative study with 10-year follow-up
van Loon J, Hoornenborg D, van der Vis HM, Sierevelt IN, Opdam KT, Kerkhoffs GM, Haverkamp D
- 24 Hello, can you hear me? Orthopaedic clinic telephone consultations in the COVID-19 era- a patient and clinician perspective
Vusirikala A, Ensor D, Asokan AK, Lee AJX, Ray R, Tsekes D, Edwin J

META-ANALYSIS

- 35 Thirty-day mortality of patients with hip fracture during COVID-19 pandemic and pre-pandemic periods: A systematic review and meta-analysis
Tripathy SK, Varghese P, Panigrahi S, Panda BB, Velagada S, Sahoo SS, Naik MA, Rao SK

CASE REPORT

- 51 Hallux rigidus treated with adipose-derived mesenchymal stem cells: A case report
Braile A, Toro G, De Cicco A, Cecere AB, Zanchini F, Schiavone Panni A

ABOUT COVER

Editorial Board Member of *World Journal of Orthopedics*, Maciej Płaszewski, MEd, PT, PhD, is a university professor and Head of the Department of Rehabilitation, Faculty in Biała Podlaska, Józef Piłsudski University of Physical Education, Warsaw, Poland. He has a background in physiotherapy and health policy, and also in physical education and occupational therapy. His research interests are primarily focused on evidence-based practice, secondary research and research synthesis methods. He has been engaged in research synthesis studies addressing management and recommendations for school screening and treatment interventions for people with idiopathic scoliosis, as well as other, especially physiotherapeutic, interventions, including physical modalities and pulmonary physiotherapy. Currently, he is leading an international project from the Polish Ministry of Science and Higher Education, focusing on the implementation and dissemination of evidence-based physiotherapy practice in Poland. (L-Editor: Filipodia)

AIMS AND SCOPE

The primary aim of *World Journal of Orthopedics* (*WJO*, *World J Orthop*) is to provide scholars and readers from various fields of orthopedics with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

WJO mainly publishes articles reporting research results and findings obtained in the field of orthopedics and covering a wide range of topics including arthroscopy, bone trauma, bone tumors, hand and foot surgery, joint surgery, orthopedic trauma, osteoarthropathy, osteoporosis, pediatric orthopedics, spinal diseases, spine surgery, and sports medicine.

INDEXING/ABSTRACTING

The *WJO* is now abstracted and indexed in PubMed, PubMed Central, Emerging Sources Citation Index (Web of Science), China National Knowledge Infrastructure (CNKI), China Science and Technology Journal Database (CSTJ), and Superstar Journals Database.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Yan-Xia Xing; Production Department Director: Xiang Li; Editorial Office Director: Jin-Lai Wang.

NAME OF JOURNAL

World Journal of Orthopedics

ISSN

ISSN 2218-5836 (online)

LAUNCH DATE

November 18, 2010

FREQUENCY

Monthly

EDITORS-IN-CHIEF

Massimiliano Leigheb

EDITORIAL BOARD MEMBERS

<http://www.wjnet.com/2218-5836/editorialboard.htm>

PUBLICATION DATE

January 18, 2021

COPYRIGHT

© 2021 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Hallux rigidus treated with adipose-derived mesenchymal stem cells: A case report

Adriano Braile, Giuseppe Toro, Annalisa De Cicco, Antonio Benedetto Cecere, Fabio Zanchini, Alfredo Schiavone Panni

ORCID number: Adriano Braile 0000-0001-5366-5733; Giuseppe Toro 0000-0002-8560-721X; Annalisa De Cicco 0000-0003-0548-5767; Antonio Benedetto Cecere 0000-0001-6531-2521; Fabio Zanchini 0000-0001-6438-3395; Alfredo Schiavone Panni 0000-0003-3262-3483.

Author contributions: Braile A, Toro G, and Panni AS conceived the study; Cecere AB, and De Cicco AD collected clinical data; Cecere AB, De Cicco AD, and Zanchini F interpreted the obtained data; Braile A and Toro G wrote the article; Panni AS supervised the entire process; all authors revised and approved the final version of the manuscript.

Informed consent statement: As routinely performed, the patient signed a written informed consent in which he authorized the surgical procedure and data collection for research and audit purposes. According to Italian law no formal ethical approval is required for this type of study as it includes routinely performed clinical evaluations.

Conflict-of-interest statement: The authors declare that they had no conflict of interest.

CARE Checklist (2016) statement: The authors have read the CARE

Adriano Braile, Giuseppe Toro, Annalisa De Cicco, Antonio Benedetto Cecere, Fabio Zanchini, Alfredo Schiavone Panni, Department of Medical and Surgical Specialties and Dentistry, University of Campania “Luigi Vanvitelli”, Naples 80138, Italy

Corresponding author: Giuseppe Toro, MD, Surgeon, Department of Medical and Surgical Specialties and Dentistry, University of Campania “Luigi Vanvitelli”, Via L. De Crecchio 4, Naples 80138, Italy. giusep.toro@gmail.com

Abstract

BACKGROUND

First metatarsophalangeal joint arthritis (FMTPA), also known as hallux rigidus, is the most frequent degenerative disease of the foot. Diagnosis is made through both clinical and radiological evaluation. Regenerative medicine showed promising results in the treatment of early osteoarthritis. The aim of the present study was to report the results of a case of FMTPA treated with the injection of autologous adipose-derived mesenchymal stem cells.

CASE SUMMARY

A gentleman of 50 years of age presented with a painful hallux rigidus grade 2 resistant to any previous conservative treatment (including nonsteroidal anti-inflammatory drugs and hyaluronic acid injections). An injection of autologous adipose-derived mesenchymal stem cells into the first metatarsophalangeal joint was performed. No adverse events were reported, and both function and pain scales improved after 9 mo of follow-up.

CONCLUSION

The FMTP joint injection of mesenchymal stem cells improved symptoms and function in our patient with FMTPA at 9 mo of follow-up.

Key Words: First metatarsophalangeal joint arthritis; Hallux rigidus; Stem cells; Regenerative medicine; Early osteoarthritis; Adipose derived-mesenchymal stem cells; Case report

©The Author(s) 2021. Published by Baishideng Publishing Group Inc. All rights reserved.

Checklist (2016), and the manuscript was prepared and revised according to the CARE Checklist (2016).

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Unsolicited manuscript

Specialty type: Orthopedics

Country/Territory of origin: Italy

Peer-review report's scientific quality classification

Grade A (Excellent): 0
 Grade B (Very good): B
 Grade C (Good): 0
 Grade D (Fair): 0
 Grade E (Poor): 0

Received: July 27, 2020

Peer-review started: July 27, 2020

First decision: September 21, 2020

Revised: October 5, 2020

Accepted: October 20, 2020

Article in press: October 20, 2020

Published online: January 18, 2021

P-Reviewer: Long X

S-Editor: Zhang L

L-Editor: Webster JR

P-Editor: Li JH



Core Tip: Recently, the use of intra-articular injections of stem cells has been proposed as a promising treatment in early osteoarthritis. In particular, autologous adipose-derived stem cells (aASCs) have attracted considerable attention, considering the easy access to fat tissue and the absence of adverse events registered. These characteristics make aASCs one of the most promising cell types used in regenerative medicine. Hallux rigidus, is the most frequent degenerative disease of the foot. Patients with hallux rigidus present a history of pain, gait discomfort, articular effusion, and a reduction in range of motion. Different types of treatment are available, both conservative and operative, but both are often ineffective. aASCs might overcome the gap between these two methods of treatment.

Citation: Braile A, Toro G, De Cicco A, Cecere AB, Zanchini F, Schiavone Panni A. Hallux rigidus treated with adipose-derived mesenchymal stem cells: A case report. *World J Orthop* 2021; 12(1): 51-55

URL: <https://www.wjgnet.com/2218-5836/full/v12/i1/51.htm>

DOI: <https://dx.doi.org/10.5312/wjo.v12.i1.51>

INTRODUCTION

First metatarsophalangeal joint arthritis (FMTPA), also known as hallux rigidus, is the most frequent degenerative arthritis disease of the foot^[1]. Depending on the severity of the disease, the symptoms in the case of FMTPA are pain, gait discomfort, articular effusion, and a reduction in range of motion. The diagnosis is generally completed by observing the typical findings of osteoarthritis (OA) revealed by X-rays. Coughlin and Shurnas proposed a classification for FMTPA based on both clinical features and imaging^[2] (Table 1).

Different types of treatments had been proposed, both conservative [*i.e.*, orthosis, nonsteroidal anti-inflammatory drugs, hyaluronic acid (HA) injections] and operative (*i.e.*, cheilectomy, arthroplasty, arthrodesis)^[3].

Recently, emerging evidence has supported the use of autologous adipose-derived mesenchymal stem cells (aAMSCs) for the treatment of early OA^[4-7].

The aim of the present study was to report the clinical results of a case of FMTPA treated with the injection of aAMSCs.

CASE PRESENTATION

Chief complaints

A gentleman of 50 years of age presented with a painful FMTP joint in the left foot.

History of present illness

His symptoms were resistant to both nonsteroidal anti-inflammatory drugs and HA injections.

History of past illness

The patient was already scheduled for an aAMSCs injection due to right knee Kellgren-Lawrence grade 2 OA.

Personal and family history

Personal and family history were negative for foot pathologies.

Physical examination

The clinical examination showed a positive axial grind test and joint pain, exacerbated by the dorsiflexion that impaired the patient's ability to walk.

Laboratory examinations

Laboratory tests did not support the final diagnosis.

Table 1 Coughlin and Shurnas clinical and radiographic classification of hallux rigidus

Grade	Dorsiflexion	Radiographs	Clinical
0	40°-60°	Normal	No pain, stiffness with loss of motion
1	30°-40°	Dorsal osteophytes; Minimal narrowing; Minimal flattening	Mild pain and stiffness pain with maximum dorsiflexion/plantar flexion
2	10°-30°	Global osteophytes, mild/moderate narrowing	Moderate to severe pain and stiffness relatively constant, pain near extreme ROM
3	< 10°	Cystic changes	Nearly constant pain and stiffness, no midrange pain
4	< 10°	Same as grade 3	Grade 3 + midrange pain

Imaging examinations

Antero-posterior and latero-lateral standard X-rays showed FMTPA stage 2.

FINAL DIAGNOSIS

FMTPA stage 2 was diagnosed.

TREATMENT

The patient was already scheduled for an aAMSCs injection due to right knee Kellgren-Lawrence grade 2 OA, and therefore a similar injection was proposed to treat the hallux rigidus.

Concentrated aAMSCs were obtained from the abdomen, filtered as previously reported^[5], and then injected into the FMTP joint after its distraction (**Figure 1**).

Protected weight bearing was prescribed during the first week after the procedure.

OUTCOME AND FOLLOW-UP

The patient was followed up at regular intervals. The clinical evaluation was completed using both the American Orthopedic Foot and Ankle Society (AOFAS) for hallux metatarsophalangeal-interphalangeal scale and the Visual Analog Scale (VAS).

No adverse effects were reported, excluding a transient pain in the FMTP joint in the first week after surgery. During the entire follow-up, an improvement in both VAS and AOFAS scales were observed (**Table 2**), leading to a final VAS scale of 0 and an AOFAS of 78 at 9 mo of follow-up.

DISCUSSION

Recently, the use of regenerative medicine principles has been proposed for various applications in both trauma and orthopedics, and especially for the treatment of early OA^[4-6,8-11]. While conventional therapies for early OA (*i.e.* physical therapy, glucosamine and chondroitin sulfate supplementation) showed little benefits, regenerative medicine was demonstrated to be a promising option, due to the paracrine, anti-apoptotic, anti-inflammatory, and anti-aging effects of stem cells^[12,13].

aAMSCs showed some theoretical advantages over other sources of stem cells. In fact, aAMSCs are easy to harvest, because of the wide availability of adipose tissue, and their sampling is generally associated with minimal discomfort, considering that it can be carried out using local anesthesia^[7]. Moreover, aAMSCs demonstrated a high capacity for proliferation and fibroblastic differentiation^[14]. Hass *et al*^[15] showed that adipose tissue should be considered a primary source of cells for regenerative medicine as it contains 500 times more MSCs than the same volume of bone marrow.

Emerging literature has underlined the role of aAMSCs in the treatment of early OA. Schiavone Panni *et al*^[5] conducted a study of 52 patients with early knee OA treated with arthroscopic debridement and aAMSCs injection, and showed improvement in both function and pain at an average of 15.3 mo of follow-up.

Table 2 American Orthopedic Foot and Ankle Society and Visual Analog Scale scores

	Pre-operatively	6-mo follow-up	9-mo follow-up
AOFAS score			
Pain	30	30	30
Function	35	40	40
Alignment	8	8	8
Total score	73	78	78
VAS	7	5	0

AOFAS: American Orthopedic Foot and Ankle Society; VAS: Visual analog scale.



Figure 1 Intraoperative picture showing the autologous adipose-derived mesenchymal stem cells injection into the first metatarsophalangeal joint.

Similarly, Song *et al*^[16] reported the amelioration of pain, function and cartilage volume of the knee after multiple injections of aAMSCs. The efficacy of aAMSCs in OA was recently confirmed in a systematic review conducted by McIntyre *et al*^[17].

FMTPA is a degenerative disease with an incidence of 2.5% in patients over 50 years of age. Its treatment might be frustrating for both the orthopedic and the patient, considering the conflicting outcomes reported after conservative treatment, including HA injections. Petrella *et al*^[18] in their study of 47 patients with FMTPA, described the long-term improvement in both pain and function after multiple HA injections; however, Munteanu *et al*^[19] in their randomized controlled trial did not observe any differences when HA was compared with placebo.

Pons *et al*^[20] in a randomized study compared the use of sodium hyaluronate with triamcinolone acetonide in FMTPA, and reported an improvement in pain relief and function at 3 mo after the injections. However, a high percentage of patients in both groups required subsequent surgery after 1 year of follow-up, due to further progression of the disease with worsening of both pain and function.

To the best of our knowledge, this is the first report on the use of aAMSCs for FMTPA. A single injection was effective in treating FMTPA, improving both the AOFAS and VAS score at 9 mo of follow-up.

CONCLUSION

The present case report indicates that the injection of aAMSCs might be a promising treatment for FMTPA. Obviously, larger cohorts and longer follow-up studies are needed to confirm these findings.

REFERENCES

- 1 **Ho B**, Baumhauer J. Hallux rigidus. *EFORT Open Rev* 2017; **2**: 13-20 [PMID: 28607766 DOI: 10.1302/2058-5241.2.160031]
- 2 **Coughlin MJ**, Shurnas PS. Hallux rigidus. Grading and long-term results of operative treatment. *J Bone Joint Surg Am* 2003; **85**: 2072-2088 [PMID: 14630834 DOI: 10.2106/00004623-200311000-00003]
- 3 **Herrera-Pérez M**, Andarcia-Bañuelos C, de Bergua-Domingo J, Paul J, Barg A, Valderrabano V. [Proposed global treatment algorithm for hallux rigidus according to evidence-based medicine]. *Rev Esp Cir Ortop Traumatol* 2014; **58**: 377-386 [PMID: 25088241 DOI: 10.1016/j.recot.2014.05.004]
- 4 **Damia E**, Chicharro D, Lopez S, Cuervo B, Rubio M, Sopena JJ, Vilar JM, Carrillo JM. Adipose-Derived Mesenchymal Stem Cells: Are They a Good Therapeutic Strategy for Osteoarthritis? *Int J Mol Sci* 2018; **19** [PMID: 29966351 DOI: 10.3390/ijms19071926]
- 5 **Schiavone Panni A**, Vasso M, Braile A, Toro G, De Cicco A, Viggiano D, Lepore F. Preliminary results of autologous adipose-derived stem cells in early knee osteoarthritis: identification of a subpopulation with greater response. *Int Orthop* 2019; **43**: 7-13 [PMID: 30280218 DOI: 10.1007/s00264-018-4182-6]
- 6 **Ha CW**, Park YB, Kim SH, Lee HJ. Intra-articular Mesenchymal Stem Cells in Osteoarthritis of the Knee: A Systematic Review of Clinical Outcomes and Evidence of Cartilage Repair. *Arthroscopy* 2019; **35**: 277-288. e2 [PMID: 30455086 DOI: 10.1016/j.arthro.2018.07.028]
- 7 **Nöth U**, Steinert AF, Tuan RS. Technology insight: adult mesenchymal stem cells for osteoarthritis therapy. *Nat Clin Pract Rheumatol* 2008; **4**: 371-380 [PMID: 18477997 DOI: 10.1038/ncprheum0816]
- 8 **Caggiari G**. Efficacy of platelet-rich plasma in experiential instrumented interbody spinal fusion. *EuroMediterranean Biomed J* 2016; 141-147 [DOI: 10.3269/1970-5492.2016.11.20]
- 9 **Doria C**, Mosele GR, Caggiari G, Puddu L, Ciurlia E. Treatment of Early Hip Osteoarthritis: Ultrasound-Guided Platelet Rich Plasma vs Hyaluronic Acid Injections in a Randomized Clinical Trial. *Joints* 2017; **5**: 152-155 [PMID: 29270545 DOI: 10.1055/s-0037-1605584]
- 10 **Toro G**, Moretti A, Toro G, Tirelli A, Calabrò G, Toro A, Iolascon G. Surgical treatment of neglected hip fracture in children with cerebral palsy: case report and review of the literature. *Clin Cases Miner Bone Metab* 2017; **14**: 317-323 [PMID: 29354160 DOI: 10.11138/ccmbm/2017.14.3.317]
- 11 **Toro G**, Lepore F, Calabrò G, Toro G, Rossini M, Vasso M, Schiavone Panni A. Humeral shaft non-union in the elderly: Results with cortical graft plus stem cells. *Injury* 2019; **50** Suppl 2: S75-S79 [PMID: 30846284 DOI: 10.1016/j.injury.2019.01.050]
- 12 **Wehling P**, Evans C, Wehling J, Maixner W. Effectiveness of intra-articular therapies in osteoarthritis: a literature review. *Ther Adv Musculoskelet Dis* 2017; **9**: 183-196 [PMID: 28835778 DOI: 10.1177/1759720X17712695]
- 13 **Platas J**, Guillén MI, Pérez Del Caz MD, Gomar F, Castejón MA, Mirabet V, Alcaraz MJ. Paracrine effects of human adipose-derived mesenchymal stem cells in inflammatory stress-induced senescence features of osteoarthritic chondrocytes. *Aging (Albany NY)* 2016; **8**: 1703-1717 [PMID: 27490266 DOI: 10.18632/aging.101007]
- 14 **McIntosh KR**, Frazier T, Rowan BG, Gimble JM. Evolution and future prospects of adipose-derived immunomodulatory cell therapeutics. *Expert Rev Clin Immunol* 2013; **9**: 175-184 [PMID: 23390948 DOI: 10.1586/eci.12.96]
- 15 **Hass R**, Kasper C, Böhm S, Jacobs R. Different populations and sources of human mesenchymal stem cells (MSC): A comparison of adult and neonatal tissue-derived MSC. *Cell Commun Signal* 2011; **9**: 12 [PMID: 21569606 DOI: 10.1186/1478-811X-9-12]
- 16 **Song Y**, Du H, Dai C, Zhang L, Li S, Hunter DJ, Lu L, Bao C. Human adipose-derived mesenchymal stem cells for osteoarthritis: a pilot study with long-term follow-up and repeated injections. *Regen Med* 2018; **13**: 295-307 [PMID: 29417902 DOI: 10.2217/rme-2017-0152]
- 17 **McIntyre JA**, Jones IA, Han B, Vangness CT Jr. Intra-articular Mesenchymal Stem Cell Therapy for the Human Joint: A Systematic Review. *Am J Sports Med* 2018; **46**: 3550-3563 [PMID: 29099618 DOI: 10.1177/0363546517735844]
- 18 **Petrella RJ**, Cogliano A. Intra-articular Hyaluronic Acid Treatment for Golfer's Toe: Keeping Older Golfers on Course. *Phys Sportsmed* 2004; **32**: 41-45 [PMID: 20086421 DOI: 10.3810/psm.2004.07.453]
- 19 **Munteanu SE**, Zammit GV, Menz HB, Landorf KB, Handley CJ, Elzarka A, Deluca J. Effectiveness of intra-articular hyaluronan (Synvisc, hylan G-F 20) for the treatment of first metatarsophalangeal joint osteoarthritis: a randomised placebo-controlled trial. *Ann Rheum Dis* 2011; **70**: 1838-1841 [PMID: 21791454 DOI: 10.1136/ard.2011.153049]
- 20 **Pons M**, Alvarez F, Solana J, Viladot R, Varela L. Sodium hyaluronate in the treatment of hallux rigidus. A single-blind, randomized study. *Foot Ankle Int* 2007; **28**: 38-42 [PMID: 17257536 DOI: 10.3113/FAI.2007.0007]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA
Telephone: +1-925-3991568
E-mail: bpgoffice@wjgnet.com
Help Desk: <https://www.f6publishing.com/helpdesk>
<https://www.wjgnet.com>

