

Non-standard neutrino interactions in IceCube

² Elisa Lohfink for the IceCube Collaboration

3 E-mail: elisa.lohfink@icecube.wisc.edu

Non-standard neutrino interactions (NSI) may arise in various types of new physics. Their existence would change the potential that atmospheric neutrinos encounter when traversing Earth matter and hence alter their oscillation behavior. This imprint on coherent neutrino forward scattering can be probed using high-statistics neutrino experiments such as IceCube and its low-energy extension, DeepCore. Both provide extensive data samples that include all neutrino flavors, with oscillation baselines between tens of kilometers and the diameter of the Earth. DeepCore event energies reach from a few GeV up to the order of 100 GeV - which marks the lower threshold for higher energy IceCube atmospheric samples, ranging up to 10 TeV. In DeepCore data, the large sample size and energy range allow us to consider not only flavor-violating and flavor-nonuniversal NSI in the $\mu - \tau$ sector, but also those involving electron flavor. The effective parameterization used in our analyses is independent of the underlying model and the new physics mass scale. In this way, competitive limits on several NSI parameters have been set in the past. The 8 years of data available now result in significantly improved sensitivities. This improvement stems not only from the increase in statistics but also from substantial improvement in the treatment of systematic uncertainties, background rejection and event reconstruction.

Corresponding authors: Elisa Lohfink^{1*}, Sebastian Böser¹, Thomas Ehrhardt¹, Grant Parker² ¹ Institute of Physics, University of Mainz, Staudinger Weg 7, D-55099 Mainz, Germany ² Dept. of Physics, University of Texas at Arlington, 502 Yates St., Science Hall Rm 108, Box 19059, Arlington, TX 76019, USA

* Presenter

4

*** The European Physical Society Conference on High Energy Physics (EPS-HEP2021), *** *** 26-30 July 2021 *** *** Online conference, jointly organized by Universität Hamburg and the research center DESY ***

© Copyright owned by the author(s) under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License (CC BY-NC-ND 4.0).

5 1. Introduction

6 1.1 Neutrino oscillations in IceCube and DeepCore

The IceCube Neutrino Observatory is an 7 ice Cherenkov detector located at the South 8 Pole. It consists of 5160 light detectors, sog called Digital Optical Modules (DOMs), that 10 have been deployed along 86 strings within a 11 cubic kilometer of glacial ice [1]. When neutri-12 nos interact with matter inside or close to the in-13 strumented volume, IceCube detects Cherenkov 14 light produced by charged secondary particles. 15 Being optimized for the detection of cosmic 16 neutrinos, the lowest energy at which individ-17 ual events can be reconstructed in IceCube is 18 ~100 GeV. DeepCore is IceCube's low en-19 ergy extension, consisting of a more densely 20 instrumented volume at the detector center (see 21 Fig. 1). For events occurring within this re-22 gion, the energy threshold is lowered to ap-23 proximately 5 GeV [2]. Atmospheric neutrinos 24 in DeepCore are a great probe for matter ef-25 fects on neutrino oscillation [3], as they cover 26 the energy range from approximately 5 GeV to 27 300 GeV. Depending on the angle at which they 28 traverse the Earth, oscillation baselines range 29



Figure 1: Layout of the IceCube Neutrino Observatory and its low energy extension, DeepCore (highlighted in green). Each dot represents a DOM, deployed within 1 km³ of glacier inside the Antarcic ice sheet at depths between 1450 and 2450 m

³⁰ between a few km for down-going trajectories and $O(10^4 \text{ km})$ for up-going trajectories, crossing

the entire Earth. Event samples of atmospheric neutrinos contain neutral current (NC) and charged

³² current (CC) interactions of all neutrino flavors.

1.2 Non-standard neutrino interactions

IceCube's sensitivity to neutrino oscillations allows us to probe matter effects from a number 34 of beyond the standard model (BSM) scenarios, including non-standard neutrino interactions (NSI). 35 We test for NC forward scattering between neutrinos of all flavors and charged fermions in Earth 36 matter (up- and down-quarks, electrons), assuming the existence of a new heavy mediator particle. 37 This is well motivated from theory, as NSI come up in a multitude of neutrino mass models [4]. By 38 effectively introducing new degrees of freedom to neutrino oscillations, NSI effects might resolve 39 tensions that are currently observed in measurements of standard oscillation parameters, such as 40 $\delta_{\rm CP}$ [5]. 41

In an approach that is mostly independent from the underlying model and the mediator mass, the NSI-induced change to the matter potential can be parametrized with five effective coupling parameters:

$$H_{\text{mat}} = \sqrt{2}G_F N_e(x) \begin{pmatrix} 1 + (\epsilon_{ee}^{\oplus} - \epsilon_{\mu\mu}^{\oplus}) & \epsilon_{e\mu}^{\oplus} & \epsilon_{e\tau}^{\oplus} \\ \epsilon_{e\mu}^{\oplus *} & 0 & \epsilon_{\mu\tau}^{\oplus} \\ \epsilon_{e\tau}^{\oplus *} & \epsilon_{\mu\tau}^{\oplus *} & (\epsilon_{\tau\tau}^{\oplus} - \epsilon_{\mu\mu}^{\oplus}) \end{pmatrix}$$

The three flavor violating off-diagonal parameters, $\epsilon_{e\mu}^{\oplus}$, $\epsilon_{e\tau}^{\oplus}$ and $\epsilon_{\mu\tau}^{\oplus}$, are complex-valued. Subtract-42 ing $\epsilon_{\mu\mu}^{\oplus} \times \mathbb{1}$ yields two real-valued lepton universality-violating diagonal parameters, $\epsilon_{ee}^{\oplus} - \epsilon_{\mu\mu}^{\oplus}$ and 43 $\epsilon_{\tau\tau}^{\oplus} - \epsilon_{\mu\mu}^{\oplus}$, thereby reducing dimensionality without observable consequences. 44

NSI analyses in IceCube 2. 45

IceCube NSI analyses are being performed 46 on multiple data sets, varying in choice of meth-47 ods and investigated NSI parameters. This is 48 motivated by how the sensitivity to different 49 NSI signatures depends on event types and en-50 ergies. 51

The atmospheric neutrino flux at GeV to 52 TeV energies is dominated by muon neutri-53 nos [6]. Together with IceCube's capacity to 54 discriminate v_{μ} CC events from others (see 55 Sec. 2.1), this renders our data sets most sen-56 sitive to effects on v_{μ} oscillation probabilities, 57 $P_{\mu\alpha}^{1}$. 58

The impact of individual NSI couplings on 59 single oscillation probabilities differs between 60 the energy ranges at which IceCube and Deep-61 Core can resolve individual neutrino interac-62 tions (see Fig. 2). IceCube events have ener-63 gies above 100 GeV, making them almost ex-64 clusively sensitive to $\epsilon^{\oplus}_{\mu\tau}$. Similarly, for Deep-65 Core event energies of approximately 5 GeV 66 to 100 GeV, dominant effects are observed for



Figure 2: Examples of oscillation probabilities at different NSI hypotheses. Each panel shows $P_{\mu\tau}$ for different positive (in red) and negative (in blue) values of the individual NSI parameter together with the standard interaction oscillations (SI) case in black.

 $\epsilon_{\mu\tau}^{\oplus}$ [7]. Current DeepCore analyses, however, are able to constrain all effective NSI parameters. 68

2.1 Analysis principle for DeepCore samples 69

67

In a typical DeepCore NSI analysis, binned data are compared to Monte Carlo (MC) simulated 70 expectation templates, generated for a variety of NSI hypotheses. A metric based on the differences 71 in event counts per bin is optimized or sampled to obtain the best-fitting NSI hypothesis and 72 confidence intervals. 73

The binned quantities are the reconstructed energy of the events, the cosine of their zenith angle 74 and their topology. The zenith angle corresponds to the oscillation baseline at which the neutrino 75

¹We use the term neutrinos for both neutrinos and antineutrinos and indicate arbitrary flavor as $\alpha \in [e, \mu, \tau]$

traverses the Earth. The reconstructed event topology is based on a binned quantity, subdivided 76 between more track-like and more cascade-like events. Track-like events are v_{μ} CC interactions, 77 with the secondary muon emitting light along its extended path through the detector. Any other 78 neutrino interactions result in approximately spherical, cascade-like light deposition. 79

The analysis varies the NSI hypothesis being studied, as well as nuisance parameters for signal 80 and background events, and detector characteristics [8]. Uncertainties in our understanding of 81 the detector concern the efficiency of DOMs as well as the optical properties of the surrounding 82 ice. Ice absorption and scattering characteristics differ between pristine glacial ice and the melted 83 and re-frozen ice in which modules have been deployed. Signal nuisance parameters relate to the 84 atmospheric flux of neutrinos, the oscillations they undergo and their interaction cross section. 85

Current prospects and results 3. 86

While DeepCore samples allow for resolv-87 ing the signatures of all NSI couplings at few 88 GeV, analyses of IceCube events make use of 89 the sensitivity of higher energy data to $\epsilon^{\oplus}_{\mu\tau}$ es-90 pecially. A soon to be published analysis is 91 based on the same data set as a recent ster-92 ile neutrino analysis [10], including ν_{μ} events 93 at 500 GeV to 10 TeV from 7.5 years of Ice-94 Cube data. Preliminary sensitivities to NSI im-95 ply globally competitive constraints on complex 96 $\epsilon_{\mu\tau}^{\oplus} \text{ of } -2.91 \cdot 10^{-3} \le |\epsilon_{\mu\tau}^{\oplus}| \le 2.93 \cdot 10^{-3}.$ 97

10^{0} 10^{-1} 90% allowed region 10^{-1} 10 10^{-2} 0 0 -10^{-2} -2 10 -10^{-1} -10^{-1} -10^{0} $\epsilon^{\oplus}_{\mu\mu} = \epsilon^{\oplus}_{\tau\tau} = \epsilon^{\oplus}_{\mu\mu}$ $\epsilon^{\oplus}_{e\mu}$ $\epsilon_{e\tau}^{\oplus}$ $\epsilon^{\oplus}_{\mu\tau}$

Figure 3: Limits on singly tested NSI couplings based

on 3 years of DeepCore data, compared to earlier re-

3.1 Recent DeepCore results 98

A recently published analysis [9] was for 99

the first time able to constrain all effective NSI 100

parameters from a single IceCube sample, using the approach outlined in Sec. 2.1. It was based on 101 three years of DeepCore data, including events of all flavors, from all zenith angles and with energies 102 of 5.6 GeV to 100 GeV. In Fig. 3, the limits obtained from investigating all effective NSI parameters 103 individually, while fixing all other NSI couplings to 0, can be found. This approach results in some 104 model dependence, as it excludes any hypotheses with multiple non-zero NSI parameters. The 105 reason why this approach is chosen is that keeping all NSI parameters free is not computationally 106 feasible. In order to cover the full model space, a second part of the analysis was carried out using a 107 different parametrization with three free parameters (generalized matter potential), yielding limits 108 on the scale and structure of fully free NSI. 109

sults [9].

3.2 Upcoming 8 year DeepCore analysis 110

An analysis similar to the one performed in [9] is applied to 8 years of DeepCore data with 111 an extended energy range with respect to earlier analyses of 5.6 GeV to 300 GeV. Preliminary 112 sensitivities suggest significant improvement, specifically to the limits on $\epsilon^{\oplus}_{\mu\tau}$ (see Fig. 4), while 113 sensitivity to the complex phases of flavor violating parameters remains low. 114



The improved sensitivity is not only due to the larger data set, but also to a multitude of 115 refinements of the analysis tools and technique. An improved background rejection using Boosted 116 Decision Trees (BDTs) results in a sample purity of 97% (compared to 95% in earlier DeepCore 117 samples [3]), with $\sim 3\%$ of atmospheric muon events and negligible contribution of random noise 118 hits of < 0.03% of the final sample. This is reached while keeping signal rates at ~ 1 mHz, improving 119 upon earlier samples by approximately a factor of two. 120

A new reconstruction keeps the original approach of minimizing a tabulated likelihood, but 121 reverses the process by starting at the individual modules and tracing photons back to their source. 122 For topology classification, for the first time, BDTs are used. These measures result in an at least 123 two times faster reconstruction and better resolution in all observable variables. 124

With the enhanced data set containing approximately $3 \cdot 10^5$ events, analysis techniques 125 face unprecedented requirements in speed and optimization of computational resource usage. 126 For SI analyses, these could be met through optimization of the established PISA tool [11]. 127

For an NSI analysis of these data further re-128 vision of the analysis technique is required as 129 the traditional approach through minimization 130 of the parameter space is rendered computation-131 ally challenging due to the number and behavior 132 of NSI parameters: fast and simple minimizers 133 fail to resolve symmetries e.g. in the complex 134 phases of flavor violating couplings. Additional 135 complexity is introduced through the degener-136 acy of $\epsilon_{ee}^{\oplus} - \epsilon_{\mu\mu}^{\oplus}$ with neutrino mass ordering 137 (NMO) and vacuum-like oscillations behavior 138



Figure 4: Preliminary sensitivities based on 8 years

of DeepCore data to all NSI couplings, each evaluated

at $\epsilon_{ee}^{\oplus} - \epsilon_{\mu\mu}^{\oplus} = -1$. 139 Markov Chain Monte Carlo (MCMC) sam-140

pling of the parameter space in place of mini-141

142

146



well with high dimensional space and yield the full parameter space profile and confidence limits 143 in addition to the best fitting hypothesis. We use the emcee Python package [12], which allows 144 for parallelization via openMPI. This promises to allow for simultaneous evaluation of all effective 145 NSI parameter magnitudes.

3.3 On the horizon: The IceCube Upgrade 147

An IceCube low-energy extension, the IceCube Upgrade, consists of seven densely instru-148 mented additional strings which will be deployed within the next several years [13]. Its increased 149 sensitivity to low energy neutrinos owes to the increase in event rate as well as the Upgrade's 150 capacity to observe individual events starting at $\sim 1 \text{ GeV}$, resolving NSI signatures over a wider 151 range. Calibration devices along the new strings will furthermore improve our understanding of 152 detector systematics. As a result, the IceCube Upgrade will constitute a significant improvement in 153 sensitivity to NSI effects. 154

155 References

- [1] IceCube Collaboration, R. Abbasi *et al. Astroparticle Physics* **35** no. 10, (2012) 615–624.
- [2] IceCube Collaboration, R. Abbasi *et al. Astroparticle Physics* **35** no. 10, (2012) 615–624.
- ¹⁵⁸ [3] **IceCube** Collaboration, M. G. Aartsen *et al. Phys. Rev. Lett.* **120** (Feb, 2018) 071801.
- ¹⁵⁹ [4] M. B. Gavela, D. Hernandez, T. Ota, and W. Winter *Phys. Rev. D* **79** (Jan, 2009) 013007.
- ¹⁶⁰ [5] S. S. Chatterjee and A. Palazzo *Phys. Rev. Lett.* **126** (Feb, 2021) 051802.
- [6] M. Honda, M. S. Athar, T. Kajita, K. Kasahara, and S. Midorikawa *Phys. Rev. D* 92 (Jul, 2015) 023004.
- ¹⁶³ [7] **IceCube** Collaboration, M. G. Aartsen *et al. Phys. Rev. D* **97** (Apr, 2018) 072009.
- [8] IceCube Collaboration, M. G. Aartsen *et al. Phys. Rev. D* 99 (Feb, 2019) 032007.
- ¹⁶⁵ [9] **IceCube** Collaboration, R. Abbasi *et al. Phys. Rev. D* **104** (Oct, 2021) 072006.
- ¹⁶⁶ [10] **IceCube** Collaboration, M. G. Aartsen *et al. Phys. Rev. D* **102** (Sep, 2020) 052009.
- [11] M. G. Aartsen *et al.* Nuclear Instruments and Methods in Physics Research Section A:
 Accelerators, Spectrometers, Detectors and Associated Equipment **977** (2020) 164332.
- [12] D. Foreman-Mackey, D. W. Hogg, D. Lang, and J. Goodman *Publications of the Astronomical Society of the Pacific* **125** no. 925, (Mar, 2013) 306–312.
- 171 [13] A. Ishihara *PoS* **ICRC2019** (2019) 1031.

172 Full Author List: IceCube Collaboration*

R. Abbasi¹⁷, M. Ackermann⁵⁹, J. Adams¹⁸, J. A. Aguilar¹², M. Ahlers²², M. Ahrens⁵⁰, C. Alispach²⁸, A. A. Alves Jr.³¹, N. M. Amin⁴², 173 R. An¹⁴, K. Andeen⁴⁰, T. Anderson⁵⁶, G. Anton²⁶, C. Argüelles¹⁴, Y. Ashida³⁸, S. Axani¹⁵, X. Bai⁴⁶, A. Balagopal V.³⁸, A. Barbano²⁸, 174 S. W. Barwick³⁰, B. Bastian⁵⁹, V. Basu³⁸, S. Baur¹², R. Bay⁸, J. J. Beatty^{20, 21}, K.-H. Becker⁵⁸, J. Becker Tjus¹¹, C. Bellenghi²⁷, S. 175 BenZvi⁴⁸, D. Berley¹⁹, E. Bernardini^{59, 60}, D. Z. Besson^{34, 61}, G. Binder^{8, 9}, D. Bindig⁵⁸, E. Blaufuss¹⁹, S. Blot⁵⁹, M. Boddenberg¹, 176 F. Bontempo³¹, J. Borowka¹, S. Böser³⁹, O. Botner⁵⁷, J. Böttcher¹, E. Bourbeau²², F. Bradascio⁵⁹, J. Braun³⁸, B. Brinson⁶, S. Bron²⁸, 177 J. Brostean-Kaiser⁵⁹, S. Browne³², A. Burgman⁵⁷, R. T. Burley², R. S. Busse⁴¹, M. A. Campana⁴⁵, E. G. Carnie-Bronca², C. Chen⁶, Z. Chen⁵¹, D. Chirkin³⁸, K. Choi⁵², B. A. Clark²⁴, K. Clark³³, L. Classen⁴¹, A. Coleman⁴², G. H. Collin¹⁵, J. M. Conrad¹⁵, P. Coppin¹³, P. Correa¹³, D. F. Cowen⁵⁵, ⁵⁶, R. Cross⁴⁸, C. Dappen¹, P. Dave⁶, C. De Clercq¹³, J. J. DeLaunay⁵⁴, D. Delgado López¹⁴, H. Dembinski⁴², K. Deoskar⁵⁰, A. Desai³⁸, P. Desiati³⁸, K. D. de Vries¹³, G. de Wasseige¹³, M. de With¹⁰, T. DeYoung²⁴, A. Diaz¹⁵, 178 179 180 181 J. C. Díaz-Vélez³⁸, M. Dittmer⁴¹, H. Dujmovic³¹, M. Dunkman⁵⁶, M. A. DuVernois³⁸, E. Dvorak⁴⁶, T. Ehrhardt³⁹, P. Eller²⁷, R. 182 Engel^{31, 32}, H. Erpenbeck¹, J. Evans¹⁹, P. A. Evenson⁴², K. L. Fan¹⁹, A. R. Fazely⁷, N. Feigl¹⁰, S. Fiedlschuster²⁶, A. T. Fienberg⁵⁶, 183 K. Filimonov⁸, C. Finley⁵⁰, L. Fischer⁵⁹, D. Fox⁵⁵, A. Franckowiak^{11, 59}, E. Friedman¹⁹, A. Fritz³⁹, P. Fürst¹, T. K. Gaisser⁴², J. 184 Gallagher³⁷, E. Ganster¹, A. Garcia¹⁴, S. Garrappa⁵⁹, L. Gerhardt⁹, A. Ghadimi⁵⁴, C. Glaser⁵⁷, T. Glauch²⁷, T. Glüsenkamp²⁶, J. 185 G. Gonzalez⁴², S. Goswami⁵⁴, D. Grant²⁴, T. Grégoire⁵⁶, S. Griswold⁴⁸, C. Günther¹, C. Haack²⁷, A. Hallgren⁵⁷, R. Halliday²⁴, 186 L. Halve¹, F. Halzen³⁸, M. Ha Minh²⁷, K. Hanson³⁸, J. Hardin³⁸, A. A. Harnisch²⁴, A. Haungs³¹, D. Hebecker¹⁰, K. Helbing⁵⁸, F. 187 Henningsen²⁷, E. C. Hettinger²⁴, S. Hickford⁵⁸, J. Hignight²⁵, C. Hill¹⁶, G. C. Hill², K. D. Hoffman¹⁹, R. Hoffmann⁵⁸, T. Hoinka²³, 188 B. Hokanson-Fasig³⁸, K. Hoshina^{38, 62}, F. Huang⁵⁶, M. Huber²⁷, T. Huber³¹, K. Hultqvist⁵⁰, M. Hünnefeld²³, R. Hussain³⁸, S. In⁵², 189 N. Iovine¹², A. Ishihara¹⁶, M. Jansson⁵⁰, G. S. Japaridze⁵, M. Jeong⁵², M. Jin¹⁴, B. J. P. Jones⁴, D. Kang³¹, W. Kang⁵², X. Kang⁴⁵, A. 190 Kappes⁴¹, D. Kappesser³⁹, T. Karg⁵⁹, M. Karl²⁷, A. Karle³⁸, U. Katz²⁶, M. Kauer³⁸, M. Kellermann¹, J. L. Kelley³⁸, A. Kheirandish⁵⁶, 191 K. Kin¹⁶, T. Kintscher⁵⁹, J. Kiryluk³¹, S. R. Klein^{8, 9}, R. Koirala⁴², H. Kolanoski¹⁰, T. Kontrimas²⁷, L. Köpke³⁹, C. Kopper²⁴, S. 192 Kopper⁵⁴, D. J. Koskinen²², P. Koundal³¹, M. Kovacevich⁴⁵, M. Kowalski^{10, 59}, T. Kozynets²², E. Kun¹¹, N. Kurahashi⁴⁵, N. Lad⁵⁹, 193 C. Lagunas Gualda⁵⁹, J. L. Lanfranchi⁵⁶, M. J. Larson¹⁹, F. Lauber⁵⁸, J. P. Lazar^{14, 38}, J. W. Lee⁵², K. Leonard³⁸, A. Leszczyńska³², 194 Y. Li⁵⁶, M. Lincetto¹¹, Q. R. Liu³⁸, M. Liubarska²⁵, E. Lohfink³⁹, C. J. Lozano Mariscal⁴¹, L. Lu³⁸, F. Lucarelli²⁸, A. Ludwig^{24, 35}, 195 W. Luszczak³⁸, Y. Lyu⁸, ⁹, W. Y. Ma⁵⁹, J. Madsen³⁸, K. B. M. Mahn²⁴, Y. Makino³⁸, S. Mancina³⁸, I. C. Mariş¹², R. Maruyama⁴³, K. 196 Mase¹⁶, T. McElroy²⁵, F. McNally³⁶, J. V. Mead²², K. Meagher³⁸, S. Mechbal⁵⁹, A. Medina²¹, M. Meier¹⁶, S. Meighen-Berger²⁷, J. 197 Micallef²⁴, D. Mockler¹², T. Montaruli²⁸, R. W. Moore²⁵, R. Morse³⁸, M. Moulai¹⁵, R. Naab⁵⁹, R. Nagai¹⁶, U. Naumann⁵⁸, J. Necker⁵⁹, 198 L. V. Nguyễn²⁴, H. Niederhausen²⁷, M. U. Nisa²⁴, S. C. Nowicki²⁴, A. Obertacke Pollmann⁵⁸, M. Oehler³¹, B. Oeyen²⁹, A. Olivas¹⁹, E. 199 O'Sullivan⁵⁷, H. Pandya⁴², D. V. Pankova⁵⁶, N. Park³³, G. K. Parker⁴, E. N. Paudel⁴², L. Paul⁴⁰, C. Pérez de los Heros⁵⁷, L. Peters¹, J. 200 Peterson³⁸, S. Philippen¹, D. Pieloth²³, S. Pieper⁵⁸, M. Pittermann³², A. Pizzuto³⁸, M. Plum⁴⁰, Y. Popovych³⁹, A. Porcelli²⁹, M. Prado 201 Rodriguez³⁸, P. B. Price⁸, B. Pries²⁴, G. T. Przybylski⁹, C. Raab¹², A. Raissi¹⁸, M. Rameez²², K. Rawlins³, I. C. Rea²⁷, A. Rehman⁴², 202 P. Reichherzer¹¹, R. Reimann¹, G. Renzi¹², E. Resconi²⁷, S. Reusch⁵⁹, W. Rhode²³, M. Richman⁴⁵, B. Riedel³⁸, E. J. Roberts², S. 203 Robertson^{8, 9}, G. Roellinghoff⁵², M. Rongen³⁹, C. Rott^{49, 52}, T. Ruhe²³, D. Ryckbosch²⁹, D. Rysewyk Cantu²⁴, I. Safa^{14, 38}, J. Saffer³², 204 S. E. Sanchez Herrera²⁴, A. Sandrock²³, J. Sandroos³⁹, M. Santander⁵⁴, S. Sarkar⁴⁴, S. Sarkar²⁵, K. Satalecka⁵⁹, M. Schaufel¹, H. Schieler³¹, S. Schindler²⁶, P. Schlunder²³, T. Schmidt¹⁹, A. Schneider³⁸, J. Schneider²⁶, F. G. Schröder^{31, 42}, L. Schumacher²⁷, G. 205 206 Schwefer¹, S. Sclafani⁴⁵, D. Seckel⁴², S. Seunarine⁴⁷, A. Sharma⁵⁷, S. Shefali³², M. Silva³⁸, B. Skrzypek¹⁴, B. Smithers⁴, R. Snihur³⁸, 207 J. Soedingrekso²³, D. Soldin⁴², C. Spannfellner²⁷, G. M. Spiczak⁴⁷, C. Spiering^{59, 61}, J. Stachurska⁵⁹, M. Stamatikos²¹, T. Stanev⁴², 208 R. Stein⁵⁹, J. Stettner¹, A. Steuer³⁹, T. Stezelberger⁹, T. Stürwald⁵⁸, T. Stuttard²², G. W. Sullivan¹⁹, I. Taboada⁶, S. Ter-Antonyan⁷, S. 209 Tilav⁴², F. Tischbein¹, K. Tollefson²⁴, C. Tönnis⁵³, S. Toscano¹², D. Tosi³⁸, A. Trettin⁵⁹, M. Tselengidou²⁶, C. F. Tung⁶, A. Turcati²⁷, 210 R. Turcotte³¹, C. F. Turley⁵⁶, J. P. Twagirayezu²⁴, B. Ty³⁸, M. A. Unland Elorrieta⁴¹, N. Valtonen-Mattila⁵⁷, J. Vandenbroucke³⁸, N. 211 van Eijndhoven¹³, D. Vannerom¹⁵, J. van Santen⁵⁹, S. Verpoest²⁹, C. Walck⁵⁰, T. B. Watson⁴, C. Weaver²⁴, P. Weigel¹⁵, A. Weindl³¹, 212 M. J. Weiss⁵⁶, J. Weldert³⁹, C. Wendt³⁸, J. Werthebach²³, M. Weyrauch³², N. Whitehorn^{24, 35}, C. H. Wiebusch¹, D. R. Williams⁵⁴ 213 M. Wolf²⁷, K. Woschnagg⁸, G. Wrede²⁶, J. Wulff¹¹, X. W. Xu⁷, J. P. Yanez²⁵, S. Yoshida¹⁶, S. Yu²⁴, T. Yuan³⁸, Z. Zhang⁵¹, P. Zhelnin¹⁴ 214

- 215
- ¹ III. Physikalisches Institut, RWTH Aachen University, D-52056 Aachen, Germany
- ²¹⁷ ² Department of Physics, University of Adelaide, Adelaide, 5005, Australia
- ²¹⁸ ³ Dept. of Physics and Astronomy, University of Alaska Anchorage, 3211 Providence Dr., Anchorage, AK 99508, USA
- ⁴ Dept. of Physics, University of Texas at Arlington, 502 Yates St., Science Hall Rm 108, Box 19059, Arlington, TX 76019, USA
- ⁵ CTSPS, Clark-Atlanta University, Atlanta, GA 30314, USA
- ⁶ School of Physics and Center for Relativistic Astrophysics, Georgia Institute of Technology, Atlanta, GA 30332, USA
- ²²² ⁷ Dept. of Physics, Southern University, Baton Rouge, LA 70813, USA
- ⁸ Dept. of Physics, University of California, Berkeley, CA 94720, USA
- ⁹ Lawrence Berkeley National Laboratory, Berkeley, CA 94720, USA
- ¹⁰ Institut für Physik, Humboldt-Universität zu Berlin, D-12489 Berlin, Germany
- 226 ¹¹ Fakultät für Physik & Astronomie, Ruhr-Universität Bochum, D-44780 Bochum, Germany
- ¹² Université Libre de Bruxelles, Science Faculty CP230, B-1050 Brussels, Belgium
- ¹³ Vrije Universiteit Brussel (VUB), Dienst ELEM, B-1050 Brussels, Belgium
- ¹⁴ Department of Physics and Laboratory for Particle Physics and Cosmology, Harvard University, Cambridge, MA 02138, USA
- ¹⁵ Dept. of Physics, Massachusetts Institute of Technology, Cambridge, MA 02139, USA
- ¹⁶ Dept. of Physics and Institute for Global Prominent Research, Chiba University, Chiba 263-8522, Japan

7

- ¹⁷ Department of Physics, Loyola University Chicago, Chicago, IL 60660, USA
- ¹⁸ Dept. of Physics and Astronomy, University of Canterbury, Private Bag 4800, Christchurch, New Zealand
- ¹⁹ Dept. of Physics, University of Maryland, College Park, MD 20742, USA
- ²⁰ Dept. of Astronomy, Ohio State University, Columbus, OH 43210, USA
- ²¹ Dept. of Physics and Center for Cosmology and Astro-Particle Physics, Ohio State University, Columbus, OH 43210, USA
- ²² Niels Bohr Institute, University of Copenhagen, DK-2100 Copenhagen, Denmark
- ²³ Dept. of Physics, TU Dortmund University, D-44221 Dortmund, Germany
- ²⁴ Dept. of Physics and Astronomy, Michigan State University, East Lansing, MI 48824, USA
- ²⁵ Dept. of Physics, University of Alberta, Edmonton, Alberta, Canada T6G 2E1
- ²⁴¹ ²⁶ Erlangen Centre for Astroparticle Physics, Friedrich-Alexander-Universität Erlangen-Nürnberg, D-91058 Erlangen, Germany
- ²⁷ Physik-department, Technische Universität München, D-85748 Garching, Germany
- ²⁸ Département de physique nucléaire et corpusculaire, Université de Genève, CH-1211 Genève, Switzerland
- ²⁹ Dept. of Physics and Astronomy, University of Gent, B-9000 Gent, Belgium
- ²⁴⁵ ³⁰ Dept. of Physics and Astronomy, University of California, Irvine, CA 92697, USA
- ²⁴⁶ ³¹ Karlsruhe Institute of Technology, Institute for Astroparticle Physics, D-76021 Karlsruhe, Germany
- ²⁴⁷ ³² Karlsruhe Institute of Technology, Institute of Experimental Particle Physics, D-76021 Karlsruhe, Germany
- ³³ Dept. of Physics, Engineering Physics, and Astronomy, Queen's University, Kingston, ON K7L 3N6, Canada
- ³⁴ Dept. of Physics and Astronomy, University of Kansas, Lawrence, KS 66045, USA
- ³⁵ Department of Physics and Astronomy, UCLA, Los Angeles, CA 90095, USA
- ³⁶ Department of Physics, Mercer University, Macon, GA 31207-0001, USA
- ³⁷ Dept. of Astronomy, University of Wisconsin–Madison, Madison, WI 53706, USA
- 253 ³⁸ Dept. of Physics and Wisconsin IceCube Particle Astrophysics Center, University of Wisconsin–Madison, Madison, WI 53706, USA
- ³⁹ Institute of Physics, University of Mainz, Staudinger Weg 7, D-55099 Mainz, Germany
- ⁴⁰ Department of Physics, Marquette University, Milwaukee, WI, 53201, USA
- ⁴¹ Institut für Kernphysik, Westfälische Wilhelms-Universität Münster, D-48149 Münster, Germany
- ⁴² Bartol Research Institute and Dept. of Physics and Astronomy, University of Delaware, Newark, DE 19716, USA
- ⁴³ Dept. of Physics, Yale University, New Haven, CT 06520, USA
- ⁴⁴ Dept. of Physics, University of Oxford, Parks Road, Oxford OX1 3PU, UK
- ⁴⁵ Dept. of Physics, Drexel University, 3141 Chestnut Street, Philadelphia, PA 19104, USA
- ⁴⁶ Physics Department, South Dakota School of Mines and Technology, Rapid City, SD 57701, USA
- ⁴⁷ Dept. of Physics, University of Wisconsin, River Falls, WI 54022, USA
- ⁴⁸ Dept. of Physics and Astronomy, University of Rochester, Rochester, NY 14627, USA
- ⁴⁹ Department of Physics and Astronomy, University of Utah, Salt Lake City, UT 84112, USA
- ⁵⁰ Oskar Klein Centre and Dept. of Physics, Stockholm University, SE-10691 Stockholm, Sweden
- ⁵¹ Dept. of Physics and Astronomy, Stony Brook University, Stony Brook, NY 11794-3800, USA
- ⁵² Dept. of Physics, Sungkyunkwan University, Suwon 16419, Korea
- ⁵³ Institute of Basic Science, Sungkyunkwan University, Suwon 16419, Korea
- ⁵⁴ Dept. of Physics and Astronomy, University of Alabama, Tuscaloosa, AL 35487, USA
- ⁵⁵ Dept. of Astronomy and Astrophysics, Pennsylvania State University, University Park, PA 16802, USA
- ⁵⁶ Dept. of Physics, Pennsylvania State University, University Park, PA 16802, USA
- ⁵⁷ Dept. of Physics and Astronomy, Uppsala University, Box 516, S-75120 Uppsala, Sweden
- ⁵⁸ Dept. of Physics, University of Wuppertal, D-42119 Wuppertal, Germany
- ⁵⁹ DESY, D-15738 Zeuthen, Germany
- ⁶⁰ Università di Padova, I-35131 Padova, Italy
- ⁶¹ National Research Nuclear University, Moscow Engineering Physics Institute (MEPhI), Moscow 115409, Russia
- ⁶² Earthquake Research Institute, University of Tokyo, Bunkyo, Tokyo 113-0032, Japan
- 278
- 279 *E-mail: analysis@icecube.wisc.edu