

RESEARCH OUTPUTS / RÉSULTATS DE RECHERCHE

Local Exponential Stability of Nonlinear Distributed Parameter Systems

HASTIR, Anthony; WINKIN, Joseph; Dochain, Denis

Published in:
39th Benelux Meeting on Systems and Control

Publication date:
2020

Document Version
Peer reviewed version

[Link to publication](#)

Citation for pulished version (HARVARD):
HASTIR, A, WINKIN, J & Dochain, D 2020, Local Exponential Stability of Nonlinear Distributed Parameter Systems: Application to a Nonisothermal Tubular Reactor. in *39th Benelux Meeting on Systems and Control*.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Local exponential stability of nonlinear distributed parameter systems: Application to a nonisothermal tubular reactor

Anthony Hastir*, Joseph J. Winkin* and Denis Dochain†

* Namur Institute for Complex Systems (naXys) and Department of Mathematics, University of Namur, Rempart de la vierge 8, B-5000 Namur, Belgium

† Institute of Information and Communication Technologies, Electronics and Applied Mathematics (ICTEAM),

Université Catholique de Louvain, Avenue Georges Lemaitre 4-6, B-1348 Louvain-La-Neuve, Belgium

anthony.hastir@unamur.be, joseph.winkin@unamur.be, denis.dochain@uclouvain.be

Deducing exponential stability of an equilibrium of a nonlinear distributed parameter, i.e. infinite-dimensional, system on the basis of the stability of a linear approximation of it is in general quite challenging. Some of the existing theories, see e.g. [4, 1], rely on the Fréchet differentiability of the nonlinear semigroup generated by the nonlinear operator dynamics. However, checking Fréchet differentiability for nonlinear operators defined on an infinite-dimensional space is difficult or even impossible if these are unbounded. In many cases, the general theory cannot be applied and a case-by-case study has to be performed by working directly on the semigroup instead of its generator.

The approach that is proposed here is based on an adapted concept of Fréchet differentiability which takes different spaces and norms into account. This is called the (Y, X) -Fréchet differentiability, where X is the state (Hilbert) space and Y is an auxiliary space chosen to handle more easily norm-inequalities when working in infinite-dimensions (typically L^∞ or Sobolev spaces $(H^p, p \in \mathbb{N}_0)$, which are all multiplicative algebras). The systems that are considered here are governed by the following abstract ODE:

$$\dot{x}(t) = Ax(t) + N(x(t)), \quad x(0) = x_0, \quad (1)$$

where $A : D(A) \subset X \rightarrow X$ and $N : D(N) \subset X \rightarrow X$ are linear and nonlinear operators, respectively. Let $x^e \in D(A) \cap D(N)$ be an equilibrium of (1).

Definition : Let $(Y, \|\cdot\|_Y)$ be an infinite-dimensional (possibly Banach) space such that $D(A) \cap D(N) \subset Y \subset X$. The operator N is called (Y, X) -Fréchet differentiable at x^e if there exists a bounded linear operator $dN(x^e) : X \rightarrow X$ such that for all $h \in D(A) \cap D(N)$, $N(x^e + h) - N(x^e) = dN(x^e)h + R(x^e, h)$ where $\lim_{\|h\|_Y \rightarrow 0} \|R(x^e, h)\|_X / \|h\|_X = 0$.

This allows more easily checkable adapted Fréchet differentiability conditions, provided that local exponential stability of the equilibrium of (1) holds in a weaker sense, see [3]. Based on this new concept, our approach to deduce exponential stability of the equilibrium of (1) can be summarized as in Figure 1. Let $(S(t))_{t \geq 0}$ be the nonlinear semigroup generated by the operator $A + N$ on X . The standard concept of Fréchet differentiability is needed for $(S(t))_{t \geq 0}$ on Y , with $(T_{x^e}(t))_{t \geq 0}$ as Fréchet derivative, the linear semigroup generated by the Gâteaux derivative $A + dN(x^e)$ of $A + N$. After showing that $(T_{x^e}(t))_{t \geq 0}$ satisfies some Lyapunov-type stability condition and that it is exponentially stable on X , the new concept of (Y, X) -Fréchet differentiability pops up to make the connection between Y and X , in order to deduce




Figure 1: Illustration of the new theoretical framework.

local exponential stability of the equilibrium of (1) on X , see [3]. Local means here that the Y -norm of the initial condition has to be small instead of its X -norm.

Our theoretical approach is illustrated on a nonisothermal axial dispersion tubular reactor which exhibits different numbers of equilibria depending on the diffusion coefficients: see [2] for the existence and multiplicity of the equilibria. Our main result states that, in the case of only one equilibrium profile, the latter is locally exponentially stable for the nonlinear system governing the dynamics. Moreover, in the case where the reactor exhibits three equilibria, local exponential bistability is established, that is, the pattern "stable – unstable – stable" is highlighted, see [3].

Acknowledgment

This research was conducted with the financial support of F.R.S-FNRS. Anthony Hastir is a FNRS Research Fellow under the grant FC 29535. The scientific responsibility rests with its authors.

References

- [1] R. Al Jamal and K. Morris, 2018. "Linearized stability of partial differential equations with application to stabilization of the kuramotosivashinsky equation", SIAM Journal on Control and Optimization 56 (1), 120 – 147
- [2] A. Hastir, F. Lamoline, J.J. Winkin and D. Dochain, 2019. "Analysis of the existence of equilibrium profiles in nonisothermal axial dispersion tubular reactors", IEEE Transactions on Automatic Control, to appear
- [3] A. Hastir, J.J. Winkin and D. Dochain, 2019. "Exponential stability of nonlinear infinite-dimensional systems: application to nonisothermal axial dispersion tubular reactors", submitted
- [4] N. Kato, 1995. "A principle of linearized stability for nonlinear evolution equations", Transactions of the American Mathematical Society 347 (8), 2851 – 2868.