



Ocular surface response in contact lens wearers

Aurore Muselier, Maxime Souchier, B. Mathieu, Stéphane Grégoire, F. Nicot, Françoise Baudouin, Lionel Bretilon, Alain Bron, Catherine Creuzot Garcher

► **To cite this version:**

Aurore Muselier, Maxime Souchier, B. Mathieu, Stéphane Grégoire, F. Nicot, et al.. Ocular surface response in contact lens wearers. Association for Research in Vision and Ophthalmology annual congress, May 2009, Fort Lauderdale, Florida, United States. 51, 2010, Investigative Ophthalmology and Visual Science. <hal-01136997>

HAL Id: hal-01136997

<https://hal.archives-ouvertes.fr/hal-01136997>

Submitted on 30 Mar 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Ocular Surface Response in Contact Lens Wearers

A. Muselier, Jr.¹, M. Souchier¹, B. Mathieu¹, S. Grégoire², F. Nicot¹, F. Baudouin³, L. Bretillon²,
A. M. Bron^{1,2} and C. Creuzot-Garcher^{1,2}

¹Department of Ophthalmology, University Hospital, Dijon, France

²Eye & Nutrition Research Group, INRA, University of Burgundy, Dijon, France

³Laboratory of Toxicology, Faculty of Pharmaceutical and Biological Sciences, University of Paris-5 René Descartes, Dijon, France

Commercial Relationships: A. Muselier, Jr., None; M. Souchier, None; B. Mathieu, None; S. Grégoire, None; F. Nicot, None; F. Baudouin, None; L. Bretillon, None; A.M. Bron, None; C. Creuzot-Garcher, None.

Support: None.

Abstract

Purpose: To evaluate the ocular surface changes induced by contact lens wear.

Methods: Data of 7 tolerant contact lens wearers, 7 non-tolerant contact lens wearers and 7 healthy subjects were collected. Every patient underwent a thorough ophthalmic examination and tear osmolarity test, conjunctival impression cytology and meibomian lipid sampling. Symptoms, Break-Up Time (BUT), tear osmolarity, conjunctival expression of human leucocyte antigen (HLA) DR and meibomian fatty acid composition were evaluated.

Results: Twenty one subjects (8 men and 13 women) were included (42 eyes). Symptoms reported by tolerant and non-tolerant contact lens wearers were as follows: superficial foreign body sensation (29% vs 21%), burning (14% vs 21%) and ocular dryness (28% vs 78%). No symptoms were reported by healthy subjects. There was no significant difference in tear osmolarity ($p=0.63$) and meibomian fatty acid composition ($p=0.97$) in both groups. Break-up time was significantly lower in tolerant ($p<0.05$) and non-tolerant ($p<0.0001$) contact lens wearers than in healthy subjects. The HLA-DR expression was significantly higher in non-tolerant (50%) contact lens wearers than in healthy subjects (16%) ($p<0.01$).

Conclusions: Contact lens wear is responsible for ocular surface alterations which lead to intolerance. These patterns are very similar to those reported in dry eye symptoms.

Keywords: contact lens • cornea: tears/tear film/dry eye • lipids