



Comment on "The use of BRET to study receptor-protein interactions"

Ralf Jockers

► **To cite this version:**

Ralf Jockers. Comment on "The use of BRET to study receptor-protein interactions". *Frontiers in Endocrinology, Frontiers*, 2014, pp.3. <10.3389/fendo.2014.00003>. <hal-01085046>

HAL Id: hal-01085046

<https://hal.archives-ouvertes.fr/hal-01085046>

Submitted on 20 Nov 2014

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Comment on “The use of BRET to study receptor-protein interactions”

Ralf Jockers^{1,2,3*}

¹ U1016, INSERM, Institut Cochin, Paris, France

² CNRS UMR 8104, Paris, France

³ University of Paris Descartes, Paris, France

*Correspondence: ralf.jockers@inserm.fr

Edited by:

Pierre De Meyts, Hagedorn Research Institute, Denmark

Keywords: BRET, high throughput screening assays, GPCR heteromers, PPI inhibitors, protein–protein interactions

This Research Topic assembles for the first time a comprehensive selection of articles (mini review, review articles, original research, and opinion articles) on the bioluminescence resonance energy transfer (BRET) technology. BRET is a natural phenomenon known in several marine organisms. It relies on the non-radiative transfer of energy from an appropriate energy donor to an energy acceptor, provided that both are located at a distance lower than 10 nm [see Ref. (1) for a basic and theoretical introduction]. BRET has been first applied to the detection of protein–protein interactions (PPIs) in 1999 (2). Since then, the technology constantly evolved by combining new donor and acceptor couples and developing various BRET assay formats as discussed in the Mini Review article of De et al. (3).

Many BRET assays have been developed to study the oligomerization of seven-transmembrane-spanning G protein-coupled receptors (GPCRs). The BRET technology provides an attractive way to study this phenomenon in intact cells without the need of solubilization of receptors from their natural membrane environment. The overwhelming majority of BRET-based studies conclude that GPCRs do indeed exist as dimers or higher-order oligomers when transfected into cells at physiological levels. Three of the articles of this Research Topic discuss BRET assays that have been developed to properly address the specificity of BRET signals obtained upon expression of different GPCRs (1, 4, 5). The current consensus confirms that several different BRET assays are needed to evaluate the specificity of BRET signals. The precise role of each of these assays remains a source of controversy in the field (4). Further refinement of BRET control experiments and application of new techniques like single-molecule measurements and functional *in vivo* studies are likely to provide new insights to the existence and physiological relevance of GPCR oligomerization. Not surprisingly, GPCR oligomerization is the main issue of four articles of this Research Topic (4–7) ranging from studies on class A and B GPCR homo- and heteromers using BRET or alternative approaches like time-resolved FRET measurements.

Apart from performing the proper control experiments, another important issue in the BRET field concerns the proper interpretation of stimulus-induced BRET signals. In the context of GPCR oligomerization, agonist-induced BRET signals can be generated by an agonist-driven oligomerization or agonist-induced conformational changes within preassembled oligomers. Discrimination between these two possibilities is not trivial but has become

possible due to the development of BRET donor saturation experiments in the absence and presence of receptor stimulation (8, 9). This issue, to which BRET has made a significant contribution, is obviously of general importance for the field of PPIs. This is illustrated by the articles describing the interaction of the protease-activated receptor 1 and 2 with its cognate G proteins as described by Ayoub and Pin (10) and Ayoub et al. (11).

The BRET technology has been extended toward other receptor families like tyrosine kinase receptors and cytokine receptors. This diversification demonstrates the general feature of this technique. These studies did not only address the question of receptor oligomerization but also monitored the real-time interaction of receptors with various effector molecules such as Grb2, PTP1-B, PLC- γ 1, etc. This important aspect is discussed in the review article of Siddiqui et al. (12). Receptor–effector interactions have been also monitored by BRET for two privileged GPCR interacting partners, heterotrimeric G proteins and β -arrestins, as documented in articles of this Research Topic (6, 10, 11, 13).

A more recent application of BRET concerns the development of biosensors to monitor downstream events of cellular signaling like the generation of second messengers and activation of intracellular kinases. These sensors are typically composed of the energy donor and acceptor separated by an assay-specific domain that changes its conformation upon second messenger binding or phosphorylation, thus modifying the position of the donor and acceptor and consequently the BRET signal. Similar sensors have been developed for FRET applications, which served as source of inspiration for the development of the BRET sensors. The articles from Salahpour et al. (13) and Xu et al. (2) describe the design and validation of cAMP and ERK sensors.

The high reproducibility of BRET and the robustness of the measurements make BRET an interesting option for the design of high throughput screening assays. Two applications are discussed in this Research Topic. The first concerns the design of an assay for the identification of compounds that specifically activate GPCR heteromers (6) based on the recruitment of β -arrestin to GPCR heteromers. The second case concerns the identification of PPI inhibitors (14).

Taken together, this Research Topic provides an illustrative overview of the principles and applications of the BRET technology that should be of interest for any scientist interested in monitoring PPI in intact cells.

ACKNOWLEDGMENTS

The author thanks Erika Cecon (University of Sao Paulo, Brazil) for comments. This work was supported by grants from the Agence National pour la Recherche (Grants ANR RPIB 2012 "MED-HET-REC-2," the "Who am I?" laboratory of excellence No. ANR-11-LABX-0071 funded by the French Government through its "Investments for the Future" program operated by ANR under grant No. ANR-11-IDEX-0005-01), "Fondation Recherche Médicale" (Grant FRM DEQ20130326503), and ARC (Grants ARC N°5051, ARC N°N° SFI20121205906).

REFERENCES

1. Drinovec L, Kubale V, Nøhr Larsen J, Vrecl M. Mathematical models for quantitative assessment of bioluminescence resonance energy transfer: application to seven transmembrane receptors oligomerization. *Front Endocrin* (2012) **3**:104. doi:10.3389/fendo.2012.00104
2. Xu Y, Piston DW, Johnson CH. A bioluminescence resonance energy transfer (BRET) system: application to interacting circadian clock proteins. *Proc Natl Acad Sci U S A* (1999) **96**:151–6. doi:10.1073/pnas.96.1.151
3. De A, Jasani A, Arora R, Gambhir SS. Evolution of BRET biosensors from live cell to tissue-scale in vivo imaging. *Front Endocrinol* (2013) **4**:131. doi:10.3389/fendo.2013.00131
4. Felce JH, Davis SJ. Unraveling receptor stoichiometry using BRET. *Front Endocrin* (2012) **3**:86. doi:10.3389/fendo.2012.00086
5. Roed SN, Orgaard A, Jorgensen R, Meyts PD. Receptor oligomerization in family B1 of G-protein-coupled receptors: focus on BRET investigations and the link between GPCR oligomerization and binding cooperativity. *Front Endocrin* (2012) **3**:62. doi:10.3389/fendo.2012.00062
6. Johnstone EKM, Pflieger KDG. Receptor-Heteromer Investigation Technology and its application using BRET. *Front Endocrin* (2012) **3**:101. doi:10.3389/fendo.2012.00101
7. Cottet M, Faklaris O, Maurel D, Scholler P, Doumazane E, Trinquet E, et al. BRET and time-resolved FRET strategy to study GPCR oligomerization: from cell lines toward native tissues. *Front Endocrin* (2012) **3**:92. doi:10.3389/fendo.2012.00092
8. Couturier C, Jockers R. Activation of leptin receptor by a ligand-induced conformational change of constitutive receptor dimers. *J Biol Chem* (2003) **278**:26604–11. doi:10.1074/jbc.M302002200
9. Mercier JF, Salahpour A, Angers S, Breit A, Bouvier M. Quantitative assessment of beta 1 and beta 2-adrenergic receptor homo and hetero-dimerization by bioluminescence resonance energy transfer. *J Biol Chem* (2002) **277**:44925–31. doi:10.1074/jbc.M205767200
10. Ayoub MA, Pin J-P. Interaction of protease-activated receptor 2 with G proteins and β -arrestin 1 studied by bioluminescence resonance energy transfer. *Front Endocrinol* (2013) **4**:196. doi:10.3389/fendo.2013.00196
11. Ayoub MA, Al-Senaïdy A, Pin J-P. Receptor-G protein interaction studied by bioluminescence resonance energy transfer: lessons from protease-activated receptor 1. *Front Endocrin* (2012) **3**:82. doi:10.3389/fendo.2012.00082
12. Siddiqui S, Cong WN, Daimon CM, Martin B, Maudsley S. BRET biosensor analysis of receptor tyrosine kinase functionality. *Front Endocrinol* (2013) **4**:46. doi:10.3389/fendo.2013.00046
13. Salahpour A, Espinoza S, Masri B, Lam V, Barak LS, Gainetdinov RR. BRET biosensors to study GPCR biology, pharmacology, and signal transduction. *Front Endocrin* (2012) **3**:105. doi:10.3389/fendo.2012.00105
14. Couturier C, Deprez B. Setting up a bioluminescence resonance energy transfer high throughput screening assay to search for protein/protein interaction inhibitors in mammalian cells. *Front Endocrin* (2012) **3**:100. doi:10.3389/fendo.2012.00100

Received: 08 January 2014; accepted: 08 January 2014; published online: 22 January 2014.

Citation: Jockers R (2014) Comment on "The use of BRET to study receptor-protein interactions". *Front. Endocrinol.* **5**:3. doi: 10.3389/fendo.2014.00003

This article was submitted to *Molecular and Structural Endocrinology*, a section of the journal *Frontiers in Endocrinology*.

Copyright © 2014 Jockers. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.