



UNIVERSITY OF LEEDS

This is a repository copy of *Chromatic index of graphs with no cycle with a unique chord*.

White Rose Research Online URL for this paper:

<http://eprints.whiterose.ac.uk/74348/>

Article:

Machado, RCS, de Figueiredo, CMH and Vuskovic, K (2010) Chromatic index of graphs with no cycle with a unique chord. *Theoretical Computer Science*, 411 (7-9). 1221 - 1234 . ISSN 0304-3975

<https://doi.org/10.1016/j.tcs.2009.12.018>

Reuse

See Attached

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

Chromatic index of graphs with no cycle with a unique chord

R. C. S. Machado^{*†}, C. M. H. de Figueiredo^{*}, K. Vušković[‡]

February 8, 2009

Abstract

The class \mathcal{C} of graphs that do not contain a cycle with a unique chord was recently studied by Trotignon and Vušković [26], who proved strong structure results for these graphs. In the present paper we investigate how these structure results can be applied to solve the edge-colouring problem in the class. We give computational complexity results for the edge-colouring problem restricted to \mathcal{C} and to the subclass \mathcal{C}' composed of the graphs of \mathcal{C} that do not have a 4-hole. We show that it is NP-complete to determine whether the chromatic index of a graph is equal to its maximum degree when the input is restricted to regular graphs of \mathcal{C} with fixed degree $\Delta \geq 3$. For the subclass \mathcal{C}' , we establish a dichotomy: if the maximum degree is $\Delta = 3$, the edge-colouring problem is NP-complete, whereas, if $\Delta \neq 3$, the only graphs for which the chromatic index exceeds the maximum degree are the odd order cycle-graphs and the odd order complete graphs, a characterization that solves edge-colouring problem in polynomial time. We determine two subclasses of graphs in \mathcal{C}' of maximum degree 3 for which edge-colouring is polynomial. Finally, we remark that a consequence of one of our proofs is that edge-colouring is NP-complete for r -regular tripartite graphs of degree $\Delta \geq 3$, for $r \geq 3$.

Keywords: cycle with a unique chord, decomposition, recognition, Petersen graph, Heawood graph, edge-colouring.

1 Motivation

Let $G = (V, E)$ be a simple graph. The degree of a vertex v in G is denoted $\deg_G(v)$, and the maximum degree of a vertex in G is denoted $\Delta(G)$. An

^{*}COPPE, Universidade Federal do Rio de Janeiro, Rio de Janeiro, RJ, Brazil. E-mail: {raphael, celina}@cos.ufrj.br.

[†]Instituto Nacional de Metrologia Normalização e Qualidade Industrial

[‡]School of Computing, University of Leeds, Leeds LS2 9JT, United Kingdom. Email: vuskovi@comp.leeds.ac.uk. Partially supported by EPSRC grant EP/C518225/1 and Serbian Ministry for Science and Technological Development grant 144015G.

edge-colouring of G is a function $\pi : E \rightarrow \mathbf{C}$ such that no two adjacent edges receive the same colour $c \in \mathbf{C}$. If $\mathbf{C} = \{1, 2, \dots, k\}$, we say that π is a *k-edge-colouring*. The *chromatic index* of G , denoted by $\chi'(G)$, is the least k for which G has a *k-edge-colouring*.

Vizing's theorem [27] states that $\chi'(G) = \Delta(G)$ or $\Delta(G) + 1$, defining the *classification problem*: graphs with $\chi'(G) = \Delta(G)$ are said to be *Class 1*, while graphs with $\chi'(G) = \Delta(G) + 1$ are said to be *Class 2*. The *edge-colouring problem* or *chromatic index problem* is the problem of determining the chromatic index of a graph. Edge-colouring is a challenging topic in graph theory and the complexity of the problem is unknown for several important well studied classes. Edge-colouring is NP-complete for regular graphs [13, 18] of degree $\Delta \geq 3$. The problem is NP-complete also for the following classes [5]:

- r -regular comparability (hence perfect) graphs, for $r \geq 3$;
- r -regular line graphs of bipartite graphs (hence line graphs and clique graphs), for $r \geq 3$;
- r -regular k -hole-free graphs, for $r \geq 3, k \geq 3$;
- cubic graphs of girth k , for $k \geq 4$.

Graph classes for which edge-colouring is polynomially solvable include the following:

- bipartite graphs [14];
- split-indifference graphs [21];
- series-parallel graphs (hence outerplanar) [14];
- k -outerplanar graphs [2], for $k \geq 1$.

The complexity of edge-colouring is unknown for several well-studied strong structured graph classes, for which only partial results have been reported, such as cographs [1], join graphs [10, 11, 19], cobipartite graphs [19], planar graphs [24, 28], chordal graphs, and several subclasses of chordal graphs such as indifference graphs [8], split graphs [7] and interval graphs [3].

Given a graph F , we say that a graph G *contains* F if graph F is isomorphic to an induced subgraph of G . A graph G is *F-free* if G does not contain F . A *cycle* C in a graph G is a sequence of vertices $v_1 v_2 \dots v_n v_1$, that are distinct except for the first and the last vertex, such that for $i = 1, \dots, n - 1$, $v_i v_{i+1}$ is an edge and $v_n v_1$ is an edge – we call these edges *the edges of C*. An edge of G with both endvertices in a cycle C is called a *chord* of C if it is not an edge of C . One can similarly define a path and a chord of a path. A *hole* is a chordless cycle of length at least four and an ℓ -*hole* is a hole of length ℓ . A *triangle* is a cycle of length 3 and a *square* is a 4-hole. A graph is called a *cycle-graph* (resp. *path-graph*) if it has a cycle (resp. path) that contains every vertex and has no chord.

Trotignon and Vušković [26] studied the class \mathcal{C} of graphs that do not contain a cycle with a unique chord. The main motivation to investigate this class was to find a structure theorem for it, a kind of result which is not very frequent in the literature. Basically, this structure result states that every graph in \mathcal{C} can be built starting from a restricted set of basic graphs and applying a series of known “gluing” operations. Another interesting property of this class is that it belongs to the family of the χ -bounded graphs, introduced by Gyárfás [12] as a natural extension of perfect graphs. A family of graphs \mathcal{G} is χ -bounded with χ -binding function f if, for every induced subgraph G' of $G \in \mathcal{G}$, $\chi(G') \leq f(\omega(G'))$, where $\chi(G')$ denotes the chromatic number of G' and $\omega(G')$ denotes the size of a maximum clique in G' . The research in this area is mainly devoted to understanding for what choices of forbidden induced subgraphs, the resulting family of graphs is χ -bounded, see [23] for a survey. Note that perfect graphs are a χ -bounded family of graphs with χ -binding function $f(x) = x$, and perfect graphs are characterized by excluding odd holes and their complements. Also, by Vizing’s Theorem, the class of line graphs of simple graphs is a χ -bounded family with χ -binding function $f(x) = x + 1$ (this special upper bound is known as the *Vizing bound*) and line graphs are characterized by nine forbidden induced subgraphs [29]. The class \mathcal{C} is also χ -bounded with the Vizing bound [26]. Also in [26] the following results are obtained for graphs in \mathcal{C} : an $\mathcal{O}(nm)$ algorithm for optimal vertex-colouring, an $\mathcal{O}(n + m)$ algorithm for maximum clique, an $\mathcal{O}(nm)$ recognition algorithm, and the NP-completeness of the maximum stable set problem.

In this paper we consider the complexity of determining the chromatic index of graphs in \mathcal{C} . We also investigate the subclasses obtained from \mathcal{C} by forbidding 4-holes and/or 6-holes. Tables 1 and 2 summarize the main results achieved in this work:

Class	$\Delta = 3$	$\Delta \geq 4$	regular
graphs of \mathcal{C}	NP-complete	NP-complete	NP-complete
4-hole-free graphs of \mathcal{C}	NP-complete	Polynomial	Polynomial
6-hole-free graphs of \mathcal{C}	NP-complete	NP-complete	NP-complete
{4-hole,6-hole}-free graphs of \mathcal{C}	Polynomial	Polynomial	Polynomial

Table 1: Complexity dichotomy for edge-colouring in the class of graphs with no cycle with unique chord.

Class	$k \leq 2$	$k \geq 3$
k -partite graphs	Polynomial	NP-complete

Table 2: Complexity dichotomy for edge-colouring in the class of multipartite graphs.

The results of Tables 1 and 2 show that, even for graph classes with strong structure and powerful decompositions, the edge-colouring problem may be difficult.

The class initially investigated in this work is the class \mathcal{C} of graphs with no cycle with a unique chord. Each non-basic graph in this class can be decomposed [26] by special cutsets: 1-cutsets, proper 2-cutsets or proper 1-joins. We prove that edge-colouring is NP-complete for graphs in \mathcal{C} . We consider, then, a subclass $\mathcal{C}' \subset \mathcal{C}$ whose graphs are the graphs in \mathcal{C} that do not have a 4-hole. By forbidding 4-holes we avoid decompositions by joins, which are difficult to deal with in edge-colouring [1, 10, 11]. That is, each non-basic graph in \mathcal{C}' can be decomposed of 1-cutsets and proper 2-cutsets. For this class \mathcal{C}' we establish a dichotomy: edge-colouring is NP-complete for graphs in \mathcal{C}' with maximum degree 3 and polynomial for graphs in \mathcal{C}' with maximum degree **not** 3. We determine also a necessary condition for a graph $G \in \mathcal{C}'$ of maximum degree 3 to be Class 2. This condition is having graph P^* – a subgraph of the Petersen graph – as a basic block in the decomposition tree. As a consequence, if both 4-holes and 6-holes are forbidden, the chromatic index of graphs with no cycle with unique chord can be determined in polynomial time. The results achieved in this work have connections with other areas of research in edge-colouring, as we describe in the following four observations.

The first observation refers to the complexity dichotomy result found for class \mathcal{C}' . This dichotomy presents great interest since, to the best of our knowledge, this is the first class for which edge-colouring is NP-complete for graphs with a given fixed maximum degree Δ and is polynomial for graphs with maximum degree $\Delta' > \Delta$, as the reader may verify in the NP-completeness results reviewed in the beginning of the present section. Class \mathcal{C}' is, also, the first interesting graph class for which edge-colouring is NP-complete in general, but is polynomial when restricted to regular graphs.

The second observation refers to a conjecture of Chetwynd and Hilton. An important tool to identify Class 2 graphs is the concept of *overfullness* [9]. A graph $G = (V, E)$ is *overfull* if $|E| > \Delta(G)\lfloor |V|/2 \rfloor$, and G is *subgraph overfull* if it has a subgraph of same maximum degree which is overfull.

Subgraph-overfull graphs are [9] Class 2, and it can be verified in polynomial time whether a graph is subgraph overfull [20]. For some graph classes, being subgraph overfull is equivalent to being Class 2. Examples of such classes are graphs with universal vertex [22], complete multipartite graphs [17] and split graphs with odd maximum degree [7]. A conjecture of Chetwynd and Hilton [16] states that a graph $G = (V, E)$ with $\Delta(G) > |V|/3$ is Class 2 if and only if it is subgraph overfull. In fact, for most graph classes for which the edge-colouring problem can be solved in polynomial time, the equivalence “Class 2 = Subgraph Overfull” holds. We observe that the majority of these classes is composed of graphs whose maximum degree are large – always larger than one third of the number of vertices. So, for these graph classes, the equivalence “Class 2 = Subgraph Overfull” – and the consequent polynomial time algorithm for the edge-colouring problem – would be a direct consequence of the Subgraph Overfull Conjecture, in case of its validity. In this sense, the class \mathcal{C}' investigated in this work is of great interest: for graphs in \mathcal{C}' there is no bound on the relation “number of vertices over maximum degree”, yet, if the maximum degree is not 3, it holds the equivalence “Class 2 = Subgraph Overfull”. So, the class of the graphs in \mathcal{C}' with maximum degree not 3 is a class of graphs which do not fit the assumptions of Subgraph Overfull Conjecture, but for which edge-colouring is still solvable in polynomial time since the Class 2 graphs in \mathcal{C}' are Subgraph Overfull.

The third observation is related to the study of *snarks* [25]. A *snark* is a cubic bridgeless graph with chromatic index 4. In order to avoid trivial (easy) cases, snarks are commonly restricted to have girth 5 or more and not to contain three edges whose deletion results in a disconnected graph, each of whose components is non-trivial. The study of snarks is closely related to the Four Colour Theorem. By the result of Proposition 8, the only non-trivial snark which has **no** cycle with unique chord is the Petersen graph.

Finally, the fourth observation refers to the problem of determining the chromatic index of a k -partite graph, that is, a graph whose vertices can be partitioned into k stable sets. The problem is known to be polynomial [14, 17] for $k = 2$ and for complete multipartite graphs. However, there is no explicit result in the literature regarding the complexity of determining the chromatic index of a k -partite graph for $k \geq 3$. From the proof of Theorem 2 we can observe that edge-colouring is NP-complete for k -partite r -regular graphs, for each $k \geq 3$, $r \geq 3$.

The remainder of the paper is organized as follows. In Section 2, we prove NP-completeness results regarding edge-colouring in the classes \mathcal{C} and

\mathcal{C}' . In Section 3, we review known results on the structure of graphs in \mathcal{C} and obtain stronger structure results for graphs in \mathcal{C}' . In Section 4 we show how to determine in polynomial time the chromatic index of a graph in \mathcal{C}' with maximum degree $\Delta \geq 4$. In Section 5 we further investigate graphs in \mathcal{C}' with maximum degree 3: we show that edge-colouring can be solved in polynomial time if the inputs are restricted to regular graphs of \mathcal{C}' and to 6-hole-free graphs of \mathcal{C}' .

2 NP-completeness results

In this section, we state NP-completeness results on the edge-colouring problem restricted to the class \mathcal{C} of graphs that do not contain a cycle with a unique chord and to the class \mathcal{C}' composed of the graphs in \mathcal{C} that do not contain a 4-hole. First, we prove that edge-colouring is NP-complete for regular graphs of \mathcal{C} with fixed degree $\Delta \geq 3$. We observe that it can be shown that the construction of Cai and Ellis [5] which proves the NP-completeness of r -regular k -hole-free graphs, for $r \geq 3$ and $k \neq 4$, creates a graph with no cycle with a unique chord. Nevertheless, in the present section, we give a simpler construction. Second, we prove that edge-colouring is NP-complete for graphs in \mathcal{C}' with maximum degree $\Delta = 3$. For the proof of this second result, we construct a replacement graph which is not present in any edge-colouring NP-completeness proof we could find in the literature.

We use the term $\text{CHRIND}(P)$ to denote the problem of determining the chromatic index restricted to graph inputs with property P . For example, $\text{CHRIND}(\text{graph of } \mathcal{C})$ denotes the following problem:

INSTANCE: a graph G of \mathcal{C} .
 QUESTION: is $\chi'(G) = \Delta(G)$?

The following theorem [13, 18] establishes the NP-completeness of determining the chromatic index of Δ -regular graphs of fixed degree Δ at least 3:

Theorem 1. ([13, 18]) *For each $\Delta \geq 3$, $\text{CHRIND}(\Delta\text{-regular graph})$ is NP-complete.*

Please refer to Figure 1. Graph Q_n , for $n \geq 3$, is obtained from the complete bipartite graph $K_{n,n}$ by removing an edge xy , by adding new pendant vertices u and v , and by adding pendant edges ux and vy . Graph Q'_n is obtained from Q_n by identifying vertices u and v into a vertex w . Observe that Q'_n is a graph of maximum degree n , and has $2n + 1$ vertices and $n^2 + 1$ edges. So, Q'_n is overfull and, hence [9], Class 2. Lemma 1




Figure 1: NP-complete gadget Q_n and graph Q'_n .

investigates the properties of graph Q_n , which is used as “gadget” in the NP-completeness proof of Theorem 2.

Lemma 1. *Graph Q_n is n -edge-colourable, and in any n -edge-colouring of Q_n , edges ux and vy receive the same colour.*

Proof:

We use the notation from Figure 1. First, we exhibit an n -edge-colouring of Q_n . Denote by x_0, \dots, x_{n-1} (resp. y_0, \dots, y_{n-1}) the vertices of Q_n which belong to the same partition as x (resp. y), where $x = x_0$ (resp. $y = y_0$). An n -edge-colouring of Q_n is constructed as follows: just let the colour of edge $x_i y_j$ be $(i + j \bmod n) + 1$ and let the colour of edges $x_0 u$ and $y_0 v$ be 1.

Now we prove that, in any n -edge-colouring of Q_n , edges ux and vy have the same colour. Suppose there is an n -edge-colouring π of Q_n where ux and vy have different colours. Consider the graph $Q'_n = (V', E')$ obtained from Q_n , by contracting vertices u and v into vertex w . Then we can construct an n -edge-colouring π' of Q'_n by setting $\pi'(e) = \pi(e)$ if $e \in E' \setminus \{ux, vy\}$, $\pi'(wx) = \pi(ux)$ and $\pi'(wy) = \pi(vy)$, which is a contradiction to the fact that Q'_n is Class 2. \square

We prove in Theorem 2 the NP-completeness of edge-colouring regular graphs that do not contain a cycle with a unique chord for each fixed degree $\Delta \geq 3$.

Theorem 2. *For each $\Delta \geq 3$, $\text{CHRIND}(\Delta\text{-regular graph in } \mathcal{C})$ is NP-complete.*

Proof:

Let $G = (V, E)$ be an input of the NP-complete problem $\text{CHRIND}(\Delta\text{-regular graph})$. Now, let G' be the graph obtained from G by removing each edge $pq \in E$ and adding a copy of Q_Δ , identifying vertices u and v of Q_Δ with vertices p and q of G . For each edge pq of G , denote H_{pq} the subgraph of G' isomorphic to Q_Δ whose pendant vertices are p and q . Observe that G' is also Δ -regular.

Claim 1: G' can be constructed in polynomial time from G . In fact, we make one substitution – by a copy of Q_Δ – for each edge of G , so that the construction time is linear on the number of edges of G .

Claim 2: if G is Δ -edge-colourable, then so is G' . Let π be a Δ -edge-colouring of G . We construct a Δ -edge-colouring π' of G' in the following way: for each edge pq of G , let the edges of H_{pq} in G' be coloured in such a way that the pendant edges have the colour $\pi(pq)$ – this colouring exists and is described by Lemma 1.

Claim 3: if G' is Δ -edge-colourable, then so is G . Let π' be a Δ -edge-colouring of G' . We construct a Δ -edge-colouring π of G as follows: let the colour in π of each edge pq of G be equal to the colour in π' of the pendant edges of H_{pq} (by Lemma 1, these two pendant edges must receive the same colour).

Claim 4: $G' \in \mathcal{C}$. Suppose G' has a cycle C with a unique chord $\alpha\beta$. Observe that, by construction, every edge of G' – and, in particular, chord $\alpha\beta$ – has both endvertices in the same copy of Q_Δ . Denote by $H_{p'q'}$ this copy and observe that cycle C , when restricted to $H_{p'q'}$, is a path between p' and q' , and that $\alpha\beta$ is a unique chord of this path. But there is no path with unique chord between the pendant vertices of Q_Δ , so that we have a contradiction. \square

Observe that graph G' in the proof of Theorem 2 is tripartite with vertex tripartition (P_1, P_2, P_3) determined as follows:

- P_1 is the set whose elements are the original vertices of G and the vertices denoted y_1, \dots, y_Δ in each copy of Q_Δ ;
- P_2 is the set whose elements are the vertices denoted x_0 and y_0 in each copy of Q_Δ ;
- P_3 is the set whose elements are the vertices denoted x_1, \dots, x_Δ in each copy of Q_Δ .

So, the following result holds:

Theorem 3. *For each $k \geq 3, \Delta \geq 3$, $\text{CHRIND}(\Delta\text{-regular } k\text{-partite graph})$ is NP-complete.*

We emphasize that \mathcal{C} is a class with strong structure [26], yet, it is NP-complete for edge-colouring. We manage in Section 4 to define a subclass of \mathcal{C} where edge-colouring is solvable in polynomial time. Consider the class \mathcal{C}' as the subset of the graphs of \mathcal{C} that do not contain a square. The structure of graphs in \mathcal{C}' is stronger than that of graphs in \mathcal{C} , and is described in detail in Section 3. Yet, the edge-colouring problem is still NP-complete for inputs in \mathcal{C}' , as we prove next in Theorem 4. We recall that the proof of Cai and Ellis [5] for the NP-completeness of edge-colouring cubic square-free graphs generates a graph which **has** a cycle with unique a chord. In addition, remark that the gadget Q_Δ used in the proof of the NP-completeness of edge-colouring graphs with no cycle with unique chord **has** a square. So, we need an alternative construction, which is based on the gadget \tilde{P} shown in Figure 2. Graph \tilde{P} is constructed in such a way that the identification of its pendant vertices generates a graph isomorphic to P^* , the graph obtained from the Petersen graph by removing one vertex. Graph P^* is a non-overfull Class 2 graph [6, 15]. The properties of \tilde{P} with respect to edge-colouring are described in Lemma 2.




Figure 2: 3-edge-colouring of gadget graph \tilde{P} .

Lemma 2. *Graph \tilde{P} is 3-edge-colourable, and in any 3-edge-colouring of \tilde{P} , the edges ux and vy receive the same colour.*

Proof:

Figure 2 shows a 3-edge-colouring of \tilde{P} – observe that edges ux and vy receive the same colour.

The fact that edges ux and vy always receive the same colour is a consequence of P^* being Class 2. The proof is similar to that of Lemma 1, except

that gadget \tilde{P} is used instead of Q_Δ . \square

Theorem 4. *CHRIND(graph in \mathcal{C}' with maximum degree 3) is NP-complete.*

Proof:

The proof is similar to that of Theorem 2, except that $\Delta = 3$ and gadget \tilde{P} is used instead of Q_Δ . \square

Observe that the graph G' constructed in the proof of Theorem 4 is not regular. In fact, as we prove in Section 5.1, the edge-colouring problem can be solved in polynomial time if the input is restricted to cubic graphs of \mathcal{C}' .

3 Structure of graphs in \mathcal{C} and \mathcal{C}'

The goal of the present section is to review structure results for the graphs in \mathcal{C} and obtain stronger results for the subclass \mathcal{C}' . These results are used in Section 4 to edge-colour the graphs in \mathcal{C}' with maximum degree at least 4. In the present section we review the results of Trotignon and Vušković [26] on the structure of graphs in \mathcal{C} and obtain stronger results for graphs in \mathcal{C}' .

Let \mathcal{C} be the class of the graphs that do not contain a cycle with a unique chord and let \mathcal{C}' be the class of the graphs of \mathcal{C} that do not contain a square. Trotignon and Vušković give a decomposition result [26] for graphs in \mathcal{C} and graphs in \mathcal{C}' in the following form: every graph in \mathcal{C} or in \mathcal{C}' either belongs to a basic class or has a cutset. Before we can state these decomposition theorems, we define the basic graphs and the cutsets used in the decomposition.

The *Petersen graph* is the graph on vertices $\{a_1, \dots, a_5, b_1, \dots, b_5\}$ so that both $a_1a_2a_3a_4a_5a_1$ and $b_1b_2b_3b_4b_5b_1$ are chordless cycles, and such that the only edges between some a_i and some b_i are $a_1b_1, a_2b_4, a_3b_2, a_4b_5, a_5b_3$. We denote by P the Petersen graph and by P^* the graph obtained from P by removal of one vertex. Observe that $P \in \mathcal{C}$.

The *Heawood graph* is a cubic bipartite graph on vertices $\{a_1, \dots, a_{14}\}$ so that $a_1a_2 \dots a_{14}a_1$ is a cycle, and such that the only other edges are $a_1a_{10}, a_2a_7, a_3a_{12}, a_4a_9, a_5a_{14}, a_6a_{11}, a_8a_{13}$. We denote by H the Heawood graph and by H^* the graph obtained from H by removal of one vertex. Observe that $H \in \mathcal{C}$.

A graph is *strongly 2-bipartite* if it is square-free and bipartite with bipartition (X, Y) where every vertex in X has degree 2 and every vertex in Y has degree at least 3. A strongly 2-bipartite graph is in \mathcal{C} because any chord of a cycle is an edge between two vertices of degree at least three, so every cycle in a strongly 2-bipartite graph is chordless.

For the purposes of this work, a graph G is called *basic*¹ if

1. G is a complete graph, a cycle-graph with at least five vertices, a strongly 2-bipartite graph, or an induced subgraph of the Petersen graph or of the Heawood graph; and
2. G has no 1-cutset, proper 2-cutset or proper 1-join (all defined next).

We denote by \mathcal{C}_B be the set of the basic graphs. Observe that $\mathcal{C}_B \subseteq \mathcal{C}$.

A *cutset* S of a connected graph G is a set of elements, vertices and/or edges, whose removal disconnects G . A decomposition of a graph is the removal of a cutset to obtain smaller graphs, called the *blocks* of the decompositions, by possibly adding some nodes and edges to connected components of $G \setminus S$. The goal of decomposing a graph is trying to solve a problem on the whole graph by combining the solutions on the blocks. For a graph $G = (V, E)$ and $V' \subseteq V$, $G[V']$ denotes the subgraph of G induced by V' . The following cutsets are used in the known decomposition theorems of the class \mathcal{C} [26]:

- A *1-cutset* of a connected graph $G = (V, E)$ is a node v such that V can be partitioned into sets X , Y and $\{v\}$, so that there is no edge between X and Y . We say that (X, Y, v) is a *split* of this 1-cutset.
- A *proper 2-cutset* of a connected graph $G = (V, E)$ is a pair of non-adjacent nodes a, b , both of degree at least three, such that V can be partitioned into sets X , Y and $\{a, b\}$ so that: $|X| \geq 2$, $|Y| \geq 2$; there is no edge between X and Y , and both $G[X \cup \{a, b\}]$ and $G[Y \cup \{a, b\}]$ contain an ab -path. We say that (X, Y, a, b) is a *split* of this proper 2-cutset.
- A *1-join* of a graph $G = (V, E)$ is a partition of V into sets X and Y such that there exist sets A, B satisfying:
 - $\emptyset \neq A \subseteq X$, $\emptyset \neq B \subseteq Y$;
 - $|X| \geq 2$ and $|Y| \geq 2$;
 - there are all possible edges between A and B ;
 - there is no other edge between X and Y .

¹By the definition of [26], a basic graph is not, in general, indecomposable. However, our slightly different definition helps simplifying some of our proofs.

We say that (X, Y, A, B) is a *split* of this 1-join.

A *proper 1-join* is a 1-join such that A and B are stable sets of G of size at least two.

We can now state a decomposition result for graphs in \mathcal{C} :

Theorem 5. (Trotignon and Vušković [26]) *If $G \in \mathcal{C}$ is connected then either $G \in \mathcal{C}_B$ or G has a 1-cutset, or a proper 2-cutset, or a proper 1-join.*

The *block* G_X (resp. G_Y) of a graph G with respect to a 1-cutset with split (X, Y, v) is $G[X \cup \{v\}]$ (resp. $G[Y \cup \{v\}]$).

The *block* G_X (resp. G_Y) of a graph G with respect to a 1-join with split (X, Y, A, B) is the graph obtained by taking $G[X]$ (resp. $G[Y]$) and adding a node y complete to A (resp. x complete to B). Nodes x, y are called *markers* of their respective blocks.

The *blocks* G_X and G_Y of a graph G with respect to a proper 2-cutset with split (X, Y, a, b) are defined as follows. If there exists a node c of G such that $N_G(c) = \{a, b\}$, then let $G_X = G[X \cup \{a, b, c\}]$ and $G_Y = G[Y \cup \{a, b, c\}]$. Otherwise, block G_X (resp. G_Y) is the graph obtained by taking $G[X \cup \{a, b\}]$ (resp. $G[Y \cup \{a, b\}]$) and adding a new node c adjacent to a, b . Node c is called the *marker* of the block G_X (resp. G_Y).

The blocks with respect to 1-cutsets, proper 2-cutsets and proper 1-joins are constructed in such a way that they remain in \mathcal{C} , as shown by Lemma 3.

Lemma 3. (Trotignon and Vušković [26]) *Let G_X and G_Y be the blocks of decomposition of G w.r.t. a 1-cutset, a proper 1-join or a proper 2-cutset. Then $G \in \mathcal{C}$ if and only if $G_X \in \mathcal{C}$ and $G_Y \in \mathcal{C}$.*

We reviewed results that show how to decompose a graph of \mathcal{C} into basic blocks: Theorem 5 states that each graph in \mathcal{C} has a 1-cutset, a proper 2-cutset or a proper 1-join, while Lemma 3 states that the blocks generated with respect to any of these cutsets are still in \mathcal{C} . We now obtain similar results for \mathcal{C}' . These results are not explicit in [26], but they can be obtained as consequences of results in [26] and by making minor modifications in its proofs. As we discuss in the following observation [4], for the goal of edge-colouring, we only need to consider the **biconnected** graphs of \mathcal{C}' .

Observation 1. *Let G be a connected graph with a 1-cutset with split (X, Y, v) . The chromatic index of G is $\chi'(G) = \max\{\chi'(G_X), \chi'(G_Y), \Delta(G)\}$.*

By Observation 1, if both blocks G_X and G_Y are $\Delta(G)$ -edge-colourable, then so is G . That is, once we know the chromatic index of the biconnected components of a graph, it is easy to determine the chromatic index of the whole graph. So, we may focus our investigation on the biconnected graphs of \mathcal{C}' .

Theorem 6. (Trotignon and Vušković [26]) *If $G \in \mathcal{C}'$ is biconnected, then either $G \in \mathcal{C}_B$ or G has a proper 2-cutset.*

Theorem 6 is an immediate consequence of Theorem 5: since G has no 4-hole, G cannot have a proper 1-join, and since G is biconnected, G cannot have a 1-cutset.

Next, in Lemma 4, we show that the blocks of decomposition of a biconnected graph of \mathcal{C}' w.r.t. a proper 2-cutset, are also biconnected graphs of \mathcal{C}' . The proof of Lemma 4 is similar to that of Lemma 5.2 of [26]. For the sake of completeness, the proof, which uses the result of Theorem 7 below, is included here.

Theorem 7. (Trotignon and Vušković [26]) *Let $G \in \mathcal{C}$ be a connected graph. If G contains a triangle then either G is a complete graph, or some vertex of the maximal clique that contains this triangle is a 1-cutset of G .*

Lemma 4. *Let $G \in \mathcal{C}'$ be a biconnected graph and let (X, Y, a, b) be a split of a proper 2-cutset of G . Then both G_X and G_Y are biconnected graphs of \mathcal{C}' .*

Proof:

We first prove that G is triangle-free. Suppose G contains a triangle. Then, by Theorem 7, either G is a complete graph, which contradicts the assumption that G has a proper 2-cutset, or G has a 1-cutset, which contradicts the assumption that G is biconnected. So G is triangle-free, and hence by construction, both of the blocks G_X and G_Y are triangle-free.

Now we show that G_X and G_Y are square-free. Suppose w.l.o.g. that G_X contains a square C . Since G is square-free, C contains the marker node M , which is not a real node of G , and $C = MazbM$, for some node $z \in X$. Since M is not a real node of G , we have $\deg_G(z) > 2$, otherwise, z would be a marker of G_X . Let z' be a neighbor of z distinct of a and b . Since G is triangle-free, z' is not adjacent to a nor b . Since z is not a 1-cutset, there exists a path P in $G[X \cup \{a, b\}]$ from z' to $\{a, b\}$. We choose z' and P subject to the minimality of P . So, w.l.o.g., $z'Pa$ is a chordless path. Note that b is not adjacent to the neighbor of a along P because G is triangle-free and square-free, so that z is the unique common neighbor of a and b in G .

So, by the minimality of P , vertex b does not have a neighbor in P . Now let Q be a path from a to b whose interior is in Y . So, $bzz'PaQb$ is a cycle of G with a unique chord (namely az), contradicting the assumption that $G \in \mathcal{C}$.

By Lemma 3, G_X and G_Y both belong to \mathcal{C} , and since G_X and G_Y are both square-free, it follows that G_X and G_Y both belong to \mathcal{C}' .

Finally we show that G_X and G_Y are biconnected. Suppose w.l.o.g. that G_X has a 1-cutset with split (A, B, v) . Since G is biconnected and $G[X \cup \{a, b\}]$ contains an ab -path, we have that $v \neq M$, where M is the marker of G_X . Suppose $v = a$. Then, w.l.o.g., $b \in B$, and $(A, B \cup Y, a)$ is a split of a 1-cutset of G , with possibly M removed from $B \cup Y$, if M is not a real node of G , contradicting the assumption that G is biconnected. So $v \neq a$ and by symmetry $v \neq b$. So $v \in X \setminus \{M\}$. W.l.o.g. $\{a, b, M\} \subset B$. Then $(A, B \cup Y, v)$ is a split of a 1-cutset of G , with possibly M removed from $B \cup Y$ if M is not a real node of G , contradicting the assumption that G is biconnected. \square

Observe that Lemma 3 is somehow stronger than Lemma 4. While Lemma 3 states that a graph is in \mathcal{C} **if and only if** the blocks with respect to any cutset are also in \mathcal{C} , Lemma 4 establishes only one direction: **if** a graph is a biconnected graph of \mathcal{C}' , **then** the blocks with respect to any cutset are also biconnected graph of \mathcal{C}' . For our goal of edge-colouring, there is no need of establishing the “only if” part. Anyway, it is possible to verify that, if both blocks G_X and G_Y generated with respect to a proper 2-cutset of a graph G are biconnected graphs of \mathcal{C}' , then G itself is a biconnected graph of \mathcal{C}' .

Next lemma shows that every non-basic biconnected graph in \mathcal{C}' has a decomposition such that one of the blocks is basic.

Lemma 5. *Every biconnected graph $G \in \mathcal{C}' \setminus \mathcal{C}_B$ has a proper 2-cutset such that one of the blocks of decomposition is basic.*

Proof:

By Theorem 6 G has a proper 2-cutset. Consider all possible 2-cutset decompositions of G and pick a proper 2-cutset S that has a block of decomposition B whose size is smallest possible. By Lemma 4, $B \in \mathcal{C}'$ and is biconnected. So by Theorem 6, either B has a proper 2-cutset or it is basic. We now show that in fact B must be basic.

Let (X, Y, a, b) be a split w.r.t. S . Let M be the marker node of G_X , and assume w.l.o.g. that $B = G_X$. Suppose G_X has a proper 2-cutset with split (X_1, X_2, u, v) . By minimality of $B = G_X$, $\{a, b\} \neq \{u, v\}$. Assume w.l.o.g. $b \notin \{u, v\}$. Note that since $\deg_{G_X}(u) \geq 3$ and $\deg_{G_X}(v) \geq 3$, it follows that $M \notin \{u, v\}$. Suppose $a \notin \{u, v\}$. Then w.l.o.g. $\{a, b, M\} \subseteq X_1$,

and hence $(X_1 \cup Y, X_2, u, v)$, with M removed if M is not a real node of G , is a proper 2-cutset of G whose block of decomposition G_{X_2} is smaller than G_X , contradicting the minimality of $G_X = B$. Therefore $a \in \{u, v\}$. Then w.l.o.g. $\{b, M\} \subseteq X_1$, and hence $(X_1 \cup Y, X_2, u, v)$, with M removed if M is not a real node of G , is a proper 2-cutset of G whose block of decomposition G_{X_2} is smaller than G_X , contradicting the minimality of $G_X = B$. Therefore G_X does not have a proper 2-cutset, and hence it is basic. \square

4 Chromatic index of graphs in \mathcal{C}' with maximum degree at least 4

The first NP-completeness result of Section 2 proves that edge-colouring is difficult for the graphs in \mathcal{C} . We consider, further, the subclass \mathcal{C}' and verify that the edge-colouring problem is still NP-complete when restricted to \mathcal{C}' . In the present section we apply the structure results of Section 3 to show that edge-colouring graphs in \mathcal{C}' of maximum degree $\Delta \geq 4$ is polynomial by establishing that the only Class 2 graphs in \mathcal{C}' are the odd order complete graphs. Remark that the NP-completeness holds only for 3-edge-colouring restricted to graphs in \mathcal{C}' with maximum degree 3.

We describe, next, the technique applied to edge-colour a graph in \mathcal{C}' by combining edge-colourings of its blocks with respect to a proper 2-cutset. Observe that the fact that a graph F is isomorphic to a block B obtained from a proper 2-cutset decomposition of G does **not** imply that G contains F : possibly B is constructed by the addition of a marker vertex. This is illustrated in the example of Figure 3, where G is P^* -free, yet, graph P^* appears as a block with respect to a proper 2-cutset of G .

The reader will also observe that it is not necessary that a block of decomposition of G is $\Delta(G)$ -edge-colourable in order that G itself is $\Delta(G)$ -edge-colourable: graph G in Figure 3 is 3-edge-colourable, while block P^* is not. This is an important observation: possibly, the edges adjacent to a marker vertex of a block of decomposition are not real edges of the original graph, or are already coloured by an edge-colouring of another block, so that these edges do not need to be coloured.

Observation 2. *Consider a graph $G \in \mathcal{C}'$ with the following properties:*

- (X, Y, a, b) is a split of a proper 2-cutset of G ;
- \tilde{G}_Y is obtained from G_Y by removing its marker if this marker is not a real vertex of G ;




Figure 3: Example of decomposition with respect to a proper 2-cutset $\{a, b\}$. Observe that the marker vertices and their incident edges – identified by dashed lines – do not belong to the original graph.

- $\tilde{\pi}_Y$ is a $\Delta(G)$ -edge-colouring of \tilde{G}_Y ;
- F_a (resp. F_b) is the set of the colours in $\{1, 2, \dots, \Delta\}$ not used by $\tilde{\pi}_Y$ in any edge of \tilde{G}_Y incident to a (resp. b).

If there exists a $\Delta(G)$ -edge-colouring π_X of $G_X \setminus M$, where M is the marker vertex of G_X , such that each colour used in an edge incident to a (resp. b) is in F_a (resp. F_b), then G is Δ -edge-colourable.

The above observation shows that, in order to extend a $\Delta(G)$ -edge-colouring of \tilde{G}_Y to a $\Delta(G)$ -edge-colouring of G , one must colour the edges of $G_X \setminus M$ in such a way that the colours of the edges incident to a (resp. b) are not used at the edges of \tilde{G}_Y incident to a (resp. b). This guarantees that we create no conflicts. Moreover, there is no need to colour the edges incident to the marker M of G_X : if this marker is a vertex of G , its incident edges are already coloured by $\tilde{\pi}$, otherwise, these edges are not real edges of G . In the example of Figure 3, we exhibit a 3-edge-colouring $\tilde{\pi}_Y$ of \tilde{G}_Y . In the notation of Observation 2, $F_a = \{2, 3\}$ and $F_b = \{2, 3\}$. We exhibit, also, a 3-edge-colouring of $G_X \setminus M$ such that the colours of the edges incident to a are $\{2, 3\} \subset F_a$ and the colours of the edges incident to b are $\{2, 3\} \subset F_b$.

So, by Observation 2, we can combine colourings $\tilde{\pi}_Y$ and π_X in a 3-edge-colouring of G , as it is done in Figure 3.

Before we proceed and show how to edge-colour graphs in \mathcal{C}' with maximum degree $\Delta \geq 4$, we need to introduce some additional tools and concepts. A *partial k -edge-colouring* of a graph $G = (V, E)$ is a colouring of a subset E' of E , that is, a function $\pi : E' \rightarrow \{1, 2, \dots, k\}$ such that no two adjacent edges of E' receive the same colour.

The *set of free-colours* at vertex u with respect to a partial-edge-colouring $\pi : E' \rightarrow \mathbf{C}$ is the set $\mathbf{C} \setminus \pi(\{uv \mid uv \in E'\})$. The *list-edge-colouring* problem is described next. Let $G = (V, E)$ be a graph and let $\mathcal{L} = \{L_e\}_{e \in E}$ be a collection which associates to each edge $e \in E$ a set of colours L_e called the *list* relative to e . It is asked whether there is an edge-colouring π of G such that $\pi(e) \in L_e$ for each edge $e \in E$. Theorem 8 is a result on list-edge-colouring which is applied, in this work, to edge-colour some of our basic graphs: strongly 2-bipartite graphs, Heawood graph and its subgraphs, and cycle-graphs.

Theorem 8. (Borodin, Kostochka, and Woodall [4]) *Let $G = (V, E)$ be a bipartite graph and $\mathcal{L} = \{L_e\}_{e \in E}$ be a collection of lists of colours which associates to each edge $uv \in E$ a list L_{uv} of colours. If, for each edge $uv \in E$, $|L_{uv}| \geq \max\{\deg_G(u), \deg_G(v)\}$, then there is an edge-colouring π of G such that, for each edge $uv \in E$, $\pi(uv) \in L_{uv}$.*

We investigate, now, how to $\Delta(G)$ -edge-colour a graph $G \in \mathcal{C}'$ by combining $\Delta(G)$ -edge-colourings of its blocks with respect to a proper 2-cutset. More precisely, Lemma 6 shows how this can be done if one of the blocks is basic. Subsequently, we obtain, in Theorem 9, a characterization for graphs in \mathcal{C}' of maximum degree at least 4 of its Class 2 graphs which establishes that edge-colouring is polynomial for these graphs.

Lemma 6. *Let $G \in \mathcal{C}'$ be a graph of maximum degree $\Delta \geq 4$ and let (X, Y, a, b) be a split of proper 2-cutset, in such a way that G_X is basic. If G_Y is Δ -edge-colourable, then G is Δ -edge-colourable.*

Proof:

Denote by M the marker vertex of G_X and let \tilde{G}_Y be obtained from G_Y by removing its marker if this marker is not a real vertex of G . Since \tilde{G}_Y is a subgraph of G_Y , graph \tilde{G}_Y is Δ -edge-colourable. Let π_Y be a Δ -edge-colouring of \tilde{G}_Y , i.e. a partial-edge-colouring of G , and let F_a and F_b be the sets of the free colours of a and b , respectively, with respect to the partial edge-colouring π_Y . We show how to extend the partial edge-colouring π_Y to G , as described in Observation 2, that is, by colouring the

edges of $G_X \setminus M$. Since a and b are not adjacent, G_X is not a complete graph. Moreover, the block G_X cannot be isomorphic to the Petersen graph or to the Heawood graph, because these graphs are cubic and G_X has a marker vertex M of degree 2. So, G_X is isomorphic to an induced subgraph of P^* , or to an induced subgraph of H^* , or to a strongly 2-bipartite graph, or to a cycle-graph.

Case 1. G_X is a strongly 2-bipartite graph.

Since $\deg_{G_X}(M) = 2$, vertex M belongs to the bipartition of G_X whose vertices have degree 2. So, vertices a and b belong to the bipartition of G_X whose vertices have degree larger than 2, and $|F_a| \geq \deg_{G_X \setminus M}(a) \geq 2$ and $|F_b| \geq \deg_{G_X \setminus M}(b) \geq 2$. Associate to each edge of $G_X \setminus M$ incident to a (resp. b) a list of colours equal to F_a (resp. F_b). To each of the other edges of $G_X \setminus M$, associate list $\{1, \dots, \Delta\}$. Now, to each edge uv of $G_X \setminus M$, a list of colours is associated whose size is not smaller than $\max\{\deg_{G_X \setminus M}(u), \deg_{G_X \setminus M}(v)\}$ and, by Theorem 8, there is an edge-colouring π_1 of $G_X \setminus M$ from these lists. Finally, set $\pi := \pi_1$ for the edges of $G_X \setminus M$.

Case 2. G_X is a cycle-graph.

In this case, $G_X \setminus M$ is a path. Denote the vertices of $G_X \setminus M$ by $a = x_1, x_2, \dots, x_k = b$, in such a way that $x_1x_2\dots x_k$ is a path. We now show that $k \geq 4$. Since a and b are not adjacent, $k \geq 3$. Suppose that $k = 3$. If M is a real node of G , then G_X is a square and it is an induced subgraph of G , contradicting the assumption that G is square-free. So M is not a real node of G , and hence $G_X \setminus M = X$. But, then, $|X| = 1$, contradicting the definition of a proper 2-cutset. Therefore, $k \geq 4$.

Observe that there is at least one colour c_α in F_a and one colour c_β in F_b . We construct a 3-edge-colouring π of $C_X \setminus M$ by setting $\pi(x_1x_2) := c_\alpha$ and $\pi(x_{k-1}x_k) := c_\beta$, and by colouring the other edges of $G_X \setminus M$ as follows. If $k = 4$, let $\pi(x_2x_3)$ be some colour in $\{1, 2, 3\} \setminus \{c_\alpha, c_\beta\}$, which is clearly a non-empty set. If $k \geq 5$, let $\mathcal{L}_2 = \{L_2, L_3, \dots, L_{k-2}\}$ be a collection which associates to each edge $x_i x_{i+1}$ a list of colours L_i such that:

- $L_i = \{1, 2, 3\} \setminus \{c_\alpha\}$, for $i = 2, 3, \dots, k - 3$, and
- $L_{k-2} = \{1, 2, 3\} \setminus \{c_\beta\}$.

Observe that $G_X \setminus \{M, a, b\}$ is a path, hence bipartite of maximum degree 2, and that $|L_i| \geq 2$ for each $i = 2, \dots, k - 2$, so that by, Theorem 8, there is an edge-colouring π_2 of $G_X \setminus \{M, a, b\}$ from the lists \mathcal{L}_2 . Moreover, this colouring creates no conflicts with the colours c_α of x_1x_2 and c_β of $x_{k-1}x_k$, so that we can set $\pi := \pi_2$ for edges $x_2x_3, x_3x_4, \dots, x_{k-2}x_{k-1}$.

Case 3. G_X is an induced subgraph of the Heawood graph.

Observe that a and b have only M as common neighbor in G_X , otherwise G_X has a square (recall that Heawood graph is square-free). We construct a 4-edge-colouring of $G_X \setminus M$. Denote the neighbors of a (resp. b) in $G_X \setminus M$ by a_1, \dots, a_x (resp. b_1, \dots, b_y), where $x = \deg_{G_X \setminus M}(a)$ (resp. $y = \deg_{G_X \setminus M}(b)$). Note that $x, y \in \{1, 2\}$. Observe that F_a (resp. F_b) contains at least x (resp. y) colours, which we denote by c_{a_1}, \dots, c_{a_x} (resp. $c_{b_1}, c_{b_2}, \dots, c_{b_y}$). Set the colour π of edge aa_i (resp. bb_j), for $i = 1, \dots, x$ (resp. for $j = 1, \dots, y$), to c_{a_i} (resp. c_{b_j}). Now, associate to each edge incident to a_i and different from aa_i a list of colours $\{1, 2, 3, 4\} \setminus \{c_{a_i}\}$. Similarly, associate to each edge incident to b_j and different of bb_j a list of colours $\{1, 2, 3, 4\} \setminus \{c_{b_j}\}$. Finally, associate to each of the other edges of $G_X \setminus \{M, a, b\}$ the list of colours $\{1, 2, 3, 4\}$. Observe that $G_X \setminus \{M, a, b\}$ is bipartite of maximum degree at most 3 and that each of the lists has 3 or 4 colours, so that, by Theorem 8, there is an edge-colouring π_3 of $G_X \setminus \{M, a, b\}$ from these lists, and we set $\pi := \pi_3$ for the edges of $G_Z \setminus M$.

Case 4.a: $G_X = P^*$.

Observe that there are at least two colours c_{a_1}, c_{a_2} in F_a and two colours c_{b_1}, c_{b_2} in F_b , and that exactly one of the following three possibilities holds:

- $|\{c_{a_1}, c_{a_2}\} \cap \{c_{b_1}, c_{b_2}\}| = 0$;
- $|\{c_{a_1}, c_{a_2}\} \cap \{c_{b_1}, c_{b_2}\}| = 1$; or
- $|\{c_{a_1}, c_{a_2}\} \cap \{c_{b_1}, c_{b_2}\}| = 2$.

In the three cases, it is possible to extend the Δ -edge-colouring π_Y to G by colouring the edges of $G_X \setminus M$, as it is shown on Figure 4.

Case 4.b: G_X is a proper induced subgraph of P^* .

We need to investigate which are the proper induced subgraphs of P^* . We invite the reader to verify that, except for graph P^{**} shown on the left of Figure 5, each proper induced subgraph of P^* either has a 1-cutset or a proper 2-cutset, and we do not consider it because G_X is assumed basic, or is a cycle-graph, which is already considered in Case 2.

There is only one possible choice for the marker M of $G_X = P^{**}$, in the sense that, for any other choice of marker M' , we have $G_X \setminus M' = G_X \setminus M$. As in Case 4.a, there are at least two colours c_{a_1}, c_{a_2} in F_a and two colours c_{b_1}, c_{b_2} in F_b , and $|\{c_{a_1}, c_{a_2}\} \cap \{c_{b_1}, c_{b_2}\}| = 0, 1$ or 2 . In Figure 5 we exhibit three edge-colourings for $P^{**} \setminus M$, one for each possibility. \square

Using Lemma 6 we can determine in polynomial time the chromatic index of the graphs of \mathcal{C}' , as we show in Theorem 9.




Figure 4: Extending the colouring to the edges of G_X .




Figure 5: Graph P^{**} and three edge-colourings of $P^{**} \setminus M$ subject to each possible free colour restriction.

Theorem 9. *A connected graph $G \in \mathcal{C}'$ of maximum degree $\Delta \geq 4$ is Class 2 if and only if it is an odd order complete graph.*

Proof:

If G is complete, then the theorem clearly holds. So, we prove that every connected graph of \mathcal{C}' that is not complete and whose maximum degree is at least 4 is Class 1. More precisely, we prove, by induction on the number of vertices, that, if Δ is an integer at least 4 and G is a connected graph of \mathcal{C}' with maximum degree $\Delta(G) \leq \Delta$ and is not a complete graph on $\Delta + 1$ vertices, then G is Δ -edge-colourable.

If $G = (V, E)$ has four vertices or less, then $\Delta(G) \leq 3$. Since $\Delta \geq 4$, by Vizing's theorem, graph G is Δ -edge-colourable.

Now, let $G \in \mathcal{C}'$ be a connected graph with k vertices such that $\Delta(G) \leq \Delta$ and G is not a complete graph on $\Delta + 1$ vertices. Assume, by induction

hypothesis, that every connected graph $G' \in \mathcal{C}'$ with $k' < k$ vertices such that $\Delta(G') \leq \Delta$ and G' is not a complete graph on $\Delta + 1$ vertices, is Δ -edge-colourable. By Theorem 6 either G is basic, or G has a 1-cutset, or G is biconnected and has a proper 2-cutset.

Suppose G is basic. If G is strongly 2-bipartite, then G is Δ -edge-colourable because bipartite graphs are Class 1 and $\Delta(G) \leq \Delta$. If G is not strongly 2-bipartite, then $\Delta(G) \leq \Delta - 1$ and G is Δ -edge-colourable by Vizing's theorem.

Now, suppose G has a 1-cutset with split (X, Y, v) . Note that blocks of decomposition G_X and G_Y are induced subgraphs of G and hence both belong to \mathcal{C}' . By the induction hypothesis, both G_X and G_Y are Δ -edge-colourable, and hence by Observation 1, graph G is Δ -edge-colourable.

Finally, suppose G is biconnected and has a proper 2-cutset. Let (X, Y, a, b) be a split of a proper 2-cutset such that block G_X is basic (note that such a cutset exists by Lemma 5). By Lemma 4 block G_Y is in \mathcal{C}' . By the induction hypothesis, block G_Y is Δ -edge-colourable. If $\Delta(G) \leq 3$, then since $\Delta \geq 4$, by Vizing's theorem, G is Δ -edge-colourable. So we may assume that $\Delta(G) \geq 4$, and hence by Lemma 6, G is Δ -edge-colourable. \square

5 Graphs of \mathcal{C}' with maximum degree 3

Class \mathcal{C}' has a stronger structure than \mathcal{C} , yet, edge-colouring problem is NP-complete for inputs in \mathcal{C}' . In fact, the problem is NP-complete for graphs in \mathcal{C}' with maximum degree $\Delta = 3$. In this section, we further investigate graphs in \mathcal{C}' with maximum degree $\Delta = 3$, providing two subclasses for which edge-colouring can be solved in polynomial time: cubic graphs of \mathcal{C}' and 6-hole-free graphs of \mathcal{C}' .

5.1 Cubic graphs of \mathcal{C}'

In the present section, we prove the polynomiality of the edge-colouring problem restricted to cubic graphs of \mathcal{C}' . This is a direct consequence of Lemma 7, which states that every non-biconnected cubic graph is Class 2, and Lemma 8, that states that the Petersen graph is the only biconnected cubic Class 2 graph in \mathcal{C}' .

Lemma 7. *Let G be a connected cubic graph. If G has a 1-cutset, then G is Class 2.*

Proof:

Denote by (X, Y, v) a split of a 1-cutset of G . Observe that v has degree 1 in exactly one of the blocks G_X and G_Y ; assume, w.l.o.g. that this block is G_X . Let G'_X be the graph obtained from G_X by removing vertex v . Observe that G'_X has exactly one vertex of degree 2 and each of the other vertices has degree 3. Since the sum of the degrees of the vertices is even, G'_X has an even number of vertices of degree 3, say n . So, the number of edges in G'_X is $(3n + 2)/2$. Since $3\lfloor(n + 1)/2\rfloor = 3n/2 < (3n + 2)/2$, graph G'_X is overfull, so that G is subgraph-overfull, hence Class 2. \square

Lemma 8. *Let $G \in \mathcal{C}'$ be biconnected graph. If G is cubic, then G is isomorphic to the Petersen graph or to the Heawood graph or is a complete graph on four vertices.*

Proof:

Suppose G is not basic. By Lemma 5, G has a proper 2-cutset such that one of the blocks is basic. Let (X, Y, a, b) be a split of such cutset, in such a way that G_X is basic, and denote by M the marker vertex of G_X . If $\deg_{G_X}(a) = 1$, vertex M is the only neighbor of a and, clearly, is a 1-cutset of G_X . By Lemma 4 G_X is a biconnected graph of \mathcal{C}' . Since G_X is biconnected $\deg_{G_X} \geq 2$. Let a' be a neighbor of a in G_X that is distinct from M . Since $\{M, a, b, a'\}$ cannot induce a square, b is not adjacent to a' , and hence (since G is cubic) a' has two neighbors in $G_X \setminus \{a, b, M\}$. If $\deg_{G_X}(a) = 2$ then $\{a', b\}$ is a proper 2-cutset of G , contradicting the assumption that G_X is basic. Hence $\deg_{G_X}(a) \geq 3$, and by symmetry $\deg_{G_X}(b) \geq 3$. Observe that each of the other vertices – different from a, b and M – has degree $\Delta(G)$. In other words, G_X is a graph with exactly one vertex of degree 2, and each of the other vertices has degree 3. But there is no graph in \mathcal{C}_B with this property, and we have a contradiction to the fact that G_X is basic. So, G is basic and the statement of the proposition clearly holds. \square

Theorem 10. *Let $G \in \mathcal{C}'$ be a connected cubic graph. Then G is Class 1 if and only if G is biconnected and is not isomorphic to the Petersen graph.*

Proof:

If G is not biconnected, then, by Lemma 7, G is Class 2. If G is biconnected, then, by Lemma 8, then G is isomorphic to the Petersen graph or to the Heawood graph or is a complete graph on four vertices. Hence, G is Class 2 if and only if it is isomorphic to the Petersen graph. \square

5.2 6-hole-free graphs of \mathcal{C}'

In the present section, we prove the polynomiality of the edge-colouring problem restricted to 6-hole-free graphs of \mathcal{C}' . This is a consequence of

Lemma 9, a variation for 3-edge-colouring of Lemma 6.

Lemma 9. *Let $G \in \mathcal{C}'$ be a graph of maximum degree at most 3 and (X, Y, a, b) be a split of a proper 2-cutset, in such a way that G_X is basic but not isomorphic to P^* . If G_Y is 3-edge-colourable, then G is 3-edge-colourable.*

Proof:

Assume G_Y is 3-edge-colourable. Denote by M the marker vertex of G_X and let \tilde{G}_Y be obtained from G_Y by removing its marker if this marker is not a real vertex of G . Since \tilde{G}_Y is a subgraph of G_Y , graph \tilde{G}_Y is 3-edge-colourable. Let π_Y be a 3-edge-colouring of \tilde{G}_Y , i.e. a partial-edge-colouring of G , and let F_a and F_b be the sets of the free colours of a and b , respectively, with respect to the partial edge-colouring π_Y . We show how to extend the partial edge-colouring π_Y to G , as described in Observation 2, that is, by colouring the edges of $G_X \setminus M$. Since a and b are not adjacent, G_X is not a complete graph. Moreover, the block G_X cannot be isomorphic to the Petersen graph or to the Heawood graph, because these graphs are cubic and G_X has a marker vertex M of degree 2. Also, by assumption, block G_X is not isomorphic to P^* . So, G_X is isomorphic to a proper induced subgraph of P^* , or to an induced subgraph of H^* , or to a strongly 2-bipartite graph, or to a cycle-graph.

Case 1. G_X is a strongly 2-bipartite graph.

Similar to the Case 1 of the proof of Lemma 6, which also works for $\Delta = 3$.

Case 2. G_X is a cycle-graph.

Similar to the Case 2 of the proof of Lemma 6, where at most three colours are used in the edges of $G_X \setminus M$.

Case 3. G_X is an induced subgraph of H^* .

First, observe that $\deg_{G_X \setminus M}(a) = 2$ and $\deg_{G_X \setminus M}(b) = 2$, otherwise G_X has a decomposition by a 1-cutset or a proper 2-cutset and is not basic. Observe, also, that there are at least two colours c_{a_1}, c_{a_2} in F_a and two colours c_{b_1}, c_{b_2} in F_b , and that $|\{c_{a_1}, c_{a_2}\} \cap \{c_{b_1}, c_{b_2}\}| = 1$ or 2 . We consider each case next.

If $|\{c_{a_1}, c_{a_2}\} \cap \{c_{b_1}, c_{b_2}\}| = 1$, we must exhibit a 3-edge-coloring π of $G_X \setminus M$ such that the free colors at a and b are different. If M is a real node of G , then G_X is an induced subgraph of G , and hence $\Delta(G_X) \leq 3$. If M is not a real node of G , then by definition of proper 2-cutset both a and b have a neighbor in Y , and hence $\Delta(G_X) \leq 3$. So $\Delta(G_X) \leq 3$. Since G_X is bipartite, G_X has a 3-edge-colouring π' . So, let π be the restriction of π' to $G_X \setminus M$.

If $|\{c_{a_1}, c_{a_2}\} \cap \{c_{b_1}, c_{b_2}\}| = 2$, we must exhibit a coloring of $G_X \setminus M$ such that the free colors at a and b are the same. We exhibit these colourings for each possible induced subgraph of the Heawood graph. First, consider the case $G_X = H^*$, whose coloring is given in Figure 6.




Figure 6: A 3-edge-colouring of $H^* \setminus M$ such that the sets of the colours incident to vertex a and vertex b are the same.

Now, observe that each non-basic proper subgraph of H^* is a subgraph of the graph H_1 of Figure 7, which is obtained from H^* by removing a vertex of degree 2. Graph H_2 of Figure 7 is obtained from H_1 by removing one of the four vertices of degree 2 (any choice yields the same graph up to an isomorphism). Finally, the last non-basic proper subgraph of H^* is the graph H_3 of Figure 7. Observe that there is only one possible choice M for




Figure 7: Non-basic proper induced subgraphs of H^*

the marker when $G_X = H_1$, in the sense that, for any other choice \tilde{M} , we have $G_X \setminus \tilde{M} = G_X \setminus M$. If $G_X = H_2$, there are two possible choices M' and M'' for the marker, in the sense that, for any other choice \tilde{M}' , we have $G_X \setminus \tilde{M}' = G_X \setminus M'$ or $G_X \setminus \tilde{M}' = G_X \setminus M''$. We show, in Figure 8, one

edge-colouring of $H_1 \setminus M$, and two edge-colourings of $H_2 \setminus M$, one for each possible choice of marker M . We don't consider here that case $G_X = H_3$




Figure 8: 3-edge-colourings of H_1 and H_2 , for each possible choice of marker.

because H_3 is a strongly 2-bipartite graph, considered in Case 1.

Case 4. G_X is a proper subgraph of P^* .

As we already discussed in Case 4 of Lemma 6, except for graph P^{**} shown on the left of Figure 5, each of the other proper induced subgraphs of P^* either has a 1-cutset or a proper 2-cutset, and we do not consider because G_X is basic, or is a cycle-graph, which are considered in Case 2. There is only one possible choice of marker M_1 for the case $G_X = P^{**}$, in the sense that for any other choice of marker M'_1 , we have $G_X \setminus M'_1 = G_X \setminus M_1$. Observe, also, that there are at least two colours c_{a_1}, c_{a_2} in F_a and two colours c_{b_1}, c_{b_2} in F_b , and that $|\{c_{a_1}, c_{a_2}\} \cap \{c_{b_1}, c_{b_2}\}| = 1$ or 2 . These two possibilities are considered in the first two colourings of Figure 5. \square

Remark that the NP-Complete gadget \tilde{P} of Figure 2 is constructed from P^* . The NP-completeness of edge-colouring graphs in \mathcal{C}' is obtained as a consequence of $P^* \in \mathcal{C}'$. Using Lemma 9, we can prove that if the special graph P^* does not appear as a leaf in the decomposition tree, i.e., as a basic block when we recursively apply the proper 2-cutset decomposition to a biconnected graph $G \in \mathcal{C}'$ of maximum degree 3, then G is Class 1.

Theorem 11. *Let $G \in \mathcal{C}'$ be a connected graph of maximum degree 3. If G does not contain a 6-hole all of whose nodes are of degree 3, then G is Class 1.*

Proof: Assume the theorem does not hold and let G be a counterexample with fewest number of nodes. So G is a connected graph of \mathcal{C}' of maximum degree 3, it does not contain a 6-hole all of whose nodes are of degree 3, and it is not 3-edge-colourable. By Theorem 6 either G is basic, or it has a 1-cutset, or it is biconnected and has a proper 2-cutset.

Suppose G is basic. G cannot be strongly 2-bipartite nor an induced subgraph of Heawood graph, since bipartite graphs are Class 1 [29]. G cannot be a complete graph on four vertices, because such a graph is 3-edge-colourable. G cannot be a hole since it has maximum degree 3. So G must be an induced subgraph of the Petersen graph. G cannot be isomorphic to P nor P^* , because both of these graphs contain a 6-hole all of whose nodes are of degree 3. But all the other induced subgraphs of the Petersen graph are in fact 3-edge-colourable. Therefore G cannot be basic.

Now suppose that G has a 1-cutset with split (X, Y, v) . Note that blocks of decomposition are induced subgraphs of G , and hence both are connected graphs of \mathcal{C}' that do not contain a 6-hole all of whose nodes are of degree 3. If $\Delta(G_X) = 3$ then since G is a minimum counterexample, G_X is 3-edge-colourable. If $\Delta(G_X) \leq 2$ then G_X is 3-edge-colourable by Vizing's Theorem. So G_X is 3-edge-colourable, and similarly so is G_Y . But then by Observation 1, G is also 3-edge-colourable, a contradiction.

Therefore G is biconnected and has a proper 2-cutset. Let (X, Y, a, b) be a split of a proper 2-cutset such that block G_X is basic (note that such a cutset exists by Lemma 5). By Lemma 4 both of the blocks G_X and G_Y are biconnected graphs of \mathcal{C}' . Since the marker node M is of degree 2 in both G_X and G_Y , and $G_X \setminus M$ and $G_Y \setminus M$ are both induced subgraphs of G , it follows that neither G_X nor G_Y can contain a 6-hole all of whose nodes are of degree 3. If M is a real node of G , then G_X and G_Y are both induced subgraphs of G , and hence $\Delta(G_X) \leq 3$ and $\Delta(G_Y) \leq 3$. If M is not a real node of G , then by definition of proper 2-cutset both a and b have a neighbor in both X and Y , and hence $\Delta(G_X) \leq 3$ and $\Delta(G_Y) \leq 3$. Since both G_X and G_Y have fewer nodes than G , it follows either from minimality of counterexample G or by Vizing's Theorem that both G_X and G_Y are 3-edge-colourable. Since G_X does not contain a 6-hole all of whose nodes are of degree 3, G_X is not isomorphic to P^* , and hence by Lemma 9, G is 3-edge-colourable, a contradiction. \square

Corollary 1. *Every connected 6-hole-free graph of \mathcal{C}' with maximum degree 3 is Class 1.*

A natural question in connection with Theorem 12 is whether forbidding 6-holes would make it easier to edge-colour graphs of \mathcal{C}' , and the answer is **no**. By observing graph G' of the proof of Theorem 2, one can easily verify that this graph has no 6-hole, so that the following theorem holds:

Theorem 12. *For each $\Delta \geq 3$, $\text{CHRIND}(\Delta\text{-regular 6-hole-free graph in } \mathcal{C})$ is NP-complete.*

References

- [1] M. M. Barbosa, C. P. de Mello, and J. Meidanis. *Local conditions for edge-colouring of cographs*. Congr. Numer. **133** (1998) 45–55.
- [2] H. L. Bodlaender. *Polynomial algorithms for graph isomorphism and chromatic index on partial k -trees*. J. Algorithms **11** (1990) 631–643.
- [3] V. A. Bojarshinov. *Edge and total colouring of interval graphs*. Discrete Appl. Math. **114** (2001) 23–28.
- [4] O. V. Borodin, A. V. Kostochka, and D. R. Woodall. *List edge and list total colourings of multigraphs*. J. Combin. Theory Ser. B **71** (1997) 184–204.
- [5] L. Cai and J. A. Ellis. *NP-Completeness of edge-colouring some restricted graphs*. Discrete Appl. Math. **30** (1991) 15–27.
- [6] D. Cariolaro and G. Cariolaro. *Colouring the petals of a graph*. Electron. J. Combin. **10** (2003) #R6.
- [7] B. L. Chen, H.-L. Fu, and M. T. Ko. *Total chromatic number and chromatic index of split graphs*. J. Combin. Math. Combin. Comput. **17** (1995) 137–146.
- [8] C. M. H. de Figueiredo, J. Meidanis, C. P. de Mello, and C. Ortiz. *Decompositions for the edge colouring of reduced indifference graphs*. Theoret. Comput. Sci. **297** (2003) 145–155.
- [9] C. M. H. de Figueiredo, J. Meidanis, and C. P. de Mello. *Local conditions for edge-coloring*. J. Combin. Math. Combin. Comput. **32** (2000) 79–91.
- [10] C. De Simone and C. P. de Mello. *Edge-colouring of join graphs*. Theoret. Comput. Sci. **355** (2006) 364–370.
- [11] C. De Simone and A. Galluccio. *Edge-colouring of regular graphs of large degree*. Theoret. Comput. Sci. **389** (2007) 91–99.
- [12] A. Gyárfás. *Problems from the world surrounding perfect graphs*. Zastos. Mat. **19** (1987) 413–441.
- [13] I. Holyer. *The NP-completeness of edge-colouring*. SIAM J. Comput. **10** (1982) 718–720.

- [14] D. S. Johnson. *The NP-completeness column: an ongoing guide*. J. Algorithms **6** (1985) 434–451.
- [15] A. J. W. Hilton and Cheng Zhao. *On the edge-colouring of a graph whose core has maximum degree two*. J. Combin. Math. Combin. Comput. **21** (1996) 97–108.
- [16] A. G. Chetwynd and A. J. W. Hilton. *Star multigraphs with three vertices of maximum degree*. Math. Proc. Cambridge Philos. Soc. **100** (1986) 303–317.
- [17] D. G. Hoffman and C. A. Rodger. *The chromatic index of complete multipartite graphs*. J. Graph Theory **16** (1992) 159–163.
- [18] D. Leven and Z. Galil. *NP-completeness of finding the chromatic index of regular graphs*. J. Algorithms **4** (1983) 35–44.
- [19] R. C. S. Machado and C. M. H. de Figueiredo. *Decompositions for edge-coloring join graphs and cobipartite graphs*. To appear in Discrete Appl. Math.
- [20] T. Niessen. *How to find overfull subgraphs in graphs with large maximum degree. II*. Electron. J. Combin. **8** (2001) #R7.
- [21] C. Ortiz, N. Maculan, and J. L. Szwarcfiter. *Characterizing and edge-colouring split-indifference graphs*. Discrete Appl. Math. **82** (1998) 209–217.
- [22] M. Plantholt. *The chromatic index of graphs with a spanning star*. J. Graph Theory **5** (1981) 45–53.
- [23] B. Randerath and I. Schiermeyer. *Vertex colouring and forbidden subgraphs—a survey*. Graphs Combin. **20** (2004) 1–40.
- [24] D. P. Sanders and Y. Zhao. *Planar graphs of maximum degree seven are class I*. J. Combin. Theory Ser. B **83** (2001) 201–212.
- [25] E. Steffen. *Classifications and characterizations of snarks*. Discrete Math. **188** (1998) 183–203.
- [26] N. Trotignon and K. Vušković. *A structure theorem for graphs with no cycle with a unique chord and its consequences*. To appear in Journal of Graph Theory. (Available online at <http://www.comp.leeds.ac.uk/vuskovi/chord.ps>.)

- [27] V. G. Vizing. *On an estimate of the chromatic class of a p -graph* (in russian). *Diskret. Analiz* **3** (1964) 25–30.
- [28] V. G. Vizing. *Critical graphs with given chromatic index* (in russian). *Diskret. Analiz* **5** (1965) 9–17.
- [29] D. B. West. *Introduction to Graph Theory*. Second edition. Prentice Hall (2001).