

REPRESENTING FINITE CONVEX GEOMETRIES BY RELATIVELY CONVEX SETS

KIRA ADARICHEVA*

ABSTRACT. A closure system with the anti-exchange axiom is called a convex geometry. One geometry is called a sub-geometry of the other if its closed sets form a sublattice in the lattice of closed sets of the other. We prove that convex geometries of relatively convex sets in n -dimensional vector space and their finite sub-geometries satisfy the n -Carousel Rule, which is the strengthening of the n -Carathéodory property. We also find another property, that is similar to the simplex partition property and does not follow from 2-Carousel Rule, which holds in sub-geometries of 2-dimensional geometries of relatively convex sets.

1. INTRODUCTION

A closure system $\mathbf{A} = (A, -)$, i.e. a set A with a closure operator $- : 2^A \rightarrow 2^A$ defined on A , is called a *convex geometry* (see [3]), if it is a zero-closed space (i.e. $\overline{\emptyset} = \emptyset$) and it satisfies *the anti-exchange axiom*, i.e.

$$x \in \overline{X \cup \{y\}} \text{ and } x \notin X \text{ imply that } y \notin \overline{X \cup \{x\}}$$

for all $x \neq y$ in A and all closed $X \subseteq A$.

A convex geometry $\mathbf{A} = (A, -)$ is called finite, if set A is finite.

Very often, a convex geometry can be represented by its collection of closed sets. There is a convenient description of those collections of subsets of a given finite set A , which are, in fact, the closed sets of a convex geometry on A : if $\mathcal{A} \subseteq 2^A$ satisfies

- (1) $\emptyset \in \mathcal{A}$;
- (2) $X \cap Y \in \mathcal{A}$, as soon as $X, Y \in \mathcal{A}$;
- (3) $X \in \mathcal{A}$ and $X \neq A$ implies $X \cup \{a\} \in \mathcal{A}$, for some $a \in A \setminus X$,

then \mathcal{A} represents the collection of closed sets of a convex geometry $\mathbf{A} = (A, \mathcal{A})$.

A reader can be referred to [8],[9] for the further details of combinatorial and lattice-theoretical aspects of finite convex geometries.

For convex geometries $\mathbf{A} = (A, -)$ and $\mathbf{B} = (B, \tau)$, one says that \mathbf{A} is a sub-geometry of \mathbf{B} , if there is a one-to-one map ϕ of closed sets of \mathbf{A} to closed sets of \mathbf{B} such that $\phi(X \cap Y) = \phi(X) \cap \phi(Y)$, and $\phi(\overline{X \cup Y}) = \tau(\phi(X) \cup \phi(Y))$, where $X, Y \subseteq A$, $\overline{X} = X$, $\overline{Y} = Y$. In other words, the lattice of closed subsets of \mathbf{A} is a sublattice of the lattice of closed sets of \mathbf{B} . When geometries \mathbf{A} and \mathbf{B} are defined on the same set $X = A = B$, we also call \mathbf{B} a *strong extension* of \mathbf{A} . Extensions of finite convex geometries were considered in [4] and [3], the more systematic treatment of extensions of finite lattices was given in [15].

Keywords and phrases: convex geometry, relatively convex sets, Carathéodory property.

2000 *Mathematics Subject Classification:* 05B25, 06B15, 06B05, 51D20.

*Stern College for Women, Yeshiva University, 245 Lexington Ave., New York, NY 10016, USA.

Given any class \mathcal{L} of convex geometries, we will call it *universal*, if an arbitrary finite convex geometry is a sub-geometry of some geometry in \mathcal{L} .

One of main results in [3] proves that a specially designed class of convex geometries \mathcal{AL} is universal. Namely, \mathcal{AL} consists of convex geometries of the form $Sp(A)$, each of which is built on a carrier set of an algebraic and dually algebraic lattice A and whose closed sets are all complete lower subsemilattices of A closed with respect to taking joins of non-empty chains. At the same time, a subclass of all *finite* geometries from class \mathcal{AL} cease to be universal, see [2] and [3].

In this paper, we want to consider another conveniently designed class of convex geometries, in fact, even an infinite hierarchy of classes.

Given a set of points A in Euclidean n -dimensional space \mathbb{R}^n , one defines a closure operator $- : 2^A \rightarrow 2^A$ on A as follows: for any $Y \subseteq A$, $\bar{Y} = ch(Y) \cap A$, where ch stands for *the convex hull*. One easily verifies that such an operator satisfies the anti-exchange axiom. Thus, $(A, -)$ is a convex geometry, which also will be denoted as $\mathbf{Co}(\mathbb{R}^n, A)$. We will call such convex geometry *a geometry of relatively convex sets* (assuming that these are convex sets “relative” to A). The convex geometries of relatively convex sets were studied in [13],[7] and [1].

For any geometry $C = \mathbf{Co}(\mathbb{R}^m, A)$, we will call $n \in \mathbb{N}$ *a dimension* of C , if n is the smallest number such that C could be represented as $\mathbf{Co}(\mathbb{R}^n, A)$, for appropriate $A \subseteq \mathbb{R}^n$. In particular, $n \leq m$, and $n \leq p - 1$, if A is a finite non-empty set of cardinality $p > 1$.

Let \mathcal{C}_n be the class of convex geometries of relatively convex sets of dimension $\leq n$, and let \mathcal{C} be the the class of of all convex geometries of relatively convex sets of finite dimension (thus, including \mathcal{C}_n , $n \in \mathbb{N}$, as subclasses). By \mathcal{C}_B we denote a subclass of \mathcal{C} that consists of geometries of convex sets relative to bounded sets, i.e. $\mathbf{Co}(\mathbb{R}^n, A)$, for some n and $A \subseteq B$, where B is a ball in \mathbb{R}^n . By \mathcal{C}_f we denote a subclass if *finite* convex geometries in \mathcal{C} .

It is known that none of \mathcal{C}_n is universal, due to the n -Carathéodory property that holds on any sub-geometry of geometry from \mathcal{C}_n (see, for example, [7]), but fails on any geometry of dimension $n + 1$. We introduce a stronger property called the n -Carousel Rule and show that it holds on sub-geometries of \mathcal{C}_n . It allows to build a series of finite convex geometries C_n such that C_n satisfies the n -Carathéodory property, but cannot be a sub-geometry of any geometry in \mathcal{C}_n . On the other hand, C_n is a sub-geometry of some geometry in \mathcal{C}_{n+1} . We also prove that the so-called Sharp Carousel Rule holds in all sub-geometries in \mathcal{C}_2 , a slight modification of *the simplex partition property* from [14].

It was shown in [7] that *every finite* closure system can be embedded into some geometry in the class \mathcal{C} , in particular, this class is universal for all finite convex geometries. This observation is a direct consequence of deep and complex result proved in [16] that every finite lattice is a sublattice of a finite partition lattice. Thus, class \mathcal{C} can not be considered as specific to finite convex geometries. It is worth noting that the construction in [7] uses convex sets relative to A which is the collection of lines, in particular, A is always an unbounded set.

This leaves the following open questions:

Problem 1.1. *Is class \mathcal{C}_B of geometries of convex sets relative to bounded sets universal? Is the class \mathcal{C}_f of finite geometries of relatively convex sets universal?*

Note that the second question of two is a modification of Problem 3 from [3].

2. CARATHÉODORY PROPERTY AND CAROUSEL RULE

We recall that a convex geometry $(A, -)$ satisfies the n -Carathéodory property, if $x \in \bar{S}$, $S \subseteq A$, implies $x \in \overline{\{a_0, \dots, a_n\}}$ for some $a_0, \dots, a_n \in S$. Besides, a_0 can be taken to be any pre-specified element of S .

Proposition 2.1. ([13, Lemma 3.2],[7, Proposition 25]) *For any $n \in \mathbb{N}$ and $A \subseteq \mathbb{R}^n$, convex geometry $\mathbf{Co}(\mathbb{R}^n, A)$ satisfies the n -Carathéodory property.*

Our aim is to formulate a stronger property, which we call the n -Carousel Rule, extending to arbitrary finite dimensions the 2-Carousel Rule introduced in [5].

Definition 2.2. A convex geometry $(A, -)$ satisfies the n -Carousel Rule, if $x, y \in \bar{S}$, $S \subseteq A$, implies $x \in \overline{\{y, a_1, \dots, a_n\}}$ for some $a_1, \dots, a_n \in S$.

Note that the n -Carathéodory property follows from the n -Carousel Rule. Indeed, if y is chosen among elements of S , and $x \in \bar{S}$, then, according to the n -Carousel Rule, $x \in \overline{\{y, a_1, \dots, a_n\}}$ for some $a_1, \dots, a_n \in S$, which is also a desired conclusion for the n -Carathéodory property.

Lemma 2.3. *For any $n \in \mathbb{N}$ and $A \subseteq \mathbb{R}^n$, convex geometry $\mathbf{Co}(\mathbb{R}^n, A)$ satisfies the n -Carousel Rule.*

Proof. Consider $\mathbf{G} = \mathbf{Co}(\mathbb{R}^n, A)$, and let $x, y \in \bar{S}$, for some $S \subseteq A$.

Due to the n -Carathéodory property, $x \in \{c_0, c_1, \dots, c_n\}$ and $y \in \overline{\{b_0, b_1, \dots, b_n\}}$ for some $c_0, b_0, \dots, c_n, b_n \in S$. In other words, points x, y belong to a convex polytope P in \mathbb{R}^n with the vertices among $c_0, b_0, \dots, c_n, b_n$. Suppose F_1, \dots, F_k are the faces of this polytope, i.e. they are at most $(n-1)$ -dimensional convex polytopes. For arbitrary $y \in P$, we have $P \subseteq \bigcup_{i \leq k} P_i$, where $P_i = ch(y \cup F_i)$, $i = 1, \dots, k$. Hence, $x \in \overline{y \cup F_i}$ for some $i \leq k$. Now, due to the n -Carathéodory property, $x \in \overline{\{y, f_1, \dots, f_n\}}$ for some vertices f_1, \dots, f_n of F_i , which are also elements of S . Thus, the conclusion of the n -Carousel Rule holds. \square

Our next goal is to show that the n -Carousel Rule is preserved on finite subgeometries.

Lemma 2.4. *If geometry \mathbf{H} satisfies the n -Carousel Rule, and \mathbf{G} is a finite subgeometry of \mathbf{H} , then \mathbf{G} satisfies the n -Carousel Rule.*

Proof. Suppose $\mathbf{H} = (H, -)$, $\mathbf{G} = (G, \tau)$ and ϕ is a one-to-one mapping from closed sets of \mathbf{G} to closed sets of \mathbf{H} that preserves the intersection and the closure of finite unions of sets.

Let assume that \mathbf{G} does not satisfy the n -Carousel Rule. It means that, for some $x, y \in G$ and $S \subseteq G$, we have $x, y \in \bar{S}$, but $x \notin \overline{\{y, s_1, \dots, s_n\}}$, for any $s_1, \dots, s_n \in S$. In any finite convex geometry $(A, -)$, for any $a \in A$, the subset $\bar{a} \setminus a$ is closed. Hence, set $X = \bar{x} \setminus x$ is closed in \mathbf{G} . According to our assumption, $\bar{x} \cap \overline{y \cup \bar{s}_1 \cup \dots \cup \bar{s}_n} \subseteq X$, for any $s_1, \dots, s_n \in S$.

Take $x' \in \phi(\bar{x}) \setminus \phi(X)$ and $y' \in \phi(\bar{y})$. Note that $\bar{S} = \bigcup \{\bar{s} : s \in S\}$, hence $S' = \phi(\bar{S}) = \tau(\bigcup \{\phi(\bar{s}) : s \in S\})$. Since $x', y' \in S' = \tau(\bigcup \{\phi(\bar{s}) : s \in S\})$ and \mathbf{H} satisfies the n -Carousel Rule, we have $x' \in \tau(\{y', s'_1, \dots, s'_n\})$, for some $s'_i \in \phi(\bar{s}_i)$, $s_i \in S$. It follows $x' \in \phi(\bar{x}) \cap \tau(\phi(\bar{y}) \cup \phi(\bar{s}_1) \cup \dots \cup \phi(\bar{s}_n)) = \phi(\bar{x}) \cap \phi(\overline{y \cup \bar{s}_1 \cup \dots \cup \bar{s}_n})$, which means $\phi(\bar{x} \cap \overline{y \cup \bar{s}_1 \cup \dots \cup \bar{s}_n}) \not\subseteq \phi(X)$, a contradiction. \square

3. CONVEX GEOMETRIES C_n

Using the n -Carousel Rule, it will not be difficult to built an example of a finite convex geometry that cannot be a sub-geometry of relatively convex sets of dimension $\leq n$.

Consider a point configuration in \mathbb{R}^n that consists of extreme points a_0, \dots, a_n , equivalently, the vertices of a n -dimensional polytope P , and inner points x, y of P . Besides, choose x, y so that x belongs to only one of polytopes $P_i = ch(\{y\} \cup D \setminus a_i)$ and y belongs to only one of polytopes $Q_j = ch(\{x\} \cup D \setminus a_j)$, where $D = \{a_0, \dots, a_n\}$, $i, j \leq n$.

Let $\mathbf{D}_n = \mathbf{Co}(\mathbb{R}^n, D \cup \{x, y\})$.

According to our assumption, $\{y\} \cup D \setminus a_i$ and $\{x\} \cup D \setminus a_j$ are not closed sets in convex geometry \mathbf{D}_n , for some unique $i, j \leq n, i \neq j$.

Consider closure space $\mathbf{C}_n = (D \cup \{x, y\}, \mathcal{D})$, where a family of closed sets \mathcal{D} is defined as a collection of all closed sets of convex geometry \mathbf{D}_n , plus sets $\{y\} \cup D \setminus a_i$ and $\{x\} \cup D \setminus a_j$. These are, indeed, the closed sets of a closure operator, since the intersection of any members of \mathcal{D} is again in \mathcal{D} . For this, it is enough to note that any subset of $\{y\} \cup D \setminus a_i$ and $\{x\} \cup D \setminus a_j$ is a closed set of convex geometry \mathbf{D}_n . We can claim more, namely:

Lemma 3.1. \mathbf{C}_n is a (finite) convex geometry that satisfies the n -Carathéodory property.

Proof. To show that \mathbf{C}_n is a convex geometry, one needs to demonstrate that every closed set can be extended by one point to obtain another closed set. This is true for any closed set of \mathbf{D}_n , since it is a convex geometry itself. This is also true for additional sets $\{y\} \cup D \setminus a_i$ and $\{x\} \cup D \setminus a_j$: the first can be extended by x to obtain $\{x, y\} \cup D \setminus a_i$, a closed set of \mathbf{D}_n , the second can be extended by y to obtain $\{x, y\} \cup D \setminus a_j$, another closed set of \mathbf{D}_n . \square

Lemma 3.2. \mathbf{C}_n cannot be a sub-geometry of any geometry of relatively convex sets of dimension $\leq n$.

Proof. Indeed, \mathbf{C}_n does not satisfy the n -Carousel Rule, since x is not in a closure of y with any n points from D (similarly, y is not in a closure of x with any n points from D). Hence, the claim of this lemma follows from 2.3 and 2.4. \square

On the other hand, we can show that \mathbf{C}_n is a sub-geometry of some $(n + 1)$ -dimensional geometry of relatively convex sets. Indeed, consider \mathbb{R}^{n+1} , and subspace $S_0 \subseteq \mathbb{R}^{n+1}$ of all points whose last projection is 0; correspondingly, let $S_1 \subseteq \mathbb{R}^{n+1}$ be a subspace of all points whose last projection is 1. Consider points $c_0, c_1, \dots, c_n \in S_0$ whose convex hull is n -dimensional polytope C , and take an inner point u of C . Let $b_0, b_1, \dots, b_n, v \in S_1$ be obtained from c_0, c_1, \dots, c_n, u , correspondingly, by replacing the last projection by 1. Let $K = \{c_0, b_0, c_1, b_1, \dots, c_n, b_n, u, v\}$ and $\mathbf{G}_{n+1} = \mathbf{Co}(\mathbb{R}^{n+1}, K)$.

Define a mapping ϕ from closed sets of \mathbf{C}_n to closed sets of \mathbf{G}_{n+1} : $\phi(\{a_i\}) = \{c_i, b_i\}$, $i = 0, \dots, n$, $\phi(\{x\}) = \{u\}$, $\phi(\{y\}) = \{v\}$. For any closed set $S = \{s_1, \dots, s_k\}$, $k > 1$, of \mathbf{C}_n , it is straightforward to check that $\phi(s_1) \cup \dots \cup \phi(s_k)$ is closed in \mathbf{G}_{n+1} , thus, we may define $\phi(S) = \phi(s_1) \cup \dots \cup \phi(s_k)$, for any closed S in \mathbf{C}_n . Evidently, this mapping preserves intersections. As for the closure of a union of closed sets X, Y in \mathbf{C}_n , we observe that

$$\overline{X \cup Y} = \begin{cases} X \cup Y \cup \{x, y\}, & a_0, \dots, a_n \in X \cup Y; \\ X \cup Y, & \text{otherwise.} \end{cases}$$

Similarly, in \mathbf{G}_{n+1} , u (symmetrically, v) is not in a closure of v (u) with any n sets $\{c_i, b_i\}$, $i = 0, \dots, n$, since u (v) is an inner point of n -dimensional polytope with vertices c_0, \dots, c_n (b_0, \dots, b_n). Hence, we have in \mathbf{G}_{n+1}

$$\overline{\phi(X) \cup \phi(Y)} = \begin{cases} \phi(X) \cup \phi(Y) \cup \{u, v\}, & a_0, \dots, a_n \in X \cup Y; \\ \phi(X) \cup \phi(Y), & \text{otherwise.} \end{cases}$$

Therefore, ϕ preserves the closure of the union of closed sets, too.

4. SHARPENING CARUSEL RULE

It turns out that one can slightly strengthen the n -Carusel Rule, and we are going to illustrate it in case of 2-Carusel Rule.




FIGURE 1.

First of all, let see the visual image of 2-Carusel Rule on Figure 1: if x, y are in the convex polygone generated by a_1, \dots, a_n , then x should be at least in one triangle generated by y and two points from a_1, \dots, a_n . In general, there might be multiple triangles of that sort containing x . On the other hand, if $n = 3$, i.e. x, y are inside the triangle defined by a_1, a_2, a_3 , x can belong to maximum two triangles. In this case, x will be also on the segment containing y and one of points a_1, a_2, a_3 . Indeed, if, say, $x \in \overline{\{y, a_1, a_2\}}$ and $x \in \overline{\{y, a_1, a_3\}}$, then $x \in \overline{\{y, a_1\}}$. Note that the property will hold even if y belongs to the boundary of triangle a_1, a_2, a_3 .

The version of this property under additional assumption that the points on the plane are in the *general position*, i.e. no three of them are on the same line, is called *the simplex partition property* in [14]. In this case, one would say that x can be in exactly one of triangles $\overline{\{y, a_i, a_j\}}$, $i, j \in \{1, 2, 3\}$.

It turns out we can make the similar statement in any sub-geometry of 2-dimensional geometry, as long as we assume that y is not on the boundary of a_1, a_2, a_3 .

Theorem 4.1. *Let $\mathbf{G} = (G, -)$ be any sub-geometry of 2-dimensional finite geometry $\mathbf{G}_0 = \text{Co}(\mathbb{R}^2, G_0)$. Then the following implication holds for all x, y, a, b, c in G : if $y \in \overline{\{a, b, c\}}$, $\overline{y} \cap \overline{\{a, b\}} = \overline{y} \cap \overline{\{b, c\}} = \overline{y} \cap \overline{\{a, c\}} = \overline{y} \cap \overline{x} = \emptyset$, $x \in \overline{\{y, a, b\}}$ and $x \in \overline{\{y, a, c\}}$, then $\overline{x} \cap \overline{\{y, a\}} > \emptyset$.*

To prove Theorem we will need a few auxiliary statements.

Lemma 4.2. *Let $a_1, \dots, a_i, \dots, a_j, \dots, a_k, \dots, a_s, \dots, a_n$ be a circular order of vertices of some convex polygon on the plane. If s is a point of intersection of segments $[a_1, a_j]$ and $[a_i, a_k]$, then s is in triangle $\{a_s, a_i, a_j\}$.*




FIGURE 2.

Proof. It is true for any "diagonal" of a convex polygon $[a_1, a_j]$ that all the vertices between a_j and a_1 in their circular order belong to the same semi-plane generated by the line (a_1, a_j) . In particular, $[a_1, a_j]$ and $[a_i, a_k]$, indeed, intersect at some point s , since the points a_i and a_k are separated by line (a_1, a_j) .

In order to show that s is inside triangle $\{a_s, a_i, a_j\}$, one needs to show that, for each side of a triangle, the third vertex and point s belong to the same semiplane generated by the line extending this side.

Take side $[a_i, a_j]$, then vertices a_k, a_s, a_1 are in the same semiplane generated by line (a_i, a_j) , hence, both segments $[a_1, a_k]$ and $[a_i, a_j]$ are in that semiplane, implying that their intersection point s belongs there as well.

Take another side of triangle $[a_i, a_s]$. Then a_j, a_k are in the same semiplane generated by line (a_i, a_j) . Since s is on segment $[a_i, a_k]$, it belongs to the same semiplane. Thus, s and a_j belong to the same semiplane generated by (a_i, a_s) , which is needed. Similar is true for the side $[a_j, a_s]$ and points a_i and s . \square

Lemma 4.3. *Suppose the vertices of a convex polygon M with at least 4 vertices are split into three subsets A, B, C . If the vertices of one of these subsets are separated by the vertices of the others in the circular order, then every point of convex polygon M belongs to $\overline{A \cup B} \cup \overline{A \cup C} \cup \overline{B \cup C}$.*

Proof. Assume without loss of generality that vertices $a_1, a_2 \in A$ are separated by points either from B or C in the circular order of vertices of polygon M . If points




FIGURE 3.

from B all belong to one semi-plane generated by line (a_1, a_2) , and all points from C are in the other semi-plane, then every point from M is in $\overline{A \cup B \cup A \cup C}$. Thus, assume that there are points from both B and C in one of semi-planes, and points from, say, B are located in both semi-planes. Then the only points of M that do not belong to $\overline{A \cup B \cup A \cup C}$ are the points of triangles of the form $\{b_i, c_j, s_{ij}\}$, where $b_i \in B$, $c_j \in C$, c_j immediately follows b_i in the circular order of vertices of M , and s_{ij} is the point of intersection of lines (u, c_j) , and (b_i, v) , where u is closest from $A \cup C$ preceding point to b_i in the circular order, and v is the closest in circular order point from $A \cup B$ following c_j .

According to the assumption, there is vertex $b_k \in B$ that belongs to the other semi-plane generated by (a_1, a_2) . Due to Lemma 4.2, when a_i is replaced by b_i , a_1 by u , a_j by c_j , a_k by v , s by s_{ij} and a_s by b_k , it follows that s_{ij} belongs to triangle $\{b_k, b_i, c_j\}$. In particular, $\{b_i, c_j, s_{ij}\} \subseteq \{b_k, b_i, c_j\} \subseteq B \cup C$. \square

Proof of Theorem 4.1. Due to Lemma 2.4, \mathbf{G} satisfies 2-Carousel Rule, therefore, $y \in \{x, b, c\}$. According to assumption that \mathbf{G} is a subgeometry of \mathbf{G}_0 , one can find an embedding ϕ of lattice of closed sets of \mathbf{G} into lattice of closed sets of \mathbf{G}_0 . Denote $U = \phi(\bar{u})$, for any $u \in \{\bar{a}, \bar{b}, \bar{c}, \bar{x}, \bar{y}\}$, and let $P = \phi(\emptyset)$. Then, according to conditions of theorem, $X, Y \subseteq \overline{A \cup B \cup C}$, $X \subseteq \overline{A \cup B \cup Y}$, $X \subseteq \overline{A \cup C \cup Y}$, $Y \subseteq \overline{B \cup C \cup X}$. Besides, $P = Y \cap \overline{A \cup B} = Y \cap \overline{A \cup C} = Y \cap \overline{B \cup C} = Y \cap X$.

Since points of $Y \setminus P$ are inside of convex polygon $\overline{A \cup B \cup C}$, but not in any $\overline{A \cup B}, \overline{A \cup C}, \overline{B \cup C}$, the vertices of $\overline{A \cup B \cup C}$ should appear in clusters, due to Lemma 4.3: elements from A should follow elements from C , which should follow elements of B , in their circular order. Figure 4 makes a sketch of arrangement, where a_1 and a_2 are end points of A -cluster, similarly, b_1, b_2 and c_1, c_2 are end-points of clusters B and C , correspondingly. Elements of $Y \setminus P$ are located inside triangle formed by points of intersection of lines (a_1, b_2) , (b_1, c_2) and (c_1, a_2) . We need to show that some point $x \in X \setminus P$ is in $\overline{Y \cup A}$.

Claim 1. *Let points of $\overline{A \cup C \cup Y}$ follow each other in the circular order $c_1, \dots, c_i, \dots, c_2, a_1, \dots, a_i, \dots, a_2, \dots, u, \dots, v, \dots, c_1$. Then $v \in \overline{u \cup B \cup C}$.*




FIGURE 4.

One can use Figure 4 with possible identification of u as y , and v as c_j . We assume that $u \notin W = \overline{\{b_1, \dots, b_i, \dots, b_2, c_1, \dots, c_i, \dots, c_2\}}$, since otherwise the claim is obvious.

Draw the line (u, a_2) , then v should be in the same semi-plane as c_1 . Draw the line (c_1, u) , then v should be in the semi-plane opposite to a_2 . Since u is an inner point of triangle $\{a_2, b_1, c_2\}$, line (a_2, u) crosses segment $[c_2, b_1]$ at some inner point. Hence, this line crosses another segment of convex polytope W , say, at point w (this point does not necessarily belong to configuration that generates G_0). Thus, v belongs to the convex polytope formed by u, c_1, w and all the vertices of the border of W between c_1 and w . In particular, $v \in u \cup B \cup C$, as desired. End of proof of Claim 1.

Since $\overline{A \cup C} \subset \overline{Y \cup A \cup C}$, some vertices of $\overline{Y \cup A \cup C}$ should be from $Y \setminus \overline{A \cup C} = Y \setminus P = y \setminus \overline{B \cup C}$. Let y_1 be the first element from $Y \setminus P$ that appears after a_2 in the circular order of vertices of $\overline{Y \cup A \cup C}$ given in Claim 1. According to Claim 1, no point from C can appear between a_2 and y_1 , since, otherwise, y_1 will be in $\overline{B \cup C}$. Thus, we have in the sequence from a_2 to y_1 only elements from A .

Similarly, let y_2 be the first element from $Y \setminus P = Y \setminus \overline{A \cup B}$ that appears in the circular order of vertices $b_2, \dots, b_1, a_2, \dots, a_1, \dots, b_2$ of $\overline{Y \cup A \cup B}$ between a_1 and b_2 . Then there is only elements from A in this sequence between a_1 and y_2 .

Claim 2. *The sequences of segments forming the border of $\overline{Y \cup A \cup C}$ from c_1 to a_2 containing point y_1 , and the border of $\overline{Y \cup A \cup B}$ from b_2 to a_1 containing point from y_2 intersect at some point (not necessarily the point of configuration forming G_0).*

If only one vertex in $\overline{A \cup B \cup C}$ is from A , i.e. $a_1 = a_2$, then a_1 might be the only point of intersection of $\overline{Y \cup A \cup B}$ and $\overline{Y \cup A \cup C}$. In this case we assume that this is point of intersection of borders of these two convex polygons stated in the claim.

Otherwise, points c_1 and a_2 are separated by line (a_1, b_2) . If $c_1, t, \dots, s, w, u, v, \dots, a_2$ are the vertices of $\overline{Y \cup A \cup C}$, on the path from c_1 to a_2 that has point y_1 , then one of the segments of this border, say, $[u, v]$, will cross $[a_1, b_2]$.

Now a_1 and b_2 are separated by line (u, v) , hence, by all the lines $(w, u), (s, w), \dots, (c_1, t)$. Therefore, the vertices of $\overline{Y \cup A \cup B}$ on the path from b_2 to a_1 and containing y_2 , will cross each of those lines. It should cross line (c_1, t) on the "right" ray, i.e. on the ray with endpoint c_1 that contains t . On the other hand, it can cross line (u, v) only on the "left" ray, i.e. on the ray with the endpoint v that contains u . There should be a sequence of vertices in $\overline{Y \cup A \cup C}$, say, s, w, u, v , where the sequence of segments of $\overline{Y \cup A \cup B}$ will cross (s, w) on the "right" ray, while it will cross (u, v) on the "left" ray. This implies it will cross one of segments $[s, w], [w, u], [u, v]$. End of proof of claim.

Let us call a point of intersection from Claim 2 by O . Note again that, unlike points from A, B, C or Y , point O is just a geometrical location of intersection of some segments formed by points from $A \cup Y$. There are three possibilities for positioning of points y_1, y_2 and O (see Figure 5). In all three cases, $V = \overline{Y \cup A \cup B} \cap \overline{Y \cup A \cup C}$ is a convex polytope formed by points from A , point O and all the points prior to O on the path from a_1 to b_2 and on the path from a_2 to c_1 . According to the assumption, $X \subseteq V$ and $Y \subseteq \overline{X \cup B \cup C}$. We need to show that some point from $X \setminus P$ belongs to $\overline{Y \cup A}$.




FIGURE 5.

(I) On the path from a_1 to b_2 , point O occurs prior to y_2 , but on the path from a_2 to c_1 point O occurs after y_1 . Evidently, O belongs to $\overline{Y \cup A}$, since O is on a segment connecting two points from $A \cup Y$. We want to show that any vertex of V between O and y_1 (which is also a vertex of $\overline{Y \cup A \cup C}$) cannot be from C . Indeed, if one vertex would be $c \in C$, then we can apply Claim 1 to vertex c in place of u , and any vertex of $\overline{Y \cup A \cup C}$ on the path from c to c_1 in place of v . Then $v \in \overline{B \cup C}$. In particular, O is in $\overline{B \cup C}$. We can apply now a symmetric statement of Claim 1 to the points on the border of $\overline{Y \cup A \cup B}$, identifying u with O and v with y_2 . Then $y_2 \in \overline{B \cup C}$, a contradiction.

Thus, all the vertices of V must be in $\overline{Y \cup A}$, which proves $X \subseteq \overline{Y \cup A}$.

(II) Both y_1, y_2 occur after O on the corresponding paths. Then all the vertices of V are in $\overline{Y \cup A}$, which is needed.

(III) Both y_1, y_2 occur prior to O on the corresponding paths. According to Claim 1, points of the path from a_2 to c_1 that appear after y_1 belong to $\overline{y_1 \cup B \cup C}$, in particular, $O \in \overline{y_1 \cup y_2 \cup B \cup C}$, thus, the part of polytope V formed by O, y_1, y_2

and all the vertices of both paths between y_1 and O , and y_2 and O , correspondingly, belong to $\overline{y_1 \cup y_2 \cup B \cup C}$. If all the points from $X \setminus P$ would be in that part of V , we would have $X \subseteq \overline{y_1 \cup y_2 \cup B \cup C}$. At least one of y_1, y_2 should be a vertex of $\overline{y_1 \cup y_2 \cup B \cup C}$. Then it can be in $\overline{X \cup B \cup C}$ only when it belongs to $X \cup B \cup C$. But then this points would be in P due to $Y \cap X = P = Y \cap \overline{B \cup C}$, a contradiction. It follows that at least one point from $X \setminus P$ should be in the part of V formed by points from $A \cup \overline{y_1 \cup y_2}$. Thus, $X \cap \overline{Y \cup A} > P$. *End of proof of Theorem 4.1*

It follows from the proof of the theorem that the following property always holds in any geometry $G_0 = \mathbf{Co}(\mathbb{R}^2, A)$, hence, in any of its subgeometry:

$$\begin{aligned} & \text{For all closed sets } X, Y, A, B, C, \\ & \text{if } Y \subseteq \overline{A \cup B \cup C} \\ & Y \cap \overline{A \cup B} = Y \cap \overline{B \cup C} = Y \cap \overline{A \cup C} = Y \cap X = P < Y, X \\ & X \subseteq \overline{Y \cup A \cup B}, X \subseteq \overline{Y \cup A \cup C} \text{ and } Y \subseteq \overline{X \cup B \cup C} \\ & \text{then } X \cap \overline{\{A \cup Y\}} > P. \end{aligned}$$

We will refer to this property as *the Sharp 2-Carousel Rule*.

In conclusion of this section, we give an example of the convex geometry that satisfies 2-Carousel Rule, but does not satisfy the Sharp 2-Carousel Rule.

Let $A = \{a, b, c, x, y\}$ and the collection of closed sets of $(A, -)$ include all one-element and two-element subsets; besides, three-element subsets are $\{x, a, b\}$, $\{x, a, c\}$, $\{y, b, c\}$, $\{x, y, w\}$, for $w \in \{a, b, c\}$, and four-element are $\{a, b, x, y\}$, $\{b, c, x, y\}$, $\{a, c, x, y\}$. This implies $x, y \in \overline{\{a, b, c\}}$, $x \in \overline{\{y, a, b\}}$, $\overline{\{y, a, c\}}$, and $y \in \overline{\{x, b, c\}}$. The Sharp 2-Carousel Rule fails since $x \notin \overline{\{y, a\}}$. Hence, $(A, -)$ is not a sub-geometry of any geometry of relatively convex sets.

5. CONCLUDING REMARKS

Problem 1.1 asks whether any of classes $\mathcal{C}_B, \mathcal{C}_f$ is universal for *all* finite convex geometries. In fact, it is enough to check whether every finite *atomistic* convex geometry is a sub-geometry in one of those classes. Recall that a closure system $\mathbf{A} = (A, -)$ is called *atomistic*, if all one-element subsets of A are closed. This follows from the result proved in [3] (a different proof was given in [4]):

Proposition 5.1. *Every finite convex geometry has a strong atomistic extension. In particular, every finite convex geometry is a sub-geometry of some atomistic convex geometry.*

On the other hand, for the description of sub-geometries of class \mathcal{C}_n , the proposition above is not of great help, due to the fact the strong atomistic extension might not preserve the n -Carousel Rule.

Indeed, it is enough to give an example of an atomistic extension that does not preserve the n -Carathéodory Property.

Consider finite geometry $\mathbf{G} = (\{a, b, c, d, x\}, -)$ given by its collection of closed sets $\mathcal{G} = \{\emptyset, a, b, d, ab, ad, bd, cd, abd, acd, abx, adx, bcd, bcdx, acdx, abdx, abcdx\}$. In this convex geometry, for any closed sets $\overline{U} = U, \overline{V} = V$,

$$\overline{U \cup V} = \begin{cases} U \cup V \cup \{x\}, & a, b, c \in U \cup V; \\ U \cup V, & \text{otherwise.} \end{cases}$$

In particular, this convex geometry satisfies the 3-Carathéodory Property.

Let $\mathbf{H} = (\{a, b, c, d, x\}, \tau)$ be another convex geometry on $\{a, b, c, d, x\}$, whose closed sets are all subsets of $\{a, b, c, d, x\}$, except $abcd$. One easily verifies that \mathbf{G} is a sub-geometry of \mathbf{H} , therefore, \mathbf{H} is an atomistic extension of \mathbf{G} . On the other hand, 3-Carathéodory property fails in \mathbf{H} , since $x \in \tau(abcd)$, but $x \notin \tau(abc) \cup \tau(abd) \cup \tau(acd) \cup \tau(bcd)$.

It would be interesting to describe necessary and sufficient properties of finite geometries which are sub-geometries of n -dimensional geometries of relatively convex sets. One of main results in [6] states that if a finite atomistic convex geometry with k extreme points a_1, \dots, a_k and points x, y in the closure of a_1, \dots, a_k , satisfies the so-called Carousel Rule and Splitting Rule then it can be represented as $\mathbf{Co}(\mathbb{R}^2, A)$, with $A = \{a_1, \dots, a_k, x, y\}$ being some set of points on a plane. In this result the Carousel Rule is slightly more elaborate property than the 2-Carousel Rule (a version of the Carousel Rule was also formulated in [10], where the case of one point x in the closure of a_1, \dots, a_k was investigated).

At the moment we are not aware of any example of a finite convex geometry satisfying the 2-Carousel Rule and the Sharp 2-Carousel Rule but not representable by relatively convex sets on the plane. Thus, we would like to ask:

Problem 5.2. *Is every finite convex geometry that satisfies 2-Carousel Rule and the Sharp 2-Carousel Rule a sub-geometry of some (finite) geometry $\mathbf{Co}(\mathbb{R}^2, A)$?*

In [5], the 2-Carousel Rule was essential in establishing the correspondence between two problems: the representation of an atomistic convex geometry as $\mathbf{Co}(\mathbb{R}^2, A)$ and the realization of an order type by point configuration on the plane. See [11] for the definition of an order type and [12] for the recent overview and references on the topic.

Acknowledgements. The paper was inspired by Maurice Pouzet and a prospect to celebrate his achievements at ROGICS'08 conference in Mahdia, Tunisia. We are grateful to Nejib Zaguia and other organizers of the conference to make this event possible.

REFERENCES

[1] K. Adaricheva, *Join-semidistributive lattices of relatively convex sets*, Contributions to General Algebra 14, Proceedings of the Olomouc Conference 2002 (AAA 64) and the Potsdam conference 2003 (AAA 65), Verlag Johannes Heyn, Klagenfurt, 2004, 1–14. Also see on http://arxiv.org/PS_cache/math/pdf/0403/0403104v1.pdf

[2] ———, *Two embedding theorems for lower bounded lattices*, Algebra Universalis **36** (1996), 425–30.

[3] K.V. Adaricheva, V.A. Gorbunov, and V.I. Tumanov, *Join-semidistributive lattices and convex geometries*, Adv. Math. **173**(2003), 1–49.

[4] K.V. Adaricheva and J.B. Nation, *Largest extension of finite convex geometry*, Algebra Universalis **52** (2004), 185–195.

[5] K.Adaricheva and M.Wild, *Realization of abstract convex geometries by point configurations. Part I: Complexity*, submitted to European Journal of Combinatorics. Also see on http://arxiv.org/PS_cache/arxiv/pdf/0710/0710.1248v1.pdf

[6] ———, *Realization of convex geometries by point configurations on a plane. Part II: two points inside an n -gon*, manuscript.

[7] G.M. Bergman, *On lattices of convex sets in R^n* , Algebra Universalis **53** (2005), 357–395.

- [8] B. Dietrich, *A circuit characterization of antimatroids*, J. Comb. Theory B **43**(1987), 314–321.
- [9] P.H. Edelman and R. Jamison, *The theory of convex geometries*, Geom.Dedicata **19**(1985), 247-274.
- [10] P. Edelman and D. Larman, *On characterizing collections arising from N -gons in the plane*, Geom. Dedicata **33**(190), 83–89.
- [11] J.E. Goodman and P. Pollack, *Multidimensional sorting*, SIAM J.Computing **12**(1983), 484–503.
- [12] *Handbook of Discrete and Computational Geometry*, edited by J.E.Goodman and J.O'Rourke, Chapman& Hall/CRC, 2004.
- [13] A.P.Huhn, *On nonmodular n -distributive lattices.I.Lattices of convex sets*, Acta Sci.Math (Szeged) **52** (1988),35–45.
- [14] W.Morris and V.Soltan, *The Erdős-Szekeres problem on points in convex position - a survey*, Bulletin of Amer.Math.Soc. **37** (2000), 437–458.
- [15] J.B. Nation, *Closure operators and lattice extensions*, Order **21** (2004), 43–48.
- [16] P. Pudlák and J. Tůma, *Every finite lattice can be embedded in a finite partition lattice*, Algebra Universalis **10** (1980), 74–95.

STERN COLLEGE FOR WOMEN, YESHIVA UNIVERSITY, 245 LEXINGTON AVE., NEW YORK , NY 10106, USA

E-mail address: `adariche@yu.edu`