



## How environmental managers perceive and approach the issue of invasive species: the case of Japanese knotweed s.l. (Rhône River, France)

Marylise Cottet, Florence Piola, Yves-François Le Lay, Soraya Rouifed, Anne Riviere-Honegger

### ► To cite this version:

Marylise Cottet, Florence Piola, Yves-François Le Lay, Soraya Rouifed, Anne Riviere-Honegger. How environmental managers perceive and approach the issue of invasive species: the case of Japanese knotweed s.l. (Rhône River, France) . *Biological Invasions*, Springer Verlag, 2015, 17 (12), <10.1007/s10530-015-0969-1>. <halshs-01248768>

**HAL Id: halshs-01248768**

**<https://halshs.archives-ouvertes.fr/halshs-01248768>**

Submitted on 28 Dec 2015

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



1 **How environmental managers perceive and approach the issue of invasive species: the case of Japanese**  
2 **knotweed *s.l.* (Rhône River, France)**

3 *Marylise Cottet<sup>1\*</sup>, Florence Piola<sup>2</sup>, Yves-François Le Lay<sup>1</sup>, Soraya Rouifed<sup>2</sup> and Anne Rivière-Honegger<sup>1</sup>*

4  
5 \*Corresponding author:

6 Email: [marylise.cottet@ens-lyon.fr](mailto:marylise.cottet@ens-lyon.fr)

7 Phone number: +33 437 376 348

8 Postal address: ENS de Lyon, bureau R 243, 15 parvis René Descartes, 69007 LYON

9  
10 <sup>1</sup>UMR 5600 "Environnement, Ville, Société", CNRS, Université de Lyon, ENS de Lyon, 15 parvis René  
11 Descartes, BP 7000, 69342 Lyon Cedex 07, France

12 <sup>2</sup>UMR 5023 LEHNA, CNRS, Université Claude Bernard Lyon 1, ENTPE, 43 Boulevard du 11 novembre 1918,  
13 69622 Villeurbanne Cedex, France

14  
15 **ABSTRACT**

16 Studying the perceptions of stakeholders or interested parties is a good way to better understand behaviours and  
17 decisions. This is especially true for the management of invasive species such as Japanese knotweed *s.l.* This  
18 plant has spread widely in the Rhône basin, where significant financial resources have been devoted to its  
19 management. However, no control technique is recognized as being particularly effective. Many uncertainties  
20 remain and many documents have been produced by environmental managers to disseminate current knowledge  
21 about the plant and its management. This article aims at characterizing the perceptions that environmental  
22 managers have of Japanese knotweed *s.l.* A discourse analysis was conducted on the printed documentation  
23 produced about Japanese knotweeds *s.l.* by environmental managers working along the Rhône River (France). The  
24 corpus was both qualitatively and quantitatively analysed. The results indicated a diversity of perceptions  
25 depending on the type of environmental managers involved, as well as the geographical areas and scales on  
26 which they acted. Whereas some focused on general knowledge relating to the origins and strategies of  
27 colonization, others emphasized the diversity and efficacy of the prospective eradication techniques. There is a  
28 real interest in implementing targeted actions to meet local issues. To do so, however, these issues must be better  
29 defined. This is a challenging task, as it must involve all types of stakeholders.

30           KEYWORDS

31   Discourse analyses; environmental managers; invasive species; Japanese knotweed *s.l.*; management;  
32   perceptions.

33

34 1. INTRODUCTION

35 **1.1. When scientific uncertainties make management more difficult**

36 Biological invasions, which are believed to be the result of global change, are of growing interest in the  
37 biological sciences (Vitousek *et al.* 1997) because of their potential effects on biodiversity. Invaded ecosystems  
38 are generally considered as disrupted areas, where native species are strongly threatened. Thus, the control of  
39 invasive species has become a priority for many countries, and several policy engagements have already been  
40 ratified and are being implemented (Genovesi and Shine 2004; Heywood and Brunel 2011). Therefore, managers  
41 need to find efficient and feasible control methods (Delbart *et al.* 2012). However, in the case of particularly  
42 efficient invaders, managing such species presents a considerable challenge, and while ecological studies are  
43 numerous and can help to define successful management techniques (Genovesi, 2011), the complexity of the  
44 ecological processes involved in invasion often makes it difficult to develop effective control methods.

45

46 Biological invasions pose not only a critical ecological issue but also an important social issue. The social  
47 dimensions of biological invasions were first considered in the early 2000s (McNeely 2001), following  
48 initiatives in environmental economics (Perring *et al.* 2000; Pimentel *et al.* 2000). If managers want to  
49 efficiently manage invasive species when defining a management strategy, they must also consider the social  
50 dimension of the issue (Binimelis *et al.* 2007; Gobster 2011). On the one hand, the impacts of biological invasions  
51 are many and not limited solely to ecological consequences. They may also affect market or non-market goods  
52 and services produced (or no longer produced) by invaded systems (Colautti *et al.* 2006). On the other hand, what  
53 we define as a biological invasion (the degree to which the colonization of an area by a species becomes an  
54 invasion) as well as the management strategies we use to control them are widely influenced by human  
55 perceptions (Mack 2001).

56 **1.2. Human perceptions, a key factor for managing biological invasions**

57 When managing complex ecological processes, there are no standard rules defining what constitutes a good  
58 decision or action. Human perceptions that guide the definition of strategies rely sometimes on non-scientific  
59 criteria (Lévêque *et al.* 2012). For example, Starfinger *et al.* (2003) showed that when there are insufficient data  
60 available to tackle a specific issue, there is a tendency to believe in stories about the beneficial or noxious impact  
61 of an alien species. Focusing on human perceptions as well as human values associated with biological invasions  
62 is a good way to better understand behaviours and decisions. Furthermore, this focus may help to define more

63 efficient management strategies (Vanderhoeven *et al.* 2011). This research direction is also legitimized given  
64 the fact that there is a wide diversity of stakeholders or interested parties involved in the management of  
65 biological invasions, each of them having a specific perception regarding the issue and a specific point of view  
66 regarding the action to take (Simberloff *et al.* 2005). Human perceptions are at their most heterogeneous when  
67 addressing questions that are confounded by many scientific uncertainties (Pahl-Wostl 2006), as everyone tends  
68 to have his or her own perceptions depending on his or her personal experiences. With respect to invasive  
69 species, uncertainty is the norm (Williamson 1999; Horan *et al.* 2002). The definition of action relating to  
70 biological invasions must therefore take into account this diversity of perceptions (Binimelis *et al.* 2007; Garcia  
71 Llorente *et al.* 2008). This article aims to serve this objective by choosing an original angle: the study of  
72 managers' perceptions.

### 73 **1.3. The choice of studying managers' perceptions**

74 Many scientific studies until now have been interested in characterizing public perceptions of the control and  
75 eradication of invasive species (Simberloff *et al.* 2005; Hulme 2006; Bardsley and Edwards-Jones 2007; Bremner  
76 and Park 2007; Andreu *et al.* 2009), and public opposition has repeatedly caused delays in or the abandonment  
77 of control efforts (Marshall *et al.* 2011; McNeely 2011). Better knowledge of public opinion towards these  
78 actions is a first step in fostering public involvement and gaining social acceptance (Selge *et al.* 2011).  
79 Nevertheless, the perceptions of environmental managers require specific attention.

- 80 - Environmental managers are charged with defining and implementing environmental plans. They are  
81 the ones who have to act (or not) and who have to meet the challenge of overcoming uncertainties (Liu  
82 *et al.* 2011). How do these uncertainties influence their perceptions, decisions and behaviours towards  
83 the plant and its management?
- 84 - They are also the ones in charge of producing and communicating information about invasive species to  
85 other stakeholders, tasks that require certain skills and principles (Jurin *et al.* 2010). Beyond the  
86 uncertainties that they must address, when communicating on issues relating to a biological invasion,  
87 they face the challenge of correctly informing stakeholders about the invasive species. Indeed, the  
88 ambiguous and inconsistent use of terminology (Richardson *et al.* 2000; Collauti and Richardson  
89 2009) and the use of emotive and manipulative language (Gobster 2005; Larson 2005; 2008; Stromberg  
90 *et al.* 2009; Selge *et al.* 2011) have already been widely criticized.

91 **1.4. The case of Japanese knotweed *s.l***

92 *1.4.1. The importance of conducting case studies*

93 Research conducted on invasive species is instructive. In particular, these studies show that different criteria  
94 influence human perceptions regarding a biological invasion. Several researchers have listed influencing criteria  
95 (Garcia Llorente *et al.* 2008; Selgeet *et al.* 2011), and some have even proposed a model to explain human  
96 perceptions of biological invasions (Gobster 2011). The following are among the influencing criteria identified:

- 97 - The impact caused by an invasive species appears to be one of the main structuring factors of human  
98 perceptions. These impacts can be negative or positive (Shapiro 2002; Bardsley and Edward-Jones  
99 2006) and may affect ecosystems as well as social systems (Garcia Llorente *et al.* 2008). The negative  
100 impacts of invasive species on ecosystems are found to be a strong motivation for their eradication  
101 (Levine *et al.* 2003; Garcia Llorente *et al.* 2008; Selgeet *et al.* 2011). Nevertheless, ecological functions  
102 provided by species (Binimelis *et al.* 2007) or values associated with them (i.e., the aesthetic value of  
103 colourful plants such as purple loosestrife; the cultural value of feral pigs for particular ethnic groups,  
104 etc., Gobster 2011) may help to attenuate the observed negative impacts.
- 105 - Time can also influence human perceptions of a biological invasion such that the older the invasion, the  
106 less the species is identified as exotic, and the better it is valued (Bardsley and Edwards-Jones 2006;  
107 Garcia Llorente *et al.* 2008; Gobster 2011). Moreover, Starfinger *et al.* (2003) showed that perceptions of  
108 an invasive species can evolve over time, according to the scientific data available.
- 109 - Personal factors add many variations to the perceptions of an invasive species. Such factors include  
110 education, economic status, rural or urban residence, cultural or regional characteristics, house size and  
111 distance: the closer the invasion, the more concerned people are likely to be (Garcia Llorente *et al.* 2008;  
112 Ehrenfeld 2010; Gobster 2011). Knowledge and expertise also play a critical role in the way we consider  
113 issues linked to biological invasions. Consequently, different stakeholders will often have different  
114 perceptions of the issue.

115 The previous studies show that reflections about biological invasions cannot be exclusively conducted in a  
116 global framework, as human perceptions of biological invasions depend on which species is under consideration,  
117 which area has been invaded and which stakeholder is affected. Data, therefore, should be focused on the effect  
118 of a single species over a single area on a single-type stakeholder.

119           1.4.2.   *The Japanese knotweed s.l. invasion: a preoccupation of environmental managers*

120 Taxa from the hybrid complex *Fallopia* (mainly *Fallopia japonica*  
121 (Houtt.)RonseDecraene,*Fallopia sachalinensis* (F. Schmidt ex Maxim.) RonseDecraene and the hybrid *Fallopia*  
122 *x bohemica* (ChrtketChrtkovaĭ)) are widespread invaders of North America (Shaw and Seiger 2002) and Europe  
123 (Child and Wade 2000). *F. japonica* and *F.sachalinensis* are rhizomatous perennial herbaceous plants that  
124 originated from Asia and were introduced in Europe in the XIX century because of their ornamental qualities  
125 (Bailey and Conolly 2000). Today,*F. japonica* is one of the most common invasive species in Europe, having  
126 been identified in 40 countries (Lambdon *et al.* 2008), including France. Hybridization between *Fallopia* taxa  
127 produces *F. x bohemica*, which exhibits higher genetic diversity (Tiébré *et al.* 2007), higher phenotypic variation  
128 (Herpigny *et al.* 2012), and higher performances (Parepaet *et al.* 2014). *Fallopia* spp. colonize mainly in riparian  
129 habitats and disturbed areas (Bailey *et al.* 2009). In invaded sites, identified functional impacts include altered  
130 nutrient cycles (Dassonville *et al.* 2007; 2011) and reduced plant species diversity (Vanderhoeven *et al.* 2005;  
131 Gerber *et al.* 2008). Numerous mechanical, chemical or biological techniques have been tested, but none have  
132 demonstrated satisfying outcomes (for a review of these techniques and their effectiveness, see Delbart *et al.*  
133 2012). In fact, some of them may even promote further invasion, as is the case for mowing interventions  
134 (Beerling *et al.* 1994; McHugh 2006), the most frequently used technique. Little is known about the response of  
135 the different taxa to the control methods. Moreover, as they invade similar habitats and as it is sometimes  
136 difficult to distinguish between them, they are often considered together. For these reasons and for convenience,  
137 in this paper, the term Japanese knotweed *s.l.* will be used to refer to the three taxa of interest: *F. japonica*, *F.*  
138 *sachalinensis*, and *F. x bohemica* (Bailey *et al.* 2009).

139 Because of the uncertainties regarding management, there is no consensus with respect to the most effective way  
140 to control Japanese knotweed *s.l.* (Delbart *et al.* 2012) or even whether it should be controlled. Ongoing ecological  
141 field research may produce knowledge likely to help define the best management plan for this plant (Rouifed  
142 2011). Nevertheless, this research must be completed with data relating to the field of human perceptions,  
143 especially to better understand the main issues associated with the management of Japanese knotweed *s.l.* and the  
144 best action to take from the perspective of the managers.

145           1.4.3.   *A need for research relating to human perceptions of Japanese knotweed s.l.*

146 Only a few studies have considered the social dimension of Japanese knotweed *s.l.* invasion. Child *et al.* (1998)  
147 conducted a contingent valuation study to assess the socially acceptable cost of controlling the plant. Apart from  
148 this study, only factual and incomplete data relating to the human perceptions of Japanese knotweed *s.l.* are



149 available. For instance, Vanderhoeven *et al.* (2011) determined that, in Belgium, Japanese knotweed *s.l.* is one of  
150 the species mentioned by horticulture professionals and nature reserve managers during surveys when asked  
151 which species would become a problem in the next few years. Further research on human perceptions of  
152 Japanese knotweed *s.l.* is therefore required.

153 Herein, we aim to extend these initial data by characterizing the perceptions of environmental managers working  
154 along the Rhône River with Japanese knotweed *s.l.* By analysing what they have written about this plant, we aim  
155 to determine:

- 156 (1) What types of environmental managers are actually concerned about Japanese knotweed *s.l.* and produce  
157 information about it?
- 158 (2) What are the perceived issues linked with the management of the plant?
- 159 (3) What strategies do the environmental managers recommend to manage this plant, and what arguments do  
160 they use to justify their choices?

## 161 2. MATERIALS AND METHODS

### 162 2.1. Study area

163 The Rhône River, one of the main Mediterranean rivers, originates at the Furka glacier in the Swiss Alps. The  
164 river is 812 km long, with more than 500 km located in France. The Rhône flows into the Mediterranean Sea,  
165 where it terminates in a very large delta. Upstream of this delta, at Beaucaire, the mean river flow reaches 1700  
166 m<sup>3</sup>/s (Olivier *et al.* 2009). The Rhône River is a powerful unifying element of southern and eastern France  
167 because it crosses or delimits many administrative areas, including regional, infra-regional and local authorities.  
168 Many environmental managers attached to these various geographical areas are facing the challenge of how to  
169 control Japanese knotweed *s.l.* as part of their mission.

170  
171 While few data are available concerning the intensity of the Japanese knotweed *s.l.* invasion along this  
172 watercourse, according to the inventory conducted in 2001 for the regional water authority (Boyer and Laval  
173 2001), the invasion is variable, with the upper region exhibiting a far greater invasion than the lower region.  
174 However, two sectors appear to be more specifically colonized. One, in the Upper Rhône, is the 40 km section  
175 downstream from the confluence with Les Ussets River, and the second, in the Middle Rhône, is a 90 km section  
176 downstream from the city of Lyon. The Japanese knotweed *s.l.* invasion is of major concern to environmental

177 managers and has led to numerous management plans, following the measures (2010 to 2015) defined by the  
178 regional water authority to “fight against invasive exotic species” (article OF6-C).

## 179 **2.2. Constitution of the corpus**

180 We aim to gather all documents relating to Japanese knotweed *s.l.* produced by managers involved in the  
181 management of this plant along the Rhône River in France. The preliminary analysis suggests that printed  
182 documents on this plant are relevant sources for studying stakeholders’ perceptions, as each document published  
183 by a management structure is often collectively written and validated. The discourse it produces about a given  
184 issue is therefore assumed to correspond to management structure’s perceptions regarding the issue.

185

186 To be as exhaustive as possible when collecting this documentation, a multi-step, rigorous collection strategy  
187 was established. As a first step, we conducted a general search of the Internet, using the scientific and common  
188 names of the plant, place names, and words such as ‘management’, ‘control’, and ‘management practices’, in  
189 French. As a second step, we contacted by telephone different types of management bodies whose  
190 responsibilities included issues related to the Rhône River. They were asked to send us by post or e-mail a copy  
191 of any documents relating to Japanese knotweed *s.l.* that they had produced or had in their possession. For some  
192 of these institutions, we went on-site to gather the available documents. This contact procedure was pursued at  
193 three geographical levels:

- 194 - at the local level: all "communautés de communes" (a regrouping of local authorities with jurisdiction  
195 on certain matters) or individual municipalities (when not collectively organized), as well as all local  
196 water associations;
- 197 - at the regional or infra-regional level: every regional or infra-regional authority (“conseils régionaux et  
198 départementaux”) as well as all decentralized state services (“direction régionale de l’environnement, de  
199 l’aménagement et du logement”, “direction départementale des territoires”);
- 200 - at the Rhone-Mediterranean basin level: the Rhône-Mediterranean Corsica water agency (“Agence de  
201 l’eau Rhône Méditerranée Corse”).

202 Two criteria were used to justify the inclusion of a document in the corpus. First, it had to specifically focus on  
203 Japanese knotweed *s.l.* (not simply on invasive species); second, it must have been produced by an environmental  
204 manager who was responsible for the management of an area crossed or bordered by the Rhône River or one of  
205 its tributaries. We digitally scanned each document using OCR software (Omnipage professional©, Nuance

206 Communications Inc., Burlington, Massachusetts, US) and built a database to associate metadata with each of  
207 these documents. These metadata included the following:

- 208 - the publication date. In the case that the publication was in draft (i.e., not final) form, the date planned  
209 for publication was entered into the database. As a result, the corpus includes some sources listed as  
210 published in 2013;
- 211 - the type of stakeholder producing the document;
- 212 - the nature of the document.

213 The collection of the documentation for the corpus ended in April 2012 and allowed us to gather 81 documents.  
214 In spite of the systematic sampling, we may have missed certain documents. Nevertheless, since we have  
215 collected every document that the managers have archived or have heard about, we can consider that our  
216 collection is a large sample of the universe of relevant documents. One limit must however be underlined: we  
217 chose to gather only the printed documentation, which excludes online publications dealing with the question of  
218 Japanese knotweed *s.l.*, such as websites, blogs... As the environmental managers are more and more inclined to  
219 use these media as a way for communicating, this exclusion may have biased our sample.

### 220 **2.3. Corpus analysis: a statistical analysis of textual data**

221 The corpus was both qualitatively and quantitatively analysed. A statistical analysis of textual data (Lebart *et al.*  
222 1998) was performed using the open source software Iramuteq© (Ratinaud and Dejean 2009), which was  
223 recently developed (2008) and which is regularly updated. It relies on R software (R Core team, 2013) as well as  
224 the python language.

225 Iramuteq© reproduces the classification algorithm, described by Reinert (1983; 1990) and implemented in the  
226 Alceste© software, which has led to numerous publications (Brochet and Dubourdieu 2001; Dransfield *et al.*  
227 2004; Parr *et al.* 2011). It corresponds to a top-down hierarchical classification based on five stages.

228 (1) Segmentation. The corpus is segmented into two textual units: (a) the texts composing the corpus –  
229 in our case, each document produced by environmental managers and qualified by the metadata  
230 defined above (publication date, type of stakeholder and nature of the document); (b) the text  
231 segments (through an iterative process, each text is cut into a number of segments, defined  
232 according to a number of words – 40 by default).

233 (2) Lemmatization. Using a grammatical dictionary, each verb, noun, adjective, *etc.*, is reduced to its  
234 basic dictionary entry, named a lemma.

235 (3) Production of a contingency table. The matrix crosses the reduced forms (in columns) and the text  
236 segments (in rows). Among the reduced forms, only the analysable forms (nouns, verbs, adjectives,  
237 adverbs, *etc.*) are retained; supplementary forms (prepositions, pronouns, conjunctions and  
238 auxiliary verbs, *etc.*) are excluded from the table. The presence or absence of each analysable  
239 reduced form within each text segment is specified at intersections of the matrix (respectively noted  
240 as 1 vs. 0).

241 (4) Processing of a top-down hierarchical classification from the contingency table. Relying on an  
242 iterative algorithm, the software aims at defining classes that maximize the distance between two  
243 subsets (using  $\chi^2$  metrics). The iterative process stops when sub-classes are not significantly  
244 different. Specific forms of a class are then removed from the other class. This analysis is then  
245 repeated on the larger of the two classes, and so on, until the requested number of classes is reached  
246 (10 by default). Two independent analyses with different lengths of text segment are processed, and  
247 their results are statistically compared to test the stability of the classes.

248 (5) Description of the classes. The number of text segments classified in each class is specified. For  
249 each class, a list of the reduced forms associated with it is created (the degree of association with  
250 the class is indicated by the  $\chi^2$  value). The modality of the metadata variables most associated with  
251 each class is also specified. These classes are then finally interpreted as “lexical worlds” (Rouré  
252 and Reinert 1993).

253 We also performed statistical treatments relying on the co-occurrence analyses of specific lexical forms (Lebart  
254 *al.* 1998). In particular, using Iramuteq©, we conducted similarity analyses. Based on graph theory,  
255 these analyses aim to study proximity and relationships between components (in our case, the lexical forms) of a  
256 set (in our case, the corpus) using a maximum tree (Marchand and Ratinaud 2012). The plot resulting from this  
257 analysis had specific properties: (a) the greater the occurrence of a lexical form, the greater the size of the  
258 characters; (b) the greater the co-occurrence between two lexical forms, the thicker the line that linked them.  
259 Such a graph was used to analyse the relationships between all lexical forms associated with a class, which  
260 resulted from the top-down hierarchical analysis.

261 3. RESULTS

262 **3.1. Who produces information about Japanese knotweed *s.l.*, and in what form?**

263 The corpus was composed of 81 documents (195,006 words total) published between 1998 and 2013. However,  
264 the production of information about Japanese knotweed *s.l.* was marginal until 2006 (Fig1A). That year marked a  
265 strong increase in the number of documents produced, and the number continued to increase in the following  
266 years.

267 Diverse types of environmental managers were involved in this production (Fig1A). Some were attached to the  
268 regional water authority, to local or regional authorities or state services, to environmental protection  
269 associations, to environment consultancy firms, to building companies operating on the river, to regional  
270 conservation bodies (conservatoires botaniques régionaux), or to local water associations. Thus, both public and  
271 private stakeholders contributed to the production of information about Japanese knotweed *s.l.* Managers  
272 working within local water associations were the main stakeholders diffusing information about Japanese  
273 knotweed *s.l.* as they alone produced almost half of the documentation. However, their contribution to the  
274 production was relatively late and occurred mainly from 2010 onward. In other words, the strong increase in  
275 production observed over these last few years was mainly due to this one group of stakeholders. Nonetheless,  
276 local and regional authorities, state services, and environmental consultancy firms also had a significant input.

277 The documentation produced was quite diverse, consisting of management plans, syntheses of technical  
278 meetings, management guides, identification guides, information leaflets and specialized journal articles (see  
279 Table 1 for a description of the documents). The various documents addressed diverse target groups in that some  
280 - management guides and synthesis studies - were intended for environmental managers who already confronted  
281 ecological invasions and thus focused on management methods. Others - specialized journal and information  
282 leaflet - addressed the general public with potentially no knowledge of the issue and thus provided  
283 information about the plant, its colonization and its impacts. These two uses were roughly equally shared if we  
284 consider the number of produced documents per category (Fig1B).

285 Following a spatial analysis, we observed that the production of documents relating to Japanese knotweed *s.l.*  
286 also turned out to be diverse (Fig2), with the number of documents produced upstream being far higher than the  
287 number produced downstream. This longitudinal division of the river seemed to correspond to the intensity of  
288 the invasion (Boyer and Laval, 2001). That is, the more invaded the area was, the more documentation on the  
289 plant being produced by environmental managers.

## 290 3.2. What is said about Japanese knotweed *s.l.*

### 291 3.2.1. Main components of the discourses

292 The top-down hierarchical clustering classified 97.7 % of the text segments within 5 lexical classes (Fig3a). The  
293 clustering tree marked a first segmentation dividing textual segments into two parts, each having a semantic  
294 unity. The first regrouped classes 1 and 2 (including 13.2 % and 24.2 % of the textual segments, respectively), and  
295 focused on the fight against Japanese knotweed *s.l.* The second regrouped classes 3, 4 and 5 (including 19.6 %,  
296 18.7 % and 24.3 % of the textual segments, respectively), and involved information on the invasion (mechanisms  
297 of dispersion) as well as strategies for managing the plant. The most significant textual segments for each class  
298 are presented in Table 2.

299  
300 Class 3 (Fig3b) brought together all available strategies and resources for managing Japanese knotweed *s.l.* On  
301 the one hand, it listed all possible actions aimed at controlling the plant: it mentioned both preventive measures  
302 aimed at limiting its expansion, such as awareness actions (“prevention”, “awareness”, “communication”) or  
303 follow-up actions (“cartography”, “diagnostic”, “monitoring”, “inventory”), and restorative measures when the  
304 plant is already established (“eradication”, “fight”, “techniques”, “method”, “means”). The need to acquire  
305 knowledge also held a significant position within this class (“information”, “knowledge”, “to know”). On the  
306 other hand, this class took stock of all stakeholders concerned with the invasion of Japanese knotweeds *s.l.* in that  
307 it referred both to management experts (“manager”, “Rhône-Méditerranée-Corse” [water agency], “local” or  
308 “field” stakeholder”, “authority”), to economic actors (“company”), and to the general public (“public”). Many  
309 geographical areas and many spatial scales involved in the management of the plant were also listed, including  
310 “basin”, “Saône”, “Rhône”, “regional”, “*région*” (or regional authority), and “*département*” (or local authority).  
311 This inventory responds to the concerns of developing specialized stakeholder networks to better control  
312 Japanese knotweeds *s.l.* (“network”, “work[ing]” “group”).

313  
314 Classes 5 (Fig3d) and 4 (Fig3c) comprised knowledge relating to the dispersal mechanisms of Japanese  
315 knotweed *s.l.* Class 5 described the origins of the introduction of the species (“to introduce”, “exotic”, “Europe”,  
316 “Asia”, “origin”, “ornamental”, “century”, “human” “activity”), its strategies of colonization and adaptation  
317 (“hybrid”, “competition”, “reproduction”, “pioneer”), including its preferred environments for developing  
318 (“wetland”, “railway” “track”, “road”, “side”) and finally, problems it could cause (“ecological” “impact”,  
319 “nuisance”, “biodiversity”, “monospecific”, “ecosystem” “functioning”, “bank” “erosion”). This class appeared

320 to more generally describe certain dispersal mechanisms common to invasive species (“giant hogweed”,  
321 “ragweed”, “animal”). Class 4 was more specifically focused on watercourses, described as one of the main  
322 affected environments but also one of the main means of dispersal.

323

324 Eradication was at the core of classes 1 (Fig3e) and 2 (Fig3f). Class 2 listed a considerable number of the  
325 techniques used – either today or in the past – to fight Japanese knotweed *s.l.* (“to mow”, “to cut”, “chemical”  
326 “treatment”, “uprooting”, “tarpaulin”). Actions specifically focused on the main reproductive organs of the plant:  
327 its aerial (“stalk”), or its underground parts (“rhizome”) – as both are strongly involved in its reproductive  
328 capabilities in vegetative propagation. Therefore, a significant place in this class was given to the physiology of  
329 the plant. The experimental component of these actions was strongly present, as the effectiveness of the  
330 eradication techniques used was assessed experimentally (“protocol”, “experimentation”, “plot”, “counting”,  
331 “m<sup>2</sup>”). Finally, the lexicon relating to seasons and seasonality was very much present (“end”, “beginning”  
332 “May”, “June”, “season”, “period”, “month”), as was the lexicon linked to repetition (“to repeat”, “times”,  
333 “repetitive”). This finding indicated, on the one hand, that environmental managers were used to relying on a  
334 seasonal calendar to implement actions against Japanese knotweed *s.l.*, and on the other hand, that their  
335 interventions must necessarily be repeated. Class 1 focused on a specific technique that was experimentally used  
336 to eradicate the plant. This technique involved mechanical action aimed at crushing the rhizomes into small  
337 enough particles (using construction machinery) to impede their vegetative reproduction. This class insisted on  
338 the effectiveness of this method (“complete” “mortality”, “to achieve” “result”, “efficiency”) and mentioned the  
339 necessary preventive measures to implement in such experimental trials so as not to further disperse the plant  
340 (“cleaning”, “to clean”, “caterpillar” “machine”).

#### 341 3.2.2. *A technical discourse responding to a wish for effective action*

342 This classification shows that more than one-half of the discourses were dedicated to actions aimed at controlling  
343 Japanese knotweed *s.l.*, and more than one-third specifically focused on restorative methods. The discourses of  
344 environmental managers appeared to strongly promote the implementation of actions to eradicate the plant. This  
345 willingness to take action against Japanese knotweed *s.l.* was specifically obvious when we considered the  
346 “efficiency” lemma, cited as many as 121 times in the corpus. Table 3, presenting its co-occurring lemmas,  
347 highlights the fact that the “efficiency” lemma relates, above all, to the actions implemented for controlling this  
348 plant, as this wish to be “more efficient” (36 co-occurrences) led environmental managers to be creative about  
349 the methods used to destroy the plant. Figure 4 summarizes the new techniques proposed (and when) and how

350 their uses have evolved over time. We observed that while many techniques were tested, some were only very  
351 marginally cited in the documentation produced by environmental managers (thermal actions, grazing, *etc.*), and  
352 others had sometimes been abandoned (chemical treatments). On the other hand, other techniques were often  
353 mentioned. Mowing, for example, was the most cited method for controlling the plant. Uprooting was also well  
354 represented. It can be concluded that the information produced about Japanese knotweed *s.l.* was highly  
355 technical.

### 356 3.2.3. *Issues mentioned for controlling this plant*

357 In the documentation produced by environmental managers, actions against Japanese knotweed *s.l.* were strongly  
358 motivated by its impacts: there were 228 occurrences of the “impact” lemma present in the corpus. However,  
359 when we considered its co-occurring lemmas (Table 4), discourses on their possible impacts appeared to be  
360 rather weakly explained. While “environments” (n=30), “ecosystems” (n=16) and “landscapes” (n=11) were said  
361 to be affected by Japanese knotweed *s.l.*, the ways in which they were affected were unclear. The only impacts  
362 concretely qualified, albeit with very poor frequency, were related to the loss of biological diversity  
363 (“biodiversity”, n=11; “diversity”, n=7, “monospecific”, n=4) or landscape diversity (“homogenization”, n=2;  
364 “standardization”, n=2). Few references were made to the role of foliage in reducing biodiversity (“foliage”,  
365 n=4; “shade”, n=3). Reduced access (“accessibility”, n=4), and bank erosion (“undercutting”, n=2), both induced  
366 by the plant, were only marginally mentioned.

## 367 **3.3. Does every type of environmental manager have similar views about Japanese knotweed *s.l.*?**

### 368 3.3.1. *A diversity of arguments focusing on certain themes*

369 The distribution of textual segments within the different classes of the top-down hierarchical classification,  
370 according to the type of environmental manager who produced them, provided information about the diversity of  
371 proposals relating to Japanese knotweed *s.l.* Each type of stakeholder was inclined to use certain classes of  
372 discourse. The probability of occurrence of classes, for each type of stakeholder, is particularly high a low in  
373 respect to a hypothesis of independence (Pearson’s residuals; tables 5a and 5b).

374 - Class 3, which describes all strategies and stakeholders involved in the management of Japanese  
375 knotweed *s.l.*, was significantly more represented in the discourse of regional conservation bodies  
376 compared to other stakeholders.



- 377 - Class 5, which describes origins, colonization strategies and the potential impact of Japanese knotweed  
 378 *s.l.* invasion, was greatly more represented in the discourse of the regional water authority, whereas less  
 379 represented in the discourse of consultancy firms.
- 380 - Class 4, which describes the role of watercourses in dispersing the plant, was more represented in the  
 381 discourse of consultancy firms but also, to a far lesser extent, of environmental protection associations.  
 382 It was on the contrary less represented in the discourse of other stakeholders. Class 2, which describes  
 383 all the techniques used to eradicate Japanese knotweed *s.l.*, was generally more represented in the  
 384 discourse of local or regional authorities or state services, of local water associations, and, to a lesser  
 385 extent, of environmental protection associations. Conversely, it was greatly less represented in the  
 386 discourse of the regional water agency.
- 387 - Class 1, which is related to mechanical means of crushing rhizomes, was more represented in the  
 388 discourse of consultancy firms and, to a lesser extent, of building companies operating on the river. This  
 389 class is less represented in the discourses of other stakeholders, particularly the regional water agency.

390 These results show that each group had a specific position when approaching the issue of Japanese knotweed *s.l.*  
 391 Some types of stakeholders, such as the regional water agency, tended to emphasize all knowledge relating to the  
 392 Japanese knotweed *s.l.* invasion processes. On the other hand, other types of stakeholders, such as local or  
 393 regional authorities, state services, local water associations and environmental protection associations, were  
 394 more likely to diffuse information relating to the techniques used for eradicating the plant. This was also the case  
 395 for consultancy firms, who more specifically mentioned mechanical crushing techniques. Regional conservation  
 396 bodies appeared to take a particular stand, as they gave a more general view of the issue, mentioning both  
 397 strategies and the stakeholders involved in the management of Japanese knotweed *s.l.*

### 398 3.3.2. *Degrees of emotionalism linked to the Japanese knotweed s.l. invasion*

399 Four terms that were frequently used to qualify the Japanese knotweed *s.l.* invasion were “impact”, “nuisance”,  
 400 “risk” and “menace”, all of which have a negative connotation. Nevertheless, they were associated in the French  
 401 language (the original language of the corpus), with different significations, thus revealing information about the  
 402 way the environmental managers perceived the invasion. The “nuisance” and “impact” terms (in this corpus,  
 403 systematically considered as negative) characterize the invasion in a tangible, measurable way. Other terms, such  
 404 as “risk” or “menace”, on the other hand, refer to the invasion in a more intangible way: both evoke a potential  
 405 danger. The former supposes that ecosystems and/or societies (often not specified further) were vulnerable to

406 this invasion. The latter clearly evoked the danger associated with the invasion, as the terms were related to the  
407 vocabulary of fear and had a strong emotive connotation.

408 The different types of environmental managers seemed to have different uses for some of these terms (Fig5). In  
409 particular, the terms “impact” and “nuisance” were more represented in the discourse of the regional water  
410 agency compared to the whole population of managers. Conversely, the term “menace” was more represented in  
411 the discourse of local water associations. Thus, the Japanese knotweed *s.l.* invasion appeared to generate strong  
412 emotions, specifically among this group of environmental managers.

#### 413 4. DISCUSSION

##### 414 4.1. Discourses promote an integrated action and focus on the eradication stage

415 This study indicates that environmental managers of the Rhone River strongly want to take action against  
416 Japanese knotweed *s.l.* This attitude may be encouraged by the many policy engagements aimed at controlling  
417 and even eradicating priority invasive species (Delbart *et al.* 2012). As there is heavy pressure being placed on  
418 the environmental managers who are held responsible for maintaining environmental quality for the future, the  
419 uncertainties linked with the management of Japanese knotweed *s.l.* may have contributed, in several ways, to  
420 the willingness to take action. Most invasions are spreading too fast and too unpredictably to do anything other  
421 than respond immediately (Sims and Finnoff 2013). Thus, the “wait and see” approach appears to be, in many  
422 cases, inappropriate, thereby justifying the motivation for environmental managers to act. Moreover, behavioural  
423 studies have demonstrated that the threat of fearsome risks (those that induce strong emotional response, such as  
424 fear and anxiety) activate certain cognitive mechanisms that push people towards action (Loewenstein and  
425 Lerner 2003; Sunstein and Zeckhauser 2011). Such reactions are thought to be more frequent in uncertain  
426 situations (Patt and Zeckhauser 2000). The emotional response of environmental managers to a Japanese  
427 knotweed *s.l.* invasion – tangible in their discourses – may explain their willingness to take action against the  
428 plant. This tendency may also be reinforced by uncertainties relating to the effectiveness of the proposed  
429 experimental methods of control. The absence of visible results and the need for repeated interventions encourage  
430 managers to experiment with other methods. This series of successive failures and the lack of control over the  
431 plant may have resulted in frustration among the network of environmental managers (Allison 2011), thus  
432 encouraging them to pursue other strategies.

433 Which actions are considered by environmental managers? Their discourses tackle a wide diversity of modalities  
434 of intervention and promote an integrated approach towards environmental management:

435 - They take into account every stakeholder (managers, economic actors, the public) and every territorial  
436 scale (local, regional, catchment) concerned with Japanese knotweed *s.l.*

437 - They consider different actions depending on the invasion stage. During introduction stage, knowledge  
438 acquisition and information campaigns are recommended; during colonisation stage, follow-up actions  
439 are proposed; and during establishment stage, eradication measures are advanced.

440 Nevertheless, the discourses of environmental managers focus, above all, on possible actions when the plant is  
441 already established. They have tested over time a large variety of techniques aimed at stopping or at least  
442 slowing down colonization. The information produced about Japanese knotweed *s.l.* is very technical, and  
443 environmental managers mention all of their successes and failures, sometimes describing in detail the way they  
444 implemented actions and sometimes recommending certain procedures for increased efficiency. We believe that  
445 the development of discourses – and consequently of actions – relating to the invasion prevention and monitoring  
446 stages are beneficial, and we argue that it is best to act at the earliest possible stage of the invasion, that is, before  
447 it is too advanced to be reversed (Boyer, 2005). In the Rhône River case, for instance, we have found that  
448 uninvaded areas, such as the downstream section of the river, produce little information about the Japanese  
449 knotweed *s.l.* invasion. Conversely, the more heavily invaded an area is, the more documentation the  
450 environmental managers produce about these plants, and therefore, the more they worry about the presence of  
451 the plants. An efficient management system should consider implementing action before this stage.

#### 452 **4.2. Varying positions of environmental managers**

453 Japanese knotweed *s.l.* has led to widely differing proposals, according to the type of environmental manager  
454 making the proposal. In particular, there appears to be a strong difference of approach between those managers  
455 who define policy at a regional level (regional water agency) and focus on the knowledge available and those  
456 who implement policy at a regional or local level (local or regional authorities, state services, local water  
457 associations) and focus on action. Managers implementing environmental policies are highly interested in control  
458 and eradication techniques and the effectiveness of these techniques. They are also the ones more frequently  
459 using emotional language to characterize the Japanese knotweed *s.l.* invasion. This use of language may reflect  
460 their strong and increasing need for success in managing this issue given that since 2006, the majority of  
461 information produced about Japanese knotweed *s.l.* has come from these stakeholders. The pressure resulting  
462 from management policies that expect quick results in managing invasive species may be a more sensitive issue  
463 for managers in the field, as they directly experience the success or failure of their intervention and feel more  
464 directly responsible for it. From another perspective, the emotional component of their discourses may not be a

465 sign of a greater preoccupation but may only translate their willingness to bring the Japanese knotweed *s.l.* issue  
466 to the attention of funding bodies (including the regional water agency) to obtain funds to act against the spread  
467 of this plant. The use of controversial language in the field of invasion biology has already been widely  
468 observed, sometimes because its terminology connotes nativism, racism or xenophobia (Subramaniam, 2001;  
469 O'Brien, 2006), sometimes because it is militaristic (Davis et al., 2001; Larson, 2005). These criticisms have led  
470 some authors to advocate that scientists use more neutral terms for introduced species (Larson, 2005; Davis,  
471 2009). The results of our study may suggest that environmental managers do the same and be more attentive to  
472 the terminology used. We doubt that this evolution of language, if indeed possible (Larson, 2010; Simberloff,  
473 2006), will lead to a consensus in the way that Japanese knotweed *s.l.* is perceived and has to be managed: those  
474 attitudes appear to be more firmly embedded and to be influenced by heterogeneity of worldviews about relation  
475 (and definition) of human society and biota (Simberloff, 2012). Nevertheless, such an effort would enable  
476 sharing of views and discussion about actions to implement on a more neutral basis.

477 The information provided by the regional water agency (the managers defining policy at a regional level) uses  
478 less emotive terms. Moreover, this agency says little about the available methods for controlling the plant but  
479 rather focuses on the diffusion of knowledge related to the origins of the Japanese knotweed *s.l.* invasion, its  
480 colonization strategies and the nuisance it creates. The position of the regional water agency appears to be more  
481 reflective, or at least more dedicated, to the diffusion of information related to the Japanese knotweed *s.l.*  
482 invasion. Nevertheless, the attitude of the agency is far from disassociated with action, as the control of invasive  
483 species is part of the programmed measures (2010-2015) that it defined and funded<sup>1</sup>. However, as the  
484 information it produced did not focus on a way to achieve this objective, a gap exists between the two types of  
485 managers: environmental managers who implement management policies need effective methods to efficiently  
486 control the plant, as required by the regional water agency, but the agency only diffuses general knowledge  
487 related to colonization processes. Misunderstandings and frustrations may result from these heterogeneous  
488 positions. The communication strategy of the regional water agency may be more successful if their general  
489 information was complemented with more technical details regarding methods of eradication. However, even if  
490 the information produced by the regional water agency does not meet the expectations of environmental  
491 managers who have to implement these management projects, the efforts aimed at diffusing general knowledge  
492 regarding the origins, mechanisms and impacts of Japanese knotweed *s.l.* colonization should be maintained.

---

<sup>1</sup> Article OF6-C requiring to “fight against invasive exotic species”.

### 493 4.3. Towards a more targeted management

494 In the environmental managers' discourses, the management of Japanese knotweed *s.l.* is approached from a global  
495 perspective. Recommendations for action define general rules regardless of the socio-economic and  
496 environmental context of the area under consideration. These rules flow from the premise that ecosystems are  
497 bounded by frontiers within which are found, either exclusively or primarily, native species. This premise,  
498 however, is far from the reality, and defining such a reference state is delicate if not impossible (Dufour and  
499 Piégay 2009). This perception of a stable and balanced nature has long been called into question by academic  
500 ecologists (Simberloff, 2014). According to this author, the idea of a balance of nature lives on especially among  
501 conservationists and environmentalists. This may explain why this point of view is salient in the discourse of  
502 environmental managers.

503 The efficiency of management may be improved if a targeted strategy for each identifiable plant area was  
504 defined. Such a definition should then be the result of a wide collective reflection relating to local issues linked  
505 with action against invasive species. In view of this idea, environmental managers may have interest in clarifying  
506 the local issues linked with action (or inaction) against the plant in each given context and in defining the  
507 priorities related to its control. There are questions that must be addressed. For example, are there  
508 ecological issues (detrimental to other species)? aesthetic issues (size of plant and visual place in the  
509 landscape)? security issues (decrease in visibility)? economic issues (bank erosion)? Are ecological or  
510 geomorphological characteristics of the invaded area favourable to a rapid dispersion of the plant? The efficiency  
511 of the management may benefit from a more specific, spatially heterogeneous action that considers local issues  
512 (Epanchin-Niell and Hastings 2010). Certain studies have already considered this approach and merit a specific  
513 attention (Filippi and Aronson 2010).

### 514 5. CONCLUSION

515 This article studied the documentation relating to Japanese knotweed *s.l.* produced by environmental managers  
516 working along the Rhône River and characterized their views on this invasive plant. The results indicated that  
517 there is a gap between the proposals of stakeholders defining management plans at a regional level and those of  
518 stakeholders who are implementing actions locally. Whereas the former were focused on providing general  
519 knowledge about invasion processes, the latter were focused on listing technical methods for controlling and  
520 eradicating the plants and providing information regarding the effectiveness of these methods. These different  
521 approaches result, each in their own way, from uncertainties related to biological invasions and may lead to  
522 misunderstandings among stakeholders. Nevertheless, they all agreed on one point: the need to take action

523 against Japanese knotweed *s.l.* While there was an interest in conducting targeted actions to meet local issues, to  
524 do so, the issues must be better defined. This is a challenging task that must involve all types of stakeholders  
525 including environmental managers, scientists, association members, users and the public. Uncertainties relating  
526 to environmental management can only be overcome if management projects result from political projects that  
527 have been collectively discussed and validated.

## 528 ACKNOWLEDGEMENTS

529 We thank the anonymous referees for their helpful comments. We are grateful to the regional water agency (in  
530 partnership with the LTER-ZABR) and the IngECOTech programme (INEE-CNRS) for funding and supporting  
531 this research. We extend our appreciation to Céline Cordani, who collected the data during a master 1 internship  
532 and also participated in the first discussions, to Hervé Tronchère for his help with figure 2, and to John Stella for  
533 his valuable reading and comments. We finally wish to thank all environmental managers who took the time to  
534 gather the documentation included in our analysis corpus.

## 535 LIST OF FIGURES

536 **Fig1** Number of documents produced (a) over time according to the type of manager; (b) according to the  
537 document's category and the type of manager

538 **Fig2** Area under study showing the number and the repartition of the documents produced by each type of  
539 stakeholder

540 **Fig3** Results of the top-down hierarchical clustering of the textual segments: (a) segmentation tree indicating the  
541 proportion of textual segments classified in each class; (b-f) characterization of the 5 classes using similarity  
542 analyses (note: as each graph was independently produced, the size of the character cannot be compared from  
543 one graph to another).

544 **Fig4** Techniques cited over time to control Japanese knotweed *s.l.*

545 **Fig5** Occurrence of different words cited to characterize Japanese knotweed *s.l.* invasion according to the type of  
546 stakeholder

## 547 LIST OF TABLES

548 Table 1 Description of document's categories that compose the textual corpus

549 Table 2 List of the most significant textual segments composing each class resulting from the top-down  
550 hierarchical clustering

551 Table 3 List of lemmas co-occurring with the “efficiency” lemma (co-frequency threshold = 3; distance = 10  
552 lexical forms before and after the considered lemma)  
553 Table 4 List of lemmas co-occurring with the “impact” lemma (co-frequency threshold = 3; distance = 10 lexical  
554 forms before and after the lemma under consideration)  
555 Table 5 Values assessing the association between the textual segments and the classes of the top-down  
556 hierarchical classification according to the stakeholder groups having produced them; (a) metric derived from the  
557 Chi<sup>2</sup> method ( $[\text{observed value} - \text{expected value}] / \text{expected value}$ ); (b) proportion of the classified textual  
558 segments

559

## 560 REFERENCES

- 561 Allison SK (2011) The paradox of invasive species in ecological restoration: do restorationists worry about them  
562 too much or too little? In: Rotherham ID, Lambert RA (eds) *Invasive and Introduced Plants and*  
563 *Animals: Human Perceptions, Attitudes and Approaches to Management*. Routledge, pp 265–276
- 564 Andreu J, Vilà M, Hulme PE (2009) An assessment of stakeholder perceptions and management of noxious alien  
565 plants in Spain. *Environmental Management* 43:1244–1255. doi: 10.1007/s00267-009-9280-1
- 566 Bailey JP, Bímová K, Mandák B (2009) Asexual spread versus sexual reproduction and evolution in Japanese  
567 Knotweed *s.l.* sets the stage for the “Battle of the Clones.” *Biol Invasions* 11:1189–1203. doi:  
568 10.1007/s10530-008-9381-4
- 569 Bailey JP, Conolly AP (2000) Prize-winners to pariahs - a history of Japanese knotweed *s.l.* (Polygonaceae) in  
570 the British Isles. *Watsonia* 23:93–110
- 571 Bardsley D, Edwards-Jones G (2006) Stakeholders’ perceptions of the impacts of invasive exotic plant species in  
572 the Mediterranean region. *GeoJournal* 65:199–210. doi: 10.1007/s10708-005-2755-6
- 573 Bardsley DK, Edwards-Jones G (2007) Invasive species policy and climate change: social perceptions of  
574 environmental change in the Mediterranean. *Environmental Science & Policy* 10:230–242. doi:  
575 10.1016/j.envsci.2006.12.002
- 576 Beerling DJ, Bailey JP, Conolly AP (1994) *Fallopia Japonica* (Houtt.) RonseDecraene. *Journal of Ecology*  
577 82:959–979. doi: 10.2307/2261459

- 578 Bímová K, Mandák B, Pyšek P (2003) Experimental study of vegetative regeneration in four invasive  
579 *Reynoutria* taxa (Polygonaceae). *Plant Ecology* 166:1–11. doi: 10.1023/A:1023299101998
- 580 Binimelis R, Monterroso I, Rodríguez-Labajos B (2007) A social analysis of the bioinvasions of  
581 *Dreissenapolyomorpha* in Spain and *Hydrillaverticillata* in Guatemala. *Environmental Management*  
582 40:555–566. doi: 10.1007/s00267-006-0206-x
- 583 Boyer M, Laval F (2001) Cartographie des renouées du Japon sur le réseau hydrographique du bassin Rhône  
584 Méditerranée Corse (hors Saône Doubs). Agence de l'Eau Rhône Méditerranée Corse
- 585 Boyer M., (2005) L'invasion des cours d'eau par les renouées du Japon *s.l.* : réflexions et propositions pour des  
586 stratégies de lutte efficaces. *Parcs et Réserves* 60:21-29
- 587 Larson MH (2005) The war of the roses: demilitarizing invasion biology. *Frontiers in Ecology and the*  
588 *Environment* 3: 495–500
- 589 Bremner A, Park K (2007) Public attitudes to the management of invasive non-native species in Scotland.  
590 *Biological Conservation* 139:306–314. doi: 10.1016/j.biocon.2007.07.005
- 591 Brochet F, Dubourdieu D (2001) Wine descriptive language supports cognitive specificity of chemical senses.  
592 *Brain and Language* 77:187–196. doi: 10.1006/brln.2000.2428
- 593 Bruno JF, Stachowicz JJ, Bertness MD (2003) Inclusion of facilitation into ecological theory. *Trends in Ecology*  
594 *& Evolution* 18:119–125. doi: 10.1016/S0169-5347(02)00045-9
- 595 Child L, Wade PM (2000) Japanese Knotweed Manual: The Management and Control of an Invasive Alien  
596 Weed (*Fallopia Japonica*). Packard Publishing
- 597 Child LE, Wade M, Wagner M (1998) Cost effective control of *Fallopia japonica* using combination treatments.  
598 In: Starfinger U, Edwards K, Kowarik I, Williamson M (eds) *Plant Invasions: Ecological Mechanisms*  
599 *and Human Responses*. Backhuys Publishers, Leiden, The Netherlands, pp 143–154
- 600 Colautti RI, Bailey SA, Overdijk CDA van, et al. (2006) Characterised and projected costs of nonindigenous  
601 species in Canada. *Biol Invasions* 8:45–59. doi: 10.1007/s10530-005-0236-y
- 602 Dassonville N, Guillaumaud N, Piola F, et al. (2011) Niche construction by the invasive Asian



603 knotweeds(species complex *Fallopia*): impact on activity, abundance and community structure of  
604 denitrifiers and nitrifiers. *Biol Invasions* 13:1115–1133. doi: 10.1007/s10530-011-9954-5

605 Dassonville N, Vanderhoeven S, Gruber W, Meerts P (2007) Invasion by *Fallopia japonica* increases topsoil  
606 mineral nutrient concentrations. *Ecoscience* 14:230–240. doi: [http://dx.doi.org/10.2980/1195-  
607 6860\(2007\)14\[230:IBFJIT\]2.0.CO;2](http://dx.doi.org/10.2980/1195-6860(2007)14[230:IBFJIT]2.0.CO;2)

608 Davis MA, Thompson K, Grime JP (2001) Charles S. Elton and the dissociation of invasion ecology from the rest  
609 of ecology. *Diversity and Distributions* 7: 97–102

610 Davis MA (2009) *Invasion biology*. Oxford University Press, New York, 244 pp

611 Delbart E, Mahy G, Weickmans B, et al. (2012) Can land managers control Japanese knotweed? Lessons from  
612 control tests in Belgium. *Environmental Management* 50:1089–1097. doi: 10.1007/s00267-012-9945-z

613 Dransfield E, Morrot G, Martin J-F, Ngapo T. (2004) The application of a text clustering statistical analysis to  
614 aid the interpretation of focus group interviews. *Food Quality and Preference* 15:477–488. doi:  
615 10.1016/j.foodqual.2003.08.004

616 Dufour S, Piégay H (2009) From the myth of a lost paradise to targeted river restoration: forget natural  
617 references and focus on human benefits. *River Research and Applications* 24:1–14.  
618 DOI: 10.1002/rra.1239

619 Ehrenfeld JG (2010) Ecosystem consequences of biological invasions. *Annual Review of Ecology, Evolution,  
620 and Systematics* 41:59–80. doi: 10.1146/annurev-ecolsys-102209-144650

621 Epanchin-Niell RS, Hastings A (2010) Controlling established invaders: integrating economics and spread  
622 dynamics to determine optimal management. *Ecology Letters* 13:528–541

623 Filippi O, Aronson J (2010) Plantes invasives en région méditerranéenne : quelles restrictions d'utilisation  
624 préconiser pour les jardins et les espaces verts ? *Ecologia Mediterranea* 36 (2): 31-54

625 García-Llorente M, Martín-López B, González JA, et al. (2008) Social perceptions of the impacts and benefits of  
626 invasive alien species: implications for management. *Biological Conservation* 141:2969–2983. doi:  
627 10.1016/j.biocon.2008.09.003

- 628 Genovesi P, Shine C (2004) European Strategy on Invasive Alien Species: Convention on the Conservation of  
629 European Wildlife and Habitats (Bern Convention). Council of Europe
- 630 Genovesi, P (2011) Are we turning the tide? Eradications in times of crisis: how the global community is  
631 responding to biological invasions. In: Veitch, CR, Clout MN and Towns DR (eds) Island Invasives:  
632 Eradication and Management, Proceedings of the International Conference on Island Invasives, IUCN,  
633 pp 5-8
- 634 Gerber E, Krebs C, Murrell C, et al. (2008) Exotic invasive knotweeds (*Fallopia spp.*) negatively affect native  
635 plant and invertebrate assemblages in European riparian habitats. *Biological Conservation* 141:646–  
636 654. doi: 10.1016/j.biocon.2007.12.009
- 637 Gobster PH (2005) Invasive species as ecological threat. *Ecological Restoration* 23(4):261-270
- 638 Gobster PH (2011) Factors affecting people's responses to invasive species management. In: Rotherham ID,  
639 Lambert RA (eds) *Invasive and Introduced Plants and Animals: Human Perceptions, Attitudes and*  
640 *Approaches to Management*. Routledge, pp 249–264
- 641 Hacker SD, Gaines SD (1997) Some implications of direct positive interactions for community species diversity.  
642 *Ecology* 78:1990–2003. doi: 10.1890/0012-9658(1997)078[1990:SIODPI]2.0.CO;2
- 643 Herpigny B, Dassonville N, Ghysels P, Mahy G, Meerts P (2012) Variation of growth and functional traits of  
644 invasive knotweeds (*Fallopia spp.*) in Belgium. *Plant Ecology* 213:419–430
- 645 Heywood V, Brunel S (2011) Code of Conduct on Horticulture and Invasive Alien Plants, Council of Europe  
646 Publishing, pp 24–27
- 647 Horan RD, Perrings C, Lupi F, Bulte EH (2002) Biological pollution prevention strategies under ignorance: the  
648 case of invasive species. *American Journal of Agricultural Economics* 84:1303–1310
- 649 Hulme PE (2006) Beyond control: wider implications for the management of biological invasions. *Journal of*  
650 *Applied Ecology* 43:835–847. doi: 10.1111/j.1365-2664.2006.01227.x
- 651 Jurin RR, Roush DE, Danter KJ (2010) *Environmental Communication: Skills and Principles for Natural*  
652 *Resource Managers, Scientists and Engineers*. Springer, Dordrecht, London, New York

- 653 Lambdon PW, Pyšek P, Basnou C, et al. (2008) Alien flora of Europe: species diversity, temporal trends,  
654 geographical patterns and research needs. *Preslia* 80:101–149
- 655 Larson BM (2005) The war of the roses: demilitarizing invasion biology. *Frontiers in Ecology and the*  
656 *Environment* 3:495–500. doi: 10.1890/1540-9295(2005)003[0495:TWOTRD]2.0.CO;2
- 657 Larson BM (2010) Embodied realism and invasive species in: de Laplante K, Brown B, Peacock K (eds)  
658 *Handbook of the Philosophy of Science*. Elsevier, London 11: 133–50
- 659 Lebart L, Salem A, Berry L (1998) *Exploring Textual Data*. Springer
- 660 Lévêque C, Tabacchi É, Menozzi M-J (2012) Les espèces exotiques envahissantes, pour une remise en cause des  
661 paradigmes écologiques. *Sciences Eaux&Territoires* 6:2–9
- 662 Levine JM (2000) Species diversity and biological invasions: relating local process to community pattern.  
663 *Science* 288:852–854. doi: 10.1126/science.288.5467.852
- 664 Levine JM, Adler PB, Yelenik SG (2004) A meta-analysis of biotic resistance to exotic plant  
665 invasions. *Ecology Letters* 7:975–989. doi: 10.1111/j.1461-0248.2004.00657.x
- 666 Levine JM, Vilà M, Antonino CMD, et al. (2003) Mechanisms underlying the impacts of exotic plant invasions.  
667 *Proceedings of the Royal Society B* 270:775–781. doi: 10.1098/rspb.2003.2327
- 668 Liu S, Sheppard A, Kriticos D, Cook D (2011) Incorporating uncertainty and social values in managing invasive  
669 alien species: a deliberative multi-criteria evaluation approach. *Biol Invasions* 13:2323–2337. doi:  
670 10.1007/s10530-011-0045-4
- 671 Loewenstein G, Lerner JS (2003) The role of affect in decision making. In: Davidson RJ, Scherer KR,  
672 Goldsmith HH (eds) *Handbook of Affective Sciences*. Oxford University Press, New York, pp 619–642
- 673 MacArthur R, Levins R (1967) The limiting similarity, convergence, and divergence of coexisting species. *The*  
674 *American Naturalist* 101:377–385
- 675 Mack RN (2001) Motivations and consequences of the human dispersal of plants. In: McNeely JA (ed) *The*  
676 *Great Reshuffling: Human Dimensions of Invasive Alien Species*. IUCN, pp 23–34

- 677 Marchand P, Ratinaud P (2012) L'analyse de similitude appliquée aux corpus textuels: les primaires socialistes  
678 pour l'élection présidentielle française (septembre-octobre 2011). Actes des 11ème Journées  
679 Internationales d'Analyse Statistique des Données Textuelles, JADT 2012, pp 687–699
- 680 Marshall NA, Friedel M, van Klinken RD, Grice AC (2011) Considering the social dimension of invasive  
681 species: the case of buffel grass. *Environmental Science & Policy* 14:327–338. doi:  
682 10.1016/j.envsci.2010.10.005
- 683 McHugh JM (2006) A review of literature and field practices focused on the management and control of invasive  
684 knotweed. The Nature Conservancy. West Haven, Vermont
- 685 McNeely JA (2001) *The Great Reshuffling: Human Dimensions of Invasive Alien Species*. IUCN
- 686 McNeely JA (2011) Xenophobia or conservation: some human dimensions. In: Rotherham ID, Lambert RA  
687 (eds) *Invasive and Introduced Plants and Animals: Human Perceptions, Attitudes and Approaches to*  
688 *Management*. Routledge, pp 19–36
- 689 Müller S (2004) *Plantes Invasives en France: Etat des Connaissances et Propositions d'Actions*. Publications  
690 Scientifiques du Muséum National d'Histoire Naturelle
- 691 Ness JH, Morales MA, Kenison E, et al. (2013) Reciprocally beneficial interactions between introduced plants  
692 and ants are induced by the presence of a third introduced species. *Oikos* 122:695–704, doi:  
693 10.1111/j.1600-0706.2012.20212.x
- 694 O'Brien W (2006) Exotic invasion, nativism and ecological restoration: on the persistence of a contentious  
695 debate. *Ethics, place and environment* 9:63–77
- 696 Olivier J-M, Carrel G, Lamouroux N, et al. (2009) The Rhône River Basin. In: Tockner K, Uehlinger U and  
697 Robinson C (eds) *Rivers of Europe*. Elsevier, pp 247-296
- 698 Pahl-Wostl C (2006) The importance of social learning in restoring the multifunctionality of rivers and  
699 floodplains. *Ecology and society* 11(1):10
- 700 Parepa M, Fischer M, Krebs C, Bossdorf O (2014) Hybridization increases invasive knotweed success.  
701 *Evolutionary Applications* 7:413–420

- 702 Parr WV, Mouret M, Blackmore S, et al. (2011) Representation of complexity in wine: influence of expertise.  
703 Food Quality and Preference 22:647–660
- 704 Patt A, Zeckhauser R (2000) Action bias and environmental decisions. Journal of Risk and Uncertainty 21:45–72
- 705 Perrings C, Williamson MH, Dalmazzone S (eds) (2000) The Economics of Biological Invasions. Edward Elgar  
706 Publishing, 264 pp
- 707 Pimentel D, Lach L, Zuniga R, Morrison D (2000) Environmental and economic costs of nonindigenous species  
708 in the United States. BioScience 50:53–65
- 709 Popovici J, Bertrand C, Jacquemoud D, et al. (2011) An allelochemical from *Myrica gale* with strong phytotoxic  
710 activity against highly invasive *Fallopia x bohemica* taxa. Molecules 16:2323–2333. doi:  
711 10.3390/molecules16032323
- 712 R Core Team (2013) R: a Language and Environment for Statistical Computing. R Foundation for  
713 Statistical Computing, Vienna, Austria
- 714 Ratinaud P, Déjean S (2009) IRaMuTeQ : Implémentation de la Méthode ALCESTE d'Analyse de Texte dans  
715 un Logiciel Libre
- 716 Reinert A (1983) Une méthode de classification descendante hiérarchique : application à l'analyse lexicale par  
717 contexte. Cahiers de l'Analyse des Données 8:187–198
- 718 Reinert M (1990) Une méthode de classification des énoncés d'un corpus présentée à l'aide d'une application.  
719 Cahiers de l'Analyse des Données 15:21–36
- 720 Richardson DM, Pyšek P, Rejmánek M, et al. (2000) Naturalization and invasion of alien plants: concepts and  
721 definitions. Diversity and Distributions 6:93–107. doi: 10.1046/j.1472-4642.2000.00083.x
- 722 Rotherham ID, Lambert RA (2011b) Invasive and Introduced Plants and Animals: Human Perceptions, Attitudes  
723 and Approaches to Management. Routledge, 352 pp
- 724 Rouifed S (2011) Bases Scientifiques pour un Contrôle des Renouées Asiatiques : Performances du Complexe  
725 Hybride *Fallopia* en Réponse aux Contraintes Environnementales. PhD, Lyon 1 University

- 726 Rouré H, Reinert M (1993) Analyse d'un entretien à l'aide d'une méthode d'analyse lexicale. Actes du Colloque  
727 des Secondes Journées Internationales d'Analyse de Données Textuelles, JADT 1993. Paris, pp 418–  
728 428
- 729 Selge S, Fischer A, van der Wal R (2011) Public and professional views on invasive non-native species – a  
730 qualitative social scientific investigation. *Biological Conservation* 144:3089–3097. doi:  
731 10.1016/j.biocon.2011.09.014
- 732 Shapiro AM (2002) The Californian urban butterfly fauna is dependent on alien plants. *Diversity and*  
733 *Distributions* 8:31–40. doi: 10.1046/j.1366-9516.2001.00120.x
- 734 Shaw RH, Seiger LA (2002) Japanese knotweed. In: Driesche RV, Blossey B, Hoddle M, et al. (eds) *Biological*  
735 *Control of Invasive Plants in the Eastern United States*. USDA Forest Service, pp 159–166
- 736 Simberloff D, Parker IM, Windle PN (2005) Introduced species policy, management, and future research needs.  
737 *Frontiers in Ecology and the Environment* 3:12–20
- 738 Simberloff D (2006) Invasional meltdown six years later—important phenomenon, unfortunate metaphor, or  
739 both? *Ecology Letters* 9: 912–19
- 740 Simberloff D (2012) Nature, natives, nativism and management: worldviews underlying controversies in  
741 invasion biology. *Environmental Ethics* 34:5–25
- 742 Simberloff D (2014) The “balance of Nature” – evolution of a panchreston. *PLoS Biol* 12(10):e1001963.  
743 doi:10.1371/journal.pbio.1001963
- 744 Sims C, Finnoff D (2013) When is a “wait and see” approach to invasive species justified? *Resource and Energy*  
745 *Economics* 35:235–255
- 746 Starfinger U, Kowarik I, Rode M, Schepker H (2003) From desirable ornamental plant to pest to accepted  
747 addition to the flora? – the perception of an alien tree species through the centuries. *Biological Invasions*  
748 5:323–335
- 749 Stromberg JC, Chew MK, Nagler PL, Glenn EP (2009) Changing perceptions of change: the role of scientists in  
750 *Tamarix* and river management. *Restoration Ecology* 17:177–186

- 751 Subramaniam B (2001) The aliens have landed! Reflections on the rhetoric of biological invasions. *Meridians*  
752 2(1):26–40
- 753 Sunstein CR, Zeckhauser R (2011) Overreaction to fearsome risks. *Environmental and Resource Economics*  
754 48:435–449
- 755 Tiebre M-S, Bizoux J-P, Hardy OJ, Bailey JP, Mahy G (2007) Hybridization and morphogenetic variation in the  
756 invasive alien *Fallopia* (Polygonaceae) complex in Belgium. *American Journal of Botany* 94:1900–  
757 1910
- 758 Vanderhoeven S, Dassonville N, Meerts P (2005) Increased topsoil mineral nutrient concentrations under exotic  
759 invasive plants in Belgium. *Plant Soil* 275:169–179
- 760 Vanderhoeven S, Piqueray J, Halford M, et al. (2011) Perception and understanding of invasive alien species  
761 issues by Nature conservation and horticulture professionals in Belgium. *Environmental Management*  
762 47:425–442
- 763 Vitousek PM, D’Antonio CM, Loope LL, et al. (1997) Introduced species: a significant component of human-  
764 caused global change. *New Zealand Journal of Ecology* 21(1):1–16
- 765 Williamson M (1999) Invasions. *Ecography* 22:5–12
- 766