



Aging of the frictional properties induced by temperature variations

Jean-Christophe Géminard, Eric Bertin

► To cite this version:

Jean-Christophe Géminard, Eric Bertin. Aging of the frictional properties induced by temperature variations. *Physical Review E : Statistical, Nonlinear, and Soft Matter Physics*, American Physical Society, 2010, <10.1103/PhysRevE.82.056108>. <hal-01238240>

HAL Id: hal-01238240

<https://hal.archives-ouvertes.fr/hal-01238240>

Submitted on 4 Dec 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Aging of the frictional properties induced by temperature variations

Jean-Christophe Géminard and Eric Bertin

Laboratoire de Physique, Ecole Normale Supérieure de Lyon, CNRS, Université de Lyon, 46 Allée d'Italie, 69364 Lyon Cedex 07, France

(Received 16 June 2010; published 9 November 2010)

The dry frictional contact between two solid surfaces is well known to obey Coulomb friction laws. In particular, the static friction force resisting the relative lateral (tangential) motion of solid surfaces, initially at rest, is known to be proportional to the normal force and independent of the area of the macroscopic surfaces in contact. Experimentally, the static friction force has been observed to slightly depend on time. Such an aging phenomenon has been accounted for either by the creep of the material or by the condensation of water bridges at the microscopic contact points. By studying a toy model, we show that the small uncontrolled temperature changes of the system can also lead to a significant increase of the static friction force.

DOI: [10.1103/PhysRevE.82.056108](https://doi.org/10.1103/PhysRevE.82.056108)

PACS number(s): 81.40.Pq, 46.55.+d, 65.40.De, 62.20.Hg

I. INTRODUCTION

Granular materials or more generally macroscopic solids in frictional contact at mechanical equilibrium are *a priori* considered as athermal systems, meaning that thermal agitation has no significant feedback on the mechanical degrees of freedom at the grain scale. However, uncontrolled temperature variations inevitably produce dilations of materials and one can wonder whether they can alter the properties, in particular the mechanical stability, of such systems.

Several recent studies showed that temperature cycles, even of small amplitude, can induce the slow compaction of dry granular materials [1–4]. The mechanisms, in particular the role played by the confining walls, are still under debate but one can anyway conclude that the temperature variations cause *aging*: the properties of the material, here at least the density, evolve in an irreversible manner with time.

In granular matter, several sources of aging were identified. First, for “dry” granular materials in a humid atmosphere, the condensation of microscopic liquid bridges at the contacts between grains leads to a cohesive force [5]: the nucleation of the bridges being an activated process, the mechanism leads to a logarithmic increase of the angle of avalanche [5,6] and of the static frictional coefficient [7,8] with time. Second, for immersed granular materials, chemical reactions at the surface of the grains result in soldering them to each other [6]. Again, the angle of avalanche [9] and the static frictional coefficient [7,8] are observed to increase logarithmically with time.

In the solid friction [10,11], which characterizes the mechanical contact between solid surfaces, a similar aging phenomenon also takes place: the static frictional coefficient is also observed to increase logarithmically with time for contacts between various materials such as metals [12], rocks [13], and the Bristol paper [14]. Again, for dry friction in a humid atmosphere, the condensation of liquid bridges at the microcontacts between the flat but nevertheless rough surfaces can account for a part of the phenomenon [15]. However, when effects of the humidity are suppressed, aging is still observed. The phenomenon is due to the creep of the material at the microcontacts, as proven by the dependence of the aging dynamics on the overall temperature of the system [16]. Experiments even exhibited a logarithmic aging of the microcontacts themselves [17].

To our knowledge, the possibility for the temperature variations to be responsible for a part of the aging observed in solid friction was still not evaluated. We can guess that the dilation or contraction of the materials which result from a change in the temperature alter the stress distribution between the microcontacts and that, as a result, the system can evolve with time due to a pinning-depinning dynamics. Such dynamics of the microcontacts has been proposed to explain the dissipation associated with the relative motion of the surfaces and, thus, to account for the dynamic friction [18–20] but effects of dilations have not been evaluated. The study of the hysteresis cycle of the tangential force induced by a quasistatic cyclic displacement is a related but slightly different problem [21]: small displacements, even small enough not to be considered as macroscopic sliding, induce irreversible microslips and a slow evolution of the system with time [22]. The temperature variations would have a slightly different effect as they rather induce random stress variations correlated at long range.

In the present paper, we propose the study of a minimal model mimicking the frictional contact between two, nominally flat, solid surfaces, a slider on top of a horizontal substrate. We first consider that, due to roughness, the real contact between the surfaces reduces to a large, but finite, number of microcontacts themselves belonging to a small number of mesoscopic, coherent, contact regions [23]. Such coherent regions are composed of a large number of microcontacts, so that their contact with the substrate obeys the Amonton-Coulomb law for static friction. However, the number of microcontacts in a coherent region is finite [24] so that the associated static friction coefficient, μ_s , is sensitive to the local microstructures in regard and, thus, depends on the position on the substrate. In addition, coherent regions are elastically connected one to another by the materials the slider and the substrate are made of.

In order to mimic the practical situation in the simplest model, we further reduce the problem to the study of a system consisting of two sliders connected by a linear spring. Each of the sliders accounts for one of the coherent regions in frictional contact with a flat substrate whereas the spring accounts for the elastic coupling between them. We shall demonstrate that, when the system is subjected to thermal dilations, the mechanical stability of the slider, accounted for by an effective static frictional coefficient, exhibits a signifi-




FIG. 1. Sketch of the considered situation.

cant increase with time due to the distributed values of the static friction coefficient μ_s .

II. MODEL

A. Description

The system (mass M), subjected to its own weight Mg , lies on a flat and horizontal surface. In a simple approach, we further assume that each contact region sustains half the weight of the system, so that the normal force acting on one slider is $N=Mg/2$. We introduce the rest distance l_0 between the two contact regions at the temperature T_0 . Due to the elasticity of the slider material (assumed to be much softer than the substrate), the small sliders are connected by a spring of length l_0 and stiffness k and, being identical, they are associated with the same inertia, i.e., the same mass $m \equiv M/2$. With these assumptions, the system reduces to two sliders, connected by a spring, moving on a flat surface as sketched in Fig. 1. We denote by x_1 and x_2 the positions of the small sliders 1 and 2, respectively, such that the length of the spring $l=x_2-x_1$.

At rest, the contact between each slider and the substrate is characterized by the static frictional coefficient μ_s , which accounts for the value that the horizontal force must overcome to move the slider. Due to the heterogeneity of the surfaces in regard, the local static frictional coefficient might take, at random, different values, $\mu_{s,1}$ and $\mu_{s,2}$, for the two contact regions. By contrast, we characterize the dynamical friction by a single constant value of μ_d . Indeed, the surface explored by the sliders being homogeneous in average, the dissipative interaction with the substrate can reasonably be characterized, over the whole surface, by a single dynamical frictional coefficient, μ_d , which quantifies the rate of energy dissipation. We further assume, in agreement with standard observations, that $\mu_d < \mu_s$.

The dynamics of the system is induced by temperature changes which, due the thermal dilation of the materials, lead to changes in the natural length of the spring l according to

$$l(T) = l_0[1 + \kappa(T - T_0)], \quad (1)$$

where T stands for the temperature and κ stands for the thermal-expansion coefficient of the slider material. We assume, for the sake of simplicity, that the substrate does not dilate, so that the positions x_1 and x_2 are not altered by the temperature changes and that the spring constant does not depend on the temperature. Thus, due to the thermal dilation, the spring expands or contracts so that the slider 2 is subjected to the tangential force

$$F_{1 \rightarrow 2} = -k(x_2 - x_1 - l), \quad (2)$$

where we assume the x axis to be oriented from the slider 1 to the slider 2. In the same way, the slider 1 is subjected to

$F_{2 \rightarrow 1} = -F_{1 \rightarrow 2}$. If $|F_{1 \rightarrow 2}|$ exceeds one of the static friction forces, $\mu_{s,1}N$ or $\mu_{s,2}N$, the corresponding slider starts moving and the system rearranges. After the slider has stopped, the corresponding value of the static frictional coefficient has changed because the slider lies at a different position on the substrate. The new value of the frictional coefficient is drawn at random from a distribution $p(\mu_s)$. In addition, we assume for consistency that μ_s is always larger than the dynamic frictional coefficient μ_d , namely, $p(\mu_s) = 0$ for $\mu_s < \mu_d$.

The aim of this study is to account for the evolution of the system properties, thus of $\mu_{s,1}$ and $\mu_{s,2}$, as time elapses (aging).

B. System of equations

The problem reduces to the study of two sliders, connected by a spring, in frictional contact with a substrate.

First, one can write the equations governing the dynamics of the sliders when in motion. Introducing the thermal dilation $\theta \equiv \kappa(T - T_0)$, one gets

$$\begin{aligned} m\ddot{x}_1 &= -k[x_1 - x_2 + l_0(1 + \theta)] - \mu_d NS(\dot{x}_1), \\ m\ddot{x}_2 &= -k[x_2 - x_1 - l_0(1 + \theta)] - \mu_d NS(\dot{x}_2), \end{aligned} \quad (3)$$

where S denotes the sign function. The dynamics is characterized by a typical time scale $\tau_{\text{dyn}} = \sqrt{m/k}$. We remark that, in practice, $\tau_{\text{dyn}} \sim l_0/c_s$, where c_s stands for the speed of sound in the material the macroscopic solid is made of. For a typical size $l_0 \sim 10$ cm and usual values of c_s (about a few kilometers per second), we estimate $\tau_{\text{dyn}} \sim 10^{-6} - 10^{-5}$ s.

Second, the static solid friction is accounted for by the coefficients $\mu_{s,j}$ ($j=1, 2$) such that, initially at rest, the slider j starts moving and, thus, the system rearranges when

$$k|x_2 - x_1 - l_0(1 + \theta)| = \mu_{s,j}N. \quad (4)$$

C. Rearrangements

From now on, we assume that after n rearrangements, the sliders are at the positions x_1^n and x_2^n associated with the values $\mu_{s,1}^n$ and $\mu_{s,2}^n$ of the static frictional coefficients. From condition (4), one notes that the rearrangements involve the weakest slider, namely, the one associated to the smallest value of the static frictional coefficient. For convenience, we denote

$$\mu_{\text{min}}^n \equiv \min(\mu_{s,1}^n, \mu_{s,2}^n) \equiv \mu_{s,i_n}^n, \quad (5)$$

where we introduce the index i_n of the corresponding slider. With these definitions, we get the value θ_c^n of the dilation θ at which the $(n+1)$ th rearrangement occurs,

$$\theta_c^n = \frac{x_2^n - x_1^n}{l_0} - 1 + \mu_{\text{min}}^n \tilde{N}S(\dot{\theta}), \quad (6)$$

where the dimensionless normal force $\tilde{N} = N/(kl_0)$ has been introduced to lighten the notations. Equation (6) holds true when the system dilates ($\dot{\theta} > 0$) or contracts ($\dot{\theta} < 0$), with the sign $S(\dot{\theta})$ being evaluated before the rearrangement, for $\theta \rightarrow \theta_c^n$.

For $\theta = \theta_c^n$, the slider i_n moves and reaches a novel static position such that

$$\Delta x_{i_n} = 2(-1)^{i_n}(\mu_{\min}^n - \mu_d)\tilde{N}l_0S(\dot{\theta}). \quad (7)$$

One can easily show that the second slider necessarily remains at rest so that the rearrangement induces a displacement

$$\Delta x_G^n = (-1)^{i_n}(\mu_{\min}^n - \mu_d)\tilde{N}l_0S(\dot{\theta}) \quad (8)$$

of the center of mass G of the two sliders.

In conclusion, when the dilation θ reaches the critical value θ_c^n , the weakest slider i_n moves by Δx_{i_n} and reaches a novel static position whereas the other slider remains at rest. As a consequence, the static frictional coefficient μ_{s,i_n} takes a new random value, μ_{s,i_n}^{n+1} , associated with the new position, $x_{i_n}^{n+1} = x_{i_n}^n + \Delta x_{i_n}$. The position of the other slider and the corresponding static frictional coefficient remain unchanged.

III. NUMERICAL ANALYSIS

We shall consider the temporal evolution of the system when subjected to aleatory temperature changes. In Sec. III A, we describe how the temperature changes are accounted for and, then, in Sec. III B we report the behavior of the system. For practical purposes, we consider in this section a Gaussian distribution $p(\mu_s)$, namely,

$$p(\mu_s) = \frac{1}{\sqrt{2\pi\sigma_\mu^2}} \exp\left[-\frac{(\mu_s - \bar{\mu}_s)^2}{2\sigma_\mu^2}\right]. \quad (9)$$

Although this distribution does not, strictly speaking, satisfy the condition $p(\mu_s) = 0$ for $\mu_s < \mu_d$, this condition will be satisfied in practice as we restrict our study to the case $\sigma_\mu \ll (\bar{\mu}_s - \mu_d)$.

A. Temperature changes

We shall assume that the temperature T of the system fluctuates around the temperature T_0 such that θ is distributed according to a Gaussian distribution $\psi(\theta)$ with zero mean and variance σ_θ^2 . In addition, due to its thermal inertia, the system exhibits a thermal characteristic time τ_{th} which limits the dynamics of the temperature changes and, thus, of the dilations. We assume that the temperature variations occur on time scales much larger than the typical time scale of the rearrangement dynamics, that is, $\tau_{th} \gg \tau_{dyn}$.

In order to account for the finite value of τ_{th} , we consider the evolution of the system in the following way: at time $t_q = q\tau_{th}$ (q is integer), the system is described by the positions x_1^n and x_2^n , the frictional coefficients $\mu_{s,1}^n$ and $\mu_{s,2}^n$, and the dilation $\theta(t_q)$. A new value of $\theta(t_{q+1})$ is drawn at random from the distribution $\psi(\theta)$. Then, we consider $\dot{\theta}$ as a constant over the time interval $[t_q, t_{q+1}]$, namely,

$$\dot{\theta} = \frac{\theta(t_{q+1}) - \theta(t_q)}{\tau_{th}}, \quad (10)$$

and one can calculate θ_c^n from Eq. (6). The corresponding positions x_1^{n+1} and x_2^{n+1} after the rearrangement are calculated




FIG. 2. Maximum frictional coefficient μ_{\max} vs dimensionless time t/τ_{th} —the maximum μ_{\max} is observed to increase almost logarithmically with time t , the aging being more effective when the amplitude σ_θ of the temperature variations is larger ($\mu_d = 0.5$, $\bar{\mu}_s = 0.6$, $\sigma_\mu = 0.01$, and $\tilde{N} = 10^{-7}$).

according to Eq. (7), the smallest frictional coefficient among $\mu_{s,1}^n$ and $\mu_{s,2}^n$ is drawn anew from $p(\mu_s)$ and the other remains unchanged. The process is repeated while $\theta_c^n \leq \theta(t_{q+1})$ if $\dot{\theta} > 0$ or $\theta_c^n \geq \theta(t_{q+1})$ if $\dot{\theta} < 0$. When the evolution stops, the dilation between t_q and t_{q+1} has led to r rearrangements. The system has reached a new state and the process can be iterated by choosing randomly a new value $\theta(t_{q+2})$.

B. Numerical results

When the system is subjected to temperature changes, the values of the static frictional coefficients continuously change in time. As the rearrangements only involve the weakest slider, the maximum of the values $\mu_{s,1}^n$ and $\mu_{s,2}^n$,

$$\mu_{\max}^n \equiv \max(\mu_{s,1}^n, \mu_{s,2}^n), \quad (11)$$

cannot decrease and thus, on average, continuously increases with time t (Fig. 2). We indeed observe that μ_{\max} increases rather logarithmically with time t and that, for instance, the aging is faster for larger σ_θ .

In addition, due to the random motion of the center of mass associated with the rearrangements [Eq. (7)], the system diffuses. We report the mean-square displacement $\langle x_G^2 \rangle$ as a function of time t (Fig. 3) and observe that $\langle x_G^2 \rangle \sim \ln(t/\tau_{th})$, the diffusion being faster for larger σ_θ .

In Sec. IV, we shall discuss further these results theoretically, describing analytically how the aging and the diffusion depend on the parameters of the problem at long times.

IV. THEORETICAL ANALYSIS

A. Dilation and rearrangements

In the first step, we consider the number r of rearrangements that occur during the time interval τ_{th} . Considering Eqs. (6) and (7), the difference between two successive values of the dilation at which rearrangements occur can be estimated as




FIG. 3. Normalized mean-square displacement $\langle x_G^2 \rangle / \sigma_\theta^2 t_0^2$ vs dimensionless time t / τ_{th} —the mean-square displacement $\langle x_G^2 \rangle$ increases logarithmically with time t , the diffusion being faster when the amplitude σ_θ of the temperature variations is larger ($\mu_d=0.5$, $\bar{\mu}_s=0.6$, $\sigma_\mu=0.01$, and $\tilde{N}=10^{-7}$).

$$|\theta_c^{n+1} - \theta_c^n| \approx 2(\bar{\mu}_s - \mu_d)\tilde{N}S(\dot{\theta}). \quad (12)$$

Thus, a dilation or a contraction of amplitude $\Delta\theta$ leads to a typical number of rearrangements $r \approx |\Delta\theta| / |\theta_c^{n+1} - \theta_c^n|$. To average over the fluctuations of $\Delta\theta$, we define it quantitatively as the difference between two values θ and θ' randomly drawn from $\psi(\theta)$. Hence, $\Delta\theta$ is a Gaussian random variable with zero mean and variance $2\sigma_\theta^2$. We thus find for the average value $\langle r \rangle$,

$$\langle r \rangle \approx \frac{\sigma_\theta}{\sqrt{\pi}(\bar{\mu}_s - \mu_d)\tilde{N}}. \quad (13)$$

Comparison with the results of numerical simulations (Fig. 4) essentially confirms this prediction. A careful analysis of the numerical data however reveals the presence of a small shift. The average number of rearrangements is found to be described by the phenomenological form,

$$\langle r \rangle \approx \frac{\sigma_\theta - \mu_d\tilde{N}}{\sqrt{\pi}(\bar{\mu}_s - \mu_d)\tilde{N}}. \quad (14)$$

The small offset $\mu_d\tilde{N}$ is intuitively expected to result from the absence of rearrangements when $\dot{\theta}$ changes sign.




FIG. 4. (a) Number of rearrangements n vs time t —the number of rearrangements n increases, to within small fluctuations, linearly with the time t . (b) Average number of rearrangements $\langle r \rangle$ during τ_{th} vs amplitude σ_θ —in average, the slope dn/dt or equivalently $\langle r \rangle$ depends linearly on the amplitude σ_θ according to Eq. (14).

From now on, we shall consider the evolution of the system in terms of the number of rearrangements n . The temporal evolution of the system can then be assessed by considering the simple relation between n and time t .

B. Aging of the static frictional coefficient

It is particularly interesting to focus on the dynamics of μ_{\max}^n as this quantity plays an important role in the stability of the system (Sec. V). Actually, one simply needs to notice that the dynamics consists in drawing randomly two values $\mu_{s,1}^1$ and $\mu_{s,2}^2$ in the initial step $n=1$ and then drawing at each step a new value of the frictional coefficient to replace the smallest one. As a result, the value μ_{\max}^n is the maximum value of the set of $n+1$ random values drawn independently. This mechanism is actually very close to the one appearing in a standard model of glassy dynamics, namely, the Barrat-Mézard model [25,26]. The statistics of μ_{\max}^n then boils down to an extreme value problem of independent and identically distributed random variables, for which standard results are well known [27,28]. In particular, the distribution of μ_{\max}^n belongs for large n to one of the three classes of extreme value statistics, namely, the Gumbel, Weibull, or Fréchet class. However, we are here more interested in the typical value of μ_{\max}^n rather than by its relative fluctuations. This typical value can be estimated through the following simple scaling argument. We denote as $F(\mu_s)$ the complementary cumulative probability distribution

$$F(\mu_s) = \int_{\mu_s}^{\infty} p(\mu) d\mu. \quad (15)$$

Having μ_{\max}^n smaller than a given value μ is equivalent to the fact that the $n+1$ random values drawn dynamically are below the value μ . The corresponding probability is simply $[1 - F(\mu)]^{n+1}$. The typical value μ_{typ} of μ_{\max}^n then satisfies

$$[1 - F(\mu_{\text{typ}})]^{n+1} \approx \frac{1}{e}, \quad (16)$$

which for large n takes the form

$$1 - F(\mu_{\text{typ}}) \approx e^{-1/(n+1)} \approx 1 - \frac{1}{n} \quad (17)$$

so that finally $F(\mu_{\text{typ}}) \approx 1/n$. If the upper wing of the distribution $p(\mu)$ decays as $p(\mu) \sim e^{-c\mu^\alpha}$, where c and α are positive constants (for instance, $\alpha=2$ for a Gaussian distribution), then $F(\mu)$ behaves in the same way as $p(\mu)$ up to algebraic prefactors. It follows that

$$\mu_{\text{typ}} \approx \bar{\mu}_s + \sigma_\mu c^{-1/\alpha} (\ln n)^{1/\alpha} \quad (18)$$

to leading order in n . Thus, in the case of the Gaussian distribution considered in Sec. III, the simple theoretical argument presented here predicts an increase of μ_{\max} like

$$\mu_{\max} \approx \bar{\mu}_s + \sigma_\mu \sqrt{2 \ln \left(\frac{\langle r \rangle t}{\tau_{th}} \right)}, \quad (19)$$

taking into account the correspondence $n \approx \langle r \rangle t / \tau_{th}$ between the time t and the total number n of rearrangements [Eq. (14)].

C. Slowing down diffusion

In Sec. III, we obtained a logarithmic subdiffusion of the center of mass, which we now would like to interpret. We first observe that the evolution of the system can be divided in successive periods during which the strongest slider remains the same; that is, the frictional coefficient of the weakest slider is repeatedly drawn anew, without exceeding the value μ_{\max} of the strong slider. During such periods, the position of the weakest slider changes due to the rearrangements, and the net displacement ℓ is limited by the extension of the spring. One can estimate $\ell \sim \sigma_{\theta} l_0$. When the chosen frictional coefficient becomes larger than the former value μ_{\max} (which can be called a record breaking), the weakest and strongest sliders get exchanged. Hence, the mean-square displacement of the center of mass after n rearrangements should be on the order of ℓ^2 times the number of record breakings in this sequence of rearrangements.

It is a classical result from statistical record theory [29] that the mean number of records that occur when successively drawing n independent and identically distributed random values is, to leading order, equal to $\ln n$ for large n . Further using the relation $n \approx \langle r \rangle t / \tau_{\text{th}}$, one eventually finds for the mean-square displacement of the center of mass,

$$\langle x_G^2 \rangle \approx \sigma_{\theta}^2 l_0^2 \ln \left(\frac{\langle r \rangle t}{\tau_{\text{th}}} \right), \quad (20)$$

thus accounting for the logarithmic time dependence and the order of magnitude of the numerical data reported in Fig. 3.

V. DISCUSSION

We have shown that the dilations associated with the temperature changes are likely to induce the aging of the physical properties of the system (the maximum frictional coefficient increases with time and the diffusion slows down). At this point, it is particularly interesting to discuss, first, the orders of magnitude of the temperature changes necessary for the aging process to be at stake and, second, the possibility to assess the aging process experimentally.

Let us now consider the order of magnitude of the dilations likely to induce the aging. From Eq. (14), given that $\mu_d \sim 1$, we deduce that the minimal amplitude of the dilations necessary to rearrange the system is on the order of \tilde{N} which can be estimated as follows. Consider that the two contact points, sketched by the sliders 1 and 2, are located at the distance l_0 one from another. Taking into account the Young modulus, Y , the material is made of, one can estimate $k \approx YA/l_0$, where A stands for the surface area of the slider cross section in the perpendicular direction. Considering the density ρ of the material and volume $V \sim Al_0$ of the slider, one can estimate the normal force due to gravity g , $N \sim \rho Al_0 g$. Thus, we get $\tilde{N} \sim \rho g l_0 / Y$. In practice, considering $g = 10 \text{ m s}^{-2}$, $\rho \sim 10^3 \text{ kg m}^{-3}$, $Y \sim 100 \text{ GPa}$, and the typical size $l_0 = 10 \text{ cm}$, we get $\tilde{N} \sim 10^{-8}$. Thus, dilations on the order of 10^{-8} are likely to rearrange the system. Such dilations, the typical thermal-expansion coefficients being of about 10^{-5} K^{-1} correspond to temperature changes of about 10^{-3} K . Thus, in usual experimental conditions, the thermal




FIG. 5. Variation $(\mu_s^{\text{eff}} - \bar{\mu}_s) / \sigma_{\mu}$ vs number $\langle r \rangle t / \tau_{\text{th}}$ —we report data obtained for two different values of the dynamical frictional coefficient (lines: $\mu_d = 0.5$; open squares: $\mu_d = 0.4$). The values of σ_{μ} are chosen in accordance with the values of $\sigma_{\mu} / (\bar{\mu}_s - \mu_d)$ given in the figure ($\bar{\mu}_s = 0.6$, $\tilde{N} = 10^{-7}$, and $\sigma_{\theta} = 10^{-6}$).

dilations produce rearrangements at the contact scale. Even more, it is not obvious to perform a control of the temperature that insures that rearrangements will not occur.

Experimentally, one possible method to highlight the aging process consists in measuring the critical angle of stability: the substrate is slowly tilted until the system starts sliding as a whole for a critical tilt angle, α_c [15]. In this configuration, the sliders are subjected to normal and tangential forces due to gravity and one can relate α_c with an effective static frictional coefficient $\mu_s^{\text{eff}} \equiv \tan \alpha_c$. We assessed numerically the temporal evolution of μ_s^{eff} as described in the Appendix. We observe that the increase in μ_{\max}^{eff} (Fig. 2) results in an almost logarithmic increase in μ_s^{eff} with time t . As suggested in Eq. (19), we report the relative variation $(\mu_s^{\text{eff}} - \bar{\mu}_s) / \sigma_{\mu}$ as a function of the typical number of rearrangements $n \approx \langle r \rangle t / \tau_{\text{th}}$ (Fig. 5). We observe a collapse of the curves corresponding to the same value of $\sigma_{\mu} / (\bar{\mu}_s - \mu_d)$, which suggests

$$\mu_s^{\text{eff}}(t) = \bar{\mu}_s + \sigma_{\mu} \Phi \left(\frac{\sigma_{\mu}}{\bar{\mu}_s - \mu_d}, \frac{\langle r \rangle t}{\tau_{\text{th}}} \right). \quad (21)$$

From the numerical data, the function Φ depends almost logarithmically on t at long times. Asymptotically, as a function of the number of rearrangements $n(t)$, $\mu_s^{\text{eff}} \sim \bar{\mu}_s + \phi \sigma_{\mu} (\ln n)^{\beta}$, with β on the order of unity and ϕ a constant which only depends on $\sigma_{\mu} / (\bar{\mu}_s - \mu_d)$. The aging phenomenon is drastically enhanced when the width σ_{μ} is increased.

It is particularly interesting to estimate the expected order of magnitude of the effect. With $\tilde{N} \sim 10^{-8}$ and typical values $\bar{\mu}_s \sim 0.6$ and $\mu_d \sim 0.5$ of the frictional coefficients, temperature changes of about 0.01 K correspond to $\sigma_{\theta} \sim 10^{-7}$ and, thus, to typically $\langle r \rangle \sim 50$ rearrangements during the characteristic time τ_{th} . From a typical thermal conductivity λ of about 1 W/(m K) and a typical heat capacity C of about 10^6 J/m^3 , one can estimate from the diffusion coefficient λ / C that $\tau_{\text{th}} \sim l_0^2 C / \lambda$ ranges from 10^{-2} to 1 s (indeed much larger than τ_{dyn}) for a typical size of the system ranging from 1 to 10 cm . Considering further a value $\sigma_{\mu} \sim 0.02$ (about 3% of the average value $\bar{\mu}_s$), we expect from Fig. 5 an increase

of about 0.2% of the effective frictional coefficient after $10^5 \tau_{\text{th}} \sim 10^3 - 10^5$ s. Note, however, that for $\sigma_\mu \sim 0.03$ (about 5% of the average value $\bar{\mu}_s$), the effect is drastically enhanced leading to a change of about 5% after the same waiting time. Thus, small temperature variations are likely to induce, in a few hours, an increase of the effective frictional coefficient of about a few percent. Note that, from Eq. (21), we expect the aging process to mainly depend on σ_μ and σ_θ (through $\langle r \rangle$), the effect being expected to be larger for broader distribution of the frictional coefficient and for larger amplitude of the temperature variations. Finally, it is important to note that the thermal dilations are important because of the change $kl_0\theta$ they induce in the force between the two sliders and that the latter change is to be compared with the normal force N . Thus, a smaller k leads to a weaker effect of the temperature changes. This is probably the reason why no or little aging is observed with soft materials [15].

VI. CONCLUSION

Reducing the study of the frictional contact between two solid surfaces to the study of two frictional sliders, connected by a spring, in contact with a flat substrate, we demonstrated that the uncontrolled thermal dilations of the system can lead to a significant increase in the effective static frictional coefficient μ_s^{eff} with time. The evolution of μ_{max} at long times is assessed theoretically and the associated μ_s^{eff} is estimated numerically. Our simplistic model makes it possible to demonstrate that the process leads to an almost logarithmic increase in μ_s^{eff} and we account for the dependence on the parameters, in particular the distribution of the static frictional coefficient associated with the solid-solid contact and the characteristics of the temperature changes (amplitude and characteristic time). Considering typical orders of magnitude of the physical ingredients, we show that temperature changes can have significant effects, which can also contribute to the aging of the frictional coefficient observed experimentally [15].

Finally, one could wonder if the effect reported here would not be specific to the system we chose. We assumed that the contact between two flat and rough surfaces reduces to a small number of coherent regions. On the one hand, one could claim that the contact between the surfaces rather consists of a large number of microscopic contact points and that non-Amonton behavior of the microscopic contacts was revealed experimentally [30]. However, whatever the characteristics of the microscopic contacts (characterized by frictional mechanical properties or not), one can guess that the thermal dilations, which induce a change in the distribution of the force network between the latter, lead to rearrangements of the less stable ones. The process again favors the most stable contacts, and we expect the effective frictional coefficient to increase with time. On the other hand, even considering coherent regions exhibiting frictional properties, one could claim that their number is likely to be significantly larger than considered in the present study. Again, we do not expect an increase in the number of contact regions to change qualitatively the aging process. However, the quantitative effect is difficult to predict theoretically. The study of a

two-dimensional irregular network of sliders is thus pertinent and will be the subject of a forthcoming publication in which the aging and the diffusion, as well as the creep motion induced by a constant tangential force, will be considered.

APPENDIX: EFFECTIVE STATIC FRICTIONAL COEFFICIENT

In this appendix, we present the numerical method used to compute the time evolution of the effective frictional coefficient μ_s^{eff} . To this aim, we consider that the substrate on which the two sliders lie can be tilted by an adjustable angle α .

The dynamics of the system on the horizontal substrate, as described in Sec. III A, is divided in time steps of duration τ_{th} . Within each time step, the sliders evolve through a sequence of rearrangements during which the positions and static frictional coefficients change. Before each rearrangement, the values of $\mu_{s,1}$, $\mu_{s,2}$ and of the force $F_{1 \rightarrow 2}$ are copied to auxiliary variables, and the angle of avalanche corresponding to this precise set of variables is determined through an iterative algorithm that we now describe.

Due to gravity, when the substrate is tilted, the sliders on the incline are both subjected to the same positive force $f = mg \sin \alpha$. Correspondingly, the normal force now becomes $N = mg \cos \alpha$. According to Eq. (4), the slider 1 (2) starts moving if one of the two following conditions is fulfilled,

$$f - F_{1 \rightarrow 2} > \mu_{s,1} N, \quad (\text{A1a})$$

$$f + F_{1 \rightarrow 2} > \mu_{s,2} N. \quad (\text{A1b})$$

One deduces from these relations that, when α is increased, the slider 1 moves first if

$$2F_{1 \rightarrow 2} < (\mu_{s,2} - \mu_{s,1})N, \quad (\text{A2})$$

where $F_{1 \rightarrow 2}$ is given in Eq. (2). The onset of motion occurs precisely when $f \equiv mg \sin \alpha$ reaches the value $f_1 = \mu_{s,1} N + F_{1 \rightarrow 2}$. If condition (A2) is not fulfilled, the slider 2 moves first when f reaches the value $f_2 = \mu_{s,2} N - F_{1 \rightarrow 2}$.

The motion of the slider i induces the motion of the other slider if condition (4) is satisfied, thus if (after simple algebra)

$$\frac{f}{N} \geq \mu_d + \frac{1}{2}(-1)^i(\mu_{s,1} - \mu_{s,2}). \quad (\text{A3})$$

Then, if $f > \mu_d N$, the motion of the two sliders is accelerated, and a macroscopic sliding of the whole system is observed. The corresponding value of $\tan \alpha_c = f/mg$ is recorded. In the opposite case, when $f < \mu_d$, the motion is damped, leading only to a short displacement of the system (creep). Once the system is again at rest, new values of $\mu_{s,1}$ and $\mu_{s,2}$ are drawn from the Gaussian distribution $p(\mu_s)$. The value of the force $F_{1 \rightarrow 2}$ also needs to be recomputed due to the change of the positions x_1 and x_2 . Since the determination of the new positions would require integrating the equations of motion during the damped motion, which is time consuming, we

rather use a simple approximation. The new force $F_{1\rightarrow 2}$ is chosen at random, with a uniform probability, between the values $f - \mu_{s,1}N$ and $\mu_{s,2}N - f$, where $\mu_{s,1}$ and $\mu_{s,2}$ designate the new values of the static frictional coefficient. We are again led back to Eq. (A1), and the process is then iterated until a macroscopic sliding is observed.

If, however, Eq. (A3) is not satisfied, the other slider remains at rest when the slider i rearranges, and the release of the elastic energy loaded in the spring leads to a change

$$\Delta F_{1\rightarrow 2} = -2(-1)^i(\mu_{s,i} - \mu_d)N \quad (\text{A4})$$

in the force $F_{1\rightarrow 2}$ between the sliders [see Eq. (7)]. The value of $\mu_{s,i}$ needs to be drawn anew from $p(\mu_s)$, the other frictional coefficient remaining unchanged, and the process is here again iterated, leading back to Eq. (A1) as long as a macroscopic motion does not emerge.

At the end of this iteration process, the critical value $\tan \alpha_c$ has been obtained, and the evolution on the horizontal substrate resumes (we recall that the evolution on the tilted substrate is only a “virtual” test that does not influence the “real” horizontal evolution). Obviously, the obtained value of $\tan \alpha_c$ depends on the values of $\{F_{1\rightarrow 2}, \mu_{s,1}, \mu_{s,2}, \theta\}$ given at the beginning of the test, which are all stochastic variables. It is thus necessary to average $\tan \alpha_c$ over the dynamics. Since, on the other hand, we wish to obtain the time dependence of the critical angle to study aging effects, we proceed through a time scale separation as follows: $\tan \alpha_c$ is averaged over all the rearrangements occurring in a given time step of duration τ_{th} and also over a large number of independent realizations of the horizontal dynamics, with random initial conditions. Then, the time dependence on a scale larger than τ_{th} remains, and the time step averages $\langle \dots \rangle_{t_q}$ are labeled by the corresponding time $t_q = q\tau_{\text{th}}$, with q as integer, yielding the effective static frictional coefficient $\mu_s^{\text{eff}}(t_q) \equiv \langle \tan \alpha_c \rangle_{t_q}$.

-
- [1] K. Chen, A. Harris, J. Draskovic, and P. Schiffer, *Granular Matter* **11**, 237 (2009).
- [2] K. Chen, J. Cole, C. Conger, J. Draskovic, M. Lohr, K. Klein, T. Scheidemantel, and P. Schiffer, *Nature (London)* **442**, 257 (2006).
- [3] T. Divoux, I. Vassilief, H. Gayvallet, and J.-C. Géminard, in *Proceedings of the 6th International Conference on the Micro-mechanics of Granular Media*, edited by M. Nakagawa and S. Luding (AIP, Golden, CO, 2009).
- [4] T. Divoux, H. Gayvallet, and J.-C. Géminard, *Phys. Rev. Lett.* **101**, 148303 (2008).
- [5] L. Bocquet, E. Charlaix, S. Ciliberto, and J. Crassous, *Nature (London)* **396**, 735 (1998).
- [6] N. Fraysse, H. Thomé, and L. Petit, *Eur. Phys. J. B* **11**, 615 (1999).
- [7] J.-C. Géminard, W. Losert, and J. P. Gollub, *Proceedings of the 4th International Conference on the Micromechanics of Granular Media*, edited by Y. Kishino (A. A. Balkema Publishers, Netherlands, 2001).
- [8] W. Losert, J.-C. Géminard, S. Nasuno, and J. P. Gollub, *Phys. Rev. E* **61**, 4060 (2000).
- [9] H. Gayvallet and J.-C. Géminard, *Eur. Phys. J. B* **30**, 369 (2002).
- [10] F. P. Bowden and D. Tabor, *Friction and Lubrication of Solids* (Clarendon, Oxford, 1950).
- [11] E. Rabinowicz, *Friction and Wear of Material* (Wiley, New York, 1965).
- [12] E. Rabinowicz, *Proc. Phys. Soc. London* **71**, 668 (1958).
- [13] C. H. Scholz, *The Mechanics of Earthquakes and Faulting* (Cambridge University, Cambridge, England, 1990).
- [14] F. Heslot, T. Baumberger, B. Perrin, B. Caroli, and C. Caroli, *Phys. Rev. E* **49**, 4973 (1994).
- [15] J. Crassous, L. Bocquet, S. Ciliberto, and C. Laroche, *Europhys. Lett.* **47**, 562 (1999).
- [16] P. Berthoud, T. Baumberger, C. G’Sell, and J.-M. Hiver, *Phys. Rev. B* **59**, 14313 (1999).
- [17] L. Bureau, T. Baumberger, and C. Caroli, *Eur. Phys. J. E* **8**, 331 (2002).
- [18] G. A. Tomlinson, *Philos. Mag.* **7**, 905 (1929).
- [19] J.-F. Joanny and P.-G. de Gennes, *J. Chem. Phys.* **81**, 552 (1984).
- [20] C. Caroli and P. Nozières, in *Physics of Sliding Surfaces*, NATO Advanced Studies Institute Series E: Applied Science, edited by B. N. J. Persson and E. Tosatti (Kluwer Academic Publishers, Dordrecht, 1996), Vol. 311.
- [21] J. Crassous, S. Ciliberto, E. Charlaix, and C. Laroche, *J. Phys. II* **7**, 1745 (1997).
- [22] U. Olofsson, *Tribol. Int.* **28**, 207 (1995).
- [23] B. N. J. Persson, *J. Phys.: Condens. Matter* **22**, 265004 (2010).
- [24] T. Baumberger and C. Caroli, *Adv. Phys.* **55**, 279 (2006).
- [25] A. Barrat and M. Mézard, *J. Phys. I* **5**, 941 (1995).
- [26] E. Bertin, *J. Phys. A: Math. Theor.* **43**, 345002 (2010).
- [27] E. J. Gumbel, *Statistics of Extremes* (Columbia University Press, New York, 1958); *Statistics of Extremes* (Dover Publication, New York, 2004).
- [28] J. Galambos, *The Asymptotic Theory of Extreme Order Statistics* (Wiley, New York, 1987).
- [29] J. Krug, *J. Stat. Mech.: Theory Exp.* **2007**, P07001.
- [30] L. Bureau, T. Baumberger, and C. Caroli, *Eur. Phys. J. E* **19**, 163 (2006).