



Local interactions and p-best response set

Jacques Durieu, Philippe Solal

► To cite this version:

Jacques Durieu, Philippe Solal. Local interactions and p-best response set. Journal of Applied Mathematics, Hindawi Publishing Corporation, 2014, vol. 2014, 7 p. <10.1155/2014/415686>. <halshs-01002448>

HAL Id: halshs-01002448

<https://halshs.archives-ouvertes.fr/halshs-01002448>

Submitted on 6 Jun 2014

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Research Article

Local Interactions and p -Best Response Set

J. Durieu¹ and P. Solal²

¹CREG, Université Pierre-Mendès-France-de Grenoble, 38400 Saint-Martin d'Hères, France

²GATE Lyon-St-Etienne, Université de Saint-Etienne, 42023 Saint-Etienne, France

Correspondence should be addressed to J. Durieu; jacques.durieu@univ-st-etienne.fr

Received 6 December 2013; Accepted 5 February 2014; Published 26 March 2014

Academic Editor: Pu-yan Nie

Copyright © 2014 J. Durieu and P. Solal. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We study a local interaction model where agents play a finite n -person game following a perturbed best-response process with inertia. We consider the concept of minimal p -best response set to analyze distributions of actions on the long run. We distinguish between two assumptions made by agents about the matching rule. We show that only actions contained in the minimal p -best response set can be selected provided that p is sufficiently small. We demonstrate that these predictions are sensitive to the assumptions about the matching rule.

1. Introduction

There is an extensive literature on evolutionary game theory which investigates the long-run outcomes when a population of boundedly rational agents uses simple rules of adaptation to recurrently play a game in normal form. These models allow for sharp-equilibrium selection results in some classes of games with multiple equilibria. Such an equilibrium can be interpreted as a convention, that is, a pattern of behavior that is customary and expected and regulates much of economic and social life (for a survey, see, among others, Young [1]). A common assumption in this literature is that interactions among agents are local and not global. Each individual may be matched according to some matching rule with a subgroup of the overall population to play a game (for a survey, see, e.g., Weidenholzer [2]). In this paper, we assume that agents have no information about this matching rule. As a consequence, we elaborate two alternative scenarios corresponding to different assumptions made by agents about the matching rule. This point can be interpreted as a consequence of bounded rationality: agents only have an imperfect representation of their environment. We use the concept of (minimal) p -best response set introduced by Tercieux [3] to make predictions for the long-run behavior of the population of agents (this concept is used by Durieu et al. [4] in a fictitious play model with bounded memory). We establish that such predictions are possible for the whole class of finite n -person games. We

also study how assumptions about the matching rule affect these predictions.

Precisely, we consider a finite population of agents located at the vertices of a chordal ring, that is, a ring topology in which each vertex has additional links with other vertices. In particular, we focus on chordal rings of degree $2K$, $K \in \mathbb{N}$, constructed by adding at each vertex $2K - 2$ chords with its closest vertices in a ring network. In order to deal with asymmetric finite n -person games, we distinguish between n classes of agents. A specific role in the game is assigned to every class. Each agent of each class is located at exactly one vertex and two agents of the same class are located in different vertices. In this way, each vertex contains exactly n agents. The game is played recurrently. At each period, one agent of each class is drawn at random to play the game. However, players' identity is not revealed. At the beginning of every period, each agent of each class observes the actions that the agents of other classes located in vertices linked with its own vertex has chosen in the previous period. He uses this information to estimate the probability distribution on the action profiles played by his potential opponents in the current period. These estimations depend on the assumptions made by agents about the matching rule. Each agent believes that he can be matched with agents located at vertices linked with his own location. Precisely, we consider the following two scenarios. In the first scenario, each agent assumes that his potential opponents are drawn jointly by location. In

the second scenario, each agent assumes that his potential opponents are drawn independently. For both scenarios, agents may have the opportunity to choose a best response to their estimation. The concept of p -best response set allows us to study the way in which the distribution of actions that people take in the n classes evolves over time. We show that, for both assumptions about the matching rule, only actions contained in the minimal p -best response set can be selected on the long run provided that p is sufficiently small. For each assumption, an explicit bound of p is given, and we analyze how this critical value evolves when n increases.

Ellison [5, 6] considers a similar local interaction structure: agents are arranged on a ring improved by adding chords between vertices in a regular form. However, since Ellison [5, 6] focuses on symmetric two-player games, each vertex is associated with one agent and each agent has full information about the matching rule. Ellison [5] considers a symmetric 2×2 coordination game. The model predicts that the risk-dominant equilibrium is selected on the long run. Ellison [6] shows that in a symmetric $m \times m$ game, the $1/2$ -dominant equilibrium, if it exists, is selected on the long run. Alós-Ferrer and Weidenholzer [7] consider the weakened solution concept of globally pairwise risk-dominant equilibrium and show that this equilibrium, if it exists, is selected in a symmetric 3×3 coordination game played by agents arranged on a ring. However, a generalization of this result to a larger class of games is not possible. Indeed, Alós-Ferrer and Weidenholzer [7] present an example of a 4×4 coordination game in which the globally pairwise risk-dominant equilibrium is not selected on the long run. From this point of view, the concept of minimal p -best response set is interesting. Since a minimal p -best response exists in every game, it allows us to investigate the long-run outcomes in the whole class of finite n -person games. Moreover, since the concept of minimal p -best response set generalizes the concept of p -dominant equilibrium [8], the result of Ellison [6] is a particular case of our result.

The paper is structured as follows. Section 2 introduces notations. Section 3 introduces the concept of p -best response set. Section 4 introduces the learning model. In Section 5 we present the main results. Section 6 concludes.

2. Notations and Definitions

Let \subseteq denote weak set inclusion and let \subset denote proper set inclusion. We denote by $\lceil \beta \rceil$ the smallest integer greater than or equal to β . For any finite set A , $\Delta(A)$ denotes the set of all probability distributions on A .

Let Γ be a finite n -person strategic-form game. Let X_i be the finite set of pure strategies x_i available to player $i \in I = \{1, 2, \dots, n\}$. We write $\Delta(X_i)$ for the set of probability distributions q_i over X_i for each $i \in I$. Let $q_i(x_i)$ denote the probability mass on strategy x_i . Define the product set $X = \prod_{i \in I} X_i$. Let $\Delta(X)$ be the set of probability distributions on X . Let $X_{-i} = \prod_{j \neq i} X_j$ denote the set of all possible combinations of strategies for the players other than i , with generic elements $x_{-i} = (x_j)_{j \neq i}$. Let $\Delta(X_{-i})$ be the set of probability distributions on X_{-i} with generic elements q_{-i} .

We sometimes identify the element of $\Delta(X_i)$ that assigns probability one to a strategy in X_i with this strategy in X_i .

In this paper, a player's belief about others' strategies takes the form of a probability measure on the product of all opponents' strategy sets. We assume that each player i has expected payoffs represented by the function $u_i : X_i \times \Delta(X_{-i}) \rightarrow \mathbb{R}$.

For each player i and probability distribution $q_{-i} \in \Delta(X_{-i})$, let

$$Br_i(q_{-i}) = \{x_i \in X_i : u_i(x_i, q_{-i}) \geq u_i(x'_i, q_{-i}), \forall x'_i \in X_i\} \quad (1)$$

be the set of pure best responses of i against q_{-i} .

Let $Y = \prod_{i \in I} Y_i$ be a product set where each Y_i is a nonempty subset of X_i . Let $\Delta(Y_{-i})$ denote the set of probability distributions with support in Y_{-i} . Finally, $Br_i(\Delta(Y_{-i}))$ denotes the set of strategies in X_i that are pure best responses of i against some distribution q_{-i} with support in Y_{-i} ; that is,

$$Br_i(\Delta(Y_{-i})) = \bigcup_{q_{-i} \in \Delta(Y_{-i})} Br_i(q_{-i}). \quad (2)$$

3. p -Best Response Sets

We will now introduce the concept of (minimal) p -best response set. Let $p \in [0, 1]$ and let Y_{-i} be a nonempty subset of X_{-i} . We write $\Delta(Y_{-i}, p) \subseteq \Delta(X_{-i})$ for the subset of distributions $q_{-i} \in \Delta(X_{-i})$ such that $\sum_{x_{-i} \in Y_{-i}} q_{-i}(x_{-i}) \geq p$. Let $Br_i(\Delta(Y_{-i}, p))$ denote the set of strategies in X_i that are pure best responses by i to some distribution $q_{-i} \in \Delta(Y_{-i}, p)$ (regardless of probability assigned to other possible combinations of strategies); that is,

$$Br_i(\Delta(Y_{-i}, p)) = \bigcup_{q_{-i} \in \Delta(Y_{-i}, p)} Br_i(q_{-i}). \quad (3)$$

Let us recall the definition of a strict p -dominant equilibrium first introduced by Morris et al. [8] in two-person games and extended to n -person games by Kajii and Morris [9]. A profile $x^* \in X$ is a strict $\mathbf{p} = (p_1, \dots, p_n)$ -dominant equilibrium if for each player $i \in I$

$$\{x_i^*\} = Br_i(\Delta(\{x_{-i}^*\}, p_i)). \quad (4)$$

In the sequel, we focus on the case where $p_i = p$ for all $i \in I$. The concept of p -best response set extends the concept of strict p -dominant equilibrium to product sets of strategies. Formally, a (minimal) p -best response set is defined as follows.

Definition 1. Let $p \in [0, 1]$. A p -best response set is a product set $Y \subseteq X$, where for each player i

$$Br_i(\Delta(Y_{-i}, p)) \subseteq Y_i. \quad (5)$$

A p -best response set Y is a minimal p -best response set if no p -best response set is a proper subset of Y .

Let \mathcal{Q}_p be the collection of p -best response sets for some $p \in [0, 1]$. The following lemma states some properties of minimal p -best response sets.

Lemma 2. Let Γ be a finite n -person game.

- (1) Γ has a minimal p -best response set for any $p \in [0, 1]$.
- (2) Fix $p \in [0, 1]$. Then two distinct minimal p -best response sets of Γ are disjoint.
- (3) For $p \leq 1/2$, there exists a unique minimal p -best response set in Γ .
- (4) Let $p \leq p' \leq 1/2$. Let Y and Y' be the minimal p -best response set and p' -best response set, respectively. Then, $Y \supseteq Y'$.

For a proof of this Lemma, we refer the reader to Durieu et al. [4] and Tercieux [3].

4. Adaptive Processes

Following Samuelson [10], we extend the Darwinian process proposed by Kandori et al. [11] to the multipopulation case. This extension is quite natural to deal with asymmetric n -person games. We think the game Γ as having n roles. For each role $i \in I$, a nonempty class C_i of individuals is eligible to play that role. We assume that each class is composed of M identical agents, where M is a finite integer.

We consider the possibility of local interactions. Let $G(L, E)$ be a graph where $L = \{l_1, l_2, \dots, l_M\}$ is the set of vertices and E is the set of edges. We assume that vertices are located increasingly in a clockwise direction around a ring. We focus on a chordal ring: vertex $l_k, k \in \{1, 2, \dots, M\}$ is adjacent to the vertices (modulo M) $l_{k\pm 1}, l_{k\pm 2}, \dots, l_{k\pm K}$ where $2K \leq M - 1$. If $2K = M - 1$, then each vertex is adjacent to each other vertex. For each vertex $l \in L$, denote by $V(l)$ the open neighborhood of l , that is, the set of vertices adjacent to l . Since $|V(l)| = 2K$ for each $l \in L$, $G(L, E)$ is regular. The closed neighborhood of l , denoted by $\bar{V}(l)$, is defined by $V(l) \cup \{l\}$.

We assume that each class of M agents is distributed among the set of vertices L . In other words, each agent of each class $C_i, i \in I$, is located at exactly one vertex of L and two agents of the same class are located in different vertices of L . In this way, each vertex contains exactly n agents. From this point of view, each vertex can be interpreted as a location. We denote by $x(l) \in X$ the actions profile chosen by agents located in $l \in L$. The collection of action profiles chosen in all locations in L is $\Theta = X^M$.

Let $t = 1, 2, \dots$ denote successive time periods. The stage game Γ is played once in each period. In period t , one agent is drawn at random from each of the n classes and assigned to play the corresponding role. We assume that, at every period, each agent has no information about who is selected to play each role in the game, only that a given action is played by someone. This lack of information gives rise to two kinds of assumptions made by each agent about his potential opponents, that is, about the matching rule. Fix a location $l \in L$. In a first scenario, each agent in $l \in L$ assumes that his potential opponents belong to a unique location in $\bar{V}(l)$. In other words, opponents would be drawn jointly by location in the closed neighborhood of his location. In a second scenario, each agent in $l \in L$ assumes that his

potential opponents may belong to different locations in $\bar{V}(l)$. In other words, opponents would be drawn independently in the closed neighborhood of his location.

We give a formal description of both scenarios. Consider a time period t . Actions chosen in t by the whole population are described by $\theta^t = (x^t(l_1), x^t(l_2), \dots, x^t(l_M)) \in \Theta$. Fix an agent $i \in C_i$ located in l and an action profile $x \in X$. Denote by $\omega_i^t(x_{-i})$ the number of locations in $\bar{V}(l)$ such that $x_{-i}(l) = x_{-i}$ in θ^t . According to the first scenario, at the beginning of period $t + 1$, agent i believes that the probability to be matched with agents playing x_{-i} is

$$q_{-i}^L(x_{-i}) = \frac{\omega_i^t(x_{-i})}{2K + 1}. \quad (6)$$

In order to describe the second scenario, it is convenient for each class C_i to identify X_i with the set $\{1, 2, \dots, r_i\}$, where $r_i = |X_i|$. For each $j \in C_j$ such that $C_j \neq C_i$, and each $x_j \in \{1, \dots, r_j\}$, denote by $\omega_i^t(x_j)$ the number of $\bar{V}(l)$ where x_j is chosen in θ^t . Consider $x_{-i} = (x_j)_{j \neq i} \in X_{-i}$. According to the second scenario, at the beginning of period $t + 1$, agent i believes that the probability to be matched with agents playing x_{-i} is

$$q_{-i}^L(x_{-i}) = \prod_{j \neq i} \frac{\omega_i^t(x_j)}{2K + 1}. \quad (7)$$

In period t , every agent in each class C_i chooses an action $x_i \in X_i$ according to the following learning rule. Every agent might receive the opportunity to revise his choice. For the sake of simplicity, we assume that this adjustment probability does not depend on the agent nor on the actions chosen in the whole population. Whenever an agent does not receive a revision opportunity, he simply repeats the action he has taken in the past. Whenever an agent receives a revision opportunity, he switches to a myopic best response. That is, the agent assumes that the action choices of other agents will remain unchanged in the next period and adopts a pure best response against it. Precisely, agent i chooses a pure best response against the probability distribution on X_{-i} computed using formula (6) or (7). In other words, we define two myopic best-response dynamics with inertia.

If probability distributions are computed using (6), the dynamics results in a Markov chain on the state space Θ , denoted by P_L^0 . We refer to it as *best-response process with joint drawing by location*. Similarly, if probability distributions are computed using (7), the dynamics results in a Markov chain on the state space Θ , denoted by P_I^0 . We refer to it as *best-response process with independent drawing*. Let \mathcal{R}^L (resp. \mathcal{R}^I) be the collection of recurrent sets of P_L^0 (resp., P_I^0). For each $i \in I$ and each subset $\Theta' \subset \Theta$, let $S_i(\Theta')$ be the set of actions in X_i that appears in Θ' . The product set of all actions that appears in Θ' is $S(\Theta') = \prod_{i \in I} S_i(\Theta')$. Consider a minimal p -best response set Y . Denote by $\mathcal{R}_Y^L \subseteq \mathcal{R}^L$ (resp., $\mathcal{R}_Y^I \subseteq \mathcal{R}^I$) the collection of recurrent sets of P_L^0 (resp. P_I^0) such that $\Theta' \in \mathcal{R}_Y^L$ (resp., $\Theta' \in \mathcal{R}_Y^I$) if and only if $S(\Theta') \subseteq Y$. Observe that $\mathcal{R}_Y^L \neq \emptyset$ and $\mathcal{R}_Y^I \neq \emptyset$ since Y is a minimal p -best response set and thus a 1-best response set.

Following the literature (see, e.g., [11, 12]), the model is completed by adding the possibility of rare mutations or experiments on the part of agents. With fixed probability $\epsilon > 0$, independent across players and across time, each agent chooses an action at random. The processes with mutations are called *perturbed processes* and are denoted by P_L^ϵ and P_I^ϵ . With these mutations as part of the processes, each state of Θ is reachable with positive probability from every other state. Hence, the perturbed processes P_L^ϵ and P_I^ϵ are irreducible and aperiodic finite state Markov chains on Θ . Consequently, for each $\epsilon > 0$, P_L^ϵ (resp., P_I^ϵ) has a unique stationary distribution μ_L^ϵ (resp., μ_I^ϵ) satisfying $\mu_L^\epsilon P_L^\epsilon = \mu_L^\epsilon$ (resp., $\mu_I^\epsilon P_I^\epsilon = \mu_I^\epsilon$). The limit stationary distribution (as the rate of mutation tends to zero) $\mu_L^* = \lim_{\epsilon \rightarrow 0} \mu_L^\epsilon$ (resp., $\mu_I^* = \lim_{\epsilon \rightarrow 0} \mu_I^\epsilon$) exists and is a stationary distribution of the unperturbed process P_L^0 (resp., P_I^0). The states in the support of μ_L^* (resp. μ_I^*) are called stochastically stable states and form a subset of \mathcal{R}^L (resp., \mathcal{R}^I). The recurrent sets appearing in the support of μ_L^* (or μ_I^*) are those which are the easiest to reach from all other recurrent sets, with “easiest” interpreted as requiring the fewest mutations (cf., Theorem 4 in [12]).

We rely on the identification of the set of stochastically stable states developed by Ellison [6]. This identification proceeds as follows. Ellison [6] introduces sufficient conditions to have a collection of recurrent sets that contains all stochastically stable states. The analysis uses three measures: the radius, the coradius, and the modified coradius. To illustrate these concepts, consider a collection of recurrent sets $\Theta^1 \subseteq \mathcal{R}^L$. Let $D(\Theta^1)$ be the basin of attraction of Θ^1 , that is, the set of states $\theta \in \Theta$ from which the unperturbed process converges to Θ^1 with probability one. In other words, $D(\Theta^1)$ is the set of states $\theta \in \Theta$ such that it is possible without mutation to build a path, that is, a sequence of distinct states, from θ to $\theta^1 \in \Theta^1$. The radius of (the basin of attraction of) Θ^1 , denoted by $R(\Theta^1)$, is the minimum number of mutations necessary to leave $D(\Theta^1)$ from Θ^1 , that is, the minimum number of mutations contained in any path from a state $\theta^1 \in \Theta^1$ to a state $\theta^2 \notin D(\Theta^1)$. The coradius of (a basin of attraction of) Θ^1 , denoted by $CR(\Theta^1)$, is the maximum over all states $\theta^2 \notin \Theta^1$ of the minimum number of mutations necessary to reach $D(\Theta^1)$, that is, the maximum over all states $\theta^2 \notin \Theta^1$ of the minimum number of mutations contained in any path from θ^2 to a state $\theta^1 \in D(\Theta^1)$. To compute the modified coradius of (the basin of attraction of) Θ^1 , consider a state $\theta^2 \notin \Theta^1$ and a path from θ^2 to a state belonging to Θ^1 . The modified number of mutations of this path is obtained by subtracting from the number of mutations of the path the radius of the intermediate recurrent sets through which the path passes. The modified coradius of (a basin of attraction of) Θ^1 , denoted by $CR^*(\Theta^1)$, is the maximum over all states $\theta^2 \notin \Theta^1$ of the minimum modified number of mutations necessary to reach $D(\Theta^1)$, that is, the maximum over all states $\theta^2 \notin \Theta^1$ of the minimum modified number of mutations associated with any path from θ^2 to a state $\theta^1 \in D(\Theta^1)$. Note that $CR^*(\Theta^1) \leq CR(\Theta^1)$ for every $\Theta^1 \subseteq \Theta$. Ellison [6] establishes the following result.

Theorem 3 (see [6]). *Consider a perturbed process and fix $\Theta^1 \subseteq \Theta$. If $R(\Theta^1) > CR^*(\Theta^1)$, then the stochastically stable states are contained in Θ^1 .*

Since $CR^*(\Theta^1) \leq CR(\Theta^1)$, an alternative condition to have all stochastically stable states contained in Θ^1 is that $R(\Theta^1) > CR(\Theta^1)$.

5. Selection Results

Fix a minimal p -best response set Y . First, consider the perturbed best-response process with joint drawing by location. We give a sufficient condition to have all stochastically stable states associated with Y .

Theorem 4. *Let Γ be a finite n -person game and $G(L, E)$ a chordal ring of degree $2K$. Let $p \leq 1/n$ and let Y be the minimal p -best response set of Γ . If K is sufficiently large and ϵ is sufficiently small, the perturbed process P_L^ϵ puts arbitrarily high probability on a subset $\Theta^* \subseteq \Theta$ such that $S(\Theta^*) \subseteq Y$.*

Proof. Observe that if $\mathcal{R}_Y^L = \mathcal{R}^L$, then the result follows. In the sequel, we assume that $\mathcal{R}_Y^L \subset \mathcal{R}^L$. We break the proof into three parts.

(1) We give a lower bound on $R(\mathcal{R}_Y^L)$. Fix a state $\theta \in \mathcal{R}_Y^L$, a location $l \in L$, and two distinct classes C_i and C_j . Consider agent $i \in C_i$ located in l and agent $j \in C_j$ located in $l_k \in \bar{V}(l)$. Assume that agent j mutates: he chooses at random a strategy not contained in Y_j . Then, agent i believes that the probability to be matched with agents playing a vector of actions not contained in Y_{-i} is $\sum_{x_{-i} \notin Y_{-i}} q_{-i}^L(x_{-i}) = 1/(2K + 1)$. Now, assume that $\alpha \leq 2K + 1$ agents in C_j located in locations belonging to $\bar{V}(l)$ mutate and choose an action outside Y_j . Then, agent i believes that the probability to be matched with agents playing a vector of actions not contained in Y_{-i} is $\sum_{x_{-i} \notin Y_{-i}} q_{-i}^L(x_{-i}) = \alpha/(2K + 1)$. Let \bar{p} be a probability such that at least one class of agents C_i has an action $x_i \notin Y_i$ as a pure best response to q_{-i}^L , where $\sum_{x_{-i} \in Y_{-i}} q_{-i}^L(x_{-i}) = \bar{p}$. By definition of a minimal p -best response set, we have $p > \bar{p}$. A transition from $\Theta^1 \in \mathcal{R}_Y^L$ to any $\Theta^2 \notin \mathcal{R}_Y^L$ requires at least z mutations (in locations belonging to $\bar{V}(l)$ and outside Y_j), where z is such that $z/(2K + 1) \geq 1 - \bar{p}$. Otherwise, agent i 's best-response(s) to q_{-i}^L belong(s) to Y_i since $\sum_{x_{-i} \in Y_{-i}} q_{-i}^L(x_{-i}) \geq p$. Hence, $R(\mathcal{R}_Y^L)$ is such that $R(\mathcal{R}_Y^L) \geq [(1 - \bar{p})(2K + 1)]$.

(2) We give an upper bound on $CR^*(\mathcal{R}_Y^L)$. To do this, it is convenient to distinguish between two situations according to the values of p and K . Firstly, consider the cases such that $\lceil p(2K + 1) \rceil \leq K$. Fix a state $\theta \notin \mathcal{R}_Y^L$ and a class C_i . Assume that, in a location $l_k \in L$, $(n - 1)$ agents $j \notin C_i$ mutate and choose actions in Y_j . Then, agent $i \in C_i$ located in $l \in \bar{V}(l_k)$ believes that the probability to be matched with agents playing a vector of actions contained in Y_{-i} is $\sum_{x_{-i} \in Y_{-i}} q_{-i}^L(x_{-i}) = 1/(2K + 1)$. Now, consider $\alpha \leq 2K + 1$ consecutive locations in $G(L, E)$ denoted by l_k , $k \in \{1, \dots, \alpha\}$. Assume that in each of these locations $(n - 1)$ agents $j \notin C_i$ mutate and choose actions in Y_j . Then, agent $i \in C_i$ located

in l such that $l \in \bar{V}(l_k)$ for each $k \in \{1, \dots, \alpha\}$ believes that the probability to be matched with agents playing a vector of actions contained in Y_{-i} is $\sum_{x_{-i} \in Y_{-i}} q_{-i}^L(x_{-i}) = \alpha/(2K + 1)$. By definition of a minimal p -best response set, agent i has at least one strategy $x_i \in Y_i$ as a pure best response to q_{-i}^L after that all agents $j \notin C_i$ located in $\alpha^* \leq 2K + 1$ consecutive locations in $G(L, E)$ belonging to $\bar{V}(l)$ mutate and choose actions in Y_j if α^* is such that $\alpha^*/(2K + 1) \geq p$. Set $\alpha^* = \lceil p(2K + 1) \rceil$. Since $p \leq 1/n$, if in α^* consecutive locations, l_1, \dots, l_{α^*} , every agent $j \notin C_i$ mutates and chooses an action in Y_j , then every agent $i \in C_i$ located in each of these locations chooses an action in Y_i . By inertia, it is then possible to reach a state in which an action profile contained in Y is played in α^* consecutive locations, l_1, \dots, l_{α^*} . From such a state, it is possible to reach a state in \mathcal{R}_Y^L without additional mutation. Indeed, consider agents located in l_{α^*+1} . Since $\lceil p(2K + 1) \rceil \leq K$, every agent of each class, C_i , believes that the probability to be matched with agents playing a vector of actions contained in Y_{-i} is $\sum_{x_{-i} \in Y_{-i}} q_{-i}^L(x_{-i}) = \alpha^*/(2K + 1) \geq p$. Then, it is possible to reach a state in which an action profile in Y is chosen in $\alpha^* + 1$ consecutive locations. Continuing in this fashion, we can reach a state in \mathcal{R}_Y^L without additional mutation. Hence, $CR(\mathcal{R}_Y^L)$ is such that $CR(\mathcal{R}_Y^L) \leq (n - 1)\lceil p(2K + 1) \rceil$.

Secondly, consider the cases such that $\lceil p(2K + 1) \rceil > K$. Observe that since $p \leq 1/n$, we necessarily have $n = 2$. Fix a state $\theta^1 \notin \mathcal{R}_Y^L$. By the above point, we know that $\alpha^* = \lceil p(2K + 1) \rceil$ mutations allow reaching a state, denoted by θ^2 , where an action profile in Y is chosen in α^* consecutive locations, l_1, \dots, l_{α^*} . If the number of locations belonging to $L \setminus \{l_k, k \in \{1, \dots, \alpha^*\}\}$ is at most equal to K , then every agent of each class C_i , $i \in \{1, 2\}$, located in these locations believes that the probability to be matched with an agent playing an action contained in Y_{-i} is $\sum_{x_{-i} \in Y_{-i}} q_{-i}^L(x_{-i}) \geq p$. A transition from θ^2 to a state in \mathcal{R}_Y^L is possible without additional mutation and $CR(\mathcal{R}_Y^L)$ is bounded as above. It remains to deal with situations where the number of locations in $L \setminus \{l_k, k \in \{1, \dots, \alpha^*\}\}$ is strictly greater than K . In θ^2 , each agent $i \in C_i$, $i \in \{1, 2\}$, located in these locations believes that the probability to be matched with an agent playing an action contained in Y_{-i} is $\sum_{x_{-i} \in Y_{-i}} q_{-i}^L(x_{-i}) < p$. Thus, it is possible that θ^2 does not belong to \mathcal{R}_Y^L nor to the recurrent set containing θ^1 . However, an additional mutation is sufficient to reach a state, say θ^3 , in which an action profile in Y is chosen in $\alpha^* + 1$ consecutive locations. To see this, consider location l_{α^*+1} . If agent $j \in C_2$ located in l_{α^*+1} mutates and chooses an action in Y_2 , then agent $i \in C_1$ located in l_{α^*+1} believes that the probability to be matched with an agent playing an action contained in Y_{-i} is $\sum_{x_{-i} \in Y_{-i}} q_{-i}^L(x_{-i}) \geq p$. Using inertia, it is possible that an action profile in Y is chosen in location l_{α^*+1} . By a similar argument, it is possible to establish that one additional mutation may allow reaching a state θ^4 in which an action profile in Y is chosen in $\alpha^* + 2$ consecutive locations. This argument can be applied until the number of locations in which an action profile not contained in Y is equal to K . From such a state, it is possible to reach a state in \mathcal{R}_Y^L without additional mutation. Hence, the path

from $\theta^1 \notin \mathcal{R}_Y^L$ to a state in \mathcal{R}_Y^L passes through a sequence of intermediate recurrent sets whose radius is equal to 1. Thus, the modified coradius of \mathcal{R}_Y^L is such that $CR^*(\mathcal{R}_Y^L) \leq \lceil p(2K + 1) \rceil$.

Finally, since $CR(\mathcal{R}_Y^L) \geq CR^*(\mathcal{R}_Y^L)$, it holds in all cases that

$$CR^*(\mathcal{R}_Y^L) \leq (n - 1)\lceil p(2K + 1) \rceil. \quad (8)$$

(3) In order to apply Theorem 3, it is sufficient to show that

$$\lceil (1 - \bar{p})(2K + 1) \rceil > (n - 1)\lceil p(2K + 1) \rceil. \quad (9)$$

A sufficient condition to obtain inequality (9) is that

$$1 - \bar{p} > (n - 1)p + \frac{n - 1}{2K + 1}. \quad (10)$$

Inequality (10) is satisfied provided that K is sufficiently large since by hypothesis $p \leq 1/n$, and, by definition of a minimal p -best response set, we have $1 - \bar{p} > 1 - p$. \square

By definition, if a minimal p -best response set is a singleton set, then it is a strict p -dominant equilibrium. The following result is an immediate application of Theorem 4.

Corollary 5. *Let Γ be a finite n -person game and $G(L, E)$ a chordal ring of degree $2K$. Let x^* be a strict p -dominant equilibrium of Γ , where $p \leq 1/n$. If K is sufficiently large and ϵ is sufficiently small, the perturbed process P_Γ^ϵ puts arbitrarily high probability on a subset $\Theta^* \subseteq \Theta$ such that $S(\Theta^*) = \{x^*\}$.*

Observe that if $n = 2$, Corollary 5 is similar to Corollary 2 in Ellison [6] except that it also holds for the class of asymmetric two-player games with a $1/2$ -dominant equilibrium.

Second, we consider the perturbed best-response process with independent drawing. We give a sufficient condition to have all stochastically stable states associated with a minimal p -best response set Y .

Theorem 6. *Let Γ be a finite n -person game and $G(L, E)$ a chordal ring of degree $2K$. Let $p \leq n^{1-n}$ and let Y be the minimal p -best response set of Γ . If K is sufficiently large and ϵ is sufficiently small, the perturbed process P_Γ^ϵ puts arbitrarily high probability on a subset $\Theta^* \subseteq \Theta$ such that $S(\Theta^*) \subseteq Y$.*

Proof. Observe that if $\mathcal{R}_Y^I = \mathcal{R}^I$, then the result follows. In the sequel, we assume that $\mathcal{R}_Y^I \subset \mathcal{R}^I$. As in the proof of Theorem 4, we break the proof into three parts.

(1) We give a lower bound on $R(\mathcal{R}_Y^I)$. By the same reasoning as in the proof of Theorem 4, it follows that $R(\mathcal{R}_Y^I)$ is such that $R(\mathcal{R}_Y^I) \geq \lceil (1 - \bar{p})(2K + 1) \rceil$.

(2) We give an upper bound on $CR^*(\mathcal{R}_Y^I)$. Observe that if $n = 2$, both processes P_Γ^ϵ and P_L^ϵ coincide. By the same reasoning as in the proof of Theorem 4, it follows that if $n = 2$ then $CR^*(\mathcal{R}_Y^I) \leq \lceil p(2K + 1) \rceil$. Now, assume that $n > 2$. Fix a state $\theta \notin \mathcal{R}_Y^I$ and a class C_i . Assume that, in a location $l_k \in L$, $(n - 1)$ agents $j \notin C_i$ mutate and choose actions in Y_j . Then, agent $i \in C_i$ located in $l \in \bar{V}(l_k)$ believes that

the probability to be matched with agents playing a vector of actions contained in Y_{-i} is $\sum_{x_{-i} \in Y_{-i}} q_{-i}^l(x_{-i}) = (1/(2K+1))^{n-1}$. Now, consider $\alpha \leq 2K+1$ consecutive locations in $G(L, E)$ denoted by l_k , $k \in \{1, \dots, \alpha\}$. Assume that in each of these locations $(n-1)$ agents $j \notin C_i$ mutate and choose actions in Y_j . Then, agent $i \in C_i$ located in l such that $l \in \bar{V}(l_k)$ for each $k \in \{1, \dots, \alpha\}$ believes that the probability to be matched with agents playing a vector of actions contained in Y_{-i} is $\sum_{x_{-i} \in Y_{-i}} q_{-i}^l(x_{-i}) = (\alpha/(2K+1))^{n-1}$. By definition of a minimal p -best response set, agent i has at least one strategy $x_i \in Y_i$ as a pure best response to q_{-i}^l after that all agents $j \notin C_i$ located in $\alpha^* \leq 2K+1$ consecutive locations in $G(L, E)$ belonging to $\bar{V}(l)$ mutate and choose actions in Y_j if α^* is such that $(\alpha^*/(2K+1))^{n-1} \geq p$. Set $\alpha^* = \lceil p^{1/(n-1)}(2K+1) \rceil$. Since $p \leq n^{1-n}$, if in α^* consecutive locations every agent $j \notin C_i$ mutates and chooses an action in Y_j , then every agent $i \in C_i$ located in each of these locations chooses an action in Y_i . By inertia, it is then possible to reach a state in which an action profile contained in Y is played in α^* consecutive locations in $G(L, E)$. From such a state, it is possible to reach a state in \mathcal{R}_Y^I without additional mutation. Indeed, since $p \leq n^{1-n}$ and $n > 2$, we necessarily have $\alpha^* \leq K$. Hence, $CR(\mathcal{R}_Y^I)$ is such that $CR(\mathcal{R}_Y^I) \leq (n-1) \lceil p^{1/(n-1)}(2K+1) \rceil$.

(3) In order to conclude, it is sufficient to show that

$$\lceil (1-\bar{p})(2K+1) \rceil > (n-1) \lceil p^{1/(n-1)}(2K+1) \rceil. \quad (11)$$

A sufficient condition to obtain inequality (11) is that

$$1-\bar{p} > (n-1) p^{1/(n-1)} + \frac{n-1}{2K+1}. \quad (12)$$

Inequality (12) is satisfied provided K is sufficiently large since by hypothesis $p \leq n^{1-n}$, and, by definition of a minimal p -best response set, we have $1-\bar{p} > 1-p$. \square

The following result is an immediate application of Theorem 6.

Corollary 7. *Let Γ be a finite n -person game and $G(L, E)$ a chordal ring of degree $2K$. Let x^* be a strict p -dominant equilibrium of Γ , where $p \leq n^{1-n}$. If K is sufficiently large and ϵ is sufficiently small, the perturbed process P_Γ^ϵ puts arbitrarily high probability on a subset $\Theta^* \subseteq \Theta$ such that $S(\Theta^*) = \{x^*\}$.*

6. Conclusion

This paper establishes that the concept of minimal p -best response set is useful to study the long-run outcomes of a process when agents are arranged on a chordal ring and follow a myopic best response rule with inertia. In particular, it allows us to obtain results for the whole class of finite n -person games. Even if predictions are not necessarily sharp (since a minimal p -best response set may become large when p decreases), those results make easier the identification of stochastically stable states. The paper also highlights that predictions depend on the assumptions made by agents about the matching rule. From this point of view, it is possible to

establish a connection between these results and the results obtained in Durieu et al. [4]. This paper considers a fictitious play model with bounded memory and sample as in Young's [12]. Two processes are studied. On the one hand, it is assumed that each agent believes that, in every period, his opponents play independently. On the other hand, each agent believes that, in every period, the play of his opponents is correlated. Durieu et al. show that the concept of p -best response set allows establishing predictions about the long-run outcomes of both processes. Furthermore, as in the present paper, there exists a similar gap between predictions obtained for each process. This conveys the idea that sampling in memory and believing that opponents correlate their actions (play independently resp.) has the same effect as believing that players are drawn jointly (independently resp.) in neighborhood to play the game.

Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.



Acknowledgments

The authors would like to thank two anonymous referees. Financial support from the National Agency for Research (ANR)—research programs “Dynamite Matching and Interactions: Theory and Experiments” (DynaMITE) ANR BLANC and “Mathématiques de la décision pour l'ingénierie physique et sociale” (MODMAD)—is gratefully acknowledge.

References

- [1] H. P. Young, *Individual Strategy and Social Structure: An Evolutionary Theory of Institutions*, Princeton University Press, Princeton, NJ, USA, 1998.
- [2] S. Weidenholzer, “Coordination games and local interactions: a survey of the game theoretic literature,” *Games*, vol. 1, no. 4, pp. 551–585, 2010.
- [3] O. Tercieux, “ p -best response set,” *Journal of Economic Theory*, vol. 131, no. 1, pp. 45–70, 2006.
- [4] J. Durieu, P. Solal, and O. Tercieux, “Adaptive learning and p -best response sets,” *International Journal of Game Theory*, vol. 40, no. 4, pp. 735–747, 2011.
- [5] G. Ellison, “Learning, local interaction, and coordination,” *Econometrica*, vol. 61, no. 5, pp. 1047–1071, 1993.
- [6] G. Ellison, “Basins of attraction, long-run stochastic stability, and the speed of step-by-step evolution,” *Review of Economic Studies*, vol. 67, no. 1, pp. 17–45, 2000.
- [7] C. Alós-Ferrer and S. Weidenholzer, “Partial bandwagon effects and local interactions,” *Games and Economic Behavior*, vol. 61, no. 2, pp. 179–197, 2007.
- [8] S. Morris, R. Rob, and H. S. Shin, “ p -dominance and belief potential,” *Econometrica*, vol. 63, pp. 145–157, 1995.
- [9] A. Kajii and S. Morris, “The robustness of equilibria to incomplete information,” *Econometrica*, vol. 65, no. 6, pp. 1283–1309, 1997.
- [10] L. Samuelson, *Evolutionary Games and Equilibrium Selection*, MIT Press, Harvard, 1997.

- [11] M. Kandori, G. J. Mailath, and R. Rob, "Learning, mutations, and long run equilibria in games," *Econometrica*, vol. 61, no. 1, pp. 29–56, 1993.
- [12] H. P. Young, "The evolution of conventions," *Econometrica*, vol. 61, no. 1, pp. 57–84, 1993.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

