

Progress with palbociclib in breast cancer: latest evidence and clinical considerations

Andrea Rocca, Alessio Schirone, Roberta Maltoni, Sara Bravaccini, Lorenzo Ceconetto, Alberto Farolfi, Giuseppe Bronte and Daniele Andreis

Ther Adv Med Oncol

2017, Vol. 9(2) 83–105

DOI: 10.1177/
1758834016677961

© The Author(s), 2016.

Reprints and permissions:
[http://www.sagepub.co.uk/
journalsPermissions.nav](http://www.sagepub.co.uk/journalsPermissions.nav)

Abstract: Deregulation of the cell cycle is a hallmark of cancer, and research on cell cycle control has allowed identification of potential targets for anticancer treatment. Palbociclib is a selective inhibitor of the cyclin-dependent kinases 4 and 6 (CDK4/6), which are involved, with their coregulatory partners cyclin D, in the G1-S transition. Inhibition of this step halts cell cycle progression in cells in which the involved pathway, including the retinoblastoma protein (Rb) and the E2F family of transcription factors, is functioning, although having been deregulated. Among breast cancers, those with functioning cyclin D-CDK4/6-Rb-E2F are mainly hormone-receptor (HR) positive, with some HER2-positive and rare triple-negative cases. Deregulation results from genetic or otherwise occurring hyperactivation of molecules subtending cell cycle progression, or inactivation of cell cycle inhibitors. Based on results of randomized clinical trials, palbociclib was granted accelerated approval by the US Food and Drug Administration (FDA) for use in combination with letrozole as initial endocrine-based therapy for metastatic disease in postmenopausal women with HR-positive, HER2-negative breast cancer, and was approved for use in combination with fulvestrant in women with HR-positive, HER2-negative advanced breast cancer with disease progression following endocrine therapy. This review provides an update of the available knowledge on the cell cycle and its regulation, on the alterations in cyclin D-CDK4/6-Rb-E2F axis in breast cancer and their roles in endocrine resistance, on the preclinical activity of CDK4/6 inhibitors in breast cancer, both as monotherapy and as partners of combinatorial synergic treatments, and on the clinical development of palbociclib in breast cancer.

Keywords: breast neoplasms, CDK4/6 inhibitors, cell cycle, cyclin-dependent kinase 4/6, palbociclib, retinoblastoma protein

Introduction

Abnormal cell proliferation, often resulting from oncogene activation or from inactivation or deletion of tumor-suppressor genes, is the key hallmark of cancer [Hanahan and Weinberg, 2011], and the main players of cell cycle control are logical potential targets for cancer therapy.

Breast cancer is the most common malignancy in women worldwide, and the most common cause of cancer death in women in less developed regions, and the second in women in more developed regions [International Agency for Research on Cancer, 2016]. Although breast cancer death rates are decreasing in developed countries, metastatic breast cancer remains largely an incurable disease, and new treatments are needed.

Palbociclib is the first of a new class of agents targeting the cell cycle machinery.

We will review recent knowledge on the cell cycle and its regulation, on the alterations in key cell cycle molecules in breast cancer and their roles in endocrine resistance, on the preclinical activity of cyclin-dependent kinases 4 and 6 (CDK4/6) inhibitors and on clinical development of palbociclib in breast cancer. The relevant literature was retrieved through PubMed (main keywords used: CDK 4/6 inhibitor, palbociclib, PD 0332991, ribociclib, abemaciclib; time period: up to May 2016) and the proceedings of the main recent cancer congresses (ASCO and AACR annual meetings 2014–2016, San Antonio Breast Cancer Symposium 2014–2015).

Correspondence to:
Andrea Rocca, MD
Istituto Scientifico
Romagnolo per lo Studio e
la Cura dei Tumori (IRST)
IRCCS, Via Maroncelli 40,
Meldola, FC 47014, Italy
andrea.rocca@irst.emr.it
Alessio Schirone, MD
Roberta Maltoni, MD
Sara Bravaccini, PhD
Lorenzo Ceconetto, MD
Alberto Farolfi, MD
Giuseppe Bronte, MD
Daniele Andreis, MSc
Istituto Scientifico
Romagnolo per lo Studio e
la Cura dei Tumori (IRST)
IRCCS, Meldola, FC, Italy




Figure 1. The cyclin D–cyclin-dependent kinase 4/6–retinoblastoma protein–E2F transcription-factor axis. AKT, protein kinase B (also known as PKB); Cdc25, cell-division cycle 25 phosphatase family; CDK, cyclin-dependent kinase; CIP/KIP, CDK-interacting protein/kinase inhibitory protein family (including p21^{Cip1}, p27^{Kip1}, and p57^{Kip2}); E2F, E2F transcription factor; ERK, extracellular-signal-regulated kinases; E, estrogen; ER, estrogen receptor; INK4, inhibitor of CDK4 family (including p16^{INK4A}, p15^{INK4B}, p18^{INK4C}, and p19^{INK4D}); MEK, mitogen-activated protein kinase kinase; mTORC1, mammalian target of rapamycin complex 1; P, phosphate; p70S6K, ribosomal protein S6 kinases; PDK1, 3-phosphoinositide-dependent protein kinase 1; PI3K, phosphoinositide 3-kinase (p85 regulatory subunit and p110 catalytic subunit); PIP2, phosphatidylinositol 4,5-bisphosphate; PIP3, phosphatidylinositol 3,4,5 trisphosphate; PTEN, phosphatase and tensin homolog; Raf, family of serine-threonine kinases; Ras, GTP-binding proteins encoded by ras genes; Rb, retinoblastoma protein; TSC, tuberous sclerosis complex protein.

Cyclin-dependent kinases and the cell cycle

Major players of the cell cycle are the serin/threonin kinases CDKs [Malumbres, 2014], their activating regulatory subunits cyclins [Ma *et al.* 2013], the so-called pocket proteins retinoblastoma (Rb, product of the Rb tumor-suppressor gene *RB1*) and Rb-like proteins p107 (RBL1) and p130 (RBL2) [Dick and Rubin, 2013], and the E2F family of transcription factors [Chen *et al.* 2009] (Figure 1). The different cyclins, each synthesized cyclically in specific phases of the cell cycle, bind to, and activate, specific CDKs, and the cyclin-CDK complexes govern the progression of the cell throughout the cycle, a process whose main limiting step is inhibiting Rb and thus releasing its inhibitory action on E2F, allowing transcription of genes that execute the cell cycle.

Two main classes of CDK inhibitors (cyclin kinase inhibitors, CKI) exist: the inhibitor of CDK4 (INK4) family, including p16^{INK4A}, p15^{INK4B}, p18^{INK4C}, and p19^{INK4D}, specifically blocking the formation of cyclin D-CDK4/6 complexes [Sherr and Roberts, 1999], and the CDK-interacting protein/kinase inhibitory protein (CIP/KIP) family, including p21^{Cip1}, p27^{Kip1}, and p57^{Kip2}, acting on all cyclin-CDK complexes, with prevalent inhibitory, but sometimes activatory effects, in particular, activating cyclin D-CDK4/6 complexes [LaBaer *et al.* 1997]. CDK activators, such as the cell-division cycle 25 (Cdc25) phosphatase family, also exist [Shen and Huang, 2012].

In the established model of cell cycle regulation in mammals [Malumbres and Barbacid, 2009], mitogenic stimuli induce the expression of D-type

cyclins (D1, D2, and D3), that bind to the two closely related CDKs 4 and 6, collectively called CDK4/6 for their similar function, activating their kinase activity. Cyclin D–CDK4/6 complexes phosphorylate Rb (and the other pocket proteins) on specific serine and threonine residues. Rb is active when unphosphorylated, its main function being binding to and blocking the activity of transcription factors, mainly of the E2F family, responsible for the transcription of genes involved in cell cycle progression [Weinberg, 1995]. Phosphorylation of Rb leads to its inactivation, with consequent release of E2F transcription factors that start their activity, promoting progression of the cell through the division cycle. Progressive phosphorylation of Rb occurs during the G1 (gap 1) phase of the cycle, culminating at the ‘restriction point’ (R point), when an ‘all or none’ decision is made on cell fate, depending on the balance between growth-promoting and growth-inhibitory stimuli, leading to G1-S transition and progression through the subsequent phases of the cycle, or to the quiescent state, G0.

Progression through the R point is pushed by a positive feedback, elicited by E2F transcription of cyclin E and the CDK activators Cdc25 phosphatases, leading to formation of cyclin E–CDK2 complexes, which further phosphorylate the pocket proteins [Lundberg and Weinberg, 1998; Harbour *et al.* 1999]. Another forward mechanism is due to the CIP/KIP proteins, which, during G1, bind to and activate the increasing cyclin D–CDK4/6 complexes, being therefore subtracted from cyclin E–CDK2 complexes (on which they are inhibitory), favoring their activation [LaBaer *et al.* 1997]. Moreover, cyclin E–CDK2 complexes phosphorylate p27^{Kip1}, leading to its degradation [Slingerland and Pagano, 2000]. The conjunct action of cyclin D–CDK4/6 and cyclin E–CDK2 complexes leads to progression of the cell cycle beyond the R point, from the G1 to the S phase.

CKI, which are silenced during the cell cycle, are expressed in response to inhibitory signals such as transforming growth factor β [Reynisdóttir *et al.* 1995], contact inhibition [Polyak *et al.* 1994], or senescence [Sharpless and Sherr, 2015].

Emerging concepts in the cell cycle and key molecules’ roles

Recent evidences have led to a widened picture of cell cycle control and of its key molecules’ roles.

Although being major players in cell cycle, CDK4/6 and CDK2 are generally dispensable for cycling of most mammalian cells, being required only in specific cell types [Malumbres *et al.* 2004; Malumbres and Barbacid, 2009]. D-type cyclins can bind different CDKs to initiate the cell cycle [Xiong *et al.* 1992]. Only CDK1, which binds cyclin A during the G2 phase and cyclin B for entry into mitosis, is indispensable and sufficient for driving the cell cycle in all cell types during embryogenesis [Santamaria *et al.* 2007]. Nonetheless, some tumor cells may acquire dependency on other CDKs for proliferation [Malumbres and Barbacid, 2009]. In mouse models, cyclin D1 [Yu *et al.* 2001] and CDK4 [Reddy *et al.* 2005; Landis *et al.* 2006; Yu *et al.* 2006] are necessary for the development of breast cancers induced by the neu and ras oncogenes, but not for those driven by c-Myc or Wnt-1, highlighting the dependency of certain tumors from specific cyclin–CDKs, which can be exploited therapeutically [Choi *et al.* 2012]. The dependency on cyclin D1 can nonetheless be overcome by cyclin D3 overexpression, in case of cyclin D1 loss [Zhang *et al.* 2011].

The levels of CDK2 at the exit from mitosis, which are regulated by mitogens *via* p21 at the end of the cell cycle, are important in deciding cell fate, leading to further proliferation when CDK2 is elevated, or to a quiescent state when suppressed [Spencer *et al.* 2013].

Also the G0 to G1 transition requires Rb phosphorylation, resulting from cyclin C binding to and activating CDK3 [Ren and Rollins, 2004].

Cyclins and CDKs have several functions beyond their canonical role in cell cycle [Hydbring *et al.* 2016; Musgrove *et al.* 2011]. Cyclin D–CDK complexes phosphorylate several transcription factors, and regulate their function independently of their action on the pocket proteins. CDK4 and CDK2 phosphorylate and inhibit small mother against decapentaplegic 3 (SMAD3), a key mediator of the anti-proliferative activity of transforming growth factor- β , favoring cell cycle progression from G1 to S phase [Matsuura *et al.* 2004]. Cyclin D–CDK4/6 complexes phosphorylate and activate the forkhead box protein M1 that sustains the expression of cell cycle genes and protects cancer cells from senescence [Anders *et al.* 2011]. Cyclin D1/CDK4 complexes phosphorylate methylosome protein 50 leading to protein arginine methyltransferase 5-dependent histone

methylation and transcriptional repression of *CUL4*, which results in overexpression of the replication-licensing protein CDT-1, required for initiation of DNA replication [Aggarwal *et al.* 2010]. Cyclin D–CDK6 complexes, through activation of JUN and signal transducer and activator of transcription 3, induce the expression of p16^{INK4A}, exerting a negative feedback control of their own activity [Kollmann *et al.* 2013]. Indeed, the transforming effect of CDK6 overexpression may occur only upon deletion or silencing of *CDKN2A*, the gene encoding p16^{INK4A} that frequently occurs in breast cancer [Cancer Genome Atlas Network, 2012]. CDK6 cooperates with the NF-κB subunit p65 to induce the expression of inflammatory genes [Handschiek *et al.* 2014]. Cyclin D–CDK4/6 complexes are also involved in glucose metabolism [Lee *et al.* 2014] and in the processes of apoptosis and cell differentiation [Hydbring *et al.* 2016]. The relevance of all these effects in different tumor types and their implication for the activity of CDK4/6 inhibitors must be better clarified.

Cyclins and CDKs can also regulate transcription factors and exert other functions independently from cyclin–CDK complexes' kinase activity [Hydbring *et al.* 2016], and therefore not subject to inhibition by current CDK4/6 inhibitors. Among these are: a direct role of cyclin D1 in gene transcription [Bienvenu *et al.* 2010]; its involvement in DNA damage repair [Jirawatnotai *et al.* 2011]; its repression of specificity protein 1-mediated transcription [Shao and Robbins, 1995]; inhibition of cyclin D-interacting myb-like protein (DMP1, affecting, *via* p19^{ARF} and MDM2, the expression of p53) [Hirai and Sherr, 1996]; induction of vascular endothelial growth factor (VEGF) expression [Yasui *et al.* 2006]; its activating action on the estrogen receptor (ER) [Zwijsen *et al.* 1997], and inhibitory action on the androgen receptor [Reutens *et al.* 2001]. CDK4/6 also regulates the expression of VEGF [Abedin *et al.* 2010; Kollmann *et al.* 2013] and is involved in DNA repair [Dean *et al.* 2012a].

Beyond Rb, the Rb-like proteins p130 and p107 have a role in repressing cell cycle gene expression during quiescence and in coordinating gene expression during S phase and G2/M phase [Sadasivam and DeCaprio, 2013]. However, mutations are much more frequent in Rb gene than in genes of the other family proteins, and Rb has a nonredundant role in repressing E2F target

genes involved in DNA replication during cell senescence [Chicas *et al.* 2010].

Accumulating evidence also shows roles for the E2F transcription factors beyond that in the cell cycle [Johnson *et al.* 2016], including roles in promoting invasiveness [Yoon *et al.* 2006] and metastatization [Andrechek, 2015].

Cyclin–cyclin-dependent kinase–retinoblastoma protein–E2F pathway alterations in breast cancer

Genomic aberrations or altered expression of molecules of the cyclin–CDK–Rb pathway are frequent in breast cancer. According to The Cancer Genome Atlas [TCGA; Cancer Genome Atlas Network, 2012], the gene encoding for cyclin D1 (*CCND1*) is amplified in 29% and 58% of luminal A and B tumors, respectively, and in 38% of HER2-enriched tumors, whereas that encoding cyclin E (*CCNE1*) is amplified in 9% of triple-negative tumors. Cyclin D1 overexpression, detected by immunohistochemistry in up to half of breast cancers, is even more frequent than *CCND1* amplification, implying that other mechanisms can lead to deregulated expression [Bartkova *et al.* 1994; Gillett *et al.* 1994].

Amplification of the CDK4 gene is reported in about 15% of breast cancers, results in protein overexpression, and is associated with high Ki-67 labeling index [An *et al.* 1999]. It occurs in 14% of luminal A, 25% of luminal B and 24% of HER2-enriched cancers [Cancer Genome Atlas Network, 2012].

Loss of Rb function in stem or progenitor cells is often a key event in neoplastic transformation [Sage, 2012] and is accompanied by epithelial–mesenchymal transition [Arima *et al.* 2012]. According to TCGA data, loss of Rb due to *RB1* gene deletion occurs overall in 2–4% of breast cancers, and loss due to *RB1* truncating mutations in 1–3% [Johnson *et al.* 2016], depending on breast cancer subtype, with up to 20% mutation/loss rate in basal-like tumors [Cancer Genome Atlas Network, 2012]. This leads to constitutive activation of E2F and induction of cyclin E and CDK2, independently from cyclin D–CDK4/6 activation, and is frequently accompanied by upregulation of p16 [Subhawong *et al.* 2009] due to a feedback loop [Kotake *et al.* 2007]. Other authors report higher frequencies

of Rb loss of heterozygosity, correlating with low mRNA expression, particularly in triple-negative but also in luminal B breast cancers [Herschkowitz *et al.* 2008]. Rb functional inactivation may result also from cyclin D1 overexpression or p16^{INK4A} inactivation, and is frequent in breast cancer, as shown by studies of Rb-loss gene signatures identifying tumors with deregulated Rb [Ertel *et al.* 2010; Herschkowitz *et al.* 2008]. Immunohistochemical assessment confirms more frequent Rb loss in triple-negative than in other breast cancer subtypes, and highlights its relation with benefit from adjuvant chemotherapy [Treré *et al.* 2009].

CDKN2A is deleted in 3–8% of breast cancers, more frequently in the triple-negative subtype [Johnson *et al.* 2016; Cairns *et al.* 1995], and may be mutated or silenced by promoter methylation [Ruas and Peters, 1998].

E2F transcription factors are not frequently mutated, but may be involved large-scale chromosomal aberrations such as loss of 16q [Johnson *et al.* 2016].

p21^{Cip1} expression is frequently reduced as a consequence of *TP53* mutation [Musgrove *et al.* 1995] or *Myc* overexpression [Mukherjee and Conrad, 2005], and p27^{Kip1} expression is reduced as a result of *HER2* amplification [Chu *et al.* 2008].

In summary [Cancer Genome Atlas Network, 2012; Witkiewicz and Knudsen, 2014], in luminal breast cancers, ER signaling induces cyclin D1 transcription, and there is often, particularly in luminal B tumors, cyclin D1 overexpression or gene amplification and CDK4 gain; p16^{INK4A} or Rb losses are quite rare in luminal A, and somewhat more frequent in luminal B tumors. Triple-negative cancers have the highest rate of Rb inactivation and of p16^{INK4A} overexpression, and may present cyclin E1 amplification. In *HER2*-positive tumors, cyclin D1 is often amplified or activated as a result of *HER2* mitogenic signaling and there may be CDK4 gain. Aberrations of p16^{INK4A} and those of Rb are mutually exclusive, as tumors with loss of p16^{INK4A} have wild-type Rb, while tumors with mutant Rb show high expression of p16^{INK4A} [Witkiewicz and Knudsen, 2014]. Within this framework, luminal tumors and, to some extent, *HER2*-positive tumors that often have an intact or overactive Rb pathway are

those more likely to benefit from CDK4/6 inhibition.

Cell cycle and endocrine resistance

Estrogens stimulate cellular proliferation mainly by binding to nuclear ER alpha (ER α) that induces the expression of cyclin D1 and ultimately leads to Rb inactivation [Altucci *et al.* 1996]. They also inhibit the expression of CIP/KIP proteins and induce the expression of phosphatase *Cdc25A* [Foster *et al.* 2001], which further contribute to sustain cell proliferation. With a positive feedback, cyclin D1 can directly activate ER α [Zwijnsen *et al.* 1997]. Although other mechanisms are implicated [Nardone *et al.* 2015], the cyclin D1-CDK4/6-Rb-E2F axis is critical for estrogen action.

Both selective ER modulators (SERMs) such as tamoxifen [Ichikawa *et al.* 2008] and selective ER downregulators (SERD) like fulvestrant [Carroll *et al.* 2000] lead to cell cycle arrest (CCA) in G1, reducing the expression of cyclins D and B and increasing that of p21^{Cip1}, with fulvestrant also inducing accumulation of p130-E2F4 complexes characteristic of quiescence.

Aberrant expression or function of molecules involved in cell cycle regulation and in estrogen action have been implicated in endocrine resistance. Some examples are: the suppression or inhibition of CIP/KIP proteins [Chu *et al.* 2008; Pérez-Tenorio *et al.* 2006], which sometimes result from *Myc* [Mukherjee and Conrad, 2005] or cyclin D1 overexpression [Hui *et al.* 2002; Stendahl *et al.* 2004]; Rb inactivation [Bosco *et al.* 2007; Thangavel *et al.* 2011]; upregulation of CDK6, subtending resistance to fulvestrant [Alves *et al.* 2016]; hyperactivation of receptor tyrosine kinases, such as the epidermal growth factor receptor (EGFR) [McClelland *et al.* 2001], the human epidermal growth factor receptor 2 (*HER2*) [Knowlden *et al.* 2003], or the insulin-like growth factor receptor [Miller *et al.* 2009], that induce phosphorylation and activation of ER α or its coregulators; alterations of components of their downstream signaling pathways, such as MAPK/ERK and phosphoinositide 3-kinase (PI3K)/Akt [Miller *et al.* 2010]; ligand-independent, ER α -dependent activation of CDK4-Rb-E2F axis in estrogen-deprived breast cancer [Miller *et al.* 2011]; ER α gene (*ESR1*) gain-of-function mutations leading to constitutive receptor activation [Jeselsohn *et al.* 2015] that ultimately acts *via* CDK4/6-Rb [Luo *et al.* 2016].

Cyclin-dependent kinase 4/6 inhibitors: preclinical activity as single agents and mechanisms of response and resistance

First generation pan-CDK inhibitors showed modest clinical activity and considerable toxicity. More recently, selective small molecule CDK inhibitors have been developed, and CDK4/6 inhibitors are those in more advanced stage of development, with three compounds that have reached the clinical stage: palbociclib, ribociclib, and abemaciclib. Palbociclib is the one in more advanced development.

Palbociclib (PD 0332991) is an orally available pyridopyrimidine derivative, inhibiting CDK4/6 by binding to their ATP pockets with high selectivity, showing IC₅₀ values for CDK4/cyclin D1, CDK4/cyclin D3, and CDK6/cyclin D2 of 11, 9, and 15 nmol/l, respectively, with low or absent activity against other kinases [Fry *et al.* 2004]. It potently inhibits cell proliferation, preventing progression of the cell cycle from G1 into the S phase, in Rb-positive cells of different tumor types, producing Rb dephosphorylation at specific serine residues as its pharmacodynamic hallmark.

Among human breast cancer cell lines representative of the different breast cancer subtypes, the ER-positive, luminal ones were the most sensitive, along with some HER2-amplified cell lines with luminal features, whereas cell lines with basal features were the most resistant [Finn *et al.* 2009]. Both endocrine-sensitive and endocrine-resistant lines may respond to CDK4/6 inhibitors [Petrossian *et al.* 2016]. High levels of cyclin D1 and of Rb, and low levels of p16 were predictors of sensitivity to palbociclib [Finn *et al.* 2009]. This was confirmed also in *ex vivo* studies on primary human tumor cultures [Dean *et al.* 2012b]. CDK4 mutations [Young *et al.* 2014], *CDKN2A* and *CDKN2C* (encoding p18^{INK4C}) deletions [Wiedemeyer *et al.* 2010] and low E2F expression [Logan *et al.* 2013] have been shown to predict palbociclib response in other tumor types, whereas high expression of cyclin E1 [Konecny *et al.* 2011] was associated with resistance.

Breast cancer cell lines with an inactive cyclin D–CDK4/6–Rb pathway, usually because of loss of Rb and consequent upregulation of p16^{INK4A} and downregulation of cyclin D1, are resistant to palbociclib [Dean *et al.* 2010]. This typically occurs in basal cell lines. Rb-proficient cells may become temporarily resistant to palbociclib after

prolonged exposure that leads to increased p107 and CDK2 expression with or without loss of p21^{Cip1} and p27^{Kip1}, but usually remain sensitive to deferred second rounds of treatment. Loss of Rb function would instead lead to true, long-term resistance, due to the increased transcription of cyclin A and E, which activates CDK2 and can drive the cell cycle independently of CDK4/6. Rb function is necessary for the induction of senescence by palbociclib, wherein tumor cells permanently exit the cell cycle [Dean *et al.* 2010].

Rb-knockdown experiments in breast cancer cell lines confirm that Rb status plays a prominent role in acute response to palbociclib, but shows partial activity of palbociclib also in some Rb-knocked-down cell lines, due to a compensatory role of p107, which is dephosphorylated in response to palbociclib, leading to E2F inhibition [Dean *et al.* 2010]. Some activity was seen also in other Rb-deficient cells owing to this mechanism [Rivadeneira *et al.* 2010].

Rarely, Rb-positive cell lines are resistant to palbociclib, and Rb phosphorylation does not decrease after drug exposure [Finn *et al.* 2009]. CKI downregulation and CDK2 reactivation, replacing CDK4/6, has been demonstrated in such cases in models of acute myelogenous leukemia [Wang *et al.* 2007]. Recently, ER-positive breast cancer cell lines have been shown to develop early adaptation to CDK4/6 inhibitors through cyclin D1–CDK2 complexes mediating G1–S transition; this ultimately results in acquired resistance to CDK4/6 inhibitors through selection of cells with Rb loss or *CCNE1* amplification [Herrera-Abreu *et al.* 2016].

Recent studies on breast cancer cell lines have highlighted different features of clones with acquired palbociclib resistance: genomic deletion of *RB1* in some cases, and Rb retention in others, with upregulation of E2F, TGFβ, Wnt, or NF-κB pathways [Lee *et al.* 2016]; increased activity of the PI3K/AKT/mTOR and changes in p53, apoptotic regulation and Rho/Rac pathway [Lenihan *et al.* 2016]; increased levels of 3-phosphoinositide-dependent protein kinase 1 (PDK1) [Jansen *et al.* 2016].

Activity of palbociclib has also been shown in some triple-negative breast cancer cell lines, belonging to the luminal androgen receptor (LAR) and the mesenchymal stem-like (MSL) subtypes [Asghar *et al.* 2015]. Sensitivity was

associated with expression of androgen receptor and with low expression of cyclin E1.

In mouse xenograft models, palbociclib showed significant antitumor activity in breast cancer and in other tumor xenografts, inducing tumor regression *in vivo*, including some durable complete remissions, despite its cytostatic effect *in vitro* [Fry *et al.* 2004]. Palbociclib has also been shown to inhibit breast cancer cell migration and invasion, epithelial–mesenchymal transition, and metastatization, *via* inhibition of the c-Jun/COX-2 pathway [Qin *et al.* 2015].

Cyclin-dependent kinase 4/6 inhibitors: preclinical activity in combination

The combination of palbociclib and tamoxifen showed synergism in ER-positive breast cancer cell lines, likely due to the concomitant cyclin D1 inhibition by tamoxifen and CDK4/6 inhibition by palbociclib. Furthermore, palbociclib monotherapy was active in tamoxifen-resistant MCF7 cell lines, and partially restored sensitivity to tamoxifen in resistant lines [Finn *et al.* 2009].

Palbociclib has also been shown to resensitize fulvestrant-resistant cells to fulvestrant [Alves *et al.* 2016].

Estrogen deprivation, occurring during therapy with aromatase inhibitors (AIs), leads to acquired resistance through both ligand-independent, ER α -dependent activation of the CDK4-Rb-E2F axis [Miller *et al.* 2011] and PI3K hyperactivation [Miller *et al.* 2010], whereas resistance to PI3K inhibitors also involves activation of the cyclin D-CDK4-Rb-E2F axis [Vora *et al.* 2014]. CDK4/6 inhibitors have shown activity in AI-resistant and long-term estrogen-deprived (LTED) cell lines [Petrossian *et al.* 2016]. In mouse xenograft models of LTED breast cancer, the combination of fulvestrant with the pan-PI3K inhibitor BKM120 (buparlisib) was synergic [Miller *et al.* 2011], while in PI3K inhibitors' resistant, PIK3CA-mutant breast cancer xenografts, synergism was demonstrated between PI3K inhibitors (the pan-class I inhibitor pictilisib or the PI3K- α specific inhibitor apelisib) and the CDK4/6 inhibitor ribociclib [Vora *et al.* 2014].

Combined treatment with CDK4/6 inhibitors plus an endocrine agent or a PI3K inhibitor prevented cell adaptation to CDK4/6 inhibitors, and a triple combination of an endocrine agent plus a

CDK4/6 and a PI3K inhibitor was more effective than paired combinations both *in vitro*, triggering apoptosis, and in patient-derived tumor xenografts [Herrera-Abreu *et al.* 2016]. In cases of acquired resistance due to *CCNE1* amplification, treatment with a CDK2-inhibitor was able to resensitize tumor cells to CDK4/6 inhibitors. PI3K inhibition was synergistic with CDK4/6 inhibition, also in PIK3CA-mutated LAR and MSL triple-negative cell lines [Asghar *et al.* 2015].

Palbociclib, in combination with a new class of SERM/SERD hybrids, has shown activity in endocrine-resistant breast cancer cell lines and in ER gene (*ESR1*)-mutant cell lines and patient-derived tumor xenografts [Wardell *et al.* 2015; Nguyen *et al.* 2016].

Also ribociclib has shown single-agent activity in xenograft models of ER-positive breast cancer that was increased by the addition of letrozole or fulvestrant and by the PI3K inhibitors buparlisib and apelisib [O'Brien *et al.* 2014].

Palbociclib also showed synergism with trastuzumab in HER2-amplified cell lines [Finn *et al.* 2009]. Tumors progressing after anti-HER2 therapy often show cyclin D1 overexpression, and abemaciclib has shown activity against anti-HER2-resistant tumors in mouse models, synergizing with anti-HER2 therapies and restoring sensitivity to anti-HER2 drugs [Goel *et al.* 2016]. Cyclin D1-CDK4 inactivates the tuberous sclerosis complex 2 (TSC2) by phosphorylation, relieving its inhibitory activity on mTORC1; mTORC1 can then exert its feedback inhibition of upstream HER receptors [Chandarlapaty *et al.* 2011]. Inhibition of CDK4/6 then restores HER receptors' activity, resensitizing tumors to HER2 blockade [Goel *et al.* 2016].

The combination of palbociclib with the mTOR inhibitor everolimus yielded significant activity in preclinical models of non-small cell lung cancer [Gopalan *et al.* 2013]. Preclinical evidence of effectiveness of the combination of CDK4/6 inhibitors with inhibitors of the MAPK pathway also exists, particularly in melanoma [Yadav *et al.* 2014] and in colorectal cancer [Ziemke *et al.* 2016]. Additionally, the PDK1 inhibitor GSK2334470 was synergistic with CDK4/6 inhibitors in breast cancer cell lines and in xenograft models and was able to restore sensitivity to CDK4/6 inhibitors [Jansen *et al.* 2016].

The association of palbociclib with chemotherapeutic agents has often yielded antagonism, particularly in Rb-proficient cell lines [Roberts *et al.* 2012; McClendon *et al.* 2012]. Palbociclib did not resensitize paclitaxel-resistant cells to paclitaxel-induced apoptosis [Trapé *et al.* 2016]. Nonetheless, a short exposure to palbociclib to synchronize cells prior to paclitaxel resulted in increased cytotoxicity [Dean *et al.* 2012a]. The combination of concurrent palbociclib and radiation, or radiation followed by palbociclib (but not the reverse sequence), outperformed each single treatment modality in glioblastoma xenograft models, inhibiting DNA double-strand-break repair and promoting apoptosis [Hashizume *et al.* 2016].

Palbociclib clinical development

Phase I dose-finding and pharmacokinetics studies

The main phase I studies of palbociclib, enrolling patients with Rb-positive solid tumors or non-Hodgkin's lymphoma, explored two different schedules: daily treatment for 2 weeks, followed by 1 week off treatment (schedule 2/1) [Schwartz *et al.* 2011] and daily treatment for 3 weeks, followed by 1 week off (schedule 3/1) [Flaherty *et al.* 2012]. The recommended phase II doses were 200 mg daily and 125 mg daily, respectively. The dose-limiting toxicities were hematological (neutropenia and thrombocytopenia) with both schedules, and the most common nonhematological adverse events were mild or moderate fatigue, nausea, diarrhea and constipation. No clinically significant effects on QTc interval were reported. Some clues of activity were evident, also in patients with breast cancer.

Palbociclib showed a mean absolute bioavailability after a dose of 125 mg of 46%, slightly increased by food; the pharmacokinetic was linear within the clinically relevant dose range, with a mean terminal half-life of about 26 hours, and steady state was reached within 8 days. Metabolism is mainly hepatic through cytochrome P450 3A (CYP3A) and sulfotransferase 2A1 (SULT2A1), and palbociclib is a weak CYP3A inhibitor. According to a population pharmacokinetic analysis, no dose adjustment is required for patients with mild or moderate renal impairment or with mild hepatic impairment [Sun and Wang, 2014]. A physiologically based pharmacokinetic model [Yu *et al.* 2016] predicts negligible effect of

weak CYP3A inhibitors, while moderate CYP3A inhibitors may increase plasma area under the curve (AUC) by ~40% and moderate CYP3A inducers may reduce plasma AUC by ~40%.

The schedule 3/1 at a dose of 125 mg daily was selected for clinical development.

Palbociclib is a substrate of both P-glycoprotein and breast cancer resistance protein, which limits its brain distribution in preclinical models [de Gooijer *et al.* 2015]. Abemaciclib, on the contrary, has been shown to cross the blood-brain barrier [Patnaik *et al.* 2016].

Phase Ib/II studies

A single arm, phase II study of palbociclib monotherapy (schedule 3/1) enrolled 37 patients with Rb-positive (assessed by immunohistochemistry) advanced breast cancer [DeMichele *et al.* 2015]: 33 with hormone receptor (HR)-positive (HER2-positive in two cases) and four with triple-negative tumors. A clinical benefit (objective response or stable disease \geq 6 months) was achieved in 7 of 33 HR-positive patients (21%) and in none of the triple negatives (whose enrollment was stopped early), and progression-free survival (PFS) was 4.5 months and 1.5 months in the two subgroups, respectively. The two HER2-positive, HR-positive patients yielded a partial response and a stable disease lasting 5 months. The drug confirmed a good tolerability, and none of the biomarkers assessed (immunohistochemical staining for Rb, Ki67 and p16, and *CCND1* amplification by FISH) correlated with clinical benefit.

Given the key roles of the estrogen and CDK4/6 pathways as drivers of breast cancer, and the pre-clinical evidence of activity in luminal tumors and of synergism with endocrine agents, clinical trials initially explored the combination of palbociclib with endocrine agents.

A small phase Ib study in postmenopausal patients with HR-positive, HER2-negative advanced breast cancer showed a good tolerability of the combination of palbociclib 125 mg daily (schedule 3/1) and letrozole 2.5 mg daily continuously, with no pharmacokinetic interactions [Slamon *et al.* 2010].

An open label, international, randomized phase II study assessed the efficacy and safety of palbociclib plus letrozole in postmenopausal patients

with ER-positive, HER2-negative breast cancer, as first-line therapy for advanced disease (PALOMA-1/TRIO-18 trial) [Finn *et al.* 2015]. Patients could have received neoadjuvant or adjuvant endocrine treatment, but, if this included letrozole, it would need to have been stopped at least 1 year before study entry. Two separate cohorts were enrolled sequentially: cohort 1 included any patient with ER-positive, HER2-negative disease, while in cohort 2, further requirements were either amplification of *CCND1*, or loss of *CDKN2A*, or both, assessed at a central laboratory. Patients were randomly allocated 1:1 to receive either palbociclib 125 mg daily for 3 weeks of any 4-week cycle, plus letrozole 2.5 mg daily continuously, or letrozole 2.5 mg daily alone. The intention of cohort 1 was to provide preliminary safety and efficacy data, and the analysis of the primary endpoint of PFS was initially intended as based on cohort 2 only, after enrollment of 150 patients. However, an unplanned interim analysis of cohort 1 showed relevant activity of palbociclib, with PFS hazard ratio (HR) of 0.35 (95% confidence interval (CI), 0.17–0.72; $p = 0.006$) favoring the combination arm. This preliminary analysis also suggested that further patient selection based on biomarkers was unlikely to further increase the difference between treatment arms. Therefore, cohort 2 was closed in advance, and the study was amended, planning a combined analysis of cohorts 1 and 2 for the primary endpoint of PFS. After enrollment of 165 women, stratified by disease site and disease-free interval, 84 were randomized to palbociclib plus letrozole and 81 to letrozole alone, and after almost 30 months (median) follow up, the median PFS was 20.2 months for the combination arm *versus* 10.2 months for letrozole alone, with HR, 0.488; 95% CI, 0.319–0.748; $p = 0.0004$. The difference in PFS between the two arms was significant in both cohorts 1 and 2, and was consistent across patient subgroups defined by demographic, clinical and biological features [Finn *et al.* 2016a], apart from the lack of significant difference between the study arms in the subgroup of patients with disease recurrence within 12 months from the end of adjuvant therapy. Also, the rates of objective responses (ORR, 43% *versus* 33%; $p = 0.13$) and clinical benefit (CBR, 81% *versus* 58%; $p = 0.0009$) were higher in the combination arm, whereas overall survival (OS) was not different between the two arms (HR, 0.813; 95% CI, 0.492–1.345; $p = 0.42$). Treatment was well tolerated and the mean relative dose intensity for palbociclib was 94%.

Leukopenia, neutropenia and fatigue were the most common adverse events in the palbociclib arm. An evaluation of pain by means of the Brief Pain Inventory showed no significant differences in pain severity and interference with daily activities between the two arms [Bell *et al.* 2016].

On the basis of these results, in February 2015, the US Food and Drug Administration (FDA) granted accelerated approval to palbociclib for use in combination with letrozole for the treatment of postmenopausal women with ER-positive, HER2-negative advanced breast cancer, as initial endocrine-based therapy for their metastatic disease [US Food and Drug Administration, 2015].

Based on preclinical data of palbociclib synergism with paclitaxel when administered on an alternating schedule, enabling tumor cell synchronization before the administration of paclitaxel, a phase Ib trial, including a dose escalation and an expansion cohort, identified palbociclib 75 mg daily as optimal combination for 4 days of run-in, and then on days 2–4, 9–11, 16–18 of each 28-day cycle, plus paclitaxel 80 mg/m² weekly for three cycles and then on days 1, 8, 15 of 28-day cycles [Clark *et al.* 2016]. Dose-limiting toxicities were grade 3 transaminitis and febrile neutropenia, and uncomplicated grade 3 and 4 neutropenia was the most common adverse event. Among 24 evaluable patients, 14 (58%) had a partial response or stable disease longer than 6 months, supporting further development of the regimen.

Phase III studies

In the double blind, phase III PALOMA-3 trial, patients with HR-positive, HER2-negative advanced breast cancer that had relapsed or progressed during prior endocrine therapy (within 12 months from completion of adjuvant endocrine therapy, or during or within 1 month after the end of prior endocrine therapy for advanced disease) with tamoxifen if pre- or perimenopausal, or with an AI if postmenopausal, were randomized in a 2:1 ratio to palbociclib (125 mg daily for 3 weeks, followed by 1 week off over 4-week cycles) plus fulvestrant (500 mg intramuscularly on days 1, 15, 29, then every 4 weeks) or to placebo plus fulvestrant [Turner *et al.* 2015; Cristofanilli *et al.* 2016]. Pre- and perimenopausal patients received also goserelin 3.6 mg subcutaneously every 4 weeks, starting at least 4 weeks before randomization. One prior line of chemotherapy for advanced disease was allowed. Of 521 patients enrolled,

stratified by disease site, menopausal status, and sensitivity to prior endocrine therapy, 347 received palbociclib plus fulvestrant and 174, placebo plus fulvestrant. Almost 80% of the patients had tumors that had been sensitive to prior endocrine therapy. At a median follow up of 8.9 months, median PFS was 9.5 months for fulvestrant plus palbociclib *versus* 4.6 months for fulvestrant plus placebo, with HR, 0.46 (95% CI, 0.36–0.59; $p < 0.0001$). The ORR in patients with measurable disease was 25% in the palbociclib arm and 11% in the placebo arm ($p = 0.0012$), with median time to response of 112 days and 57 days respectively, and the CBR was 64% and 36%, respectively ($p < 0.0001$). Results from subgroup analyses were generally consistent with those from the overall population. In particular, the degree of endocrine sensitivity (either in terms of previous endocrine response or HR expression) did not affect benefit, nor did the menopausal status [Loibl *et al.* 2016]. Although PIK3CA mutations (evaluated on circulating cell-free DNA on 394 patients, with a mutation rate of 33%) conferred a nonsignificant worsening in PFS, the magnitude of benefit from palbociclib was independent from PIK3CA status. Likewise, while *ESR1* gene mutations conferred a marginally significant worsening of PFS compared with wild-type status, benefit from palbociclib was independent from *ESR1* status [Fribbens *et al.* 2016]. Grade 3 or 4 neutropenia occurred in 65% of patients receiving palbociclib *versus* 1% of those receiving placebo; nonetheless, febrile neutropenia was uncommon (1%) in both groups. Adverse events with >10% increased frequency in the palbociclib group were neutropenia, leucopenia, anemia, thrombocytopenia, stomatitis, alopecia, rash, infections and fatigue; these were mainly mild or moderate, and treatment discontinuation due to adverse events occurred in 4% in the palbociclib group and 2% with placebo [Verma *et al.* 2016]. The median time to onset of grade ≥ 3 neutropenia was 16 days, and the median duration of grade ≥ 3 neutropenia, 7 days. The mean relative dose intensity for palbociclib plus fulvestrant was 86%, but dose reductions due to neutropenia did not have a detrimental effect on efficacy. The combined treatment significantly delayed deterioration of global quality of life and improved pain compared with the placebo arm [Harbeck *et al.* 2016].

On the basis of PALOMA-3 results, on 19 February 2016, the FDA approved palbociclib for use in combination with fulvestrant for the

treatment of women with HR-positive, HER2-negative advanced or metastatic breast cancer with disease progression following endocrine therapy [US Food and Drug Administration, 2016].

Results of the double blind, phase III PALOMA-2 trial, designed to confirm those from PALOMA-1, were presented at the 2016 ASCO meeting [Finn *et al.* 2016b]. In the trial, 666 postmenopausal patients with HR-positive, HER2-negative advanced breast cancer, not previously treated for advanced disease, stratified for disease site, disease-free interval, and prior (neo)adjuvant hormonal therapy, were randomized 2:1 to palbociclib plus letrozole (same doses and schedule as in PALOMA-1) or placebo plus letrozole. Baseline characteristics were well balanced among groups. After a median follow up of about 23 months, median PFS was 24.8 months in the palbociclib arm and 14.5 months in the placebo arm, with HR 0.58 (95% CI, 0.46–0.72; $p < 0.000001$). The ORR was 42% in the palbociclib arm *versus* 35% with placebo ($p = 0.031$) and the CBR was 85% and 70%, respectively ($p < 0.0001$). Benefit from palbociclib was apparent across all prespecified subgroups. The median relative dose intensity for palbociclib was 93%. Common adverse events (any grade, palbociclib *versus* placebo group) were: neutropenia (79.5% *versus* 6.3%), fatigue (37.4% *versus* 27.5%), nausea (35.1% *versus* 26.1%), arthralgia (33.3% *versus* 33.8%) and alopecia (32.9% *versus* 15.8%). Febrile neutropenia was seen only with palbociclib (2.5%). Permanent discontinuation due to AEs was 9.7% with palbociclib *versus* 5.9% with placebo. The OS analysis is pending.

The low incidence of febrile neutropenia compared with the high incidence of grade 3/4 neutropenia is due to the cytostatic effect of palbociclib on bone marrow cells that is reversible upon drug withdrawal, contrary to chemotherapeutic agents that induce bone marrow progenitors' apoptosis [Hu *et al.* 2016].

Neoadjuvant studies

A phase II neoadjuvant study has been conducted in patients with clinical stage II/III ER-positive, HER2-negative breast cancer [Ma *et al.* 2016] to assess molecular changes induced by anastrozole and palbociclib. Therapy consisted of anastrozole alone for 28 days, followed by anastrozole plus palbociclib (schedule 3/1) for four 28-day cycles,

followed by 2–4 weeks with anastrozole alone before surgery. A subgroup of patients also received palbociclib for 10–12 days immediately prior to surgery. The primary endpoint was the rate of complete CCA, defined as $Ki67 \leq 2.7\%$, at an early biopsy taken on day 15 of the first cycle of combined treatment. Premenopausal patients also received goserelin. Preliminary results were reported at the 2015 San Antonio Breast Cancer Symposium: 39 of 45 evaluable patients achieved CCA (87%), including 22 of 28 patients (79%) with wild-type PIK3CA, 15 out of 15 patients with mutant PIK3CA and 2 out of 2 with undetermined PIK3CA. Among 43 patients who also had a biopsy just before starting palbociclib, anastrozole alone induced CCA in 11 cases (26%), whereas 15 days of palbociclib converted non-CCA to CCA in another 26 patients (60%). The combination was ineffective in two patients with nonluminal cancers and in a subset of luminal B cancers. Clinical responses were seen in 67% of the patients and no pathological complete response was seen. While patients who stopped palbociclib 2–4 weeks before surgery showed a rebound in Ki67 assessed at surgery, those who received palbociclib immediately before surgery did not, highlighting the cytostatic action of the drug, more apt for maintenance treatment.

In a further short-term preoperative trial, aimed at assessing the CCA rate and to identify predictive biomarkers, 100 patients with untreated breast cancer were randomized 3:1 to palbociclib 125 mg daily for 14 days until the day before surgery, *versus* no treatment. The CCA rate was 58% with palbociclib *versus* 10% in the control arm. No Ki67 response was observed in triple-negative and in HER2-positive tumors. While baseline Rb, phosphorylated Rb and p16 did not predict response, responses were associated with changes in phosphorylated Rb from baseline [Arnedos *et al.* 2016].

Studies with other cyclin-dependent kinase 4/6 inhibitors

Several clinical trials are ongoing with the other CDK4/6 inhibitors ribociclib and abemaciclib, and have been reviewed elsewhere [O’Leary *et al.* 2016; Barroso-Sousa *et al.* 2016]. Particularly relevant are studies of triple drug combinations. Preliminary results have been reported for a phase Ib/II study of triplet therapy with ribociclib, everolimus, and exemestane in 77 evaluable postmenopausal women with HR-positive,

HER2-negative advanced breast cancer [Bardia *et al.* 2016], demonstrating feasibility and a disease control rate (patients who don’t have disease progression as best response) of 73% in heavily pretreated patients. In a further phase Ib/II study, 46 patients received a combination of ribociclib, alpelisib and letrozole that showed an acceptable safety profile and a disease control rate of 70% [Juric *et al.* 2016].

Ongoing clinical trials with palbociclib

Several phase Ib, II, and III studies are ongoing with palbociclib (Table 1). Among these are: some adjuvant studies, such as the phase III PALLAS trial comparing endocrine therapy of physician choice for at least 5 years with endocrine therapy for at least 5 years plus palbociclib for 2 years, and the PENELOPE-B trial, with standard endocrine treatment plus palbociclib or placebo in patients with residual disease after neoadjuvant chemotherapy; several neoadjuvant studies, both in HER2-positive, HR-positive and in HER2-negative, HR-positive breast cancer, some of which comparing endocrine therapy plus palbociclib with chemotherapy; several trials in the metastatic setting, either comparing endocrine therapy plus palbociclib with chemotherapy, or exploring combinations of palbociclib with other new drugs such as new endocrine agents, PI3K inhibitors (the pan-class I PI3K inhibitor pictilisib and the selective inhibitor of class I PI3K α , γ , and δ isoforms, taselisib), anti-PD-1 monoclonal antibodies and anti-HER2 drugs. Studies of triple combinations of palbociclib with an endocrine agent (fulvestrant or letrozole) and the PI3K/mTOR inhibitor gedatolisib are also ongoing.

Open questions and future directions

Palbociclib received FDA-accelerated approval in HER2-negative luminal breast cancer, both as initial endocrine-based therapy for metastatic disease, in combination with letrozole, and after progression, following an endocrine therapy for advanced disease, in combination with fulvestrant. The best positioning of palbociclib in the therapeutic strategy of luminal tumors remains an open question, partly intertwining with the still-open issue of the best sequencing of endocrine agents in the metastatic setting. Although the degree of endocrine sensitivity does not seem to affect the benefit from palbociclib, some distinctions are worthy. While more aggressive tumors

Table 1. Ongoing interventional trials with palbociclib in breast cancer.

ClinicalTrials.gov identifier (trial name)	Inclusion criteria	Study design	Intervention
Adjuvant setting NCT02513394 (PALLAS)	HER2-/HR + EBC	Phase III, randomized (two arms), open label	Endocrine TPC (at least 5 years) + palbociclib (2 years)
NCT01864746 (PENELOPE-B)	HER2-/HR + EBC (with residual invasive disease following neoadjuvant treatment)	Phase III, randomized (two arms), double blind	Endocrine TPC + placebo Endocrine TPC + palbociclib
NCT02040857 (13-559)	HER2-/HR + EBC	Phase II, single arm, open label	Endocrine TPC + palbociclib
Neoadjuvant setting NCT02530424 (NA-PHER2)	HER2+/ER+ EBC or LABC	Phase II, single arm, open label	Fulvestrant + trastuzumab + pertuzumab + palbociclib
NCT02764541 (16-052, PELOPS)	HER2-/HR + EBC	Phase II ('window treatment' followed by treatment phase), two-stage sequentially randomized, open label	Tamoxifen ('window treatment') followed by endocrine TPC ± palbociclib (treatment phase)
NCT02296801 (NSABP FB-11, PALLET)	HER2-/HR + postmenopausal EBC	Phase II, randomized (four arms), open label	Letrozole; letrozole + palbociclib
NCT02760030 (OSU-15266)	HER2-/ER+ older EBC not removable by surgery	Phase II, single arm, open label	Fulvestrant + palbociclib
NCT02400567 (NeoPAL)	HER2-/HR + PAM50 defined luminal postmenopausal EBC	Phase II, randomized (two arms), open label	Letrozole + palbociclib
NCT02626507 (CL- Gedatolisib-001)	HER2-/ER+ EBC or LABC	Phase Ib, single arm, open label	Gedatolisib + fulvestrant + palbociclib
NCT02603679 (PREDIX LumB)	HER2-/ER+ luminal B type or luminal A type with lymph node metastases EBC	Phase II, randomized (two arms), open label	Tamoxifen or AI ± goserelin + palbociclib
NCT02592083 (PREDIX LumA)	HER2-/ER+ luminal A type with age older than 40 years EBC	Phase II, randomized (two arms), open label	Tamoxifen or AI ± goserelin + palbociclib Tamoxifen or AI ± goserelin
			Fluorouracile + epirubicin + cyclophosphamide followed by docetaxel Paclitaxel

Table 1. (Continued)

ClinicalTrials.gov identifier (trial name)	Inclusion criteria	Study design	Intervention
NCT02624973 (PETREMAC)	High-risk EBC with known ER, PGR, HER2 and TP53 status	Phase II, nonrandomized (eight cohorts based on ER, PGR, and HER2 expression and on TP53 mutation status), open label	Tamoxifen + goserelin + palbociclib or letrozole + palbociclib; docetaxel + cyclophosphamide; docetaxel; docetaxel + trastuzumab + pertuzumab; docetaxel + cyclophosphamide + trastuzumab + pertuzumab; olaparib + carboplatin; olaparib
NCT01709370 (OOTR-N007/LET-CDK)	HER2-/ER+ postmenopausal EBC	Phase II, single arm, open label	Letrozole + palbociclib
NCT01723774 (201301106)	HER2-/ER+ MBC	Phase II, nonrandomized (three cohorts), open label	Anastrozole + palbociclib
Metastatic setting NCT01976169 (STU 042013-042)	HER2+ with Rb protein-proficiency LABC or MBC	Phase Ib, single arm, open label	Trastuzumab emtansine + palbociclib
NCT02448420 (PATRICIA)	HER2+/ER- or HER2+/ER+ postmenopausal LABC or MBC	Phase II, single arm, open label (HER2+/ER-); randomized (two arms), open label (HER2+/ER+)	Trastuzumab + palbociclib (HER2+/ER-) Trastuzumab + palbociclib (HER2+/ER+)
NCT02774681 (NU 15B08)	HER2-/HR- or HER2+/HR-LABC or MBC	Phase II, single arm, open label	Palbociclib (± trastuzumab for HER2+/HR-)
NCT02605486 (15-207)	HER2-/HR- AR+ postmenopausal MBC	Phase I/II, single arm, open label	Bicalutamide + palbociclib
NCT02491983 (PARSIFAL)	HER2-/HR+ postmenopausal LABC or MBC	Phase II, randomized (two arms), open label	Letrozole + palbociclib
NCT02600923 (A5481053)	HER2-/HR+ postmenopausal MBC	Phase III, single arm, open label	Letrozole + palbociclib
NCT01823835 (G029642)	HER2-/ER+ LABC or MBC	Phase I/II, single arm, open label (phase I part I and part II; phase II)	GDC-0810 (phase I, part I) GDC-0810 + palbociclib and/orLHRH agonist (phase I, part II and phase II)
NCT02592746 (2015-08-042)	HER2-/HR+ premenopausal MBC	Phase II, randomized (two arms), open label	Exemestane + goserelin + palbociclib Capecitabine

(Continued)

Table 1. (Continued)

ClinicalTrials.gov identifier (trial name)	Inclusion criteria	Study design	Intervention
NCT02684032 (B2151009)	HER2-/ER+ MBC	Phase Ib, nonrandomized (two cohorts), open label	Gedatolisib + letrozole + palbociclib
NCT02384239 (147522)	HER2-/HR+ MBC previously exposed to mTOR or PI3K inhibitors	Phase II, sequentially randomized, open label	Fulvestrant or tamoxifen + palbociclib 100 mg
NCT02630693 (MA38)	HER2-/ER+ MBC	Phase II, randomized (two arms), open label	Fulvestrant or tamoxifen or AI + palbociclib 100 mg
NCT02389842 (PIPA)	Advanced solid tumor (part I); HER2-/ER+ or HER2-/ER- or HER2+ MBC or other advanced solid tumor with a PIK3CA mutation (part II)	Phase Ib, nonrandomized (two cohorts), open label	Taselisib + palbociclib
NCT02549430 (TREnd)	HER2-/ER+ postmenopausal MBC progressed on endocrine therapy	Phase II, randomized (two arms), open label	Palbociclib
NCT02806050 (201600066)	HER2-/ER+ postmenopausal MBC	Phase II, single arm, open label	Letrozole + palbociclib, with FES PET at baseline
NCT01037790 (UPCC 03909)	Advanced solid tumor, including HER2+, ER+, HER2-/HR- LRBC or MBC	Phase II, nonrandomized (five cohorts), open label	Palbociclib (± trastuzumab for HER2+)
NCT02778685 (16058)	HER2-/ER+ postmenopausal MBC with stable disease but not responding to letrozole and palbociclib	Phase II, single arm, open label	Letrozole + palbociclib + pembrolizumab
NCT02679755 (A5481037)	HER2-/ER+ postmenopausal MBC (first-line treatment)	Phase IV, single arm, open label	Letrozole + palbociclib
NCT02692755 (PALINA)	Black, African or African-American HER2-/HR+ LRBC or MBC	Phase II, single arm, open label	Letrozole + palbociclib
NCT02028507 (PEARL)	HER2-/ER+ postmenopausal MBC	Phase III, randomized (two cohorts), open label	Exemestane + palbociclib (cohort I) Fulvestrant + palbociclib (cohort II)
			Capecitabine (cohort I) Capecitabine (cohort II)

Table 1. (Continued)

ClinicalTrials.gov identifier (trial name)	Inclusion criteria	Study design	Intervention
NCT02599714 (PASTOR)	HER2-/ER+ postmenopausal LRBC or MBC	Phase I/II, single arm, open label (phase I); randomized (two arms), double blind (phase II)	Fulvestrant + AZD2014 + palbociclib (phase I) Fulvestrant + AZD2014 + palbociclib (phase II) Fulvestrant + placebo + palbociclib (phase II)
NCT02499146 (A5481019)	Chinese HER2-/ER+ LRBC or MBC	Phase I, single arm, open label	Letrozole + palbociclib
NCT02738866 (J15212)	HER2-/HR+ LABC or MBC progressed on palbociclib and AI	Phase II, single arm, open label	Fulvestrant + palbociclib
NCT02690480 (FLIPPER)	HER2-/HR+ postmenopausal MBC (first-line treatment)	Phase II, randomized (two arms), double blind	Fulvestrant + palbociclib
NCT02536742 (PYTHIA)	HER2-/HR+ postmenopausal LRBC or MBC	Phase II, randomized (two arms), double blind (AURORA molecular screening program)	Fulvestrant + palbociclib
NCT02448771 (15-060)	HER2-/HR+ LABC or MBC	Phase Ib/II, single arm, open label	Bazedoxifene + palbociclib
NCT02686666 (BTCRC-BRE15-016)	HER2-/HR+ LRBC or MBC (first-line treatment)	Phase II, single arm, open label	Tamoxifen + palbociclib
NCT01684215 (A5481010)	Japanese advanced solid tumor (phase I, part II); HER2-/ER+ postmenopausal LRBC or MBC (phase I, part II and phase II, first-line treatment)	Phase I/II, single arm, open label (phase I, part I and part II; phase II)	Palbociclib (phase I part I) Letrozole + palbociclib (phase I, part I and phase II)
NCT02297438 (A5481027 - PALOMA4)	Asian HER2-/ER+ postmenopausal LRBC or MBC (first-line treatment)	Phase III, randomized (two arms), double blind	Letrozole + palbociclib
NCT02142868 (A5481034)	HER2-/HR+ postmenopausal LRBC or MBC	Phase III, single arm, open label (expanded access)	Letrozole + palbociclib
HER2, human epidermal growth factor receptor 2; AR, androgen receptor; ER, estrogen receptor; HR, hormone receptor (estrogen e/o progesterone receptor); EBC, early breast cancer; LABC, locally advanced breast cancer; LRBC, locoregionally recurrent breast cancer; MBC, metastatic breast cancer; TPC, therapy of physician choice; AI, aromatase inhibitor; LHRH, luteinizing hormone-releasing hormone; PAM50, Prediction Analysis of Microarray 50-gene classifier; PGR, Progesterone Receptor; TP53, p53 tumor suppressor gene; mTOR, mammalian target of rapamycin; PI3K, phosphoinositide 3-kinase; PIK3CA, Phosphatidylinositol-4,5-Bisphosphate 3-Kinase Catalytic Subunit Alpha; AZD2014, a dual mammalian target of rapamycin (mTOR) Complex 1 and 2 inhibitor. Data retrieved from ClinicalTrials.gov (accessed on 21 July 2016).			

certainly deserve the use of palbociclib from the first line of treatment, some indolent or highly endocrine-sensitive tumors may also have long PFS with endocrine therapy alone. Therefore, the trade-off between improved efficacy and added side effects may differ, depending on patient and tumor characteristics and line of treatment. On the other hand, everolimus, another agent that improves the clinical outcome when added to endocrine therapy, was studied in patients progressing on, or shortly after, therapy with a non-steroidal AI, and will more likely, but not necessarily, be used after therapy with palbociclib, although the efficacy of everolimus plus exemestane after progression to CDK4/6 inhibitors (or *vice versa*) has not been addressed in clinical trials. A further open issue that is being addressed in clinical trials is the efficacy of continuing CDK4/6 inhibition beyond progression to palbociclib, for example, maintaining palbociclib while changing the endocrine agent, or changing the CDK4/6 inhibitor.

The studies of triple blockade are very intriguing, combining an endocrine agent with a CDK4/6 inhibitor with another drug, like a PI3K, a MAPK, or an mTOR inhibitor, or an anti-HER2, ideally chosen based on tumor genomic profile. Should these drug combinations prove tolerable, they could have the potential to be not only cytostatic, but to induce tumor cell apoptosis, with greater potential activity. Very active targeted drug combinations would be worth exploring in the first-line setting, even as an alternative to chemotherapy in aggressive tumors, as well as in the neoadjuvant setting.

Balancing benefits with toxicity and costs will be critical in the adjuvant setting [Carey and Perou, 2015], should palbociclib demonstrate efficacy in this context, given that a relevant fraction of patients with luminal tumors would be cured with surgery alone. In this regard, studies in the neoadjuvant setting may help to identify predictors of response. The potential of 3-deoxy-3-[(18)F] fluorothymidine (FLT) positron emission tomography in early response assessment has been explored in lymphomas [Leonard *et al.* 2012] and deserves study in breast cancer too, but further genomic and proteomic studies are likely needed for more accurate prediction.

Conclusion

Palbociclib is the first member of the CDK4/6 inhibitors entering the clinical arena. With 10-month

improvement in PFS in first-line metastatic HR-positive, HER2-negative breast cancer, when added to letrozole, it represents one of the best steps forward in the treatment of luminal breast cancer. Nonetheless, an impact on OS is still unproven, given the limited follow up of the phase III trials available. It also shows promise in other breast cancer subtypes, especially in a subset of HER2-positive tumors. Its single-agent activity is basically cytostatic, and combination with other agents is usually required to induce tumor cell senescence or apoptosis. Identifying response predictors will be essential for a rational use of the drug, to avoid unnecessary toxicity and costs [Matter-Walstra *et al.* 2016]. Given the complexity of the pathways regulating cell cycle, no single biomarker has emerged as predictor of response, beyond the presence of HRs. Further, deeper efforts in the characterization of a single tumor's biomarkers are ongoing, and a systemic approach will likely be necessary to better identify biomarkers' constellations associated with response or resistance, as well as the optimal combination therapies for any single tumor.

Funding

The author(s) received no financial support for the research, authorship, and/or publication of this article.

Conflict of interest statement

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

References

- Abedin, Z., Ma, Z. and Reddy, E. (2010) Increased angiogenesis in Cdk4 (R24C/R24C): Apc(+/-Min) intestinal tumors. *Cell Cycle* 9: 2456–2463.
- Aggarwal, P., Vaites, L., Kim, J., Mellert, H., Gurung, B., Nakagawa, H. *et al.* (2010) Nuclear cyclin D1/CDK4 kinase regulates CUL4 expression and triggers neoplastic growth via activation of the PRMT5 methyltransferase. *Cancer Cell* 18: 329–340.
- Altucci, L., Addeo, R., Cicatiello, L., Dauvois, S., Parker, M., Truss, M. *et al.* (1996) 17beta-Estradiol induces cyclin D1 gene transcription, p36D1-p34cdk4 complex activation and p105Rb phosphorylation during mitogenic stimulation of G(1)-arrested human breast cancer cells. *Oncogene* 12: 2315–2324.
- Alves, C., Elias, D., Lyng, M., Bak, M., Kirkegaard, T., Lykkesfeldt, A. *et al.* (2016) High CDK6 protects cells from fulvestrant-mediated apoptosis and is a predictor of resistance to fulvestrant in estrogen

- receptor-positive metastatic breast cancer. *Clin Cancer Res* 2016 June 1. [Epub ahead of print]
- An, H., Beckmann, M., Reifemberger, G., Bender, H. and Niederacher, D. (1999) Gene amplification and overexpression of CDK4 in sporadic breast carcinomas is associated with high tumor cell proliferation. *Am J Pathol* 154: 113–118.
- Anders, L., Ke, N., Hydbring, P., Choi, Y., Widlund, H., Chick, J. *et al.* (2011) A systematic screen for CDK4/6 substrates links FOXM1 phosphorylation to senescence suppression in cancer cells. *Cancer Cell* 20: 620–634.
- Andrechek, E. (2015) HER2/Neu tumorigenesis and metastasis is regulated by E2F activator transcription factors. *Oncogene* 34: 217–225.
- Arima, Y., Hayashi, H., Sasaki, M., Hosonaga, M., Goto, T., Chiyoda, T. *et al.* (2012) Induction of ZEB proteins by inactivation of Rb protein is key determinant of mesenchymal phenotype of breast cancer. *J Biol Chem* 287: 7896–7906.
- Arnedos, M., Cheaib, B., Bayar, M., Michiels, S., Scott, V., Adam, J. *et al.* (2016) Anti-proliferative response and predictive biomarkers to palbociclib in early breast cancer: the Preoperative Palbociclib (POP) randomized trial. *Cancer Res* 76: Abstract CT041.
- Asghar, U., Herrera-Abreu, M., Cutts, R., Babina, I., Pearson, A. and Turner, N. (2015) Identification of subtypes of triple negative breast cancer (TNBC) that are sensitive to CDK4/6 inhibition. *J Clin Oncol* 33: Abstract 11098.
- Bardia, A., Modi, S., Oliveira, M., Campone, M., Ma, B., Dirix, L. *et al.* (2016). Triplet therapy with ribociclib, everolimus, and exemestane in women with HR+/HER2- advanced breast cancer. *Cancer Res* 76: Abstract P6-13-01.
- Barroso-Sousa, R., Shapiro, G. and Tolaney, S. (2016) Clinical development of the CDK4/6 inhibitors ribociclib and abemaciclib in breast cancer. *Breast Care* 11: 167–173.
- Bartkova, J., Lukas, J., Müller, H., Lützhøft, D., Strauss, M. and Bartek, J. (1994) Cyclin D1 protein expression and function in human breast cancer. *Int J Cancer* 57: 353–361.
- Bell, T., Crown, J., Lang, I., Bhattacharyya, H., Zanotti, G., Randolph, S. *et al.* (2016) Impact of palbociclib plus letrozole on pain severity and pain interference with daily activities in patients with estrogen receptor-positive/human epidermal growth factor receptor 2-negative advanced breast cancer as first-line treatment. *Curr Med Res Opin* 32: 959–965.
- Bienvenu, F., Jirawatnotai, S., Elias, J., Meyer, C., Mizeracka, K., Marson, A. *et al.* (2010) Transcriptional role of cyclin D1 in development revealed by a genetic-proteomic screen. *Nature* 463: 374–378.
- Bosco, E., Wang, Y., Xu, H., Zilfou, J., Knudsen, K., Aronow, B. *et al.* (2007) The retinoblastoma tumor suppressor modifies the therapeutic response of breast cancer. *J Clin Invest* 117: 218–228.
- Cairns, P., Polascik, T., Eby, Y., Tokino, K., Califano, J., Merlo, A. *et al.* (1995) Frequency of homozygous deletion at p16/CDKN2 in primary human tumours. *Nat Genet* 11: 210–212.
- Cancer Genome Atlas Network. (2012) Comprehensive molecular portraits of human breast tumours. *Nature* 490: 61–70.
- Carey, L. and Perou, C. (2015) Palbociclib—taking breast-cancer cells out of gear. *N Engl J Med* 373: 273–274.
- Carroll, J., Prall, O., Musgrove, E. and Sutherland, R. (2000) A pure estrogen antagonist inhibits cyclin E-CDK2 activity in MCF-7 breast cancer cells and induces accumulation of p130-E2F4 complexes characteristic of quiescence. *J Biol Chem* 275: 38221–38229.
- Chandarlapaty, S., Sawai, A., Scaltriti, M., Rodrik-Outmezguine, V., Grbovic-Huezo, O., Serra, V. *et al.* (2011) Akt inhibition relieves feedback suppression of receptor tyrosine kinase expression and activity. *Cancer Cell* 19: 58–71.
- Chen, H., Tsai, S. and Leone, G. (2009) Emerging roles of E2Fs in cancer: an exit from cell cycle control. *Nat Rev Cancer* 9: 785–797.
- Chicas, A., Wang, X., Zhang, C., McCurrach, M., Zhao, Z., Mert, O. *et al.* (2010) Dissecting the unique role of the retinoblastoma tumor suppressor during cellular senescence. *Cancer Cell* 17: 376–387.
- Choi, Y., Li, X., Hydbring, P., Sanda, T., Stefano, J., Christie, A. *et al.* (2012) The requirement for cyclin D function in tumor maintenance. *Cancer Cell* 22: 438–451.
- Chu, I., Hengst, L. and Slingerland, J. (2008) The Cdk inhibitor p27 in human cancer: prognostic potential and relevance to anticancer therapy. *Nat Rev Cancer* 8: 253–267.
- Clark, A., O'Dwyer, P., Troxel, A., Lal, P., Feldman, M., Gallagher, M. *et al.* (2016) Palbociclib and paclitaxel on an alternating schedule for advanced breast cancer: results of a phase Ib trial. *Cancer Res* 76: Abstract P6-13-08.
- Cristofanilli, M., Turner, N., Bondarenko, I., Ro, J., Im, S., Masuda, N. *et al.* (2016) Fulvestrant plus palbociclib versus fulvestrant plus placebo for treatment of hormone-receptor-positive, HER2-negative metastatic breast cancer that progressed on previous endocrine therapy (PALOMA-3): final analysis of

- the multicentre, double-blind, phase III randomised controlled trial. *Lancet Oncol* 17: 425–439.
- De Gooijer, M., Zhang, P., Thota, N., Mayayo-Peralta, I., Buil, L., Beijnen, J. *et al.* (2015) P-glycoprotein and breast cancer resistance protein restrict the brain penetration of the CDK4/6 inhibitor palbociclib. *Invest New Drugs* 33: 1012–1019.
- Dean, J., McClendon, A., Hickey, T., Butler, L., Tilley, W., Witkiewicz, A. *et al.* (2012a) Therapeutic response to CDK4/6 inhibition in breast cancer defined by *ex vivo* analyses of human tumors. *Cell Cycle* 11: 2756–2761.
- Dean, J., McClendon, A. and Knudsen, E. (2012b) Modification of the DNA damage response by therapeutic CDK4/6 inhibition. *J Biol Chem* 287: 29075–29087.
- Dean, J., Thangavel, C., McClendon, A., Reed, C. and Knudsen, E. (2010) Therapeutic CDK4/6 inhibition in breast cancer: key mechanisms of response and failure. *Oncogene* 29: 4018–4032.
- DeMichele, A., Clark, A., Tan, K., Heitjan, D., Gramlich, K., Gallagher, M. *et al.* (2015) CDK 4/6 inhibitor palbociclib (PD0332991) in Rb+ advanced breast cancer: phase II activity, safety, and predictive biomarker assessment. *Clin Cancer Res* 21: 995–1001.
- Dick, F. and Rubin, S. (2013) Molecular mechanisms underlying RB protein function. *Nat Rev Mol Cell Biol* 14: 297–306.
- Ertel, A., Dean, J., Rui, H., Liu, C., Witkiewicz, A., Knudsen, K. *et al.* (2010) Rb-pathway disruption in breast cancer: differential association with disease subtypes, disease-specific prognosis and therapeutic response. *Cell Cycle* 9: 4153–4163.
- Finn, R., Crown, J., Ettl, J., Schmidt, M., Bondarenko, I., Lang, I. *et al.* (2016a) Efficacy and safety of palbociclib in combination with letrozole as first-line treatment of ER-positive, HER2-negative, advanced breast cancer: expanded analyses of subgroups from the randomized pivotal trial PALOMA-1/TRIO-18. *Breast Cancer Res* 18: 67.
- Finn, R., Crown, J., Lang, I., Boer, K., Bondarenko, I., Kulyk, S. *et al.* (2015) The cyclin-dependent kinase 4/6 inhibitor palbociclib in combination with letrozole *versus* letrozole alone as first-line treatment of oestrogen receptor-positive, HER2-negative, advanced breast cancer (PALOMA-1/TRIO-18): a randomised phase II study. *Lancet Oncol* 16: 25–35.
- Finn, R., Dering, J., Conklin, D., Kalous, O., Cohen, D., Desai, A. *et al.* (2009) PD 0332991, a selective cyclin D kinase 4/6 inhibitor, preferentially inhibits proliferation of luminal estrogen receptor-positive human breast cancer cell lines *in vitro*. *Breast Cancer Res* 11: R77.
- Finn, R., Martin, M., Rugo, H., Jones, S., Im, S., Gelmon, K. *et al.* (2016b) PALOMA-2: primary results from a phase III trial of palbociclib (P) with letrozole (L) compared with letrozole alone in postmenopausal women with ER+/HER2– advanced breast cancer (ABC). *J Clin Oncol* 34: Abstract 507.
- Flaherty, K., Lorusso, P., DeMichele, A., Abramson, V., Courtney, R., Randolph, S. *et al.* (2012) Phase I, dose-escalation trial of the oral cyclin-dependent kinase 4/6 inhibitor PD 0332991, administered using a 21-day schedule in patients with advanced cancer. *Clin Cancer Res* 18: 568–576.
- Foster, J., Henley, D., Bukovsky, A., Seth, P. and Wimalasena, J. (2001) Multifaceted regulation of cell cycle progression by estrogen: regulation of CDK inhibitors and CDC25A independent of cyclin D1-CDK4 function. *Mol Cell Biol* 21: 794–810.
- Fribbens, C., O’Leary, B., Kilburn, L., Hrebien, S., Garcia-Murillas, I., Beaney, M. *et al.* (2016) Plasma ESR1 mutations and the treatment of estrogen receptor-positive advanced breast cancer. *J Clin Oncol* 34: 2961–2968.
- Fry, D., Harvey, P., Keller, P., Elliott, W., Meade, M., Trachet, E. *et al.* (2004) Specific inhibition of cyclin-dependent kinase 4/6 by PD 0332991 and associated antitumor activity in human tumor xenografts. *Mol Cancer Ther* 3: 1427–1438.
- Gillett, C., Fantl, V., Smith, R., Fisher, C., Bartek, J., Dickson, C. *et al.* (1994) Amplification and overexpression of cyclin D1 in breast cancer detected by immunohistochemical staining. *Cancer Res* 54: 1812–1817.
- Goel, S., Wang, Q., Watt, A., Tolaney, S., Dillon, D., Li, W. *et al.* (2016) Overcoming therapeutic resistance in HER2-Positive breast cancers with CDK4/6 inhibitors. *Cancer Cell* 29: 255–269.
- Gopalan, P., Gordillo-Villegas, A., Zajac-Kaye, M. and Kaye, F. (2013) Inhibitory effect of the CDK4/6 inhibitor, PD 0332991, is enhanced by mTOR inhibition in non-small cell lung cancer (NSCLC). *Cancer Res* 73: Abstract 693.
- Hanahan, D. and Weinberg, R. (2011) Hallmarks of cancer: the next generation. *Cell* 144: 646–674.
- Handschiek, K., Beuerlein, K., Jurida, L., Bartkuhn, M., Müller, H., Soelch, J. *et al.* (2014) Cyclin-dependent kinase 6 is a chromatin-bound cofactor for NF- κ B-dependent gene expression. *Mol Cell* 53: 193–208.
- Harbeck, N., Iyer, S., Turner, N., Cristofanilli, M., Ro, J., André, F. *et al.* (2016) Quality of life with palbociclib plus fulvestrant in previously treated hormone receptor-positive, HER2-negative metastatic breast cancer: patient-reported outcomes from the PALOMA-3 trial. *Ann Oncol* 27: 1047–1054.

- Harbour, J., Luo, R., Dei Santi, A., Postigo, A. and Dean, D. (1999) CDK phosphorylation triggers sequential intramolecular interactions that progressively block Rb functions as cells move through G1. *Cell* 98: 859–869.
- Hashizume, R., Zhang, A., Mueller, S., Prados, M., Lulla, R., Goldman, S. *et al.* (2016) Inhibition of DNA damage repair by the CDK4/6 inhibitor palbociclib delays irradiated intracranial atypical teratoid rhabdoid tumor and glioblastoma xenograft regrowth. *Neuro Oncol* 18: 1519–1528.
- Herrera-Abreu, M., Palafox, M., Asghar, U., Rivas, M., Cutts, R., Garcia-Murillas, I. *et al.* (2016) Early Adaptation and acquired resistance to CDK4/6 inhibition in estrogen receptor-positive breast cancer. *Cancer Res* 76: 2301–2313.
- Herschkowitz, J., He, X., Fan, C. and Perou, C. (2008) The functional loss of the retinoblastoma tumour suppressor is a common event in basal-like and luminal B breast carcinomas. *Breast Cancer Res* 10: R75.
- Hirai, H. and Sherr, C. (1996) Interaction of D-type cyclins with a novel myb-like transcription factor, DMP1. *Mol Cell Biol* 16: 6457–6467.
- Hu, W., Sung, T., Jessen, B., Thibault, S., Finkelstein, M., Khan, N. *et al.* (2016) Mechanistic investigation of bone marrow suppression associated with palbociclib and its differentiation from cytotoxic chemotherapies. *Clin Cancer Res* 22: 2000–2008.
- Hui, R., Finney, G., Carroll, J., Lee, C., Musgrove, E. and Sutherland, R. (2002) Constitutive overexpression of cyclin D1 but not cyclin E confers acute resistance to antiestrogens in T-47D breast cancer cells. *Cancer Res* 62: 6916–6923.
- Hydbring, P., Malumbres, M. and Sicinski, P. (2016) Non-canonical functions of cell cycle cyclins and cyclin-dependent kinases. *Nat Rev Mol Cell Biol* 17: 280–292.
- Ichikawa, A., Ando, J. and Suda, K. (2008) G1 arrest and expression of cyclin-dependent kinase inhibitors in tamoxifen-treated MCF-7 human breast cancer cells. *Hum Cell* 21: 28–37.
- International Agency for Research on Cancer. (2016) Cancer fact sheets: breast cancer. *Cancer Today*. Available at: <http://gco.iarc.fr/today/fact-sheets-cancers?cancer=15&type=0&sex=2> (accessed 26 July 2016).
- Jansen, V., Bhola, N., Bauer, J., Formisano, L., Moore, P., Koch, J. *et al.* (2016) Inhibition of 3-phosphoinositide dependent protein kinase 1 (PDK1) synergizes with CDK4/6 inhibitors against ER-positive breast cancer. *Cancer Res* 76: Abstract PD2–06.
- Jeselsohn, R., Buchwalter, G., De Angelis, C. and Brown, M. (2015) Schiff R. ESR1 mutations—a mechanism for acquired endocrine resistance in breast cancer. *Nat Rev Clin Oncol* 12: 573–583.
- Jirawatnotai, S., Hu, Y., Michowski, W., Elias, J., Becks, L., Bienvenu, F. *et al.* (2011) A function for cyclin D1 in DNA repair uncovered by protein interactome analyses in human cancers. *Nature* 474: 230–234.
- Johnson, J., Thijssen, B., McDermott, U., Garnett, M., Wessels, L. and Bernardis, R. (2016) Targeting the RB-E2F pathway in breast cancer. *Oncogene* 35: 4829–4835.
- Juric, D., Ismail-Khan, R., Campone, M., García a-Esté vez, L., Becerra, C., De Boer, R. *et al.* (2016) Phase Ib/II study of ribociclib and alpelisib and letrozole in ER+, HER2– breast cancer: safety, preliminary efficacy and molecular analysis. *Cancer Res* 76: Abstract P3–14–01.
- Knowlden, J., Hutcheson, I., Jones, H., Madden, T., Gee, J., Harper, M. *et al.* (2003) Elevated levels of epidermal growth factor receptor/c-erbB2 heterodimers mediate an autocrine growth regulatory pathway in tamoxifen-resistant MCF-7 cells. *Endocrinology* 144: 1032–1044.
- Kollmann, K., Heller, G., Schneckenleithner, C., Warsch, W., Scheicher, R., Ott, R. *et al.* (2013) A kinase-independent function of CDK6 links the cell cycle to tumor angiogenesis. *Cancer Cell* 24: 167–181.
- Konecny, G., Winterhoff, B., Kolarova, T., Qi, J., Manivong, K., Dering, J. *et al.* (2011) Expression of p16 and retinoblastoma determines response to CDK4/6 inhibition in ovarian cancer. *Clin Cancer Res* 17: 1591–1602.
- Kotake, Y., Cao, R., Viatour, P., Sage, J., Zhang, Y. and Xiong, Y. (2007) pRB family proteins are required for H3K27 trimethylation and Polycomb repression complexes binding to and silencing p16INK4alpha tumor suppressor gene. *Genes Dev* 21: 49–54.
- LaBaer, J., Garrett, M., Stevenson, L., Slingerland, J., Sandhu, C., Chou, H. *et al.* (1997) New functional activities for the p21 family of CDK inhibitors. *Genes Dev* 11: 847–862.
- Landis, M., Pawlyk, B., Li, T., Sicinski, P. and Hinds, P. (2006) Cyclin D1-dependent kinase activity in murine development and mammary tumorigenesis. *Cancer Cell* 9: 13–22.
- Lee, N., Eisele, K., Chionis, J., Yuan, J., Zhu, Z., Liu, C. *et al.* (2016) Mechanisms of resistance to CDK4/6 inhibition in ER+ breast cancer. *Cancer Res* 76: Abstract P3–06–01.
- Lee, Y., Dominy, J., Choi, Y., Jurczak, M., Tolliday, N., Camporez, J. *et al.* (2014) Cyclin D1-CDK4 controls glucose metabolism independently of cell cycle progression. *Nature* 510: 547–551.

- Lenihan, C., Bouchekioua-Bouzaghrou, K., Shia, A., Wilkes, E., Casado-Izquierdo, P., Cutillas, P. *et al.* (2016) Characterization of resistance to the selective CDK4/6 inhibitor palbociclib in ER positive breast cancer. *Cancer Res* 76: Abstract P3-06-02.
- Leonard, J., LaCasce, A., Smith, M., Noy, A., Chiriac, L., Rodig, S. *et al.* (2012) Selective CDK4/6 inhibition with tumor responses by PD0332991 in patients with mantle cell lymphoma. *Blood* 119: 4597-4607.
- Logan, J., Mostofizadeh, N., Desai, A., Von Euw, E., Conklin, D., Konkankit, V. *et al.* (2013) PD-0332991, a potent and selective inhibitor of cyclin-dependent kinase 4/6, demonstrates inhibition of proliferation in renal cell carcinoma at nanomolar concentrations and molecular markers predict for sensitivity. *Anticancer Res* 33: 2997-3004.
- Loibl, S., Turner, N., Ro, J., Cristofanilli, M., Iwata, H., Im, S. *et al.* (2016) Palbociclib (PAL) in combination with fulvestrant (F) in pre-/peri-menopausal (PreM) women with metastatic breast cancer (MBC) and prior progression on endocrine therapy – results from Paloma-3. *J Clin Oncol* 34: Abstract 524.
- Lundberg, A. and Weinberg, R. (1998) Functional inactivation of the retinoblastoma protein requires sequential modification by at least two distinct cyclin-CDK complexes. *Mol Cell Biol* 18: 753-761.
- Luo, F., Le, X., Jeselsohn, R., Brown, M. and Garraway, L. (2016) Estrogen-independent growth of ESR1-mutant breast cancer cells requires CDK4/6. *Cancer Res* 76: Abstract P3-05-13.
- Ma, C., Gao, F., Northfelt, D., Goetz, M., Forero, A., Naughton, M. *et al.* (2016) A phase II trial of neoadjuvant palbociclib, a cyclin-dependent kinase (CDK) 4/6 inhibitor, in combination with anastrozole for clinical stage 2 or 3 estrogen receptor positive HER2 negative (ER+HER2-) breast cancer (BC). *Cancer Res* 76: Abstract S6-05.
- Ma, Z., Wu, Y., Jin, J., Yan, J., Kuang, S., Zhou, M. *et al.* (2013) Phylogenetic analysis reveals the evolution and diversification of cyclins in eukaryotes. *Mol Phylogenet Evol* 66: 1002-1010.
- Malumbres, M. (2014) Cyclin-dependent kinases. *Genome Biol* 15: 122.
- Malumbres, M. and Barbacid, M. (2009) Cell cycle, CDKs and cancer: a changing paradigm. *Nat Rev Cancer* 9: 153-166.
- Malumbres, M., Sotillo, R., Santamaría, D., Galán, J., Cerezo, A., Ortega, S. *et al.* (2004) Mammalian cells cycle without the D-type cyclin-dependent kinases CDK4 and CDK6. *Cell* 118: 493-504.
- Matsura, I., Denissova, N., Wang, G., He, D., Long, J. and Liu, F. (2004) Cyclin-dependent kinases regulate the antiproliferative function of Smads. *Nature* 430: 226-231.
- Matter-Walstra, K., Ruhstaller, T., Klingbiel, D., Schwenkglens, M. and Dedes, K. (2016) Palbociclib as a first-line treatment in oestrogen receptor-positive, HER2-negative, advanced breast cancer not cost-effective with current pricing: a health economic analysis of the Swiss Group for Clinical Cancer Research (SAKK). *Breast Cancer Res Treat* 158: 51-57.
- McClelland, R., Barrow, D., Madden, T., Dutkowski, C., Pamment, J., Knowlden, J. *et al.* (2001) Enhanced epidermal growth factor receptor signaling in MCF7 breast cancer cells after long-term culture in the presence of the pure antiestrogen ICI 182,780 (Faslodex). *Endocrinology* 142: 2776-2788.
- McClendon, A., Dean, J., Rivadeneira, D., Yu, J., Reed, C., Gao, E. *et al.* (2012) CDK4/6 inhibition antagonizes the cytotoxic response to anthracycline therapy. *Cell Cycle* 11: 2747-2755.
- Miller, T., Balko, J., Fox, E., Ghazoui, Z., Dunbier, A., Anderson, H. *et al.* (2011) ER α -dependent E2F transcription can mediate resistance to estrogen deprivation in human breast cancer. *Cancer Discov* 1: 338-351.
- Miller, T., Hennessy, B., González-Angulo, A., Fox, E., Mills, G., Chen, H. *et al.* (2010) Hyperactivation of phosphatidylinositol-3 kinase promotes escape from hormone dependence in estrogen receptor-positive human breast cancer. *J Clin Invest* 120: 2406-2413.
- Miller, T., Pérez-Torres, M., Narasanna, A., Guix, M., Stål, O., Pérez-Tenorio, G. *et al.* (2009) Loss of Phosphatase and Tensin homologue deleted on chromosome 10 engages ErbB3 and insulin-like growth factor-I receptor signaling to promote antiestrogen resistance in breast cancer. *Cancer Res* 69: 4192-4201.
- Mukherjee, S. and Conrad, S. (2005) c-Myc suppresses p21WAF1/CIP1 expression during estrogen signaling and antiestrogen resistance in human breast cancer cells. *J Biol Chem* 280: 17617-17625.
- Musgrove, E., Caldon, C., Barraclough, J., Stone, A. and Sutherland, R. (2011) Cyclin D as a therapeutic target in cancer. *Nat Rev Cancer* 11: 558-572.
- Musgrove, E., Lilischkis, R., Cornish, A., Lee, C., Setlur, V., Seshadri, R. *et al.* (1995) Expression of the cyclin-dependent kinase inhibitors p16INK4, p15INK4B and p21WAF1/CIP1 in human breast cancer. *Int J Cancer* 63: 584-591.
- Nardone, A., De Angelis, C., Trivedi, M., Osborne, C. and Schiff, R. (2015) The changing role of ER in endocrine resistance. *Breast* 24: S60-S66.
- Nguyen, M., Buchwalter, G., Luo, F., Garraway, L., Brown, M. and Jeselsohn, R. (2016) Exploring

- bazedoxifene and palbociclib as potential therapeutic strategies for overcoming ESR1-mediated endocrine resistance. *Cancer Res* 74: Abstract P3–05–09.
- O'Brien, N., Di Tomaso, E., Ayala, R., Tong, L., Issakhanian, S., Linnartz, R. *et al.* (2014) *In vivo* efficacy of combined targeting of CDK4/6, ER and PI3K signaling in ER+ breast cancer. *Cancer Res* 74: Abstract 4756.
- O'Leary, B., Finn, R. and Turner, N. (2016) Treating cancer with selective CDK4/6 inhibitors. *Nat Rev Clin Oncol* 13: 417–430.
- Patnaik, A., Rosen, L., Tolaney, S., Tolcher, A., Goldman, J., Gandhi, L. *et al.* (2016) Efficacy and Safety of abemaciclib, an inhibitor of CDK4 and CDK6, for patients with breast cancer, non-small cell lung cancer, and other solid tumors. *Cancer Discov* 6: 740–753.
- Pérez-Tenorio, G., Berglund, F., Esguerra Merca, A., Nordenskjöld, B., Rutqvist, L., Skoog, L. *et al.* (2006) Cytoplasmic p21WAF1/CIP1 correlates with Akt activation and poor response to tamoxifen in breast cancer. *Int J Oncol* 28: 1031–1042.
- Petrossian, K., Lo, C., Yuan, Y., Kanaya, N., Wu, S., Synold, T. *et al.* (2016) Molecular characterization of CDK 4/6 inhibitors (palbociclib, abemaciclib, and ribociclib) in hormone receptor-positive (HR+) breast cancer cell lines. *Cancer Res* 74: Abstract P5–03–07.
- Polyak, K., Kato, J., Solomon, M., Sherr, C., Massague, J., Roberts, J. *et al.* (1994) p27Kip1, a cyclin-CDK inhibitor, links transforming growth factor-beta and contact inhibition to cell cycle arrest. *Genes Dev* 8: 9–22.
- Qin, G., Xu, F., Qin, T., Zheng, Q., Shi, D., Xia, W. *et al.* (2015) Palbociclib inhibits epithelial-mesenchymal transition and metastasis in breast cancer via c-Jun/COX-2 signaling pathway. *Oncotarget* 6: 41794–41808.
- Reddy, H., Mettus, R., Rane, S., Graña, X., Litvin, J. and Reddy, E. (2005) Cyclin-dependent kinase 4 expression is essential for neu-induced breast tumorigenesis. *Cancer Res* 65: 10174–10178.
- Ren, S. and Rollins, B. (2004) Cyclin C/CDK3 promotes Rb-dependent G0 exit. *Cell* 117: 239–251.
- Reutens, A., Fu, M., Wang, C., Albanese, C., McPhaul, M., Sun, Z. *et al.* (2001) Cyclin D1 binds the androgen receptor and regulates hormone-dependent signaling in a p300/CBP-associated factor (P/CAF)-dependent manner. *Mol Endocrinol* 15: 797–811.
- Reynisdóttir, I., Polyak, K., Iavarone, A. and Massagué, J. (1995) Kip/Cip and INK4 CDK inhibitors cooperate to induce cell cycle arrest in response to TGF-beta. *Genes Dev* 9: 1831–1845.
- Rivadeneira, D., Mayhew, C., Thangavel, C., Sotillo, E., Reed, C., Graña, X. *et al.* (2010) Proliferative suppression by CDK4/6 inhibition: complex function of the retinoblastoma pathway in liver tissue and hepatoma cells. *Gastroenterology* 138: 1920–1930.
- Roberts, P., Bisi, J., Strum, J., Combest, A., Darr, D., Usary, J. *et al.* (2012) Multiple roles of cyclin-dependent kinase 4/6 inhibitors in cancer therapy. *J Natl Cancer Inst* 104: 476–487.
- Ruas, M. and Peters, G. (1998) The p16INK4a/CDKN2A tumor suppressor and its relatives. *Biochim Biophys Acta* 1378: F115–F177.
- Sadasivam, S. and DeCaprio, J. (2013) The DREAM complex: master coordinator of cell cycle-dependent gene expression. *Nat Rev Cancer* 13: 585–595.
- Sage, J. (2012) The retinoblastoma tumor suppressor and stem cell biology. *Genes Dev* 26: 1409–1420.
- Santamaria, D., Barrière, C., Cerqueira, A., Hunt, S., Tardy, C., Newton, K. *et al.* (2007) CDK1 is sufficient to drive the mammalian cell cycle. *Nature* 448: 811–815.
- Schwartz, G., Lo Russo, P., Dickson, M., Randolph, S., Shaik, M., Wilner, K. *et al.* (2011) Phase I study of PD 0332991, a cyclin-dependent kinase inhibitor, administered in 3-week cycles (schedule 2/1). *Br J Cancer* 104: 1862–1868.
- Shao, Z. and Robbins, P. (1995) Differential regulation of E2F and Sp1-mediated transcription by G1 cyclins. *Oncogene* 10: 221–228.
- Sharpless, N. and Sherr, C. (2015) Forging a signature of in vivo senescence. *Nat Rev Cancer* 15: 397–408.
- Shen, T. and Huang, S. (2012) The role of Cdc25A in the regulation of cell proliferation and apoptosis. *Anticancer Agents Med Chem* 12: 631–639.
- Sherr, C. and Roberts, J. (1999) CDK inhibitors: positive and negative regulators of G1-phase progression. *Genes Dev* 13: 1501–1512.
- Slamon, D., Hurvitz, S., Applebaum, S., Glaspy, J., Allison, M., DiCarlo, B. *et al.* (2010) Phase I study of PD 0332991, cyclin-D kinase (CDK) 4/6 inhibitor in combination with letrozole for first-line treatment of patients with ER-positive, HER2-negative breast cancer. *J Clin Oncol* 28: Abstract 3060.
- Slingerland, J. and Pagano, M. (2000) Regulation of the CDK inhibitor p27 and its deregulation in cancer. *J Cell Physiol* 183: 10–17.
- Spencer, S., Cappell, S., Tsai, F., Overton, K., Wang, C. and Meyer, T. (2013) The proliferation-quiescence decision is controlled by a bifurcation in CDK2 activity at mitotic exit. *Cell* 155: 369–383.

- Stendahl, M., Kronblad, A., Rydén, L., Emdin, S., Bengtsson, N. and Landberg, G. (2004) Cyclin D1 overexpression is a negative predictive factor for tamoxifen response in postmenopausal breast cancer patients. *Br J Cancer* 90: 1942–1948.
- Subhawong, A., Subhawong, T., Nassar, H., Kouprina, N., Begum, S., Vang, R. *et al.* (2009) Most basal-like breast carcinomas demonstrate the same Rb-/p16+ immunophenotype as the HPV-related poorly differentiated squamous cell carcinomas which they resemble morphologically. *Am J Surg Pathol* 33: 163–175.
- Sun, W. and Wang, D. (2014) A population pharmacokinetic (pk) analysis of palbociclib (pd-0332991) in patients (pts) with advanced solid tumors. *Ann Oncol* 25: Abstract 462P.
- Thangavel, C., Dean, J., Ertel, A., Knudsen, K., Aldaz, C., Witkiewicz, A. *et al.* (2011) Therapeutically activating RB: reestablishing cell cycle control in endocrine therapy-resistant breast cancer. *Endocr Relat Cancer* 18: 333–345.
- Trapé, A., Liu, S., Cortes, A., Ueno, N. and Gonzalez-Angulo, A. (2016) Effects of CDK4/6 Inhibition in hormone receptor-positive/human epidermal growth factor receptor 2-negative breast cancer cells with acquired resistance to paclitaxel. *J Cancer* 7: 947–956.
- Treré, D., Brighenti, E., Donati, G., Ceccarelli, C., Santini, D., Taffurelli, M. *et al.* (2009) High prevalence of retinoblastoma protein loss in triple-negative breast cancers and its association with a good prognosis in patients treated with adjuvant chemotherapy. *Ann Oncol* 20: 1818–1823.
- Turner, N., Ro, J., André, F., Loi, S., Verma, S., Iwata, H. *et al.* (2015) PALOMA3 Study Group. Palbociclib in Hormone-Receptor-Positive Advanced Breast Cancer. *N Engl J Med* 373: 209–219.
- US Food and Drug Administration. (2015) *Palbociclib*. Available at: <http://www.fda.gov/Drugs/InformationOnDrugs/ApprovedDrugs/ucm432886.htm>
- US Food and Drug Administration. (2016) *Palbociclib*. Available at: <http://www.fda.gov/Drugs/InformationOnDrugs/ApprovedDrugs/ucm487080.htm>
- Verma, S., Bartlett, C., Schnell, P., DeMichele, A., Loi, S., Ro, J. *et al.* (2016) Palbociclib in combination with fulvestrant in women with hormone receptor-positive/her2-negative advanced metastatic breast cancer: detailed safety analysis from a multicenter, randomized, placebo-controlled, phase III study (PALOMA-3). *Oncologist* 21: 1–11.
- Vora, S., Juric, D., Kim, N., Mino-Kenudson, M., Huynh, T., Costa, C. *et al.* (2014) CDK 4/6 inhibitors sensitize PIK3CA mutant breast cancer to PI3K inhibitors. *Cancer Cell* 26: 136–149.
- Wang, L., Wang, J., Blaser, B., Duchemin, A., Kusewitt, D., Liu, T. *et al.* (2007) Pharmacologic inhibition of CDK4/6: mechanistic evidence for selective activity or acquired resistance in acute myeloid leukemia. *Blood* 110: 2075–2083.
- Wardell, S., Ellis, M., Alley, H., Eisele, K., VanArsdale, T., Dann, S. *et al.* (2015) Efficacy of SERD/SERM Hybrid-CDK4/6 inhibitor combinations in models of endocrine therapy-resistant breast cancer. *Clin Cancer Res* 21: 5121–5130.
- Weinberg, R. (1995) The retinoblastoma protein and cell cycle control. *Cell* 81: 323–330.
- Wiedemeyer, W., Dunn, I., Quayle, S., Zhang, J., Chheda, M., Dunn, G. *et al.* (2010) Pattern of retinoblastoma pathway inactivation dictates response to CDK4/6 inhibition in GBM. *Proc Natl Acad Sci U S A* 107: 11501–11506.
- Witkiewicz, A. and Knudsen, E. (2014) Retinoblastoma tumor suppressor pathway in breast cancer: prognosis, precision medicine, and therapeutic interventions. *Breast Cancer Res* 16: 207.
- Xiong, Y., Zhang, H. and Beach, D. (1992) D type cyclins associate with multiple protein kinases and the DNA replication and repair factor PCNA. *Cell* 71: 505–514.
- Yadav, V., Burke, T., Huber, L., Van Horn, R., Zhang, Y., Buchanan, S. *et al.* (2014) The CDK4/6 inhibitor LY2835219 overcomes vemurafenib resistance resulting from MAPK reactivation and cyclin D1 upregulation. *Mol Cancer Ther* 13: 2253–2263.
- Yasui, M., Yamamoto, H., Ngan, C., Damdinsuren, B., Sugita, Y., Fukunaga, H. *et al.* (2006) Antisense to cyclin D1 inhibits vascular endothelial growth factor-stimulated growth of vascular endothelial cells: implication of tumor vascularization. *Clin Cancer Res* 12: 4720–4729.
- Yoon, S., Shin, S. and Mercurio, A. (2006) Ras stimulation of E2F activity and a consequent E2F regulation of integrin alpha6beta4 promote the invasion of breast carcinoma cells. *Cancer Res* 66: 6288–6295.
- Young, R., Waldeck, K., Martin, C., Foo, J., Cameron, D., Kirby, L. *et al.* (2014) Loss of CDKN2A expression is a frequent event in primary invasive melanoma and correlates with sensitivity to the CDK4/6 inhibitor PD0332991 in melanoma cell lines. *Pigment Cell Melanoma Res* 27: 590–600.
- Yu, Q., Geng, Y. and Sicinski, P. (2001) Specific protection against breast cancers by cyclin D1 ablation. *Nature* 411: 1017–1021.

Yu, Q., Sicinska, E., Geng, Y., Ahnström, M., Zagozdzon, A., Kong, Y. *et al.* (2006) Requirement for CDK4 kinase function in breast cancer. *Cancer Cell* 9: 23–32.

Yu, Y., Loi, C., Hoffman, J. and Wang, D. (2016) Physiologically-based pharmacokinetic (PBPK) modeling of palbociclib. *J Clin Pharmacol.* doi:10.1002/jcph.792 [Epub ahead of print].


Zhang, Q., Sakamoto, K., Liu, C., Triplett, A., Lin, W., Rui, H. *et al.* (2011) Cyclin D3 compensates for the loss of cyclin D1 during ErbB2-induced mammary

tumor initiation and progression. *Cancer Res* 71: 7513–7524.

Ziemke, E., Dosch, J., Maust, J., Shettigar, A., Sen, A., Welling, T. *et al.* (2016) Sensitivity of KRAS-mutant colorectal cancers to combination therapy that cotargets MEK and CDK4/6. *Clin Cancer Res* 22: 405–414.

Zwijsen, R., Wientjens, E., Klompmaker, R., van der Sman, J., Bernards, R. and Michalides, R. (1997) CDK-independent activation of estrogen receptor by cyclin D1. *Cell* 88: 405–415.

Visit SAGE journals online
<http://tam.sagepub.com>

 SAGE journals