

Opportunistic predation of the Roadside Hawk (*Rupornis magnirostris*) on alates subterranean termites (*Coptotermes testaceus*) in southwestern Brazilian Amazon

Depredación oportunista del halcón caminero (*Rupornis magnirostris*) sobre termitas subterráneas aladas (*Coptotermes testaceus*) en el suroeste de la Amazonía brasileña

Abstract

Raul Afonso Pommer Barbosa¹
<https://orcid.org/0000-0002-1914-5184>
raulpommer@hotmail.com

Jessica Fernanda Teodoro Reis²
<https://orcid.org/0000-0001-9910-7374>
jessicaf19@outlook.com

Marcela Alvares Oliveira^{* 3}
<https://orcid.org/0000-0002-4129-993X>
marcela.mugrabe@gmail.com

*Corresponding author

1 Clube de Astronomia e Ciências de Rondônia, Avenida Presidente Dutra, 2965, Centro, Porto Velho-RO, 76801-059, Brazil.

2 Centro Universitário São Lucas (UNISL), Porto Velho, Brazil.

3 Centro Universitário Aparício Carvalho (FIMCA), Curso de Ciências Biológicas. Programa de Pós Graduação em Biodiversidade e Biotecnologia pela Universidade Federal de Rondônia, Porto Velho, Brazil.

Citación

Barbosa RAP, Reis JFT, Oliveira MA. 2022. Opportunistic predation of the Roadside Hawk (*Rupornis magnirostris*) on the alates subterranean termites (*Coptotermes testaceus*) in southwestern Brazilian Amazon. Revista peruana de biología 29(2): e21528 001-004 (Mayo 2022). doi: <http://dx.doi.org/10.15381/rpb.v29i2.21528>

Presentado: 05/11/2022
Aceptado: 28/02/2022
Publicado online: 30/05/2022

Editor: Cesar Arana

We report the opportunistic predation of Roadside Hawk (*Rupornis magnirostris* - Accipitridae) on subterranean termite (*Coptotermes testaceus* - Rhinotermitidae). This predatory behaviour was observed during an avifauna survey in the municipality of Candeias do Jamari, Rondônia, Brazil, in the southwestern Brazilian Amazon. There was a great abundance of *C. testaceus* alates because of heavy rainfall. This is the season when *C. testaceus* colonies produce alates in large numbers to found new colonies. The Roadside Hawk stayed on its perch and captured winged termites with little energy expenditure. The bird in question appeared to take advantage of the termite mating flight to feed opportunistically on the abundant alates. Although feeding on insects by *R. magnirostris* is well known, this is the first documentation of feeding on termite alates.

Resumen

Informamos de la depredación oportunista de *Coptotermes testaceus* (Rhinotermitidae) por *Rupornis magnirostris* (Accipitridae). Este comportamiento depredador fue observado durante un estudio de la avifauna en el municipio de Candeias do Jamari, Rondônia, Brasil, en el suroeste de la Amazonía brasileña. Había una gran abundancia de formas aladas de *C. testaceus* debido a las fuertes lluvias. Es el momento en que las colonias de *C. testaceus* producen formas aladas en gran número para fundar nuevas colonias. El aguilucho permaneció en su perchal y capturó termitas aladas con poco gasto de energía. El ave en cuestión parecía aprovechar el vuelo de apareamiento de las termitas para alimentarse de forma oportunista de las abundantes termitas aladas durante los vuelos de apareamiento. Aunque la alimentación de insectos por parte de *R. magnirostris* es bien conocida, ésta es la primera documentación de alimentación de termitas aladas.

Keywords:

Roadside Hawk; Feeding; Termite; Amazon; Rondônia.

Palabras clave:

Taguató Comum; Alimentación; termitas; Amazonia; Rondônia.

Introduction

The Roadside Hawk, *Rupornis magnirostris* (Gmelin 1788) (Accipitriformes: Accipitridae), occurs from Mexico to Argentina, including Brazil (Santos & Rosado 2009). It is separated into 12 subspecies differing mainly in plumage coloration throughout its range. These birds are diurnal and inhabit a wide variety of environments, including urbanized areas, riverbanks, lakes shores, open fields, forest edges and scrublands, being found up to 3000 m altitude in the Andes. These birds can weigh between 200 and 400 g with males being sma-

ller than females (Sick 1997, Tortato 2009, Barros et al. 2010). The Roadside Hawk lives in pairs and builds nests in the tops of large trees (Sick 1997). In recent decades, this hawk has become more common in urban centers, successfully adapting to this environment. In cities, the supply of prey is greater and natural predators (larger birds of prey) are scarce (Sick 2001, Athiê & Dias 2010, Silva & Machado 2015).

The Roadside Hawk is a generalist, opportunistic predator. Common prey items include small snakes, lizards, large insects, domestic birds and occasionally bats (Santos & Rosado 2009, Sick 1997, Montani et al. 2016, Lima et al. 2020). Tortato (2009) described the predation of Water Opossum, *Chironectes minimus* (Didelphidae), a small mammal widely found in the Neotropics. The objective of this study was to present a record of the consumption of *Coptotermes testaceus* (Rhinotermitidae) by *Rupornis magnirostris* (Accipitridae).

Material and methods

We conducted field observations in the rural area of the municipality of Candeias do Jamari, Rondônia state, Brazil, in the southwestern Brazilian Amazon. The locality comprises a cattle ranch with a pasture area with several

isolated Brazil nut trees *Bertholletia excelsa* (Humboldt & Bonpland) (Lecythidaceae). Photographic register was realized with a Nikon P900 camera, lens 24-2000 mm.

Results and discussion

During a general avifauna survey, predation of *Coptotermes testaceus* (L.) (Blattodea: Rhinotermitidae) alates by an individual *R. magnirostris* was observed at coordinates $8^{\circ}44'15.6''$ S, $63^{\circ}28'19.7''$ W, at 08:13 AM on December 6, 2020. The location was along a rural road, in a deforested area. In a Brazil nut tree, *Bertholletia excelsa*, we noted the opportunistic feeding of an individual *R. magnirostris* on abundant, swarming alate termites (Fig. 1a, b). The observation was carried about 50 m of distance approximately. The bird remained there consuming the insects for about 25 minutes. After this period, even though there were still individuals of *C. testaceus*, *R. magnirostris* left the site. Samples of the winged alate termites were captured and returned to the laboratory. They were determined to be *C. testaceus* by the wing venation (Scheffrahn et al. 2015). This observation was made during a heavy rainfall. There was a great abundance of *C. testaceus* alates (Fig. 1c), as under these conditions colonies of this species produce swarms of reproductive for breeding and formation of new colonies.




Figure 1. Photographic evidence of opportunistic predation; a. *Rupornis magnirostris*; b. *Rupornis magnirostris* feeding on *Coptotermes testaceus* alates; c. *Coptotermes testaceus*; d. *Coptotermes testaceus* wing under the dissecting microscope (Magnification 10x) for species identification.

Coptotermes testaceus is a xylophagous Neotropical subterranean termite that can be found in both urban and rural environments. It prefers high relative humidity (near 100%) and medium temperatures (20 °C) (Chouvenç et al. 2016, Pozo-Santiago et al. 2020). The swarming of colonies of the family Rhinotermitidae occurs mainly at the end of the dry season and beginning of the rainy season (Medeiros et al. 1999). This swarming is characterized by the appearance of large numbers of winged reproductive and the founding of new colonies (Ferreira et al. 2011). Termites in general are part of the diet of different species of birds (Faria 2007, Fernandes et al. 2007, Vasconcelos et al. 2015), with some species being specialist on certain genera (Vries et al. 2011).

The study by Vasconcelos et al. (2015) on termite consumption by Neotropical birds indicated that only 1.8% of the recorded termites were identified to the species level. There was no mention of termites in the diet of *R. magnirostris*. Records of the diet composition of *R. magnirostris* have been mainly documented through opportunistic records (Botero-Delgadillo & Escudero-Páez 2012, Zocche et al. 2018, Lima et al. 2020). In the case of insect consumption records, prey identification has not been possible (Baladrón et al. 2011). Termites are considered agricultural pests (Constantino 2002, Rouland-Lefèvre 2010, Costa et al. 2020) and the use of pesticides for their control can compromise the occurrence and maintenance of this food resource. These factors highlight the importance of this work for understanding the feeding habits of *R. magnirostris* and the potential importance of *C. testaceus* and other termite alates in bird diets.

Rupornis magnirostris uses the sit-and-wait technique as its hunting strategy. It typically settles on high perches waiting for its prey (Baladrón et al. 2011). This technique is highly opportunistic (Santos & Rosado 2009). According to Panasci & Whitacre (2000) its main hunting method consists of leaving the perch and to meet the stopped target. It rarely captures birds in flight. In the present report, the bird stayed on the perch and captured the insects with the least energy expenditure possible. This opportunistic feeding was possible due to the momentary availability of large numbers of termite alates in flight. This report documents the opportunistic and generalist feeding behavior of *R. magnirostris*. The bird took advantage of the presence of a *C. testaceus* reproductive swarm to feed. This is the first confirmed record of *C. testaceus* feeding by this bird.

Literature cited

- Athiê S, Dias MM. 2010. Avian community composition in an urban park in central São Paulo state, southeastern Brazil. Revista Brasileira de Zoociências 12(1): 67-73.
- Barros FMD, Granzinelli MAM, Pereira RJG, Motta-Junior JC. 2010. Home range and habitat use by the roadside hawk, *Rupornis magnirostris* (Gmelin, 1788) (Aves: Falconiformes) in southeastern Brazil. Journal of Natural History 45(1-2): 65-75. <https://doi.org/10.1080/00222933.2010.520756>
- Baladrón AV, Bó MS, Malizia AI, Bechard MJ. 2011. Food habits of the Roadside Hawk (*Buteo magnirostris*) during the nonbreeding season in the southeastern Pampas of Argentina. Journal of Raptor Research 45(3): 257-261. <https://doi.org/10.3356/JRR-10-108.1>
- Baladrón AV, Cavalli M, Pretelli MG, Bó MS. 2016. Time-activity budgets and hunting behavior of the Roadside Hawk (*Rupornis magnirostris*) and the Long-winged Harrier (*Circus buffoni*). Revista Brasileira de Ornitologia 24(3): 197-203.
- Botero-Delgadillo E, Escudero-Páez S. 2012. Roadside Hawk *Rupornis magnirostris* predating a Bogota Rail *Rallus semiplumbeus*, an endemic and endangered species of Colombia. Cotinga 34: 94-95.
- Constantino R. 2002. The pest termites of South America: taxonomy, distribution and status. Journal of Applied Entomology 126(7-8): 355-365. <https://doi.org/10.1046/j.1439-0418.2002.00670.x>
- Costa JG, Santos ICDL, Peres Filho O, Zanetti R, Santos AD. 2020. Characterization of injury caused by *Coptotermes testaceus* in *Tectona grandis* plantations in Brazil. Cerne 26(3): 310-314. <https://doi.org/10.1590/010477602026032739>
- Chouvenç T, Li HF, Austin J, Bordereau C, Bourguignon T, Cameron SL, Cancello EM, Constantino R, Costa-Leonardo AM, Eggleton P, et al. 2016. Revisiting *Coptotermes* (Isoptera: Rhinotermitidae): a global taxonomic road map for species validity and distribution of an economically important subterranean termite genus. Systematic Entomology 41: 299-306. <https://doi.org/10.1111/syen.12157>
- Faria IP. 2007. Peach-fronted Parakeet (*Aratinga aurea*) feeding on arboreal termites in the Brazilian Cerrado. Revista Brasileira de Ornitologia 15(3): 457-458.
- Fernandes FR, Cruz LD, Rodrigues AAF. 2007. Diet of the Gray-Breasted Martin (Hirundinidae: *Progne chalybea*) in a wintering area in Maranhão, Brazil. Revista Brasileira de Ornitologia 15(3): 445-447.
- Ferreira EVDO, Martins V, Inda Junior AV, Giasson E, Nascimento PCD. 2011. Ação dos térmitas no solo. Ciência Rural 41: 804-811. <https://doi.org/10.1590/S0103-84782011005000044>
- Lima NDS, Napiwoski SJ, Oliveira MA. 2020. Human-wildlife conflict in the southwestern amazon: Poaching and its motivations. Nature Conservation Research 5(1): 109-114. <https://dx.doi.org/10.24189/ncr.2020.006>
- Medeiros GSL, Bandeira AG, Martius C. 1999. Termite swarming in the northeastern Atlantic rain forest of Brazil. Studies on Neotropical Fauna and Environment 34(2): 76-87. <https://doi.org/10.1076/snfe.34.2.76.2103>
- Montani ME, Saigo G, Romano M. 2016. Predación del murciélagos moloso común (*Tadarida brasiliensis*) por el taguató común (*Rupornis magnirostris*) en la ciudad de Rosario (Santa Fe, Argentina). Natura Neotropicális, 47(1): 51-56. <https://doi.org/10.14409/natura.v1i47.5987>
- Panasci T, Whitacre D. 2000. Diet and foraging behavior of nesting Roadside Hawks in Petén, Guatemala. The Wilson Journal of Ornithology 112(4): 555-558.
- Pozo-Santiago CO, La Cruz MP, La Cruz MT, Cruz-Pérez A, Cabello-García S, Hernández-Gallegos MA, Velázquez-Martínez JR. 2020. Survival of *Coptotermes testaceus* (Isoptera: Rhinotermitidae) to environmental conditions (relative humidity and temperature) and Preference to different substrates. Sociobiology 67(3): 425-432. <https://doi.org/10.13102/sociobiology.v67i3.5141>

- Rouland-Lefèvre C. 2010. Termites as pests of agriculture. In Biology of termites: a modern synthesis. Dordrecht: Springer; p. 499-517. https://doi.org/10.1007/978-90-481-3977-4_18
- Santos WM, Rosado FR. 2009. Dados preliminares da biologia do Gavião-Carijó (*Rupornis magnirostris*, Gmelin, 1788) na região noroeste do Paraná. Revista em Agro-negócio e Meio Ambiente 2(3): 421-430. <https://doi.org/10.17765/2176-9168.2009v2n3p421-430>
- Sick H. 1997. Ornitologia brasileira: uma introdução. Rio de Janeiro: Editora Nova Fronteira.
- Sick H. 2001. Ornitologia brasileira. 3nd edn. Rio de Janeiro: Editora Nova Fronteira.
- Silva JC, Machado CA. 2016. Desmatamento e adaptações de aves de rapina na área urbana de Araguaína (TO). Revista Tocantinense de Geografia 5(7): 120-141.
- Scheffrahn RH, Carrijo TF, Křeček J, Su NY, Szalanski AL, Austin JW, Chase JA, Mangold J. R. 2015. A single endemic and three exotic species of the termite genus *Coptotermes* (Isoptera, Rhinotermitidae) in the New World. Arthropod systematics & phylogeny 73(2): 333-348.
- Tortato MA. 2009. Predação de cuíca-d'água (*Chironectes minimus*: Mammalia, Didelphidae) por gaviao-carijó (*Rupornis magnirostris*: Aves, Accipitridae). Mastozoología neotropical 16(2): 491-493.
- Vasconcelos MF, Hoffmann D, Araújo MC, Vasconcelos PN. 2015. Bird-termite interactions in Brazil: a review with perspectives for future studies. Biota Neotropica 15: e20140035. <https://doi.org/10.1590/1676-06032014003514>
- Vries JL, Pirk CWW, Bateman PW, Cameron EZ, Dalerum F. 2011. Extension of the diet of an extreme foraging specialist, the aardwolf (*Proteles cristata*). African Zoology 46(1): 194-196. <https://doi.org/10.1080/15627020.2011.11407494>
- Zocche JJ, Ceron K, Gonzaga GS, Plucenio GP, Carvalho F. 2018. Predation on *Chironius bicarinatus* (Wied, 1820) (Serpentes: Colubridae) by Roadside Hawk *Rupornis magnirostris* (Gmelin, 1788) (Aves: Accipitridae). Herpetology Notes 11: 129-131.

Agradecimientos / Acknowledgments:

We thank the Clube de Astronomia e Ciências de Rondônia and the "Projeto Sagui-de-Rondônia" for the financial support. We thank the Centro de Cultura e Formação Kanindé for the logistical support. We thank the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior- CAPES for the financial support.

Conflictos de intereses / Competing interests:

The authors declare no conflict of interest.

Rol de los autores / Authors Roles:

RAPM: conceptualization, formal analysis, writing: preparation of the original project, writing: revision and editing.

JFTR: conceptualization, formal analysis, writing: preparation of the original project, writing: revision and editing.

MAO: writing: preparation of the original project, writing: revision and editing.

Fuentes de financiamiento / Funding:

This work was funded by Clube de Astronomia e Ciências de Rondônia, "Projeto Sagui-de-Rondônia" and the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior – CAPES.

Aspectos éticos / legales; Ethics / legal:

Authors declare that they did not violate or omit ethical or legal norms in this research.