



OPEN

Toxicity thresholds of nine herbicides to coral symbionts (Symbiodiniaceae)

Magena Marzoni^{1,2}, Florita Flores^{1,2✉}, Nora Sadoun¹, Marie C. Thomas¹, Anais Valada-Mennuni¹, Sarit Kaserzon³, Jochen F. Mueller³ & Andrew P. Negri^{1,2}

Over 30 herbicides have been detected in catchments and waters of the Great Barrier Reef (GBR) and their toxicity to key tropical species, including the coral endosymbiotic algae Symbiodiniaceae, is not generally considered in current water quality guideline values (WQGVs). Mutualistic symbionts of the family Symbiodiniaceae are essential for the survival of scleractinian corals. We tested the effects of nine GBR-relevant herbicides on photosynthetic efficiency ($\Delta F/F_m'$) and specific growth rate (SGR) over 14 days of cultured coral endosymbiont *Cladocopium goreau* (formerly *Symbiodinium* clade C1). All seven Photosystem II (PSII) herbicides tested inhibited $\Delta F/F_m'$ and SGR, with toxicity thresholds for SGR ranging between 2.75 and 320 $\mu\text{g L}^{-1}$ (no effect concentration) and 2.54–257 $\mu\text{g L}^{-1}$ (EC_{10}). There was a strong correlation between EC_{50} s for $\Delta F/F_m'$ and SGR for all PSII herbicides indicating that inhibition of $\Delta F/F_m'$ can be considered a biologically relevant toxicity endpoint for PSII herbicides to this species. The non-PSII herbicides haloxyfop and imazapic did not affect $\Delta F/F_m'$ or SGR at the highest concentrations tested. The inclusion of this toxicity data for Symbiodiniaceae will contribute to improving WQGVs to adequately inform risk assessments and the management of herbicides in tropical marine ecosystems.

Pesticides in tropical marine ecosystems. Pesticide contamination from coastal agriculture affects tropical and sub-tropical nearshore marine environments globally^{1–5}. In Australia, large-scale monitoring and reporting programs for the Great Barrier Reef (GBR) aim to inform the management of pesticide loads that enter the GBR and its catchments⁶. There have been at least 55 pesticides detected in the GBR catchment area (GBRCA)⁷ and 99.8% of water samples retrieved from 2011 to 2015 contained detectable concentrations of pesticides and pesticide mixtures⁸. Herbicides are the most frequently detected pesticides (>30)⁷ and reach peak loads and concentrations during the summer wet season^{9,10}. Due to their persistence, some herbicides are detected year-round, resulting in chronic exposure to freshwater and marine organisms¹¹. The five most frequently detected herbicides in the GBRCA (termed priority herbicides) are the Photosystem II (PSII) inhibitors diuron, ametryn, atrazine, tebuthiuron, and hexazinone^{12,13}. This herbicide class acts by binding to the D1 protein in PSII, causing oxidative damage, reducing photosynthetic capacity and leading to the chronic impairment of cellular function¹⁴. Regulations have been implemented to reduce the environmental concentrations of these priority herbicides, leading to increased application of 'alternative' PSII and non-PSII herbicides in coastal agriculture^{15,16}. While alternative herbicides are detected frequently in the environment, there are still limited data available on their transport, persistence and toxicity to non-target species¹⁷.

Herbicide impacts on tropical marine organisms. PSII herbicides can negatively impact a wide range of tropical marine phototrophs including seagrasses^{18,19}, crustose coralline algae²⁰, *Halimeda*²¹, foraminifera²² and marine microalgae^{23,24}. Scleractinian corals are foundational invertebrates that provide key habitat and structure for tropical reefs and residing organisms. Corals are susceptible to herbicides as they rely on endosymbiotic dinoflagellates, Symbiodiniaceae (that reside in the coral host tissue), to provide up to 90% of their nutritional requirements through photosynthetic pathways²⁵. The blockage of electron transport in endosymbiotic algae by PSII herbicide exposure results in some of the energy that would normally drive electron transport

¹Australian Institute of Marine Science, PMB No. 3, Townsville, QLD 4810, Australia. ²AIMS@JCU: Australian Institute of Marine Science and College of Marine and Environmental Sciences, James Cook University, Townsville, QLD 4811, Australia. ³Queensland Alliance for Environmental Health Sciences (QAEHS), The University of Queensland, Woolloongabba, QLD 4102, Australia. ✉email: f.flores@aims.gov.au

for primary production instead being emitted as fluorescence^{26,27}. Consequently, pulse amplitude modulation (PAM) fluorometry has been used in several studies as a highly sensitive and non-destructive technique to quantify sub-lethal effects of PSII herbicides on photosynthetic efficiency (as effective quantum yield: $\Delta F/F_m'$) in light adapted corals²⁸. The inhibition of $\Delta F/F_m'$ by PSII herbicides is reversible^{27,28}, but high concentrations or chronic exposures can lead to the breakdown of symbiosis in corals (bleaching)^{27,29,30} and decreased reproductive output and mortality²⁹.

While there has been much focus on the effects of herbicides on Symbiodiniaceae within their coral hosts, these dinoflagellates also have a free-living phase (0.1% of global phytoplankton reads in the open ocean³¹), which act as a critical reservoir for uptake into newly settled aposymbiotic coral recruits and corals recovering from thermal bleaching events^{32,33}. Therefore, effects of herbicide contamination on the health or abundance of free-living Symbiodiniaceae could have important flow-on effects to the establishment of symbiotic associations which are critical for coral health. Several studies have demonstrated inhibition of $\Delta F/F_m'$ by PSII herbicides in freshly isolated or cultured Symbiodiniaceae and that sensitivity differs between species and increases with temperature^{34–36}. Impairment of $\Delta F/F_m'$ by PSII herbicides is highly correlated with reduced growth in several tropical marine microalgae^{23,37,38}; however, it is unclear whether inhibition of $\Delta F/F_m'$ translates to similar whole organism effects in Symbiodiniaceae.

Improving the applicability of water quality guideline values for herbicides in tropical marine ecosystems.

Reliable water quality guideline values (WQGVs) are required to assess the potential risks posed by herbicides and mixtures of herbicides, frequently detected in tropical marine waters including the GBR^{8,12}. Conventionally, concentrations of individual herbicides detected in the field are compared against the relevant WQGV for that herbicide. For example, time averaged concentrations of up to 0.778 $\mu\text{g L}^{-1}$ diuron, obtained over a month-long passive sampler deployment, have been detected in the latest GBR monitoring program¹², a value that exceeds the proposed PC99 WQGV for diuron (0.43 $\mu\text{g L}^{-1}$; protection concentration for 99% of species)³⁹. Importantly, measured concentrations of herbicide mixtures can be assessed against a risk metric which predicts the proportion of species potentially affected by that mixture. The metric applied in GBR herbicide monitoring is the multi-substance Potentially Affected Fraction (ms-PAF)⁴⁰, and more exceedances of WQGVs are reported using this approach which accounts for all herbicides present in the environment¹². The ms-PAF method requires high quality WQGVs for all herbicides detected in the environment; however, several alternative herbicides do not have reliable WQGVs and their contribution to risk can be difficult to assess. The Australian and New Zealand guidelines currently include WQGVs for nine herbicides: the five priority PSII herbicides and four alternative herbicides (2,4-D, bromacil, MCPA, and simazine), but many of these are of low reliability due to lack of toxicity data⁴¹. Their relevance to tropical marine habitats is also uncertain as the majority of these WQGVs have been derived from toxicity data from freshwater temperate species^{23,41}. Marine proposed guideline values (PGVs) have recently been proposed for 20 herbicides: 2,4-D, ametryn, bromacil, diuron, fluometuron, fluroxypyr, haloxyfop, hexazinone, imazapic, isoxaflutole, metribuzin, MCPA, simazine, tebuthiuron, pendimethalin, prometryn, propazine, terbutryn, triclopyr, terbuthylazine^{39,42,43} but the majority are of low to moderate reliability due to lack of suitable marine toxicity data.

More toxicity data are required to improve WQGVs for alternative herbicides detected in tropical marine environments, especially for species of critical ecological value such as corals and their symbionts. Here, we developed a 14-day chronic growth inhibition test for cultures of the free-living coral symbiont *Cladocopium goreaui* to determine the toxicity thresholds for nine herbicides detected in the GBR and the GBRCA. *C. goreaui* was chosen as a suitable test species as it is common in tropical benthic environments and has a relatively rapid growth rate in comparison to many other Symbiodiniaceae^{35,44}. The PSII herbicides diuron, bromacil, hexazinone, metribuzin, propazine, simazine and tebuthiuron, as well as the non-PSII herbicides haloxyfop and imazapic were chosen in consultation with the Water Quality and Investigation Team at the Queensland Department of Environment and Science and indicate current toxicity data gaps for the improvement of WQGVs. The effects on specific growth rate (SGR) and $\Delta F/F_m'$ were investigated^{23,37}. The chronic SGR toxicity thresholds (10% effect concentrations (EC₁₀) and no effect concentrations (NEC)) represent ecologically relevant endpoints required for inclusion in deriving WQGVs⁴⁵. Correlations between $\Delta F/F_m'$ and SGR effect concentrations (EC₅₀) would further validate the application of PAM fluorometry as a sensitive tool to measure sub-lethal stress in this ecologically important coral symbiont.

Results

Physico-chemical test conditions. Physico-chemical parameters were consistent for each test: dissolved oxygen ($7.8 \pm 0.3 \text{ mg L}^{-1}$), pH (7.8 ± 0.5) and salinity ($32.5 \pm 0.7 \text{ psu}$) (mean \pm SD, $n = 152$), temperature ($27 \pm 0.6 \text{ }^\circ\text{C}$) and light intensity ($71 \pm 8 \text{ } \mu\text{mol photons m}^{-2} \text{ s}^{-1}$) (mean \pm SD, $n = 7$). All data can be found in Table S2.

Bioassay performance. *Cladocopium goreaui* exhibited consistent exponential growth in control treatments ranging from 0.0877 to 0.163 SGR day⁻¹ among experiments (which were all conducted on separate occasions) (Table 1). There was no effect of carrier solvent (DMSO and ethanol) on SGR (t-test: $F_{\text{IMK-DMSO}}$ (1, 3) = 1.185; $p = 0.356$ and $F_{\text{IMK-EtOH}}$ (1, 4) = 0.529; $p = 0.507$). The effective quantum yield $\Delta F/F_m'$ was also consistent across control treatments ($\Delta F/F_m' = 0.35 \pm 0.04$). The reference toxicant diuron ($6 \text{ } \mu\text{g L}^{-1}$) applied in each toxicity test inhibited SGR by $64.9\% \pm 3.6\%$ (mean \pm SD, $n = 44$) and $\Delta F/F_m'$ by $92.8\% \pm 1.1\%$ (mean \pm SD, $n = 22$).

Toxicity of PSII herbicides: specific growth rate. The inhibition of SGR by PSII herbicides increased with concentration (Fig. 1). Concentrations of each herbicide that inhibited 10% and 50% of SGR (EC₁₀ and

Herbicide	Specific growth rate (SGR day ⁻¹)		Effective quantum yield ($\Delta F/F_m'$)	
	Control	Ref. Inh. (%)	Control	Ref. Inh. (%)
Diuron	0.148 ± 0.0046	72 ± 3.4	0.329 ± 0.049	96 ± 2.4
Bromacil	0.123 ± 0.011	44 ± 3.1	0.386 ± 0.016	94 ± 4.3
Hexazinone	0.123 ± 0.0066	85 ± 5.8	0.354 ± 0.027	NA
Metribuzin	0.127 ± 0.013	77 ± 5.0	0.356 ± 0.032	NA
Propazine	0.115 ± 0.0050	36 ± 0.7	0.395 ± 0.010	91 ± 2.1
Simazine	0.113 ± 0.011	100 ± 12	0.342 ± 0.053	100 ± 0.00
Tebuthiuron	0.152 ± 0.0064	46 ± 4.6	0.358 ± 0.023	95 ± 5.0
Haloxyfop	0.112 ± 0.0043	49 ± 3.2	0.357 ± 0.043	82 ± 0.1
Imazapic	0.0952 ± 0.0098	38 ± 4.9	0.371 ± 0.020	95 ± 5.3

Table 1. Assay performance of solvent control and reference toxicants (6 $\mu\text{g L}^{-1}$ diuron) for each of the nine herbicides tested over 14 days (SGR, $n = 4\text{--}6$ reps; $\Delta F/F_m'$, $n = 3$). Mean \pm SD. Ref. Inh. (%) denotes percent inhibition of reference toxicant relative to control. NA denotes data not measured.

EC_{50} , respectively), as well as no effect concentrations (NEC) in *C. goreau* are listed in Table 2. All PSII herbicides were toxic but exhibited a broad range of potencies. *C. goreau* was most sensitive to the reference herbicide diuron ($EC_{50} = 4.45 \mu\text{g L}^{-1}$) and least sensitive to simazine ($EC_{50} = 387 \mu\text{g L}^{-1}$) (Table 2). All relative potencies were proportionally lower than diuron ranging from bromacil (ReP = 0.16) to simazine (ReP = 0.012). Slopes for fitted concentration–response curves were similar for the PSII herbicides ranging between 2.5 and 5.4 ($R^2 \geq 0.83$), while hexazinone had a much greater slope (Slope = 9.1, $R^2 = 0.97$) (Fig. 1). The order of toxicity based on EC_{50} values were: diuron > bromacil > metribuzin > propazine > hexazinone > tebuthiuron > simazine (Table 2). The EC_{10} and NEC toxicity thresholds for SGR followed a similar order of toxicity (Table 2, Figs. 1, 2).

Toxicity of PSII herbicides: effective quantum yield. The order of toxicity of PSII herbicides on $\Delta F/F_m'$ in *C. goreau* was the same as per SGR (Table 2). However, $\Delta F/F_m'$ was a more sensitive endpoint: diuron $EC_{50} = 1.20 \mu\text{g L}^{-1}$ and simazine $EC_{50} = 93.3 \mu\text{g L}^{-1}$ (Table 2). Fitted concentration–response curves (Fig. 1) had similar shapes, with slopes for all PSII herbicides ranging from 1.2 to 1.9 ($R^2 \geq 0.86$). Relative potencies of each herbicide relative to diuron ranged from 0.013 to 0.14 (Table 2). The EC_{10} and NEC thresholds for $\Delta F/F_m'$ inhibition followed a similar order of toxicity to inhibition of SGR (Table 2, Figs. 1, 2).

Toxicity of non-PSII herbicides. Both non-PSII herbicides, imazapic and haloxyfop, failed to inhibit growth or $\Delta F/F_m'$ at the highest concentrations of 165,000 $\mu\text{g L}^{-1}$ and 2980 $\mu\text{g L}^{-1}$, respectively (Table 2; Fig. 3). Higher concentrations of imazapic and haloxyfop were unable to be tested as they altered the pH of the IMK media below an acceptable range.

Relationship between growth and effective quantum yield inhibition. There was a linear relationship between growth inhibition and $\Delta F/F_m'$ inhibition of PSII herbicides on *C. goreau* (slope = 4.45; $R^2 = 0.83$; Fig. 4). SGR: $\Delta F/F_m'$ ratios ranged between 2.96 and 4.63, with the exception of tebuthiuron (SGR : $\Delta F/F_m' = 8.10$) (Table 2).

Discussion

Effects of PSII herbicides on growth. Each of the PSII herbicides inhibited SGR and $\Delta F/F_m'$ in the cultures of the coral symbiont *C. goreau* over 14 days. This was expected as the PSII herbicides all bind to the D1 protein site in PSII which is common across phototrophs⁴⁶. Despite the shared mode of action there was a > 80-fold difference in potency between the most toxic herbicide diuron and the least toxic herbicide simazine (comparison of SGR EC_{50} s). There was no relationship between toxicity and chemical class. For example, diuron and tebuthiuron are both phenylureas but the toxicity of diuron was > 70-fold greater than tebuthiuron. Substantial differences in toxicity were also evident among the triazines (simazine and propazine) and triazinones (metribuzin and hexazinone). There was also no relationship between EC_{50} values (Table 2) and water solubility or log K_{OW} (Table S1), indicating that transport across cell walls and membranes and/or accumulation within the cells did not have a major influence on relative potency. Instead, the toxicity of each herbicide is most likely related to the specific affinity (steric compatibility and binding properties) between each herbicide and the Q_B binding site on the D1 protein in PSII⁴⁷.

Direct comparisons of toxicity between species should be made with caution due to differences in experimental conditions; however, comparing the EC_{50} s against datasets with several species' studies carried out under similar conditions can provide insights into the relative sensitivity of *C. goreau* compared with other species. Diuron, applied in this study as a reference toxicant, is the most studied PSII herbicide with respect to effects on marine microalgal growth. The growth inhibition EC_{50} values for 17 species range between 3.4 and 110 $\mu\text{g L}^{-1}$ diuron^{23,38,39,43,48}, indicating cultured *C. goreau* is among the most sensitive. The toxicity of herbicides to two other microalgal species *Rhodomonas salina*²³ and *Chaetoceros muelleri*³⁸ were tested in the same laboratory under similar conditions (Table 3). Comparisons between EC_{50} values reveal broadly similar sensitivities




Figure 1. Concentration–response curves for *Cladocopium goreauii* on the relative percent inhibition of 14-day specific growth rate (SGR; closed circle, mean \pm SD) and effective quantum yield ($\Delta F/F_m'$; open circle, mean \pm SD) following herbicide exposure to (a) diuron; (b) bromacil; (c) hexazinone; (d) metribuzin; (e) propazine; (f) simazine; and (g) tebuthiuron at increasing concentrations. Sigmoidal, 4-parameter curve fit (solid line). All concentrations in $\mu\text{g L}^{-1}$ ($n = 4\text{--}6$ per test; error bars not visible are smaller than symbol).

Herbicide	Endpoint	SGR	$\Delta F/F_m'$	SGR: $\Delta F/F_m'$ ratio
Diuron	NEC	2.75 (2.56–2.93)		
	EC ₁₀	2.54 (2.34–2.75)	0.29 (0.26–0.33)	8.76
	EC ₅₀	4.45 (4.31–4.59)	1.20 (1.15–1.26)	3.71
	ReP	<i>1</i>	<i>1</i>	
Bromacil	NEC	16.6 (15.4–20.6)		
	EC ₁₀	18.3 (16.9–19.9)	2.54 (2.29–2.82)	7.20
	EC ₅₀	27.7 (26.7–28.7)	8.36 (8.01–8.69)	3.31
	ReP	<i>0.16</i>	<i>0.14</i>	
Hexazinone	NEC	71.7 (63.4–91.0)		
	EC ₁₀	78.7 (57.8–92.0)	8.36 (7.14–9.80)	9.41
	EC ₅₀	100 (96.1–141)	33.8 (30.6–37.6)	2.96
	ReP	<i>0.045</i>	<i>0.036</i>	
Metribuzin	NEC	23.6 (21.3–27.5)		
	EC ₁₀	22.3 (16.2–25.9)	2.31 (2.08–2.56)	9.65
	EC ₅₀	33.5 (30.2–50.4)	8.75 (8.39–9.12)	3.83
	ReP	<i>0.13</i>	<i>0.14</i>	
Propazine	NEC	45.1 (37.0–51.1)		
	EC ₁₀	50.8 (44.8–57.4)	5.42 (4.94–5.95)	9.37
	EC ₅₀	86.5 (83.0–90.1)	18.7 (18.0–19.5)	4.63
	ReP	<i>0.052</i>	<i>0.064</i>	
Simazine	NEC	320 (234–452)		
	EC ₁₀	257 (226–294)	28.8 (23.9–35.3)	8.92
	EC ₅₀	387 (361–416)	93.3 (84.6–102)	4.15
	ReP	<i>0.012</i>	<i>0.013</i>	
Tebuthiuron	NEC	107 (84.6–136)		
	EC ₁₀	138 (108–173)	6.37 (4.79–8.50)	21.7
	EC ₅₀	331 (300–NA)	41.0 (36.3–46.3)	8.07
	ReP	<i>0.013</i>	<i>0.029</i>	
Haloxypop	NEC	> 2980		
	EC ₁₀	> 2980	> 2980	NA
	EC ₅₀	> 2980	> 2980	NA
	ReP	NA	NA	
Imazapic	NEC	> 165,000		
	EC ₁₀	> 165,000	> 165,000	NA
	EC ₅₀	> 165,000	> 165,000	NA
	ReP	NA	NA	

Table 2. No effect concentration values (NEC, Fig. 2) and effect concentration values (EC₁₀ and EC₅₀, Fig. 1) for growth inhibition (SGR) and photoinhibition ($\Delta F/F_m'$) of *Cladocodium goreauii* over 14 days to each herbicide. Concentrations are reported in $\mu\text{g L}^{-1}$ (95% confidence intervals). Relative equivalent potencies (ReP) of each herbicide are derived by comparing SGR EC₅₀ of each herbicide to the reference herbicide diuron. Calculated ReP values in italics to differentiate from modelled no effect and effect concentration values. NA denotes data not measured.

to each PSII herbicide (within threefold differences) except for hexazinone, which was > tenfold more toxic to the cryptophyte *R. salina* than *C. goreauii*.

Toxicity of non-PSII herbicides on growth. The two non-PSII herbicides haloxypop and imazapic had no observable effect on SGR in *C. goreauii*. This insensitivity to the non-PSII herbicides is consistent with other marine microalgae *R. salina*²³, *C. muelleri*³⁸ (Table 3) and *Navicula* spp.⁴⁹. The phenoxy herbicide haloxypop inhibits the enzyme acetyl-CoA carboxylase (ACCase inhibitor) and blocks production of lipids and fatty acids^{42,50}. ACCase inhibitors such as haloxypop act on homomeric (eukaryotic) ACCases and sequences for homomeric ACCases have been reported in dinoflagellates⁵¹, indicating the potential for activity. However, the herbicidal activity of haloxypop-p-methyl also depends on its hydrolysis within a plant and this may not occur within *C. goreauii*⁴². Furthermore, Thomas et al.²³ proposed that the active hydrolyzed form of haloxypop which contains a carboxyl moiety may bind calcium (Ca^{2+}) and/or magnesium (Mg^{2+}) ions in seawater⁵² which could affect activity directly or help to stabilize the herbicide at the surface water:air interface⁵³. Like haloxypop, imazapic also has a carboxyl group which may explain its low toxicity to *C. goreauii* cultured in seawater. Further-




Figure 2. Proportional decline of specific growth rate (SGR) for *Cladocopium goreauii* relative to control treatment (solid black line) and 95% confidence intervals (black dashed line) to derive no effect concentration (NEC) (red line) and 95% confidence interval (red dashed line) of (a) diuron; (b) bromacil; (c) hexazinone; (d) metribuzin; (e) propazine; (f) simazine; and (g) tebuthiuron from Bayesian non-linear gaussian model fits. All concentrations in $\mu\text{g L}^{-1}$ ($n = 4\text{--}6$ per test).

more, acetohydroxy acid synthase (AHAS or ALS), the target enzyme for imazapic, has not yet been reported in dinoflagellates⁵⁴.




Figure 3. Response of *Cladocopium goreauui* to alternative non-PSII herbicides, haloxyfop and imazapic. Boxplots of the specific growth rate (SGR day⁻¹) and effective quantum yields ($\Delta F/F_m'$) in response to haloxyfop (a,c) and imazapic (b,d). Note: all concentrations here are in mg L⁻¹.




Figure 4. Linear relationship (black line) between inhibition of effective quantum yield ($\Delta F/F_m'$) and inhibition of specific growth rate (SGR) in *Cladocopium goreauui* by PSII herbicides (dashed red lines 95% confidence bands). Error bars are 95% CI for each EC₅₀ value. Blue line indicates 1:1 relationship. *Diu* - diuron, *Brom* - bromacil, *Hex* - hexazinone, *Met* - metribuzin, *Pro* - propazine, *Sim* - simazine, *Teb* - tebutiuron.

Effects of herbicides on photosynthetic efficiency. All PSII herbicides inhibited effective quantum yield ($\Delta F/F_m'$), which is proportional to photosynthetic efficiency at a given light intensity⁵⁵, in cultured Symbiodiniaceae. Inhibition of $\Delta F/F_m'$ by the seven PSII herbicides was on average 4.4 times more sensitive than SGR based on EC₅₀ ratios (Table 2) and the linear regression slope (Fig. 4). The ratio of inhibition of SGR: $\Delta F/F_m'$ is comparable to other tropical marine microalgae: *R. salina* (average SGR: $\Delta F/F_m' = 4.3$ for 7 PSII herbicides)²³; *Chaetoceros muelleri* (3.0 for 3 PSII herbicides)³⁸; *Navicula* spp. (1.5 for 3 herbicides) and *Nephroselmis*

Species	<i>Cladocodium goreau</i>	<i>Rhodomonas salina</i>	<i>Chaetoceros muelleri</i>
Phylum	Dinoflagellata	Cryptophyta	Bacillariophyta
Diuron	4.45 (4.31–4.59)	6.27 (6.02–6.54)	12.4 (11.8–13.0)
Bromacil	27.7 (26.7–28.7)	19.3 (17.7–21.0)	
Hexazinone	100 (96.1–141)	8.50 (7.99–9.06)	
Metribuzin	33.5 (30.2–50.4)	13.4 (12.3–14.5)	
Propazine	86.5 (83.0–90.1)	188 (177–201)	98.2 (91.7–105)
Simazine	387 (361–416)	184 (173–195)	
Tebuthiuron	331 (300–NA)	112 (106–119)	187 (179–195)
Haloxypop	> 2980	> 3700	> 4570
Imazapic	> 165,000	790,000 (760,000–825,000)	

Table 3. Comparison among toxicity thresholds of *Cladocodium goreau*, *Rhodomonas salina* and *Chaetoceros muelleri*. Effect concentration values (SGR EC₅₀) with 95% confidence intervals derived for *Cladocodium goreau*, *Rhodomonas salina* and *Chaetoceros muelleri* exposed to herbicides. All experiments were conducted under similar conditions in the same laboratory. Concentrations are reported in µg L⁻¹. EC₅₀ values for *Rhodomonas salina* obtained from Thomas et al.²³ and *Chaetoceros muelleri* from Thomas et al.³⁸.

pyriformis (1.3 for 3 herbicides)⁴⁹. Although growth in microalgae is directly dependent on photosynthesis the relationship with inhibition of $\Delta F/F_m'$ by PSII herbicides may not necessarily be 1:1 since microalgae may draw upon their current resources to continue growth, and some (including Symbiodiniaceae) can adjust to reduced photosynthesis by increasing reliance on heterotrophy⁵⁶. Differences in $\Delta F/F_m'$: SGR between studies and species may also be due to light intensity which can affect both $\Delta F/F_m'$ and SGR⁵⁷, and the nutritional status of cells which is dependent on the composition of the growth media and may change depending on the duration of exposure⁵⁸. Interestingly, when based on EC_{10s} the average ratio of SGR: $\Delta F/F_m'$ was 10.7, indicating an even greater sensitivity of $\Delta F/F_m'$ to herbicides relative to SGR at lower herbicide concentrations (also reflected by the differences in slopes of the concentration response curves for both parameters (Fig. 1)). The reason for this is unclear; however, relative sensitivities between these endpoints may change with exposure duration. For example, inhibition of $\Delta F/F_m'$ by PSII herbicides is relatively consistent after it reaches a rapid maxima²⁷, while effects of PSII herbicides on SGR may continue to decline over the course of the exposure period (and the nature of this relationship is likely to change with light intensity⁵⁷). The insensitivity of $\Delta F/F_m'$ in *C. goreau* to the non-PSII herbicides haloxypop and imazapic was expected as their modes of action do not block electron transport in PSII, hence there would be no increase in chlorophyll *a* fluorescence⁵⁷. This result was consistent with other tropical marine microalgae *R. salina*²³, *C. muelleri*³⁸ (Table 3) and *Navicula* spp.⁴⁹.

The effects of PSII herbicides on $\Delta F/F_m'$ have been assessed for both cultured and symbiotic (*in hospite*) forms of Symbiodiniaceae. EC₅₀ values have been reported for Symbiodiniaceae *in hospite* with corals for the PSII herbicides: ametryn, diuron, hexazinone, atrazine, simazine, tebuthiuron and irgarol 1051^{20,27,28,59}, diuron and hexazinone in a jellyfish⁶⁰ and diuron in an anemone⁶¹. However, the species of Symbiodiniaceae *in hospite* was only identified in one instance²⁰. A comparison of the EC_{50s} for diuron among cultured and *in hospite* Symbiodiniaceae (Table 4) shows relatively consistent values among and between these groups. Given that experimental conditions, especially light intensity^{22,57}, can affect inhibition of $\Delta F/F_m'$ in microalgae, direct comparisons are difficult between studies. However, one study found *Durusdinium trenchii* to be twice as sensitive to *C. goreau* under identical culture and measurement conditions³⁵. It was hypothesized that differences in sensitivity to diuron among Symbiodiniaceae types may be due to the reported differences in capacity to repair damaged PSII reaction centres affected by herbicide-mediated photoinhibition⁶². The diuron EC₅₀ for *C. goreau* in the current study was the lowest reported for this species and, while this may be due to differences in experimental conditions, the comparatively long herbicide exposures (14 days) may also play a role, allowing for a build-up of damage to PSII (chronic photoinhibition) and leading to a depression of $\Delta F/F_m'$ ⁶³. Nevertheless, the strong and consistent relationships between $\Delta F/F_m'$ and SGR and between Symbiodiniaceae species for multiple PSII herbicides supports the measurement of $\Delta F/F_m'$ inhibition in marine microalgae as a valuable and biologically relevant toxicity endpoint for PSII herbicides (but should not be applied to herbicides with other modes of action).

Relevance of herbicide sensitivity of free-living Symbiodiniaceae. The apparent lack of differences in sensitivity to diuron between cultured, freshly isolated and *in hospite* Symbiodiniaceae, along with the rapid onset of $\Delta F/F_m'$ inhibition reported for symbiotic corals²⁷, indicates the multiple membrane layers of the coral host provide little barrier to diuron reaching intracellular Symbiodiniaceae. There were also similarities in the sensitivity of cultured *C. goreau* to EC_{50s} reported for coral-hosted Symbiodiniaceae: hexazinone 8.8 µg L⁻¹²⁸ and 14 µg L⁻¹²⁰; simazine 150 µg L⁻¹ and tebuthiuron 175 µg L⁻¹²⁷. These comparisons demonstrate a consistency of access and binding of PSII herbicides to the D1 protein within Symbiodiniaceae, regardless of whether the dinoflagellate is in its symbiotic or free-living form and supports the application of cultured *C. goreau* for assessing the toxicity of PSII herbicides.

The exposure of PSII herbicides to Symbiodiniaceae within corals can lead to coral bleaching (breakdown of symbiosis), reduced translocation of autotrophically-derived nutrients to the host, reduced reproduction and mortality^{29,30,65}, and similar responses are possible in symbiotic foraminifera²², jellyfish⁶⁴ and ascidians⁶¹.

Symbiodiniaceae species	Culture or symbiotic host (exposure duration)	EC ₅₀	References
Isolated cells			
<i>Cladocopium goreau</i> (formerly <i>Symbiodinium</i> Clade C1)	Culture (14 days)	1.2	This study
	Culture (2 days)	7.4	³⁴
	Culture (1 day)	1.4	¹¹
	Culture (1 day)	2.1	³⁵
<i>Durusdinium trenchii</i> (formerly <i>Symbiodinium</i> Clade D)	Culture (1 day)	1.1	³⁵
Unknown, isolated from coral <i>Stylophora pistillata</i>	Freshly isolated (10 min)	5.5	²⁷
In hospite			
<i>Cladocopium</i> C2 (formerly <i>Symbiodinium</i> Clade C2)	Coral: <i>Acropora millepora</i> (4 days)	2.9	²⁰
Unknown	Coral: <i>Seriatopora hystrix</i> (10 h)	2.3	²⁸
Unknown	Coral: <i>Acropora formosa</i> (10 h)	2.7	²⁸
Unknown	Coral: <i>Acropora formosa</i> (10 h)	5.1	²⁷
Unknown	Coral: <i>Montipora digitata</i> (10 h)	5.9	²⁷
Unknown	Coral: <i>Porites cylindrica</i> (10 h)	4.3	²⁷
Unknown	Anemone: <i>Exaiptasia pallida</i> (2 days)	8	⁶¹
Unknown	Jellyfish: <i>Cassiopea maremetens</i> (7 days)	1.4	⁶⁴

Table 4. Comparison of EC₅₀ values derived for $\Delta F/F_m'$ in Symbiodiniaceae exposed to diuron. EC₅₀ values for $\Delta F/F_m'$ in Symbiodiniaceae in cultured and freshly isolated cells and in symbiosis (*in hospite*). EC₅₀ in $\mu\text{g L}^{-1}$.

The free-living Symbiodiniaceae in culture and in the environment alternate between motile and non-motile forms⁶⁶. This mobility facilitates dispersal and infection of host recruits, which are initially symbiont free for most coral species⁶⁷. Impacts of PSII herbicides on populations of free-living Symbiodiniaceae could limit the onset of mutualistic endosymbiosis in corals and other symbiotic invertebrates that have a critical reliance on their symbionts for autotrophic energy acquisition. Free-living Symbiodiniaceae also represent a reservoir of symbionts for bleached coral hosts that are needed for rapid recovery and survival following thermal bleaching events^{31,68}. Indeed, free-living Symbiodiniaceae are widespread in the ocean, with a recent study detecting this family in over 90% of sampling sites and making up 0.1% of total eukaryotic reads in tropical and sub-tropical waters³¹. Impacts by herbicides on these highly diverse and abundant free-living populations of Symbiodiniaceae may therefore have ecological consequences beyond the risk to their role in invertebrate symbiosis.

Implications for water quality guideline values. The Australian and New Zealand marine WQGVs for all herbicides tested here (apart from the reference herbicide diuron) are of low reliability due to lack of appropriate marine toxicity threshold data^{39,41–43}. The SGR toxicity thresholds for cultured *C. goreau* represent the first toxicity dataset for Symbiodiniaceae that is suitable for application to water quality guideline derivation. The comparatively slow growth rate of *C. goreau* (common for all Symbiodiniaceae) meant that the chronic growth assay for this species needed to be longer than the standard 3- or 4-day exposure applied for most algal species. Consequently, the toxicity thresholds identified here could be considered relatively conservative and appropriate for comparison against long in situ exposure durations. NEC and EC₁₀ toxicity thresholds are the preferred data for inclusion in Species Sensitivity Distributions (SSDs) used to derive WQGVs⁴⁵ and these are compared in Table 5 against existing and proposed WQGVs for each of the herbicides. Australian and New Zealand WQGVs are currently derived to protect 99, 95, 90 and 80% (PC99, 95, 90 and 80, respectively) of species in marine and freshwater ecosystems⁴¹, and Table 5 compares SGR endpoints rather than those for $\Delta F/F_m'$, as inhibition of growth is currently considered to better represent an ecologically relevant impact⁴⁵.

The SGR NECs and EC₁₀s for all nine herbicides were greater than the proposed WQGVs (Table 5), indicating the proposed WQGVs would be protective of *C. goreau* growth. The current WQGVs⁴¹ were proposed three decades ago and current guideline values for bromacil and hexazinone would not be protective of *C. goreau*. However, the SGR toxicity thresholds for *C. goreau*, in combination with other similar data for the tropical marine cryptophyte *R. salina*²³, the diatom *C. muelleri*³⁸ and the coral *Acropora tenuis*⁶⁹ will contribute to improving the reliability of WQGVs and their relevance to tropical marine ecosystems such as the GBR. The SGR toxicity thresholds reported here for *C. goreau* were all higher than concentrations detected in the GBR of up to 0.778 $\mu\text{g diuron L}^{-1}$ over time averaged month-long passive sampler deployments¹². Additionally, the $\Delta F/F_m'$ toxicity thresholds for *C. goreau*, except for diuron, were also higher than concentrations reported in the GBR¹². However, improving WQGVs for alternative pesticides (such as the herbicides in this study) will improve confidence in the application of ms-PAF to predict the total toxicity of all pesticides detected in water quality monitoring programs. For example, over 80% of water samples taken in the GBRCA between 2011 and 2015 contained between 2 and 20 quantifiable pesticides for samples collected in the GBRCA⁸. Including the contribution of all pesticides to ecological risk using ms-PAF results in more reported exceedances of WQGVs¹², and is the preferred method in monitoring programs that inform agricultural management practices adjacent to sensitive marine habitats such as the GBR⁶.

Herbicide	This study		Proposed WQGV				Current WQGV			
	NEC	EC ₁₀	99%	95%	90%	80%	99%	95%	90%	80%
Diuron	2.75	2.54	0.43	0.67	0.86	1.2	0.2			
Bromacil	16.6	18.3	0.23	1.1	2.2	4.8	180			
Metribuzin	23.6	22.3	2.0	2.7	3.1	3.9	NA			
Propazine	45.1	50.8	2.2	4.6	6.4	9.2	NA			
Hexazinone	71.7	78.7	1.8	2.5	3.1	4.0	75			
Simazine	320	257	28	63	84	130	0.2	3.2	11	35
Tebuthiuron	107	138	4.7	11	17	26	0.02	2.2	20	160
Haloxfop	> 2980	590	2000	3400	6100	NA				
Imazapic	> 165,000	0.049	0.44	1.2	3.6	NA				

Table 5. Comparison of toxicity thresholds derived here vs Australian WQGVs. Comparisons of toxicity thresholds of EC₁₀ and NEC for specific growth rate (SGR) of *Cladocopium goreau* in response to nine herbicides with current guideline values⁴¹ and proposed guideline values^{39,42,43} derived for 99, 95, 90 and 80% species protection (PC99, PC95, PC90, PC80 respectively). All concentrations are in µg L⁻¹. NA denotes no current WQGV.

Conclusion

The toxicity growth thresholds for alternative herbicides to the free-living form of the common coral symbiont *C. goreau* were reported for the first time. The seven PSII herbicides all inhibited photosynthetic efficiency ($\Delta F/F_m'$), leading to reduced SGR over the 14-d exposure period. The sensitivity of $\Delta F/F_m'$ to PSII herbicides was on average 4.4-fold greater than for SGR, but the clear mechanistic link and consistent relationship between inhibition of $\Delta F/F_m'$ and SGR indicates that inhibition of $\Delta F/F_m'$ should be considered a biologically relevant toxicity endpoint for PSII herbicides to marine microalgae^{23,37,38}. The non-PSII herbicides haloxyfop and imazapic did not affect SGR in *C. goreau* at very high concentrations indicating these individual herbicides do not pose a significant risk to this species. The effects of PSII herbicides on free-living *C. goreau* occurred at similar concentrations as shown for Symbiodiniaceae *in hospite*, indicating cultures of this species are relevant proxies for both free-living and symbiotic forms of this species. As mutualistic endosymbiotic partners with scleractinian corals, inclusion of toxicity data for Symbiodiniaceae to relevant contaminants such as alternative herbicides will improve our confidence that future WQGVs are adequate to inform risk assessments for tropical marine ecosystems. While this study targeted some of the most frequently detected alternative herbicides in GBR waters, there remains several pesticides, including insecticides and fungicides, with no current WQGVs and further testing is needed to address this.

Materials and methods

Test species and culture conditions. A monoclonal strain of *Cladocopium goreau* (formerly *Symbiodinium* clade C1⁷⁰) was isolated in 2010 from coral *Acropora tenuis* near Magnetic Island in Queensland, Australia. In house cultures (Australian Institute of Marine Science ID: SCF 055-01.10) were maintained in sterile 75-mL culture flasks in IMK growth media prepared with 0.2 µm-filtered seawater (pH = 7.8 ± 0.5; salinity = 32.5 ± 0.7 psu; Wako Chemicals USA, Richmond) and incubated in Steridium environmental chambers at 14:10 h light:dark cycles with an irradiance of 60–75 µmol quanta m⁻² s⁻¹ (Sylvania FHO24W/T5/865) at 27 ± 0.45 °C. Fortnightly 10% inoculations were performed on cultures to consistently use 14-days old cultures in toxicity bioassays.

Herbicide stock preparation. Herbicide stock solutions were prepared with PESTANAL analytical grade products (HPLC ≥ 98%) and purchased from Sigma-Aldrich (NSW, Australia), including: diuron (CAS 330-54-1), bromacil (CAS 314-40-9), haloxyfop-p-methyl (CAS 72619-32-0), hexazinone (CAS 51235-04-2), imazapic (CAS 104098-48-8), metribuzin (CAS 21087-64-9), propazine (CAS 139-40-2), simazine (CAS 122-34-9), and tebuthiuron (CAS 34014-18-1). Stock solutions (5–600 mg L⁻¹) were prepared in Milli-Q water or filtered seawater (FSW) in 500 mL or 1 L autoclaved Schott bottles. Diuron and metribuzin were dissolved using the carrier solvent ethanol (EtOH; ≤ 0.002% v/v in all exposure treatments). Haloxyfop and simazine were dissolved in the carrier solvent dimethyl sulfoxide (DMSO; ≤ 0.006% v/v in all exposure treatments). No solvent carrier was used in the preparation of the remaining herbicide stock solutions.

Bioassay protocol. Cultures of 14-days old *C. goreau* in exponential growth phase at a starting density of 1.7–2.7 × 10⁴ cells mL⁻¹ were used in all bioassays. The required algae inoculum was transferred to 50 mL polypropylene conical centrifuge tubes (3–6 replicates per treatment) containing 30 mL of IMK media for 14-days exposure tests. Each 50 mL centrifuge tube was dosed with a range of herbicide concentrations. Treatments included control (no herbicide) and herbicide reference (diuron, 6 µg L⁻¹). *C. goreau* cells were incubated for 14 days at 130 rpm in a refrigerated incubator shaker (Thermoline Scientific) at 27 ± 1 °C on a 14:10 h light:dark cycle under 60–75 µmol quanta m⁻² s⁻¹. Samples were randomized every 1–2 days to ensure similar light conditions for all replicate samples.

Cell density counts and specific growth rate. Aliquots (1 mL) of treatment samples were fixed in glutaraldehyde (0.5% v/v final concentration) and surfactant (Pluronic F68; 0.1% v/v final concentration) to reduce clumping of cells as per Marie et al.⁴⁴. A flow cytometer (BD Accuri C6, BD Biosciences, CA, USA) equipped with one red and blue laser (488 nm, 50 mW maximum solid state; 640 nm, 30 mW diode) was used to quantify cell density as per Trenfield et al.⁷¹ at several timepoints during each bioassay. A lag phase of 3–4 days was typically observed for *C. goreau*; therefore, day 4 was considered the first timepoint for measuring cell density over the logarithmic growth phase. Subsequent timepoints (typically, day 4, 7, 10 and 14) were chosen due to availability of instrument. Flow rate was set to 35 $\mu\text{L min}^{-1}$, 16- μm core size and sample volume of 50 μL . A fixed gating was used to exclusively measure *C. goreau*, minimizing quantification of microbes and degraded chloroplasts of senescing cells. Viable *C. goreau* cells represented 70–90% of all particles identified in samples over 14 days. Specific growth rates (SGR) were expressed as the logarithmic increase in cell density from day i (t_i) to day j (t_j) as per Eq. (1), where SGR_{i-j} is the specific growth rate from time i to j , X_j is the cell density at day j and X_i is the cell density at day i ⁵⁸. SGR relative to the control treatment was used to derive modelled chronic effect concentration values for growth inhibition. Test acceptability was confirmed if the SGR of control replicates was $\geq 0.1 \text{ day}^{-1}$ (1 doubling/week) and had a coefficient of variation (CV) $\leq 10\%$ for control SGR^{58,72–75}.

$$\text{SGR}_{i-j} = \frac{\ln X_j - \ln X_i}{t_j - t_i} (\text{day}^{-1}) \quad (1)$$

Chlorophyll fluorescence. Chlorophyll fluorescence was used as a proxy to measure photosynthetic health of *C. goreau* using pulse amplitude modulation fluorometry (microscopy imaging PAM, Walz, Germany) to determine effective quantum yield ($\Delta F/F_m'$) after 14 days herbicide exposure tests. $\Delta F/F_m'$ was calculated from light adapted minimum fluorescence (F) and maximum fluorescence (F_m') measurements, following Eq. (2) from Schreiber et al.⁷⁶. $\Delta F/F_m'$ was measured from three replicates per treatment, including the control and reference toxicant. A minimum of three cells was measured with acceptable control treatment values. The observed control measurements for $\Delta F/F_m'$ were within acceptable limits ($\Delta F/F_m' > 0.30$) as per Hennige et al.⁷² and Karim et al.⁷⁷.

$$\Delta F/F_m' = \frac{F_m' - F}{F_m'} \quad (2)$$

Percent inhibition was calculated relative to controls as per Eq. (3). Inhibition of $\Delta F/F_m'$ suggests reduced energy conversion in PSII of the organism under a known light intensity. These measurements were used to compare photoinhibition between the controls and respective treatments. Three to six replicates at 6 $\mu\text{g L}^{-1}$ diuron were included in each bioassay to estimate the consistency of inhibitory responses of cells. Microscopy iPAM settings were: actinic light = 1, measuring light = 10–12, gain = 3, damp = 2, actinic light width = 180 s and saturation pulse intensity = 2.

$$\% \text{ Inhibition} = \frac{X_{\text{control}} - X_{\text{treatment}}}{X_{\text{control}}} \times 100 \quad (3)$$

Physico-chemical and herbicide analyses. Physical and chemical characteristics of each treatment were measured at 0 days and 14 days including pH and salinity (LAQUAact-PC110 Meter, HORIBA Scientific) and dissolved oxygen (HQ30D Portable Meter, HACH) as per Thomas et al.²³. Temperature was logged in 10-min intervals over the duration of the test (HOBO, Onset). Herbicide analytical samples were taken at 0 day and 14 days. Aliquots (1 mL) for chemical analysis were transferred into 1.5 liquid chromatography amber glass vials and spiked with surrogate standards (i.e. diuron-d6, hexazinone-d6, metribuzin-d3, simazine-D10, propazine-D6, bromacil-D3, haloxyfop-D4, and imazapic D7) with a 10 ng mL^{-1} final concentration of the surrogate standard. The herbicide and degradation product concentrations were determined by HPLC–MS/MS using an SCIEX Triple Quad 6500 QTRAP mass spectrometer (SCIEX, Concord, Ontario, Canada) equipped with an IonDrive Turbo V ion source using a TurboIonSpray probe^{23,78,79}. Data acquisition and quantification were performed on MultiQuant 3.0 software by Sciex. Compound identification and confirmation were achieved using retention times and via comparison of SRM transition intensity ratios between the sample and isotopically labelled standard in the same run. The geometric mean from start and end concentrations (time-weighted average) was assigned as the ‘measured’ concentration in that sample. The average loss from these measured concentrations was then applied to all nominal concentrations as per Thomas et al.²³.

Data analyses. Measured concentrations were used for all concentration–response modelling and toxicity threshold estimates (Table S1). Specific growth rate and photosynthetic inhibition were calculated as percent inhibition relative to the solvent control or IMK media control (in bioassays when no solvent carrier was used). Herbicide concentrations that inhibited 10 and 50% of growth or $\Delta F/F_m'$ relative to controls (EC_{10} and EC_{50} , respectively) were interpolated from modelled mean values using GraphPad Prism V 7.0 (GraphPad Software Inc., San Diego, USA). Relative potencies (ReP) were calculated for each herbicide ($\text{ReP} = \text{EC}_{50} \text{ diuron} / \text{EC}_{50} \text{ herbicide}$) against the reference toxicant diuron. The estimation of no effect concentrations (NEC) for SGR was calculated in R (V 3.6.1) as per Thomas et al.²³. Proportional decline in SGR (1-inhibition) was modelled as a function of log concentration of each herbicide using a Bayesian non-linear gaussian model with the R package jagsNEC⁸⁰. This model has been specifically developed to derive no effect concentrations (NECs) but also allows

the estimation of EC₁₀ and EC₅₀ values and is adapted from Fox⁸¹. See Thomas et al.²³ for further details. The linear regression of EC₅₀s SGR vs $\Delta F/F_m'$ was performed in SigmaPlot V14 (Systat Software, San Jose, CA).

Received: 14 July 2020; Accepted: 14 October 2021

Published online: 04 November 2021

References

- Castillo, L. E., de la Cruz, E. & Ruepert, C. Ecotoxicology and pesticides in tropical aquatic ecosystems of Central America. *Environ. Toxicol. Chem.* **16**, 41–51. <https://doi.org/10.1002/etc.5620160104> (1997).
- Moreno-González, R. & León, V. Presence and distribution of current-use pesticides in surface marine sediments from a Mediterranean coastal lagoon (SE Spain). *Environ. Sci. Pollut. Res.* **24**, 8033–8048. <https://doi.org/10.1007/s11356-017-8456-0> (2017).
- Hernández-Romero, A. H., Tovilla-Hernández, C., Malo, E. A. & Bello-Mendoza, R. Water quality and presence of pesticides in a tropical coastal wetland in southern Mexico. *Mar. Pollut. Bull.* **48**, 1130–1141. <https://doi.org/10.1016/j.marpolbul.2004.01.003> (2004).
- Wurl, O. & Obbard, J. P. Organochlorine pesticides, polychlorinated biphenyls and polybrominated diphenyl ethers in Singapore's coastal marine sediments. *Chemosphere* **58**, 925–933. <https://doi.org/10.1016/j.chemosphere.2004.09.054> (2005).
- Carvalho, F. P. et al. Organic contaminants in the marine environment of Manila Bay, Philippines. *Arch. Environ. Contam. Toxicol.* **57**, 348–358. <https://doi.org/10.1007/s00244-008-9271-x> (2009).
- Australian Government and Queensland Government. *Reef 2050 Water Quality Improvement Plan, Monitoring Program*. (Australian and Queensland Governments, 2018). <https://www.reefplan.qld.gov.au/tracking-progress/paddock-to-reef/modelling-and-monitoring>.
- O'Brien, D. et al. Spatial and temporal variability in pesticide exposure downstream of a heavily irrigated cropping area: Application of different monitoring techniques. *J. Agric. Food Chem.* **64**, 3975–3989. <https://doi.org/10.1021/acs.jafc.5b04710> (2016).
- Warne, M. St. J., Smith, R. & Turner, R. Analysis of pesticide mixtures discharged to the lagoon of the Great Barrier Reef, Australia. *Environ. Pollut.* **265**, 114088. <https://doi.org/10.1016/j.envpol.2020.114088> (2020).
- Shaw, M. et al. Monitoring pesticides in the Great Barrier Reef. *Mar. Pollut. Bull.* **60**, 113–122. <https://doi.org/10.1016/j.marpolbul.2009.08.026> (2010).
- Kennedy, K. et al. The influence of a season of extreme wet weather events on exposure of the World Heritage Area Great Barrier Reef to pesticides. *Mar. Pollut. Bull.* **64**, 1495–1507. <https://doi.org/10.1016/j.marpolbul.2012.05.014> (2012).
- Mercurio, P. et al. Degradation of herbicides in the tropical marine environment: Influence of light and sediment. *PLoS ONE* **11**, e0165890. <https://doi.org/10.1371/journal.pone.0165890> (2016).
- Gallen, C. et al. Marine Monitoring Program: Annual report for inshore pesticide monitoring 2017–18. Report for the Great Barrier Reef Marine Park Authority. <http://elibrary.gbrmpa.gov.au/jspui/handle/11017/3489>. (2019).
- Smith, R. et al. Large-scale pesticide monitoring across Great Barrier Reef catchments—paddock to reef integrated monitoring, modelling and reporting program. *Mar. Pollut. Bull.* **65**, 117–127. <https://doi.org/10.1016/j.marpolbul.2011.08.010> (2012).
- Oettmeier, W. Herbicide resistance and supersensitivity in photosystem II. *Cell. Mol. Life Sci.* **55**, 1255–1277. <https://doi.org/10.1007/s000180050370> (1999).
- Davis, A., Lewis, S., Brodie, J. & Benson, A. The potential benefits of herbicide regulation: A cautionary note for the Great Barrier Reef catchment area. *Sci. Total Environ.* **490**, 81–92. <https://doi.org/10.1016/j.scitotenv.2014.04.005> (2014).
- King, J., Alexander, F. & Brodie, J. Regulation of pesticides in Australia: The Great Barrier Reef as a case study for evaluating effectiveness. *Agr. Ecosyst. Environ.* **180**, 54–67. <https://doi.org/10.1016/j.agee.2012.07.001> (2013).
- Devlin, M. et al. *Advancing our Understanding of the Source, Management, Transport and Impacts of Pesticides on the Great Barrier Reef 2011–2015. Report for the Queensland Department of Environment and Heritage Protection*. (Tropical Water & Aquatic Ecosystem Research (TropWATER) Publication, James Cook University, 2015). <https://www.qld.gov.au/environment/assets/documents/agriculture/sustainable-farming/reef/rp104c-pesticide-report.pdf>.
- Flores, F., Collier, C. J., Mercurio, P. & Negri, A. P. Phytotoxicity of four photosystem II herbicides to tropical seagrasses. *PLoS ONE* **8**, e75798. <https://doi.org/10.1371/journal.pone.0075798> (2013).
- Haynes, D. et al. The occurrence and impact of herbicides in the Great Barrier Reef, Australia. *Reef Res.* **10**, 3–5 (2000).
- Negri, A. P., Flores, F., Röthig, T. & Uthicke, S. Herbicides increase the vulnerability of corals to rising sea surface temperature. *Limnol. Oceanogr.* **56**, 471–485. <https://doi.org/10.4319/lo.2011.56.2.0471> (2011).
- Marques, J. A., Flores, F., Bianchini, A., Uthicke, S. & Negri, A. P. Acclimation history modulates effect size of calcareous algae (*Halimeda opuntia*) to herbicide exposure under future climate scenarios. *Sci. Total Environ.* **736**, 140308. <https://doi.org/10.1016/j.scitotenv.2020.140308> (2020).
- van Dam, J. W., Negri, A. P., Mueller, J. F. & Uthicke, S. Symbiont-specific responses in foraminifera to the herbicide diuron. *Mar. Pollut. Bull.* **65**, 373–383. <https://doi.org/10.1016/j.marpolbul.2011.08.008> (2012).
- Thomas, M. C., Flores, F., Kaserzon, S., Fisher, R. & Negri, A. P. Toxicity of ten herbicides to the tropical marine microalgae *Rhodomonas salina*. *Sci. Rep.* **10**, 1–16. <https://doi.org/10.1038/s41598-020-64116-y> (2020).
- Magnusson, M., Heimann, K. & Negri, A. P. Comparative effects of herbicides on photosynthesis and growth of tropical estuarine microalgae. *Mar. Pollut. Bull.* **56**, 1545–1552. <https://doi.org/10.1016/j.marpolbul.2008.05.023> (2008).
- Muscatine, L. The role of symbiotic algae in carbon and energy flux in reef corals. *Coral Reefs* **25**, 1–29 (1990).
- Oettmeier, W. Herbicides of photosystems II. In *Structure, Function and Molecular Biology* (ed. Barber, J.) 349–408 (Elsevier, 1992).
- Jones, R. J., Muller, J., Haynes, D. & Schreiber, U. Effects of herbicides diuron and atrazine on corals of the Great Barrier Reef, Australia. *Mar. Ecol. Prog. Ser.* **251**, 153–167. <https://doi.org/10.3354/meps251153> (2003).
- Jones, R. J. & Kerswell, A. P. Phytotoxicity of Photosystem II (PSII) herbicides to coral. *Mar. Ecol. Prog. Ser.* **261**, 149–159. <https://doi.org/10.3354/meps261149> (2003).
- Cantin, N. E., Negri, A. P. & Willis, B. L. Photoinhibition from chronic herbicide exposure reduces reproductive output of reef-building corals. *Mar. Ecol. Prog. Ser.* **344**, 81–93. <https://doi.org/10.3354/meps07059> (2007).
- Negri, A. et al. Effects of the herbicide diuron on the early life history stages of coral. *Mar. Pollut. Bull.* **51**, 370–383. <https://doi.org/10.1016/j.marpolbul.2004.10.053> (2005).
- Decelle, J. et al. Worldwide occurrence and activity of the reef-building coral symbiont *Symbiodinium* in the open ocean. *Curr. Biol.* **28**, 3625–3633. <https://doi.org/10.1016/j.cub.2018.09.024> (2018).
- Baker, A. C. Reef corals bleach to survive change. *Nature* **411**, 765–766. <https://doi.org/10.1038/35081151> (2001).
- Muller-Parker, G., D'elia, C. F. & Cook, C. B. *Coral Reefs in the Anthropocene* 99–116 (Springer, 2015). <https://pdfs.semanticscholar.org/191e119/119ba111eab744a4054c4068f4057a4003bb4058bd4001b9628.pdf>.
- Chakravarti, L. J., Negri, A. P. & Oppen, M. J. Thermal and herbicide tolerances of chromerid algae and their ability to form a symbiosis with corals. *Front. Microbiol.* **10**, 173. <https://doi.org/10.3389/fmicb.2019.00173> (2019).

35. van Dam, J., Uthicke, S., Beltran, V., Mueller, J. & Negri, A. Combined thermal and herbicide stress in functionally diverse coral symbionts. *Environ. Pollut.* **204**, 271–279. <https://doi.org/10.1016/j.envpol.2015.05.013> (2015).
36. Mercurio, P. *et al.* Contribution of transformation products towards the total herbicide toxicity to tropical marine organisms. *Sci. Rep.* **8**, 1–12. <https://doi.org/10.1038/s41598-018-23153-4> (2018).
37. Magnusson, M., Heimann, K., Ridd, M. & Negri, A. P. Pesticide contamination and phytotoxicity of sediment interstitial water to tropical benthic microalgae. *Water Res.* **47**, 5211–5221. <https://doi.org/10.1016/j.watres.2013.06.003> (2013).
38. Thomas, M. C., Flores, F., Kaserzon, S., Reeks, T. & Negri, A. P. Toxicity of the herbicides diuron, propazine, tebuthiuron, and haloxyfop to the diatom *Chaetoceros muelleri*. *Sci. Rep.* **10**, 19592. <https://doi.org/10.1038/s41598-020-76363-0> (2020).
39. Warne, M. St. J., King, O. & Smith, R. Ecotoxicity thresholds for ametryn, diuron, hexazinone and simazine in fresh and marine waters. *Environ. Sci. Pollut. Res.* **25**, 3151–3169. <https://doi.org/10.1007/s11356-017-1097-5> (2018).
40. Traas, T. P. *et al.* In *Species Sensitivity Distributions in Ecotoxicology* (eds Posthuma, L. *et al.*) 315–344 (CRC Press, 2002).
41. ANZG. *Australian and New Zealand Guidelines for Fresh and Marine Water Quality*. 1–103 (Australian and New Zealand Governments and Australian State and Territory Governments, 2018). <http://waterquality.gov.au/anz-guidelines>.
42. King, O., Smith, R., Mann, R. & Warne, M. St. J. *Proposed Aquatic Ecosystem Protection Guideline Values for Pesticides Commonly Used in the Great Barrier Reef catchment Area: Part 2—Bromacil, Chlorothalonil, Fipronil, Fluometuron, Fluoroxypyr, Haloxyfop, MCPA, Pendimethalin, Prometryn, Propazine, Propiconazole, Terbutryn, Triclopyr and Terbutylazine*. (Department of Environment and Science, 2017). <https://www.publications.qld.gov.au/dataset/proposed-guideline-values-27-pesticides-used-in-the-gbr-catchment>.
43. King, O., Smith, R., Mann, R. & Warne, M. St. J. *Proposed Aquatic Ecosystem Protection Guideline Values for Pesticides Commonly Used in the Great Barrier Reef Catchment Area: Part 1–2, 4-D, Ametryn, Diuron, Glyphosate, Hexazinone, Imazapic, Imidacloprid, Isoxaflutole, Metolachlor, Metribuzin, Metsulfuron-methyl, Simazine and Tebuthiuron* 296 (Department of Environment and Science, 2017). <https://www.publications.qld.gov.au/dataset/proposed-guideline-values-27-pesticides-used-in-the-gbr-catchment>.
44. Marie, D., Rigaut-Jalabert, F. & Vaultot, D. An improved protocol for flow cytometry analysis of phytoplankton cultures and natural samples. *Cytom. Part A* **85**, 962–968. <https://doi.org/10.1002/cyto.a.22517> (2014).
45. Warne, M. St. J. *et al.* *Revised Method for Deriving Australian and New Zealand Water Quality Guideline Values for Toxicants: Update of 2015 Version. Prepared for the Revision of the Australian and New Zealand Guidelines for Fresh and Marine Water Quality 48* (Australian and New Zealand Governments and Australian State and Territory Governments, 2018). <https://www.waterquality.gov.au/sites/default/files/documents/warne-wqg-derivation2018.pdf>.
46. Vinyard, D. J., Ananyev, G. M. & Charles Dismukes, G. Photosystem II: The reaction center of oxygenic photosynthesis. *Annu. Rev. Biochem.* **82**, 577–606. <https://doi.org/10.1146/annurev-biochem-070511-100425> (2013).
47. Haworth, P. & Steinback, K. E. Interaction of herbicides and quinone with the qb-protein of the diuron-resistant *Chlamydomonas reinhardtii* mutant Dr2. *Plant Physiol.* **83**, 1027–1031. <https://doi.org/10.1104/pp.83.4.1027> (1987).
48. USEPA. *ECOTOX User Guide: ECOTOXicology Database System. Version 5.0*. (United States Environmental Protection Agency, 2019) <http://cfpub.epa.gov/ecotox/>.
49. Magnusson, M. *Effects of Priority Herbicides and Their Breakdown Products on Tropical, ESTUARINE Microalgae of the Great Barrier Reef Lagoon*. PhD thesis, James Cook University (2009).
50. MacBean, C. *The Pesticide Manual: A World Compendium* (British Crop Protection Council, 2012).
51. Haq, S., Bachvaroff, T. R. & Place, A. R. Characterization of acetyl-CoA carboxylases in the basal dinoflagellate *Amphidinium carterae*. *Mar. Drugs* **15**, 149. <https://doi.org/10.3390/md15060149> (2017).
52. Tang, C. Y., Huang, Z. & Allen, H. C. Interfacial water structure and effects of Mg²⁺ and Ca²⁺ binding to the COOH headgroup of a palmitic acid monolayer studied by sum frequency spectroscopy. *J. Phys. Chem. B* **115**, 34–40. <https://doi.org/10.1021/jp1062447> (2011).
53. Brzozowska, A., Duits, M. H. & Mugele, F. Stability of stearic acid monolayers on artificial sea water. *Colloid Surf. A* **407**, 38–48 (2012).
54. McCourt, J. & Duggleby, R. Acetohydroxyacid synthase and its role in the biosynthetic pathway for branched-chain amino acids. *Amino Acids* **31**, 173–210. <https://doi.org/10.1007/s00726-005-0297-3> (2006).
55. Genty, B., Briantais, J.-M. & Baker, N. R. The relationship between the quantum yield of photosynthetic electron transport and quenching of chlorophyll fluorescence. *BBA* **990**, 87–92. [https://doi.org/10.1016/S0304-4165\(89\)80016-9](https://doi.org/10.1016/S0304-4165(89)80016-9) (1989).
56. Jeong, H. J. *et al.* Heterotrophic feeding as a newly identified survival strategy of the dinoflagellate *Symbiodinium*. *Proc. Natl. Acad. Sci. U.S.A.* **109**, 12604–12609. <https://doi.org/10.1073/pnas.1204302109> (2012).
57. Ralph, P., Smith, R., Macinnis-Ng, C. & Seery, C. Use of fluorescence-based ecotoxicological bioassays in monitoring toxicants and pollution in aquatic systems. *Toxicol. Environ. Chem.* **89**, 589–607. <https://doi.org/10.1080/02772240701561593> (2007).
58. OECD. *Test No. 201: Freshwater Alga and Cyanobacteria, Growth Inhibition Test, OECD Guidelines for the Testing of Chemicals, Section 2* (OECD Publishing, 2011).
59. Kamei, M., Takayama, K., Ishibashi, H. & Takeuchi, I. Effects of ecologically relevant concentrations of Irgarol 1051 in tropical to subtropical coastal seawater on hermatypic coral *Acropora tenuis* and its symbiotic dinoflagellates. *Mar. Poll. Bull.* **150**, 110734. <https://doi.org/10.1016/j.marpolbul.2019.110734> (2020).
60. McKenzie, M. R., Templeman, M. A. & Kingsford, M. J. Detecting effects of herbicide runoff: The use of *Cassiopea maretensis* as a biomonitor to hexazinone. *Aquat. Toxicol.* **221**, 105442. <https://doi.org/10.1016/j.aquatox.2020.105442> (2020).
61. Howe, P. L., Reichelt-Brushett, A. J., Clark, M. W. & Seery, C. R. Toxicity estimates for diuron and atrazine for the tropical marine cnidarian *Exaiptasia pallida* and in-hospite *Symbiodinium* spp. using PAM chlorophyll-a fluorometry. *J. Photochem. Photobiol. B* **171**, 125–132. <https://doi.org/10.1016/j.jphotobiol.2017.05.006> (2017).
62. Takahashi, S., Whitney, S. M. & Badger, M. R. Different thermal sensitivity of the repair of photodamaged photosynthetic machinery in cultured *Symbiodinium* species. *Proc. Natl. Acad. Sci. U.S.A.* **106**, 3237–3242. <https://doi.org/10.1073/pnas.0808363106> (2009).
63. Jones, R. The ecotoxicological effects of Photosystem II herbicides on corals. *Mar. Pollut. Bull.* **51**, 495–506. <https://doi.org/10.1016/j.marpolbul.2005.06.027> (2005).
64. Rowen, D. J., Templeman, M. A. & Kingsford, M. J. Herbicide effects on the growth and photosynthetic efficiency of *Cassiopea maretensis*. *Chemosphere* **182**, 143–148. <https://doi.org/10.1016/j.chemosphere.2017.05.001> (2017).
65. Cantin, N. E., van Oppen, M. J., Willis, B. L., Mieog, J. C. & Negri, A. P. Juvenile corals can acquire more carbon from high-performance algal symbionts. *Coral Reefs* **28**, 405. <https://doi.org/10.1007/s00338-009-0478-8> (2009).
66. Fitt, W. & Trench, R. The relation of diel patterns of cell division to diel patterns of motility in the symbiotic dinoflagellate *Symbiodinium microadriaticum* Freudenthal in culture. *New Phytol.* **94**, 421–432 (1983).
67. Randall, C. J. *et al.* Sexual production of corals for reef restoration in the Anthropocene. *Mar. Ecol. Prog. Ser.* **635**, 203–232. <https://doi.org/10.3354/meps13206> (2020).
68. Baird, A. H., Bhagooli, R., Ralph, P. J. & Takahashi, S. Coral bleaching: The role of the host. *Trends Ecol. Evol.* **24**, 16–20. <https://doi.org/10.1016/j.tree.2008.09.005> (2009).
69. Flores, F., Kaserzon, S., Elisei, G., Ricardo, G. & Negri, A. P. Toxicity thresholds of three insecticides and two fungicides to larvae of the coral *Acropora tenuis*. *PeerJ* **8**, e9615. <https://doi.org/10.7717/peerj.9615> (2020).
70. LaJeunesse, T. C. *et al.* Systematic revision of Symbiodiniaceae highlights the antiquity and diversity of coral endosymbionts. *Curr. Biol.* **28**, 2570–2580. <https://doi.org/10.1016/j.cub.2018.07.008> (2018).

71. Trenfield, M. A. *et al.* Aluminium, gallium, and molybdenum toxicity to the tropical marine microalga *Isochrysis galbana*. *Environ. Toxicol. Chem.* **34**, 1833–1840. <https://doi.org/10.1002/etc.2996> (2015).
72. Hennige, S., Suggett, D., Warner, M., McDougall, K. & Smith, D. Photobiology of *Symbiodinium* revisited: Bio-physical and bio-optical signatures. *Coral Reefs* **28**, 179–195. <https://doi.org/10.1007/s00338-008-0444-x> (2009).
73. Kluetter, A., Trapani, J., Archer, F. I., McLroy, S. E. & Coffroth, M. A. Comparative growth rates of cultured marine dinoflagellates in the genus *Symbiodinium* and the effects of temperature and light. *PLoS ONE* **12**, e0187707. <https://doi.org/10.1371/journal.pone.0187707> (2017).
74. Rogers, J. E. & Davis, R. H. Application of a new micro-culturing technique to assess the effects of temperature and salinity on specific growth rates of six *Symbiodinium* isolates. *Bull. Mar. Sci.* **79**, 113–126 (2006).
75. Sakami, T. Effects of temperature, irradiance, salinity and inorganic nitrogen concentration on coral zooxanthellae in culture. *Fish. Res.* **66**, 1006–1013. <https://doi.org/10.1046/j.1444-2906.2000.00162.x> (2000).
76. Schreiber, U., Müller, J. F., Haugg, A. & Gademann, R. New type of dual-channel PAM chlorophyll fluorometer for highly sensitive water toxicity biotests. *Photosynth. Res.* **74**, 317–330. <https://doi.org/10.1023/A:1021276003145> (2002).
77. Karim, W., Nakaema, S. & Hidaka, M. Temperature effects on the growth rates and photosynthetic activities of *Symbiodinium* cells. *J. Mar. Sci. Eng.* **3**, 368–381. <https://doi.org/10.3390/jmse3020368> (2015).
78. Mercurio, P., Mueller, J. F., Eaglesham, G., Flores, F. & Negri, A. P. Herbicide persistence in seawater simulation experiments. *PLoS ONE* **10**, e0136391. <https://doi.org/10.1371/journal.pone.0136391> (2015).
79. Mercurio, P. *Herbicide Persistence and Toxicity in the Tropical Marine Environment*. PhD thesis, The University of Queensland (2016).
80. Fisher, R., Ricardo, G. & Fox, D. *jags NEC: A Bayesian No Effect Concentration (NEC) Package*. <https://github.com/AIMS/NEC-estimation>. (2019).
81. Fox, D. R. A Bayesian approach for determining the no effect concentration and hazardous concentration in ecotoxicology. *Ecotoxicol. Environ. Saf.* **73**, 123–131. <https://doi.org/10.1016/j.ecoenv.2009.09.012> (2010).

Acknowledgements

We thank Carlos Alvarez-Roa from the Symbiont Culture Facility at the Australian Institute of Marine Science (AIMS) for providing expertise for algae cultivation and Rebecca Fisher (AIMS) for advice regarding statistical analysis. This research was supported by the Australian Government's National Environmental Science Program (NESP) Tropical Water Quality Hub Project 3.1.5 Ecotoxicology of pesticides on the Great Barrier Reef for guideline development and risk assessments. M.M. was supported by AIMS@JCU: Australian Institute of Marine Science, College of Marine and Environmental Sciences, James Cook University. We acknowledge the Bindal and Wulgurukaba people as the Traditional Owners of the land where this work took place. We pay our respects to their Elders past, present and emerging and we acknowledge their continuing spiritual connection to their land.

Author contributions

A.P.N., F.F., N.S. and M.M. designed the experiments. M.M., N.S. and F.F. ran preliminary tests to develop the bioassay. M.M., A.V.-M., F.F. and M.C.T. performed experiments. F.F., M.M. and M.C.T. analyzed data. S.K. performed herbicide analyses. All authors reviewed the manuscript.

Competing interests

The authors declare no competing interests.

Additional information

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1038/s41598-021-00921-3>.

Correspondence and requests for materials should be addressed to F.F.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2021