

Beyond vegetables: effects of indoor LED light on specialized metabolite biosynthesis in medicinal and aromatic plants, edible flowers, and microgreens

Elisa Appolloni,^a Giuseppina Pennisi,^a Ilaria Zauli,^a Laura Carotti,^a Ivan Paucek,^a Stefania Quaini,^b Francesco Orsini^{a*}  and Giorgio Gianquinto^a



Abstract

Specialized metabolites from plants are important for human health due to their antioxidant properties. Light is one of the main factors modulating the biosynthesis of specialized metabolites, determining the cascade response activated by photoreceptors and the consequent modulation of expressed genes and biosynthetic pathways. Recent developments in light emitting diode (LED) technology have enabled improvements in artificial light applications for horticulture. In particular, the possibility to select specific spectral light compositions, intensities and photoperiods has been associated with altered metabolite content in a variety of crops. This review aims to analyze the effects of indoor LED lighting recipes and management on the specialized metabolite content in different groups of crop plants (namely medicinal and aromatic plants, microgreens and edible flowers), focusing on the literature from the last 5 years. The literature collection produced a total of 40 papers, which were analyzed according to the effects of artificial LED lighting on the content of anthocyanins, carotenoids, phenols, tocopherols, glycosides, and terpenes, and ranked on a scale of 1 to 3. Most studies applied a combination of red and blue light (22%) or monochromatic blue (23%), with a 16 h day⁻¹ photoperiod (78%) and an intensity greater than 200 μmol m⁻² s⁻¹ (77%). These treatment features were often the most efficient in enhancing specialized metabolite content, although large variations in performance were observed, according to the species considered and the compound analyzed. The review aims to provide valuable indications for the definition of the most promising spectral components toward the achievement of nutrient-rich indoor-grown products. © 2021 The Authors. *Journal of The Science of Food and Agriculture* published by John Wiley & Sons Ltd on behalf of Society of Chemical Industry.

Supporting information may be found in the online version of this article.

Keywords: horticultural light emitting diodes; indoor farming; vertical farms; secondary metabolites; nutraceuticals; antioxidant

INTRODUCTION

Unlike primary metabolites involved in the basic metabolic functions and development of all plants, secondary or specialized metabolites are not essential for plant life, and their distribution and biological effects can vary between species depending on the specific roles that they play.¹ Nevertheless, specialized metabolites participate in plant ecological interactions by providing protection against biotic and abiotic stresses^{1,2} and promoting environmental adaptation processes.³ Bioactive compounds may also have a role in human nutrition, being used as flavor sources, food additives, or medicines, thanks to their health-promoting properties.⁴

Specialized metabolites are usually classified in groups according to their biosynthetic pathways.¹ Polyphenols are one of the most abundant groups, highly diversified and subdivided into further subgroups such as phenolic acids, flavonoids, and anthocyanins.⁵ These compounds exhibit antioxidant and anti-

inflammatory properties as well as free radicals scavenging or antimicrobial activities.^{6,7} Carotenoids, another important group of metabolites, not only exhibit important properties for human use and health, but also have photo-protective roles in plant cells.⁸ Many specialized metabolites such as anthocyanins, betalains, flavonoids, and carotenoids are also responsible for the pigmentation of flowers, fruits, or other plant tissues,⁹ which determine the attraction or repulsion of animals, thus favoring

* Correspondence to: F Orsini, DISTAL – Department of Agricultural and Food Sciences, Alma Mater Studiorum University of Bologna, viale Fanin 44, 40127 Bologna, Italy. E-mail: f.orsini@unibo.it

a DISTAL – Department of Agricultural and Food Sciences, Alma Mater Studiorum University of Bologna, Bologna, Italy

b FEEM—Foundation Eni Enrico Mattei, Milan, Italy

reproductive success and conservation of the species. The biosynthesis and accumulation of specialized metabolites is mainly affected by light-induced responses.¹⁰

The ability of plants to respond to light is determined by the presence of multiple photoreceptors, proteins capable of sensing different light intensity, quality, direction, and photoperiod,^{11,12} triggering signals that regulate multiple physiological and metabolic responses.¹³ Five classes of photoreceptors have been identified,¹⁴ which enable plants to respond to a broad spectrum of light, from ultraviolet B (UV-B) to far-red wavelengths.¹¹ Phytochromes are the main receptors for red and far-red light spectral regions (600–750 nm), which are involved in plant growth and development, and also promote seed germination and seedling de-etiolation.¹⁵ Conversely, blue wavelengths (400–500 nm) can be absorbed by several photoreceptor classes, namely cryptochromes, phototropins, and zeitelu.¹² Cryptochromes, which are also responsible for the absorption of green light,¹⁶ are involved in plant growth and development and regulate several functions, such as the photoperiodic control of flowering, de-etiolation, or circadian clock adjustment.¹⁷ Phototropins, which are also responsive to ultraviolet A (UV-A) wavelengths (315–400 nm), are primarily involved in plant phototropism, but also mediate fundamental processes such as chloroplast movement and stomatal opening.¹⁸ Finally, the zeitelu class is involved in photoperiodic flowering and circadian clock regulation.¹² Although they are outside the so-called photosynthetic active radiation (PAR) region (400 to 700 nm), UV-B wavelengths (280–315 nm) are also absorbed by a specific photoreceptor called UV resistance locus 8 (UVR8),¹² which is involved in plant survival due to the arrangement of a series of protective gene expression responses.¹⁹

Besides the morphological and physiological responses described above, photoreceptors play a crucial role in stimulating the biochemical pathways of secondary compounds by modulating the expression of specific genes. The biosynthesis of specialized metabolites is strongly related to the wavelength absorbed by the photoreceptor. For example, blue light photoreceptors have been associated with stimulation of photo-protective compounds, such as carotenoids and anthocyanins.²⁰ It has been hypothesized that this plays a specific role in the feedback regulation of photosynthesis and non-photochemical quenching (NPQ), an energy-dissipative response against light stress.^{21,22} On the other hand, phytochrome has been observed to have less impact generally on the production of secondary compounds when subjected to monochromatic red light.²⁰ This limit may be associated with the excessive stimulation of the photoreceptor, caused by far-red unbalancing, with consequences on normal plant development and photosynthesis,^{20,23} probably leading to an uneven distribution of resources among plant metabolic pathways. When plants are exposed to UV-A radiation, an increase in phenolic compounds, especially flavonoids, as a protective mechanism, may be observed.^{24,25} In the case of UV-B, the acclimation of plants to stressing conditions may also induce a specific enhancement of some compounds such as tocopherols or glucosinolates.^{26,27} However, specialized metabolite biosynthetic responses to UV exposure can vary widely depending not only on environmental factors (such as temperature) but also on plant genetic features, even within the green and red phenotypes of the same species.^{25,28}

The specialized metabolism of plants is mainly triggered by stress conditions. Stress can be related to different factors including temperature, nutrition, or drought.^{29–31} Light is also a significant player in the mechanisms of the stress-induced production

of bioactive compounds. Accordingly, plants tend to absorb more energy than necessary for the normal photosynthetic process and carbon fixation. The excess energy must be dissipated efficiently to avoid the production of reactive oxygen species capable of initiating reactions involving radicals that damage proteins, lipids, pigments, or other fundamental bioactive molecules.²¹ However, surplus energy is closely dependent on CO₂ input, which is, in turn, related to the degree of stomatal opening.^{30,31} The imbalance between the amount of irradiated light and energy consumption due to the limited supply of CO₂ is the cause of the increased rate of specialized metabolites in a plant. This phenomenon can be observed easily in plants grown in semi-arid areas, where light, water, and heat stress induce stomatal closure and a consequent higher aromatic value than equivalent plants grown in moderate climates.^{30,31} Plants have developed several methods to dissipate excess energy and produce compounds protecting themselves from photo damage, including the NPQ mentioned above, the re-oxidation of reduced compounds, and the repair of damaged components.²¹

Nowadays, artificial lighting can enable light features to be controlled, with potential effects on plants' primary and secondary metabolic responses.²³ Among the diverse artificial lighting systems available on the market, light-emitting diodes (LEDs) are becoming increasingly popular due to several advantages that make them particularly suitable for horticultural lighting.³² These include the possibility to customize the spectral composition according to plant or photoreceptor characteristics, the relatively high efficiency in converting electrical energy to light, long operating lifespan, and possible integration with digital control systems.^{32–34} Accordingly, LED lamps are today considered the best option for indoor and vertical farming. These farming systems are characterized by tightly controlled and closed environments that require dynamic adjustment of lighting inputs to achieve the best performance in terms of energy consumption, yield, and product quality.³⁵ Concerning qualitative traits of plants, LEDs can affect the biosynthesis of specialized metabolites and antioxidant compounds directly by providing the opportunity to design specific lighting recipes characterized by optimal light spectrum, daily light exposure and intensity.^{36,37} These light features contribute to the development of nutraceuticals and functional foods that promote human health and represent a valuable opportunity for the food industry.³⁸

A relevant body of literature has already demonstrated the potential of LEDs to enhance the production of specialized metabolites in various plants categories,^{39,40} although most research traditionally targeted leafy vegetable crops.^{41,42} The present review paper aims to provide a systematic analysis of the most recent research in this field from 2016 to date. It investigates the effect of different indoor LED treatments on the accumulation of specialized metabolites in different species, going beyond vegetable production to focus on valuable species, namely edible flowers, microgreens, medicinal plants, and aromatic herbs.

MATERIALS AND METHODS

Articles were collected through the Scopus and Web of Science databases, combining specific criteria that include:

- (1) articles published from 2016 to the present;
- (2) articles that referred to four crop categories: edible flowers, microgreens, medicinal plants, and aromatic herbs;
- (3) articles that included LED treatments.

Accordingly, articles that applied other lamp typologies, such as high pressure sodium (HPS) or fluorescent lamps, were excluded from the study. However, if other lamp typologies were used in addition to LED light (e.g. HPS + LED or fluorescent + LED) or only as control treatments, the study was included in the analysis. Moreover, only studies conducted in indoor environments were considered. At the same time, articles that used LEDs as a supplemental lighting sources for natural sunlight were not examined, to focus on indoor and vertical farming possibilities. Furthermore, 14 papers that addressed *in vitro* culture were excluded from the inventory because they presented lab-scale growing conditions that differed from the horticultural approach commonly applied in indoor farms.

The search was conducted in two phases using specific search strings in order to collect a large number of papers. The first phase consisted of a general search by crop categories (edible flowers, microgreens, medicinal plants, aromatic herbs), using the following search string: 'crop category name' AND (light emitting diodes OR indoor), resulting in, e.g., *microgreens AND (light emitting diodes OR indoor)*. After the preliminary search and the screening of the papers retrieved, the investigation was deepened by selecting the most common species for each crop category and replacing them in the search string, e.g., *Brassica oleracea AND (light emitting diodes OR indoor)*. Once relevant articles were selected, information on lighting treatment characteristics and effects on specialized metabolite content was listed in a table for each crop category (see Table S1 in the supplementary material). Although medicinal and aromatic plants are often considered part of the same crop category, commonly referred to as medicinal and aromatic plants (MAPs),⁴³ for the sake of the present study it was preferred to make a clear distinction between the two categories. Medicinal plants were therefore considered to be those plants with a main or unique pharmaceutical and cosmetic function, while aromatic plants are those herbs that are mainly used for flavoring purpose in a general culinary sense, although they have possible medicinal applications. The main information on light treatment included light intensity, light spectrum, and photoperiod. The treatment effect was evaluated for the content of various specialized metabolites, focusing on anthocyanins, carotenoids, phenols, tocopherols, terpenoids, and glycosides. Light treatment features (intensity, spectrum, photoperiod) and phytochemicals (anthocyanins, carotenoids, phenols, tocopherols, terpenoids, glycosides) were evaluated in terms of frequencies to understand the prevalence of cases. With regard to lighting features, given that each study included more than one light treatment with specific light characteristics, each treatment was evaluated as an individual observation. The photoperiod was evaluated grouping the observations into three classes (<12, 12–16, >16 h day⁻¹) and the intensity was divided into two classes (<200, ≥200 μmol m⁻² s⁻¹). Spectral frequencies were assessed by counting each time a spectrum occurred individually (red, blue, green, yellow, orange, far-red, ultraviolet (UV), white) or in combination.

The effects of the different lighting treatments were evaluated within the same study by assigning a score based on the statistical significance of the results. Specifically, the treatment with the highest positive significance received the highest score, and the treatment with the lowest significance received the lowest one. For instance, if a study presented four treatments (T) with three different significance levels (e.g., a, b, c), where T1 = a, T2 = b, and T3 and T4 = c, the best treatment received a score of 3 (T1 = a), the second received 2 (T2 = b), and the third and fourth

treatments received 1 (T3 and T4 = c). If the article presented more than three significance levels (e.g., a, b, c, d, e), scores were assigned according to the number of levels (e.g., 1 to 5), with the best result receiving the highest score. In the case of double letters of significance (e.g., ab), the half score was attributed. When all values were assigned, scores were converted to a scale of 1 to 3. Accordingly, for studies with scores higher than 3, the values were normalized, dividing by the highest score assigned and multiplying by 3. For instance, with five significance levels, each value was divided by 1.66 (e.g., divided by 5 and multiplied by 3) to obtain the score 3 for the treatment showing the higher significance. If treatments in a study had no statistically significant differences, all treatments received the maximum score. If an article reported no significance letters but only the average scores of the treatments, the scores were compared with the results of the treatments and classified from best to worst performance. Some studies analyzed multiple compounds for the same specialized metabolite category (e.g., β-carotene, lutein, and neoxanthin for the carotenoids category). In these cases, the final score was assigned by averaging the scores of the compounds analyzed. The intensity of treatment effects on a 1–3 scale is given in the supplementary materials. Figure 1 shows a flow diagram of the methodological process applied to elaborate the review.

RESULTS AND DISCUSSION

The selection criteria resulted in the selection of 40 papers, ten of which referred to the microgreen crop category, one to edible flowers, 12 to medicinal plants and 17 to the category of aromatic herbs (Fig. 1). Regarding specialized metabolites, phenols were the compounds that were investigated most, being evaluated in 30 articles. Carotenoids and anthocyanins were analyzed in 13 and 10 studies respectively, whereas terpenoids were evaluated in five articles, tocopherols in two articles and glycosides in only one study (Fig. 1). Of the 40 articles, 34 focused on evaluating different spectra, two on photoperiod, two on light intensity, and two on pulsed light (Fig. 1).

Among the selected papers, 389 observations or treatments (plus relative controls) reported the photoperiod. Articles that did not report the adopted photoperiod were excluded from the frequency analysis. Most trials applied a photoperiod ranging from 12 to 16 h day⁻¹ (n = 369, 95%), with 16 h day⁻¹ as the most commonly employed photoperiod (n = 304, 78%). On the other hand, photoperiods lasting more than 16 h day⁻¹ or less than 12 h day⁻¹, were barely used, occurring 14 (4%) and 6 (1%) times, respectively (Table S1). Light intensity was detailed in 391 observations, excluding control treatments. The majority of observations (n = 301, 77%) employed a light intensity equal to or higher than 200 μmol m⁻² s⁻¹, whereas 90 cases (23%) applied an intensity lower than 200 μmol m⁻² s⁻¹ (Table S1). The intensities ranged from a minimum of 7.3 μmol m⁻² s⁻¹ to a maximum of 330 μmol m⁻² s⁻¹.⁴⁵ Spectral treatments were either monochromatic or occurred in several combinations and ratios. In total, 410 observations were counted for light quality features, both considering monochromatic and combined spectra. Counting each spectrum color separately, red and blue were used most, occurring 253 (34%) and 289 (38%) times, respectively. Other wavelengths were observed to a lesser extent: far-red n = 85 (11%), green n = 60 (8%), white n = 28 (4%), yellow n = 24 (3%), UV n = 10 (1%), and orange n = 9 (1%) (Table S1 in the supplementary material). Finally, nearly half of 410 observations (47%) applied a combination of red and blue wavelengths (n = 91,





Figure 1. Flow diagram showing the methodological process of review elaboration.

22%) or red and blue combined with other spectra ($n = 103$, 25%) (Fig. 2). The remaining percentage (53%) mainly applied monochromatic lighting ($n = 208$, 51%), especially blue, or other polychromatic combinations ($n = 8$, 2%), not including red and blue (Fig. 2).

In general, red and blue are the most common light spectral regions used in indoor agriculture, often applied in different ratios to achieve higher crop performance.^{46,47} In fact, although a red wavelength is associated with maximum absorption of chlorophyll pigments, blue light induces stomata opening, leading to greater CO_2 fixation and biomass accumulation,⁴⁸ with optimal consequences on yield. Concerning specialized metabolites, the ability of blue light to induce stomata opening or, conversely, the inability of red wavelengths to induce this effect, may play an interesting role in the modulation of specialized metabolite

content. Moreover, the suboptimal amount of CO_2 due to closed stomata can determine specific stress-induced metabolic responses in plants, leading to an increase in specialized compounds^{30,31,49} and leading to the possible conclusion that blue-less artificial lighting may help to achieve higher accumulations. However, monochromatic red light was observed to have a limited role in specialized metabolites biosynthesis probably due to a phytochrome over-stimulation linked to lack of far-red radiation,²⁰ highlighting the importance of combining red with other wavelengths, especially blue, to optimize specific metabolisms. Indeed, the phytochrome can also be stimulated by blue light, although with lower efficiency compared to far-red,⁵⁰ explaining why plants under pure blue light suffer less than those under monochromatic red, which, however, is ineffective in stimulating cryptochrome chain reactions.⁵¹







	MICROGREENS																		
	ANTHOCYANINS	-	62	-	20	-	22	-	3	24	3	-	5	6	-	-	-	-	-
	○○○	-	47%	-	55%	-	27%	-	33%	25%	0%	-	40%	0%	-	-	-	-	-
	CAROTENOIDS	-	2	-	-	-	2	-	4	33	9	-	26	6	3	3	3	-	3
	○○○	-	0%	-	-	-	50%	-	0%	9%	11%	-	23%	50%	33%	33%	0%	-	0%
	PHENOLS	1	61	1	20	-	21	1	2	6	3	-	20	6	-	-	-	-	-
	○○○	100%	33%	0%	35%	-	29%	0%	0%	17%	0%	-	45%	50%	-	-	-	-	-
	TOCOPHEROLS	-	-	-	-	-	-	-	-	-	-	-	14	6	-	-	-	-	3
	○○○	-	-	-	-	-	-	-	-	-	-	-	29%	33%	-	-	-	-	33%
	TOTAL	1	63	1	20	-	23	1	5	33	9	-	41	6	3	3	3	-	3
	FLOWERS																		
	CAROTENOIDS	-	-	-	-	-	-	-	-	6	-	-	-	-	-	-	-	-	-
	○○○	-	-	-	-	-	-	-	-	100%	-	-	-	-	-	-	-	-	-
	TOTAL	-	-	-	-	-	-	-	-	6	-	-	-	-	-	-	-	-	-
	MEDICINALS																		
	ANTHOCYANINS	-	1	-	-	-	1	-	-	5	-	-	-	-	-	-	-	-	-
	○○○	-	100%	-	-	-	0%	-	-	40%	-	-	-	-	-	-	-	-	-
	CAROTENOIDS	-	2	-	-	-	2	-	3	4	1	-	-	-	-	-	-	-	-
	○○○	-	50%	-	-	-	0%	-	33%	50%	100%	-	-	-	-	-	-	-	-
	PHENOLS	-	8	4	-	1	8	-	8	18	6	-	-	-	-	-	-	-	4
	○○○	-	38%	0%	-	0%	13%	-	25%	39%	50%	-	-	-	-	-	-	-	50%
	TERPENOIDS	-	-	-	-	-	-	-	-	4	-	-	-	-	-	-	-	-	-
	○○○	-	-	-	-	-	-	-	-	25%	-	-	-	-	-	-	-	-	-
	GLYCOSIDES	-	1	-	-	-	1	-	-	3	-	-	-	-	-	-	-	-	-
	○○○	-	0%	-	-	-	0%	-	-	33%	-	-	-	-	-	-	-	-	-
	TOTAL	-	10	4	-	1	10	-	9	22	6	-	-	-	-	-	-	-	4
	AROMATICS																		
	ANTHOCYANINS	-	2	-	-	-	2	-	2	4	6	-	-	-	-	-	-	-	-
	○○○	-	0%	-	-	-	0%	-	50%	25%	17%	-	-	-	-	-	-	-	-
	CAROTENOIDS	-	1	1	1	-	2	-	1	3	6	-	-	-	-	-	-	-	-
	○○○	-	0%	0%	0%	-	0%	-	100%	33%	67%	-	-	-	-	-	-	-	-
	PHENOLS	-	13	2	-	-	8	-	7	19	6	5	1	-	17	-	-	3	-
	○○○	-	15%	0%	-	-	13%	-	0%	26%	67%	0%	0%	-	29%	-	-	67%	-
	TERPENOIDS	-	8	3	-	-	8	-	5	8	-	2	-	-	-	-	-	-	-
	○○○	-	0%	100%	-	-	63%	-	0%	25%	-	20%	-	-	-	-	-	-	-
	TOTAL	-	22	6	1	-	18	-	13	30	6	5	1	-	17	-	-	3	-
	TOTAL OBSERVATIONS	1	95	11	21	1	51	1	27	91	21	5	42	6	20	3	3	3	7

Figure 2. Graphical summary of reviews outcomes showing the absolute frequencies of applied light spectra and percentage of those spectra with a significant positive effect on specialized metabolites (anthocyanins, carotenoids, phenols, tocopherols, glycosides and terpenes) (○○○) in each crop category: microgreens, edible flowers, medicinal plants, and aromatic herbs. (UV = ultraviolet; R = red; B = blue; G = green; Y = yellow; O = orange; FR = far-red; TOTAL OBSERVATIONS = number of treatments reporting a certain spectrum considering 410 observations).

When evaluating the relationship between the spectrum and specialized metabolites for each crop category, a large variability among the spectral combinations that optimized nutritional contents was observed (Fig. 2). This consideration can be associated with a species-specific response across the analyzed crops, and the different reactions of bioactive compounds to light treatment. For instance, anthocyanins may have a higher accumulation under yellow/green/blue light, absorbing these wavelengths to protect chloroplasts,^{20,52} as is clearly visible in Fig. 2 in the microgreens category. In polyphenols such as flavonoids, however, red and blue monochromatic lighting seems to determine an enhancement due to the antioxidant property of these compounds against light-induced ROS accumulation.^{20,53} Polyphenols, and other bioactive compounds of different crop categories seemed, on the other hand, to be fostered when a combination of red and blue light was applied (Fig. 2). However, generalizations could be equivocal and, therefore, need to be avoided. Accordingly, the following paragraphs report an in-depth overview of the main results of the different lighting treatments for each crop category, with an effort toward considering each species and each metabolite separately.

Microgreens

Microgreens are a class of specialty crops characterized by tender and immature greens,^{54,55} which can be obtained from seeds of almost any crop,⁵⁶ including wild species.^{57,58} The edible part of microgreen crops consists of the hypocotyls with cotyledons and first true leaves,⁵⁹ harvested when the cotyledons are fully unfolded and the plant has a minimum height of 5 cm.⁶⁰ The concept of this specialty crop originated in the late 1980s in San Francisco, USA,⁵⁴ and has since been gaining popularity due to its elevated nutritional and sensory properties.⁶¹ Microgreens are appreciated by consumers for their appearance, taste, and texture,⁶²⁻⁶⁴ and for their high levels of health-promoting phytochemicals.^{62,65} Microgreens are also classified as 'functional food'^{63,66} due to their higher content of vitamins, minerals, and bioactive compounds compared with adult plants,^{62,63} and their lower nitrate content.⁶⁷ These characteristics, and the ease with which microgreens can be cultivated, mean that microgreens can be a valuable nutrient-rich food source even in remote locations with difficult access to fresh vegetables.⁶⁸ Although this crop can be grown in a variety of scenarios, including outdoor, greenhouse, and indoor conditions, cultivation in controlled environments with artificial lighting represents the most common situation.^{55,63} Indoor farming technology is suggested for the production of continuous and uniform high-quality microgreens using LED light as a specific factor to modify their metabolism and obtain tailored crops rich in bioactive compounds.^{48,67}

The wide application and beneficial effects of combined red and blue light seem to be accurate, particularly in the cultivation of microgreens, presenting significant positive responses when applied together and combined with other spectra (Fig. 2). However, the effect of red and blue light on the synthesis of specialized metabolites can vary depending on plant species. Concerning the effects on carotenoids, Lobiuc *et al.*⁴⁸ found that total carotenoid content (TCC) in *Ocimum basilicum* microgreens was not affected by different red and blue ratios compared with white light, as also observed by Meas and colleagues⁴⁷ in *Amaranthus cruentus*. In *B. oleracea* microgreens (Table 1), the carotenoid content increased under a red and blue combination supplemented with far-red, in contrast with *Brassica rapa*, where carotenoids were increased when a green or yellow irradiation

was added to a red, blue, and far-red spectrum.⁵⁶ However, it should be noted that light quality is not the only parameter that affects TCC. Craver and colleagues⁶³ found that TCC in microgreens of *B. rapa* var. Japonica and *Brassica juncea* (Table 1) may also depend on light intensity, showing a lower TCC value when light intensity is increased from 105 to 315 $\mu\text{mol m}^{-2} \text{s}^{-1}$. Indeed, it is likely that carotenoid biosynthesis is enhanced at higher light intensities due to their photoprotective functions. However, such effects may be hindered by pigment photodegradation and the dilution effect of increased plant growth,⁶⁹ possibly resulting in a higher concentration of carotenoids in plant tissues exposed to lower light intensities.

In contrast with the observations for carotenoids, Lobiuc *et al.*⁴⁸ showed that contents of anthocyanins and phenolic acids in *O. basilicum* microgreens are primarily influenced by the proportion of red and blue. In particular, the content of phenolic acids in a red cultivar (Red Rubin) of *O. basilicum* was higher under a light spectrum consisting of a red:blue ratio (R:B) of 0.5:1. The effectiveness of blue light in increasing phenol content might be determined by a protective mechanism produced by cytochrome P450 against the accumulation of reactive oxygen species (ROS).⁷⁰ On the other hand, anthocyanin content showed the best results under a R:B = 2:1. However, in microgreens of *Brassica oleracea* var. *Gongylodes* (Table 1), the total phenolic content (TPC), seemed to be enhanced when far-red was added to a red and blue background (R:FR:B = 84:7:9).⁶³ Finally, in *A. cruentus* microgreens, red and blue light (R:B = 2.3:1) appeared to increase anthocyanin content compared with monochromatic red or blue light, probably due to the ability of the combined wavelengths to induce the expression of anthocyanin regulatory genes.^{48,71}

Monochromatic lighting was, however, shown to increase specialized metabolite content in some microgreen species. Zhang and colleagues⁵⁹ observed that monochromatic blue and UV-A light could be optimal lighting sources to produce soybean (*Glycine max*) microgreens rich in phenolic compounds. In fact, the absorption of blue and UV-A light, and the consequent metabolic responses, are regulated by the same receptor, the phototropin,¹⁸ which has been linked to the production of phenols as photoprotective barriers against excess light exposition and free radicals, mainly when applied monochromatically.^{20,72} Ultraviolet light potentialities were also observed,⁷³ with significant increases in carotenoids in *B. juncea* when UV was added to a red, blue and far-red spectrum (Table 1). Carotenoid biosynthesis seems to be regulated by the physiological response triggered by different blue and UV photoreceptors (e.g., UVR8). In contrast, polyphenol biosynthesis is mainly related to the plant response to UV radiation, acting as antioxidants and UV-absorbing compounds.⁷⁴ In fact, UV-A stimulates the expression of UV-protective genes and the associated accumulation of phenols.⁷³ In the work by Brazaitytė and colleagues,⁷³ the same trend observed for carotenoids was also observed for tocopherols and phenols, while the anthocyanin content did not show a significant increase after UV addition. Samuolienė *et al.*⁴⁵ suggested a mixed red and far-red spectrum as the most efficient for enhancing tocopherols in *B. juncea* (Table 1), while in *Beta vulgaris* and *Petroselinum crispum* the highest tocopherol content was associated with a red and far-red light supplemented with a blue component (resulting in an increase of +16% with R:B = 5:1 and + 33% with R:B = 2:1). Further details of the cases analyzed are reported in Table S1, while Fig. 2 summarizes the main light spectra promoting specialized metabolites in microgreens.

Table 1. Within-studies comparison of anthocyanins, carotenoids, phenols and tocopherols content in microgreens of *Brassica* grown under different LED light features (light spectrum, intensity, and photoperiod)

REFERENCES	SPECIES	LIGHTING TREATMENT		SPECIALIZED METABOLITES					
		Standard LED light features	Specific LED light features	Anthocyanins	Carotenoids	Phenols	Tocopherols		
Brazaitytė et al., 2019 ⁷³	<i>Brassica juncea</i> (Red Lion)	10 h day ⁻¹ , 300 μmol m ⁻² s ⁻¹ , R ⁸ :B ⁸ :FR ^a	UV ^a -A (366 nm) = 12.4 μmol m ⁻² s ⁻¹	○ ^b	○	○	○	○	○
			UV (390 nm) = 12.4 μmol m ⁻² s ⁻¹	○	○	○	○	○	○
			UV (402 nm) = 12.4 μmol m ⁻² s ⁻¹	○	○	○	○	○	○
Craver et al., 2017 ⁶³	<i>Brassica oleracea</i> (Gongylodes)	16 h day ⁻¹ , 300 μmol m ⁻² s ⁻¹ , R:B:FR	Standard light (Control)	○	○	○	○	○	○
			UV-A (366 nm) = 12.4 μmol m ⁻² s ⁻¹	○	○	○	○	○	○
			UV (390 nm) = 12.4 μmol m ⁻² s ⁻¹	○	○	○	○	○	○
			UV (402 nm) = 12.4 μmol m ⁻² s ⁻¹	○	○	○	○	○	○
			Standard light (Control)	○	○	○	○	○	○
			105 μmol m ⁻² s ⁻¹	○	○	○	○	○	○
			210 μmol m ⁻² s ⁻¹	○	○	○	○	○	○
			315 μmol m ⁻² s ⁻¹	○	○	○	○	○	○
			105 μmol m ⁻² s ⁻¹	○	○	○	○	○	○
			210 μmol m ⁻² s ⁻¹	○	○	○	○	○	○
B. oleracea (Gongylodes)	16 h day ⁻¹ , R:FR:B = 84:7:9	105 μmol m ⁻² s ⁻¹	○	○	○	○	○		
		315 μmol m ⁻² s ⁻¹	○	○	○	○			
B. oleracea (Gongylodes)	16 h day ⁻¹ , R:G ^b :B = 74:18:8	105 μmol m ⁻² s ⁻¹	○	○	○	○	○		
		210 μmol m ⁻² s ⁻¹	○	○	○	○			
<i>Brassica rapa</i> (Japonica)	16 h day ⁻¹ , R:B = 87:13	105 μmol m ⁻² s ⁻¹	○	○	○	○	○		
		210 μmol m ⁻² s ⁻¹	○	○	○	○			
B. rapa (Japonica)	16 h day ⁻¹ , R:FR:B = 84:7:9	105 μmol m ⁻² s ⁻¹	○	○	○	○	○		
		210 μmol m ⁻² s ⁻¹	○	○	○	○			
B. rapa (Japonica)	16 h day ⁻¹ , R:G:B = 74:18:8	105 μmol m ⁻² s ⁻¹	○	○	○	○	○		
		210 μmol m ⁻² s ⁻¹	○	○	○	○			
<i>B. juncea</i> (Garnet Giant)	16 h day ⁻¹ , R:B = 87:13	105 μmol m ⁻² s ⁻¹	○	○	○	○	○		
		210 μmol m ⁻² s ⁻¹	○	○	○	○			
<i>B. juncea</i> (Garnet Giant)	16 h day ⁻¹ , R:FR:B = 84:7:9	105 μmol m ⁻² s ⁻¹	○	○	○	○	○		
		210 μmol m ⁻² s ⁻¹	○	○	○	○			
<i>B. juncea</i> (Garnet Giant)	16 h day ⁻¹ , R:G:B = 74:18:8	105 μmol m ⁻² s ⁻¹	○	○	○	○	○		
		210 μmol m ⁻² s ⁻¹	○	○	○	○			
Samuoliėnė et al., 2019 ⁵⁶	<i>B. rapa</i> (Japonica)	16 h day ⁻¹ , 300 μmol m ⁻² s ⁻¹ , R: B = 5:1	FR = 4 μmol m ⁻² s ⁻¹	○	○	○	○	○	
			FR = 4 μmol m ⁻² s ⁻¹ , G = 15 μmol m ⁻² s ⁻¹	○	○	○	○	○	
			FR = 4 μmol m ⁻² s ⁻¹ , Y = 15 μmol m ⁻² s ⁻¹	○	○	○	○	○	

REFERENCES	SPECIES	LIGHTING TREATMENT			SPECIALIZED METABOLITES				
		Standard LED light features	Specific LED light features		Anthocyanins	Carotenoids	Phenols	Tocopherols	
	<i>B. oleracea</i> (Green)	16 h day ⁻¹ , 300 μmol m ⁻² s ⁻¹ , R; B = 5:1	FR = 4 μmol m ⁻² s ⁻¹			○○○			
			FR = 4 μmol m ⁻² s ⁻¹ , G = 15 μmol m ⁻² s ⁻¹			○○○			
			FR = 4 μmol m ⁻² s ⁻¹ , Y = 15 μmol m ⁻² s ⁻¹			○			
	<i>B. oleracea</i> (Delivery purple)	16 h day ⁻¹ , 300 μmol m ⁻² s ⁻¹ , R; B = 5:1	FR = 4 μmol m ⁻² s ⁻¹ , O = 15 μmol m ⁻² s ⁻¹			○○○			
			FR = 4 μmol m ⁻² s ⁻¹			○○○			
			FR = 4 μmol m ⁻² s ⁻¹ , G = 15 μmol m ⁻² s ⁻¹			○○○			
Samuoliené <i>et al.</i> , 2017 ⁴⁵	<i>B. juncea</i> ^c (Red Lion)	16 h day ⁻¹ , R (660 nm) = 170 μmol m ⁻² s ⁻¹ , FR = 2.5 μmol m ⁻² s ⁻¹	FR = 4 μmol m ⁻² s ⁻¹ , Y = 15 μmol m ⁻² s ⁻¹			○			
			FR = 4 μmol m ⁻² s ⁻¹ , O = 15 μmol m ⁻² s ⁻¹			○			
			R (638 nm) = 130 μmol m ⁻² s ⁻¹ B = 25 μmol m ⁻² s ⁻¹ , R (638 nm) = 105 μmol m ⁻² s ⁻¹			○○○			○○○
			B = 50 μmol m ⁻² s ⁻¹ , R (638 nm) = 80 μmol m ⁻² s ⁻¹			○○			○○
			B = 75 μmol m ⁻² s ⁻¹ , R (638 nm) = 55 μmol m ⁻² s ⁻¹			○○○			○○
			B = 100 μmol m ⁻² s ⁻¹ , R (638 nm) = 30 μmol m ⁻² s ⁻¹			○○			○

^a UV = ultraviolet; R = red; B = blue; G = green; Y = yellow; O = orange; FR = far-red.

^b Three dots represent the best performance among all treatments of the same study, one dot represents the worst performance among all treatments of the same study. In case of study treatments with similar effects, three dots were assigned to all treatments.

^c Confronted with *Beta vulgaris* and *Petroselinum crispum*.

Edible flowers

In addition to microgreens, edible flowers are another specialty crop that has recently gained interest on the market, as they are appreciated for their appearance and are therefore mainly employed in high-end cuisine.⁷⁵ In general, the whole flower can be consumed, although only some parts should be used in some species. For instance, in *Tulipa* or *Chrysanthemum*, only the petals are edible, while in *Bellis perennis* only the buds can be consumed.⁷⁶ According to Lu *et al.*,⁷⁷ there are 97 families and about 180 species of edible flowers. Beyond being used as ornamental plants, edible flowers also have potential beneficial effects for human health due to their elevated content of specialized metabolites, especially antioxidant compounds.⁷⁷ Several studies investigated various edible flower species as sources of beneficial bioactive compounds from either nutritional or health perspectives, including *Sambucus nigra*, *Cichorium intybus* L., *Hedysarum coronarium*,⁷⁸ *Tropaeolum majus*, *Tagetes erecta*, *Spilanthes oleracea*,⁷⁹ *Antirrhinum majus*, and *Viola wittrockiana*.⁸⁰

Although many studies in recent years have focused on assessing the content of specialized metabolites in edible flowers, only a few studies have investigated the effects of artificial indoor LED lighting on the accumulation of bioactive compounds. In particular, Kopsell *et al.*⁸¹ aimed to understand the potential enhancement of carotenoids in petal tissue of two different cultivars of *Tagetes tenuifolia* under different light treatments (Table S1). The experimental lighting included a fluorescent treatment and different blue and red LED compositions (R:B in the range 10:90, 20:80, and 40:60).⁸¹ Results showed that LED lighting treatments positively affected carotenoid content in both cultivars. In particular, carotenoid content in the 'Lemon Gem' cultivar had the best performance under R:B = 1:4 (Fig. 2), while lower accumulation was associated with fluorescent lighting.⁸¹ Moreover, the same positive trend was observed in 'Tangerine Gem' cultivar for carotenoid accumulation, although no relevant differences were found specifically for α -carotene, β -carotene and lutein among the different light treatments.⁸¹

Medicinal plants

Specialized metabolites are considered to be the key source of pharmaceutical and curative properties of medicinal plants.³ The commercial value of these plants is associated with their elevated bioactive compound content,⁸² which determines the effects on human health, including antimalarial, antidiabetic, hepato-protective, anti-ulcer, anti-inflammatory, and antimicrobial properties.⁸³ The concentration of specialized metabolites can be improved by growing them in a controlled environment, resulting in lower costs for extraction of the active principles. Furthermore, their facilitated crop management and harvesting suggest a higher profit margin when cultivated in vertical farms.^{84,85} Several studies revealed an association between the accumulation of specialized metabolites and artificial LED treatments in medicinal plants, although they showed different responses depending on species and experimental characteristics (Table S1 and Fig. 2). For instance, *Crepidiastrum denticulatum* is a medicinal plant the extracts of which have anti-oxidative properties.⁸⁶ According to Park *et al.*,⁸⁷ the *C. denticulatum* phenolic compound content did not show significant differences under monochromatic LED treatments compared with fluorescent light as a control. However, an increased accumulation of phenolic compounds in the shoot was observed under certain spectral compositions (e.g. a R:G:B spectrum with ratios of 8:1:1).⁸⁷ Similarly, Bae *et al.*⁸⁶ analyzed *C. denticulatum* phenolic content in response to different

ratios of far-red combined with red LED light and observed the highest phenolic levels when a red:far-red (R:FR) ratio between 0.7:1 and 1.2:1 was provided.

Medicago sativa, another medicinal plant known for its phyto-pharmacological potential,⁸⁸ was analyzed by Fiutak *et al.*,⁸⁹ demonstrating a higher accumulation of phenolic compounds and carotenoids under red, green, and blue LED lighting compared with white LED or natural sunlight conditions. Accordingly, these spectral wavelengths play a significant role in plant growth phases, with blue particularly affecting the maturity of chloroplasts and photosynthetic activity.⁸⁹ Similarly, the total amount of phenols and anthocyanins in *Lepidium sativum*, a small medicinal plant belonging to the Cruciferae family, was significantly increased under mixed red and blue LED light compared with the sunlight control.⁹⁰ Thus, it showed an increase in phenol and anthocyanin content by up to +47% and +33% under R: B = 1.5:1 and R:B = 9:1, respectively. A study on *Digitalis purpurea* also highlighted that a red and blue LED light (R:B = 1:4) increased the carotenoid content, while anthocyanins were instead higher under monochromatic blue and red lights.⁹¹ The study also investigated glycoside content in *D. purpurea*, where the highest concentrations were observed under R:B = 1:4,⁹¹ probably due to the role of blue photoreceptors in the biosynthesis and accumulation of these metabolites.⁹² Red and blue light (with R:B of 9:1, 2.3:1, and 0.4:1) also increased the phenol content in *Salvia miltiorrhiza*,⁹³ compared with monochromatic white, red or blue LED light. The combination of red and blue light was shown to upregulate the expression of many genes encoding key enzymes in the phenylpropanoid pathway and revealed transcription levels of genes consistent with the accumulation of some phenols compounds (e.g., rosmarinic acid).⁹³ Accumulation of carotenoids and phenols were also monitored in *Perovskia* sp.⁹⁴ with an observed increase in carotenoids under a red and blue light (R:B = 2.3:1), whereas phenols were mainly enhanced by monochromatic blue or red lights. It may therefore be advanced that monochromatic light may also become a stress factor for plants affecting the accumulation of metabolites.⁹⁵

Finally, the use of *Astragalus membranaceus* has been investigated in several studies with different LED spectra. Choi *et al.*⁴⁴ cultivated *A. membranaceus* under monochromatic LEDs with different irradiances, and obtained the highest phenol content under the blue treatment at $7.3 \mu\text{mol m}^{-2} \text{s}^{-1}$, although blue light presented the lowest intensity used, as compared with red ($28.7 \mu\text{mol m}^{-2} \text{s}^{-1}$), green ($24.9 \mu\text{mol m}^{-2} \text{s}^{-1}$), and white ($49.8 \mu\text{mol m}^{-2} \text{s}^{-1}$) light. However, while in Choi *et al.*,⁴⁴ white LED proved to be the least efficient in increasing the phenol content of *A. membranaceus*, the same plant showed higher values of phenols under the white treatment than when monochromatic blue and red LEDs were used in Jeong *et al.*,⁹⁶ considering an intensity around $30\text{--}40 \mu\text{mol m}^{-2} \text{s}^{-1}$ for all spectra. Similarly, white LED appeared to be most effective in increasing phenol content in *Agastache rugosa* compared to red or blue monochromatic light,⁹⁷ possibly due to the higher penetration ability of the white spectrum within the canopy, leading to increased expression of genes involved in phenylpropanoids biosynthesis.⁹⁸ In Livadariu *et al.*,⁹⁹ different monochromatic LED lights showed similar effects on phenol content in *Cannabis sativa*. This plant was also investigated by Namdar *et al.*,¹⁰⁰ who applied LED treatment in addition to fluorescent or HPS light throughout the crop cycle (in vegetative or flowering stage). According to these results, R:B = 4:1 treatment during the vegetative stage or for 35 days during flowering mainly increased phenols concentration in

Table 2. Within studies comparison of anthocyanin, carotenoid, phenol, and tocopherol content in *Ocimum basilicum* grown under different LED light features (light spectrum, intensity, and photoperiod)

REFERENCES	SPECIES	LIGHTING TREATMENT		SPECIALIZED METABOLITES				
		Standard LED light features	Specific LED light features	Anthocyanins	Carotenoids	Phenols	Terpenoids	
Bantis <i>et al.</i> , 2016 ¹¹¹	<i>O. basilicum</i> (Lettuce leaf)	14 h day ⁻¹ , 200 μmol m ⁻² s ⁻¹	UV ^a :B ^a :G ^a :R ^a :FR ^a = 0:12:19:61:8	○	○	○	○	
			UV:B:G:R:FR = 0:8:2:65:25	○	○	○	○	
<i>O. basilicum</i> (Red rubin)		14 h day ⁻¹ , 200 μmol m ⁻² s ⁻¹	UV:B:G:R:FR = 0:14:16:53:17	○	○	○	○	
			UV:B:G:R:FR = 1:20:39:35:5	○	○	○	○	
			Fluorescent (Control)	○	○	○	○	
			UV:B:G:R:FR = 0:12:19:61:8	○	○	○	○	
			UV:B:G:R:FR = 0:8:2:65:25	○	○	○	○	
Hosseini <i>et al.</i> , 2018 ¹⁰⁶	<i>O. basilicum</i> (Green)	16 h day ⁻¹ , 250 μmol m ⁻² s ⁻¹	UV:B:G:R:FR = 0:14:16:53:17	○	○	○	○	
			UV:B:G:R:FR = 0:14:16:53:17	○	○	○	○	
			UV:B:G:R:FR = 1:20:39:35:5	○	○	○	○	
			Fluorescent (Control)	○	○	○	○	
			UV:B:G:R:FR = 1:20:39:35:5	○	○	○	○	
Lin <i>et al.</i> , 2021 ¹⁰⁵	<i>O. basilicum</i> (Purple)	16 h day ⁻¹ , 250 μmol m ⁻² s ⁻¹	Monochromatic R	○	○	○	○	
			Monochromatic B	○	○	○	○	
			Monochromatic W	○	○	○	○	
			R:B = 1:1	○	○	○	○	
			R:B = 2.3:1	○	○	○	○	
Naznin <i>et al.</i> , 2019 ¹¹³	<i>Ocimum × africanum</i> (Lemon Basil)	16 h day ⁻¹ , 200 μmol m ⁻² s ⁻¹	Monochromatic R	○	○	○	○	
			Monochromatic B	○	○	○	○	
			Monochromatic W	○	○	○	○	
			R:B = 1:1	○	○	○	○	
			R:B = 2.3:1	○	○	○	○	
Pennisi <i>et al.</i> , 2020 ¹¹⁰	<i>O. basilicum</i> (Superbo)	16 h day ⁻¹ , R:B = 3:1	R:G:B = 4:1:1	○	○	○	○	
			R:G:B = 2:1:1	○	○	○	○	
			R:G:B = 1:1:1	○	○	○	○	
			R:G:B = 4:1:1	○	○	○	○	
			R:G:B = 2:1:1	○	○	○	○	
Pennisi <i>et al.</i> , 2019b ⁴⁶	<i>O. basilicum</i> (Superbo)	16 h day ⁻¹ , 215 μmol m ⁻² s ⁻¹	R:G:B = 1:1:1	○	○	○	○	
			R:B = 83:17	○	○	○	○	
			R:B = 91:9	○	○	○	○	
			R:B = 95:5	○	○	○	○	
			R = 100	○	○	○	○	

Table 2. Continued

REFERENCES	SPECIES	LIGHTING TREATMENT		SPECIALIZED METABOLITES				
		Standard LED light features	Specific LED light features	Anthocyanins	Carotenoids	Phenols	Terpenoids	
Rihan et al., 2020 ¹²¹	<i>O. basilicum</i> (Maggie)	16 h day ⁻¹ , 300 μmol m ⁻² s ⁻¹	R:B = 4:1 R:B = 1:1.5 R:B = 1:1.4 R:B = 1:1			○	○○○○	
Taulavuori et al., 2016 ¹⁰⁷	<i>O. basilicum</i> (Genovese Gigante)	16 h day ⁻¹ , 300 μmol m ⁻² s ⁻¹	Natural light + HPS (Control) 48 days under enhanced B 36 days under enhanced B 24 days under enhanced B 12 days under enhanced B 0 days under enhanced B			○○○○	○○○○	
Zotov et al., 2020 ¹³²	<i>O. basilicum</i> (Cinnamom)	14 h day ⁻¹ , 120 μmol m ⁻² s ⁻¹ , 30 days	R:B:G:FR = 33:20:44:3 R:B:G:FR = 18:57:24:1 R:B:G:FR = 65:9:23:3 R:B:G:FR = 57:23:18:2 R:B:G:FR = 33:20:44:3 R:B:G:FR = 18:57:24:1 R:B:G:FR = 65:9:23:3 R:B:G:FR = 57:23:18:2			○○○○	○○○○	
	<i>O. basilicum</i> (Cinnamom)	14 h day ⁻¹ , 120 μmol m ⁻² s ⁻¹ , 50 days				○○○○	○○○○	

^a UV, ultraviolet; R, red; B, blue; G, green; Y, yellow; O, orange; FR, far-red.

^b Three dots represent the best performance among all treatments of the same study, one dot represents the worst performance among all treatments of the same study. In case of study treatments with similar effects, three dots were assigned to all treatments.

inflorescence tissues. Terpenoids in *C. sativa* were also highest when R:B = 4:1 was supplied at the vegetative stage.¹⁰⁰

Aromatic herbs

Specialized metabolites play a fundamental role in the aromatic function of several herbs, promoting both the characteristic odors and flavors that are particularly appreciated not only for culinary purposes but also for industrial purposes. The industrial applications of aromatic plants include the production of fragrances and preservatives for the food processing industry, and perfumes and skin products in the cosmetics sector.¹⁰¹ Important uses also include pharmaceutical and livestock feeds due to their antioxidant, antimicrobial, and anticoccidial properties favoring both human and animal health.^{102,103}

Among the aromatic herbs, sweet basil (*O. basilicum*) is one of the most cultivated, presenting not only a wide application in cooking but also remarkable medicinal use.¹⁰⁴ Several studies from 2016 to date have examined LED lighting effects on specialized metabolites of different cultivars of basil, and overall confirmed a significant effect of artificial lighting treatments to increase various bioactive compounds (Table 2). According to Lin *et al.*,¹⁰⁵ cv. 'Dark opal' and 'Caesar' increased their anthocyanin, phenol, and carotenoid content under an equal combination of red, green, and blue LED light (R:G:B = 1:1:1). Moreover, comparing to 'Dark opal', 'Caesar' cultivar also demonstrated an enhanced production of phenols and carotenoids by augmenting the red proportion in the LED combination (R:G:B = 4:1:1)¹⁰⁵ (Table 2). In purple *O. basilicum*, both white LED light and a combination of red and blue (R:B = 2.3:1) significantly boosted the anthocyanin content.¹⁰⁶ Similarly, R:B = 2.3:1 increased the phenol content in green basil (Table 2). On the other hand, white LED light did not show the same effect for the green cultivar, where a monochromatic blue light was more efficient in fostering phenol content. In fact, no effect of 16 h day⁻¹ exposure of blue light for different periods (0 to 48 days) on the phenol content of basil was observed¹⁰⁷ (Table 2). The authors reported that the selected light intensity for basil in the treatments (300 $\mu\text{mol m}^{-2} \text{s}^{-1}$) may also be responsible for the unvaried results and non-reactive responses, as the light saturation of basil was evaluated considerably higher (e.g., more than 1000 $\mu\text{mol m}^{-2} \text{s}^{-1}$).¹⁰⁸ In Pennisi *et al.*,⁴⁶ the highest phenol concentration in basil in response to red and blue light was associated with R:B = 2:1 or 3:1, supporting the assumption that an increased red component in the mixture may enhance antioxidants production.¹⁰⁹ In contrast, light intensity (e.g. in the range of 100 to 300 $\mu\text{mol m}^{-2} \text{s}^{-1}$) did not increase phenols in basil leaves under red and blue irradiation (R:B = 3:1)¹¹⁰ (Table 2). Bantis *et al.*¹¹¹ evaluated the total phenol content under different UV, blue, green, red, and far-red combinations in two cultivars of *O. basilicum*, 'Lettuce leaf' and 'Red rubin', and observed the best performances for both cultivars in the case of UV:B:G:R:FR = 1:20:39:35:5. Using a broad spectrum of light with a higher blue percentage could possibly affect phenol production positively, due to stimulation of the PAL enzyme (Phenylalanine Ammonia Lyase), a key enzyme in the phenylpropanoid pathway.¹¹² However, the phenol content in the 'Red Rubin' cultivar showed relevant enhancements also under increased red and far-red portions (e.g., UV:B:G:R:FR = 0:14:16:53:17 and 0:12:19:61:8) (Table 2). Finally, Naznin *et al.*¹¹³ studied the carotenoid accumulation in another basil species, known as Lemon Basil (*Ocimum x africanum*), and observed a considerable increase after application of red and blue light (R:B = 5:1) instead of a monochromatic red light (Table 2).

The effect of LED light treatments on specialized metabolite concentration has also been evaluated in other aromatic plants, with consistent beneficial responses (Table S1). For instance, in *Allium fistulosum*, white LED proved to be the most effective treatment for carotenoid accumulation compared with monochromatic light treatments.¹¹⁴ In contrast, *Allium sativum* showed a significantly higher accumulation of phenols under monochromatic red light applied with a photoperiod of 16 h day⁻¹ compared to monochromatic blue, green and white light.¹¹⁵ In the case of *Coriandrum sativum*, monochromatic blue allowed the highest increase in phenol content compared with monochromatic red and green, as well as with red and blue, and red, blue, and far-red combinations.¹¹⁶ Both in *A. sativum* and *C. sativum*, monochromatic green light seems to have a limited effect on plant phenol accumulation. The poor effects of monochromatic green light have already been observed by other authors, although there is insufficient literature investigating this specific wavelength, which prevents general conclusions until there has been further investigation.²⁰ In lemon balm (*Melissa officinalis*), phenolic compounds were increased by the simultaneous application of drought stress and a red and blue light (R:B = 2.3:1).¹¹⁷ It is possible that the drought stress and the consequent ROS formation¹¹⁸ could be balanced by an increase in antioxidant production, particularly stimulated by the R:B treatment.¹¹⁷ Alternatively, *Anethum graveolens* presented similar levels of phenol content regardless of the light treatment with different ratios of red, blue, orange, and green light, suggesting that the changes in the red:blue ratios among combinations may have been too narrow to alter phenol accumulation.¹¹⁹ However, in the same study, a greater increase in terpenoids occurred when the red component was highest (70% of the spectrum) and the blue component was lowest (10%).¹¹⁹ In two *Thymus* species, namely *T. carmanicus* and *T. migricus*, terpenoids were enhanced by monochromatic red light.¹²⁰ Only in *T. migricus*, a combination of red and blue light (R:B = 2.3:1) also increased terpenoid concentration, suggesting that metabolite biosynthesis may be associated with increased stress from red light exposure.¹²⁰ Similarly, a red and blue combination (R:B = 1:1.4, with blue at 435 nm) proved to be efficient in enhancing terpenoid content in basil.¹²¹ Finally, in *Mentha spicata*, monochromatic red and green were more efficient in increasing terpenoids than monochromatic blue light.¹²² The main light spectra that enhance anthocyanins, carotenoids, phenols, and terpenoids are listed in Fig. 2.

CONCLUSIONS

This review highlighted the potential of indoor LED treatments to enhance specialized metabolite content in microgreens, edible flowers, medicinal plants, and aromatic herbs. Half of the studies applied a combination of light spectra, which contained red and blue light in different proportions. In some species, lighting combinations, particularly the addition of far-red or green to the lighting mixture, resulted in an improved synthesis of specialized metabolites. However, in other cases, best performances were also observed under monochromatic red or blue lights. Most trials adopted a photoperiod of between 12 and 16 h day⁻¹, often applying 16 h day⁻¹ of lighting duration. Among specialized metabolites, more than half of the articles analyzed phenols content, a large number of studies investigated carotenoids and anthocyanins. At the same time, terpenoids, tocopherols and glycosides were evaluated only in few studies. The results of each crop category highlights not only a species-specific effect

of the lighting treatments, but also a different response depending on the analyzed specialized metabolites within the same species. For this reason, giving specific recommendations on the best lighting treatment to be used in each crop category may be equivocal. In general, red and blue light, alone or combined, performed well on different crops and metabolites. However, the application of indoor LED lighting on a commercial scale to enhance product quality should consider the specific responses of different crops and compounds. In this framework, data summarized in Table S1 in the supplementary material may represent a useful tool for producers to identify the best species-light treatment combination depending on the specialized metabolite to be enhanced.

Beside its commercial usefulness, the present research also aimed to provide suggestions for future research developments. To the best of our knowledge, only one article from 2016 evaluated the phytochemical content in edible flower tissues in response to LED light. Edible flower cultivation using artificial lighting should be further investigated, not only because of the potential nutraceutical improvements that can be achieved with this technology but also because of the high economic value that these products have in their expanding market.¹²³ Similarly, although some studies have observed that the application of supplemental UV light enhances bioactive compounds,^{59,73} its application in indoor farming is still under-researched (mainly due to safety issues and regulations), so further investigation is needed for future sector development. *In vitro* cultivation with LED lighting may also represent an opportunity for the sector, especially for medicinal plants. Moreover, *in vitro* cultivation could also allow for the rapid and large-scale production of specialized metabolites to be applied in the pharmaceutical and cosmetic industries.¹²⁴⁻¹³¹

ACKNOWLEDGEMENTS

The research leading to this publication has received funding from the European Union's Horizon 2020 research and innovation programme under grant agreement No. 862663. The publication reflects the authors' view. The Research Executive Agency (REA) is not liable for any use that may be made of the information contained therein. The authors also gratefully acknowledge the financial support provided by the Foundation Eni Enrico Mattei (FEEM) and the contribution of Dario Migliorini to the data collection. The research leading to this publication has been supported by collaboration between the University of Bologna Alma Mater Studiorum and the Fondazione Eni Enrico Mattei. Open Access Funding provided by Università di Bologna within the CRUI-CARE Agreement. [Correction added on 19 May 2022, after first online publication: CRUI-CARE funding statement has been added.]

AUTHOR CONTRIBUTIONS

E.A. made contributions to the conception and design, analysis, and interpretation of the data and revised the manuscript critically. I.Z. and L.C. made substantial contributions to the acquisition of data and manuscript drafting. G.P. and F.O. made substantial contributions to the conception and design of the research and critically revised the manuscript. I.P. has critically revised the manuscript. S.Q. and G.P.G. supervised the research and reviewed the manuscript.

SUPPORTING INFORMATION

Supporting information may be found in the online version of this article.

REFERENCES

- Croteau R, Kutchan TM and Lewis NG, Natural products (secondary metabolites), in *Biochemistry and Molecular Biology of Plants*, ed. by Buchanan BB, Gruissem W and Russel RL. American Society of Plant Physiologists, Rockville, MD, pp. 1250–1319 (2000).
- Pagare S, Bhatia M, Tripathi N, Pagare S and Bansal YK, Secondary metabolites of plants and their role: overview. *Curr Trends Biotechnol Pharm* **9**:293–304 (2015).
- Jamwal K, Bhattacharya S and Puri S, Plant growth regulator mediated consequences of secondary metabolites in medicinal plants. *J Appl Res Med Aromat Plants* **9**:26–38 (2018).
- Ramakrishna A and Ravishankar GA, Influence of abiotic stress signals on secondary metabolites in plants. *Plant Signal Behav* **6**:1720–1731 (2011).
- Tsao R, Chemistry and biochemistry of dietary polyphenols. *Nutrients* **2**:1231–1246 (2010).
- Ouzounis T, Razi Parjikolaei B, Fretté X, Rosenqvist E and Ottosen CO, Predawn and high intensity application of supplemental blue light decreases the quantum yield of PSII and enhances the amount of phenolic acids, flavonoids, and pigments in *Lactuca sativa*. *Front Plant Sci* **6**:19 (2015).
- Zhang H and Tsao R, Dietary polyphenols, oxidative stress and antioxidant and anti-inflammatory effects. *Curr Opin Food Sci* **8**:33–42 (2016).
- Rosas-Saavedra C and Stange C, Biosynthesis of carotenoids in plants: enzymes and color. *Subcell Biochem* **79**:35–69 (2016).
- Tanaka Y, Sasaki N and Ohmiya A, Biosynthesis of plant pigments: anthocyanins, betalains and carotenoids. *Plant J* **54**:733–749 (2008).
- Thoma F, Somborn-Schulz A, Schlehuber D, Keuter V and Deerberg G, Effects of light on secondary metabolites in selected leafy greens: a review. *Front Plant Sci* **11**:497 (2020).
- Fankhauser C and Chory J, Light control of plant development. *Annu Rev Cell Dev Biol* **13**:203–229 (1997).
- Christie JM, Blackwood L, Petersen J and Sullivan S, Plant flavoprotein photoreceptors. *Plant Cell Physiol* **56**:401–413 (2015).
- Ouzounis T, Rosenqvist E and Ottosen CO, Spectral effects of artificial light on plant physiology and secondary metabolism: a review. *HortScience* **50**:1128–1135 (2015).
- Paik I and Huq E, Plant photoreceptors: multi-functional sensory proteins and their signaling networks. *Semin Cell Dev Biol* **92**:114–121 (2019).
- Franklin KA and Quail PH, Phytochrome functions in *Arabidopsis* development. *J Exp Bot* **61**:11–24 (2010).
- Bouly JP, Schleicher E, Dionisio-Sese M, Vandenbussche F, Van Der Straeten D, Bakrim N et al., Cryptochrome blue light photoreceptors are activated through interconversion of flavin redox states. *J Biol Chem* **282**:9383–9391 (2007).
- Liu H, Liu B, Zhao C, Pepper M and Lin C, The action mechanisms of plant cryptochromes. *Trends Plant Sci* **16**:684–691 (2011).
- Christie JM, Phototropin blue-light receptors. *Annu Rev Plant Biol* **58**: 21–45 (2007).
- Brown BA, Cloix C, Jiang GH, Kaiserli E, Herzyk P, Kliebenstein DJ et al., A UV-B-specific signaling component orchestrates plant UV protection. *Proc Natl Acad Sci U S A* **102**:18225–18230 (2005).
- Landi M, Zivcak M, Sytar O, Brestic M and Allakhverdiev SI, Plasticity of photosynthetic processes and the accumulation of secondary metabolites in plants in response to monochromatic light environments: a review. *Biochim Biophys Acta Bioenerg* **2020**:148131 (1861).
- Wilhelm C and Selmar D, Energy dissipation is an essential mechanism to sustain the viability of plants: the physiological limits of improved photosynthesis. *J Plant Physiol* **168**:79–87 (2011).
- Petroustos D, Tokutsu R, Maruyama S, Flori S, Greiner A, Magneschi L et al., A blue-light photoreceptor mediates the feedback regulation of photosynthesis. *Nature* **537**:563–566 (2016).
- Darko E, Heydarizadeh P, Schoefs B and Sabzalian MR, Photosynthesis under artificial light: the shift in primary and secondary metabolism. *Philos Trans R Soc Lond B Biol Sci* **369**:20130243 (2014).
- Winkel-Shirley B, Biosynthesis of flavonoids and effects of stress. *Curr Opin Plant Biol* **5**:218–223 (2002).
- Sytar O, Zivcak M, Bruckova K, Brestic M, Hemmerich I, Rauh C et al., Shift in accumulation of flavonoids and phenolic acids in lettuce attributable to changes in ultraviolet radiation and temperature. *Sci Hortic* **239**:193–204 (2018).
- Hectors K, Prinsen E, De Coen W, Jansen MA and Guisez Y, *Arabidopsis thaliana* plants acclimated to low dose rates of ultraviolet B radiation

- show specific changes in morphology and gene expression in the absence of stress symptoms. *New Phytol* **175**:255–270 (2007).
- 27 Topcu Y, Dogan A, Kasimoglu Z, Sahin-Nadeem H, Polat E and Erkan M, The effects of UV radiation during the vegetative period on antioxidant compounds and postharvest quality of broccoli (*Brassica oleracea* L.). *Plant Physiol Biochem* **93**:56–65 (2015).
 - 28 Zivcak M, Brückova K, Sytar O, Brestic M, Olsovska K and Allakhverdiev SI, Lettuce flavonoids screening and phenotyping by chlorophyll fluorescence excitation ratio. *Planta* **245**:1215–1229 (2017).
 - 29 Falk KL, Tokuhisa JG and Gershenzon J, The effect of sulfur nutrition on plant glucosinolate content: physiology and molecular mechanisms. *Plant Biol* **9**:573–581 (2007).
 - 30 Selmar D and Kleinwächter M, Stress enhances the synthesis of secondary plant products: the impact of stress-related over-reduction on the accumulation of natural products. *Plant Cell Physiol* **54**:817–826 (2013).
 - 31 Kleinwächter M and Selmar D, New insights explain that drought stress enhances the quality of spice and medicinal plants: potential applications. *Agron Sustain Dev* **35**:121–131 (2015).
 - 32 Morrow RC, LED lighting in horticulture. *HortScience* **43**:1947–1950 (2008).
 - 33 Yeh N and Chung JP, High-brightness LEDs—energy efficient lighting sources and their potential in indoor plant cultivation. *Renew Sustain Energy Rev* **13**:2175–2180 (2009).
 - 34 Kozai T, Why LED lighting for urban agriculture? in *LED Lighting for Urban Agriculture*, ed. by Kozai T, Fujiwara K and Runkle ES. Singapore, Springer, pp. 3–18 (2016).
 - 35 Kozai T, Niu G and Takagaki M, *Plant Factory: An Indoor Vertical Farming System for Efficient Quality Food Production*, 2nd edn. Academic Press, Cambridge, US (2019).
 - 36 Dou H, Niu G, Gu M and Masabni JG, Effects of light quality on growth and phytonutrient accumulation of herbs under controlled environments. *Horticulturae* **3**:36 (2017).
 - 37 Alrifai O, Hao X, Marcone MF and Tsao R, Current review of the modulatory effects of LED lights on photosynthesis of secondary metabolites and future perspectives of microgreen vegetables. *J Agric Food Chem* **67**:6075–6090 (2019).
 - 38 El Sohaimy SA, Functional foods and nutraceuticals—modern approach to food science. *World Appl Sci J* **20**:691–708 (2012).
 - 39 Wu MC, Hou CY, Jiang CM, Wang YT, Wang CY, Chen HH *et al.*, A novel approach of LED light radiation improves the antioxidant activity of pea seedlings. *Food Chem* **101**:1753–1758 (2007).
 - 40 Manivannan A, Soundararajan P, Halimah N, Ko CH and Jeong BR, Blue LED light enhances growth, phytochemical contents, and antioxidant enzyme activities of *Rehmannia glutinosa* cultured in vitro. *Hortic Environ Biotechnol* **56**:105–113 (2015).
 - 41 Samuolienė G, Brazaitytė A, Sirtautas R, Viršilė A, Sakalauskaitė J, Sakalauskienė S *et al.*, LED illumination affects bioactive compounds in romaine baby leaf lettuce. *J Sci Food Agric* **93**:3286–3291 (2013).
 - 42 Pennisi G, Orsini F, Blasioli S, Cellini A, Crepaldi A, Braschi I *et al.*, Resource use efficiency of indoor lettuce (*Lactuca sativa* L.) cultivation as affected by red:blue ratio provided by LED lighting. *Sci Rep* **9**:1–11 (2019).
 - 43 Lubbe A and Verpoorte R, Cultivation of medicinal and aromatic plants for specialty industrial materials. *Ind Crops Prod* **34**:785–801 (2011).
 - 44 Choi JH, Seong ES, Yoo JH, Choi SK, Lee JG, Lim JD *et al.*, Enhancement of growth characteristics and biological activities in *Astragalus membranaceus* using artificial light sources. *Russ J Plant Physiol* **65**:732–739 (2018).
 - 45 Samuolienė G, Viršilė A, Brazaitytė A, Jankauskienė J, Sakalauskienė S, Vaštakaitė V *et al.*, Blue light dosage affects carotenoids and tocopherols in microgreens. *Food Chem* **228**:50–56 (2017).
 - 46 Pennisi G, Blasioli S, Cellini A, Maia L, Crepaldi A, Braschi I *et al.*, Unraveling the role of red:blue LED lights on resource use efficiency and nutritional properties of indoor grown sweet basil. *Front Plant Sci* **10**:305 (2019).
 - 47 Meas S, Luengwilai K and Thongket T, Enhancing growth and phytochemicals of two amaranth microgreens by LEDs light irradiation. *Sci Hortic* **265**:109204 (2020).
 - 48 Lobiuc A, Vasilache V, Pintilie O, Stoleru T, Burducea M, Oroian M *et al.*, Blue and red LED illumination improves growth and bioactive compounds contents in acyanic and cyanic *Ocimum basilicum* L microgreens. *Molecules* **22**:2111 (2017).
 - 49 Sytar O, Zivcak M, Neugart S, Toutouchi PM and Brestic M, Precultivation of young seedlings under different color shades modifies the accumulation of phenolic compounds in *Cichorium* leaves in later growth phases. *Environ Exp Bot* **165**:30–38 (2019).
 - 50 Shinomura T, Nagatani A, Hanzawa H, Kubota M, Watanabe M and Furuya M, Action spectra for phytochrome a and B specific photoinduction of seed germination in *Arabidopsis thaliana*. *PNAS USA* **93**:8129–8133 (1996).
 - 51 Casal JJ, Phytochromes, cryptochromes, phototropin: photoreceptor interactions in plants. *Photochem Photobiol* **71**:1–11 (2000).
 - 52 Gould K, Davies KM and Winefield C, *Anthocyanins: Functions, and Applications*. Springer Science & Business Media, New York (2009).
 - 53 Agati G and Tattini M, Multiple functional roles of flavonoids in photoprotection. *New Phytol* **186**:786–793 (2010).
 - 54 Kyriacou MC, Rouphael Y, Di Gioia F, Kyrtatzis A, Serio F, Renna M *et al.*, Micro-scale vegetable production and the rise of microgreens. *Trends Food Sci Technol* **57**:103–115 (2016).
 - 55 Ying Q, Kong Y and Zheng Y, Growth and appearance quality of four microgreen species under light-emitting diode lights with different spectral combinations. *HortScience* **55**:1399–1405 (2020).
 - 56 Samuolienė G, Brazaitytė A, Viršilė A, Millauskienė J, Vaštakaitė-Kairienė V and Duchovskis P, Nutrient levels in Brassicaceae microgreens increase under tailored light-emitting diode spectra. *Front Plant Sci* **10**:1475 (2019).
 - 57 Lenzi A, Orlandini A, Bulgari R, Ferrante A and Bruschi P, Antioxidant and mineral composition of three wild leafy species: a comparison between microgreens and baby greens. *Foods* **8**:487 (2019).
 - 58 Harakotr B, Srijunteuk S, Rithichai P and Tabunhan S, Effects of light-emitting diode light irradiance levels on yield, antioxidants and antioxidant capacities of indigenous vegetable microgreens. *Sci Technol Asia* **24**:59–66 (2019).
 - 59 Zhang X, Bian Z, Li S, Chen X and Lu C, Comparative analysis of phenolic compound profiles, antioxidant capacities, and expressions of phenolic biosynthesis-related genes in soybean microgreens grown under different light spectra. *J Agric Food Chem* **67**:13577–13588 (2019).
 - 60 Kong Y, Kamath D and Zheng Y, Blue versus red light can promote elongation growth independent of photoperiod: a study in four brassica microgreens species. *HortScience* **54**:1955–1961 (2019).
 - 61 Tan L, Nuffer H, Feng J, Kwan SH, Chen H, Tong X *et al.*, Antioxidant properties and sensory evaluation of microgreens from commercial and local farms. *Food Sci Hum Wellness* **9**:45–51 (2020).
 - 62 Xiao Z, Lester GE, Luo Y and Wang Q, Assessment of vitamin and carotenoid concentrations of emerging food products: edible microgreens. *J Agric Food Chem* **60**:7644–7651 (2012).
 - 63 Craver JK, Gerovac JR, Lopez RG and Kopsell DA, Light intensity and light quality from sole-source light-emitting diodes impact phytochemical concentrations within brassica microgreens. *J Am Soc Hort Sci* **142**:3–12 (2017).
 - 64 Jones-Baumgardt C, Llewellyn D and Zheng Y, Different microgreen genotypes have unique growth and yield responses to intensity of supplemental PAR from light-emitting diodes during winter greenhouse production in southern Ontario, Canada. *HortScience* **55**:56–163 (2020).
 - 65 Gerovac JR, Craver JK, Boldt JK and Lopez RG, Light intensity and quality from sole-source light-emitting diodes impact growth, morphology, and nutrient content of brassica microgreens. *HortScience* **51**:497–503 (2016).
 - 66 Vaštakaitė V, Viršilė A, Brazaitytė A, Samuolienė G, Jankauskienė J, Novičkovas A *et al.*, Pulsed light-emitting diodes for a higher phytochemical level in microgreens. *J Agric Food Chem* **65**:6529–6534 (2017).
 - 67 Rajan P, Lada RR and MacDonald MT, Advancement in indoor vertical farming for microgreen production. *Am J Plant Sci* **10**:1397–1408 (2019).
 - 68 Singh N, Rani S and Mishra A, Cruciferous microgreens: growing performance and their scope as super foods at high altitude locations. *Prog Hortic* **51**:41–44 (2019).
 - 69 Lefsrud MG, Kopsell DA, Kopsell DE and Curran-Celentano J, Irradiance levels affect growth parameters and carotenoid pigments in kale and spinach grown in a controlled environment. *Physiol Plant* **127**:624–631 (2006).
 - 70 Iwai M, Ohta M, Tsuchiya H and Suzuki T, Enhanced accumulation of caffeic acid, rosmarinic acid and luteolin-glucoside in red perilla

- cultivated under red diode laser and blue LED illumination followed by UV-A irradiation. *J Funct Foods* **2**:66–70 (2010).
- 71 Cominelli E, Gusmaroli G, Allegra D, Galbiati M, Wade HK, Jenkins GI et al., Expression analysis of anthocyanin regulatory genes in response to different light qualities in *Arabidopsis thaliana*. *J Plant Physiol* **165**:886–894 (2008).
 - 72 Yang L, Wen KS, Ruan X, Zhao YX, Wei F and Wang Q, Response of plant secondary metabolites to environmental factors. *Molecules* **23**:762 (2018).
 - 73 Brazaitytė A, Viršilė A, Samuolienė G, Vaštakaitė-Kairienė V, Jankauskienė J, Miliauskienė J et al., Response of mustard microgreens to different wavelengths and durations of UV-A LEDs. *Front Plant Sci* **10**:1153 (2019).
 - 74 Neugart S and Schreiner M, UVB and UVA as eustressors in horticultural and agricultural crops. *Sci Hortic* **234**:370–381 (2018).
 - 75 Nowicka P and Wojdyło A, Anti-hyperglycemic and anticholinergic effects of natural antioxidant contents in edible flowers. *Antioxidants* **8**:308 (2019).
 - 76 Fernandes L, Casal S, Pereira JA, Saraiva JA and Ramalhosa E, Edible flowers: a review of the nutritional, antioxidant, antimicrobial properties and effects on human health. *J Food Compos Anal* **60**:38–50 (2017).
 - 77 Lu B, Li M and Yin R, Phytochemical content, health benefits, and toxicology of common edible flowers: a review (2000–2015). *Crit Rev Food Sci Nutr* **56**:S130–S148 (2016).
 - 78 Loizzo MR, Pugliese A, Bonesi M, Tenuta MC, Menichini F, Xiao J et al., Edible flowers: a rich source of phytochemicals with antioxidant and hypoglycemic properties. *J Agric Food Chem* **64**:2467–2474 (2016).
 - 79 Navarro-González I, González-Barrio R, García-Valverde V, Bautista-Ortín AB and Periago MJ, Nutritional composition and antioxidant capacity in edible flowers: characterisation of phenolic compounds by HPLC-DAD-ESI/MSn. *Int J Mol Sci* **16**:805–822 (2015).
 - 80 González-Barrio R, Periago MJ, Luna-Recio C, García-Alonso FJ and Navarro-González I, Chemical composition of the edible flowers, pansy (*Viola wittrockiana*) and snapdragon (*Antirrhinum majus*) as new sources of bioactive compounds. *Food Chem* **252**:373–380 (2018).
 - 81 Kopsell D, Belisle C, Lowery H and Whitlock C, Genotype and lighting environment impact petal tissue pigmentation in *Tagetes tenuifolia*. *Acta Hort* **1134**:103–110 (2016).
 - 82 Wagner H, Hikino H and Farnsworth NR, *Economic and Medicinal Plant Research*. Academic Press, Cambridge, US (2012).
 - 83 Sharma A, Gene expression analysis in medicinal plants under abiotic stress conditions, in *Plant Metabolites and Regulation under Environmental Stress*, ed. by Ahmad P, Ahanger MA, Singh VP, Tripathi DK, Alam P and Alyemeni MN. Academic Press, Cambridge, US, pp. 407–414 (2018).
 - 84 Giurgiu RM, Morar GA, Dumitraş A, Boancă P, Duda BM and Doldovan C, Study regarding the suitability of cultivating medicinal plants in hydroponic systems in controlled environment. *Res J Agric Sci* **46**:84–92 (2014).
 - 85 Appolloni E, Vitali C, Petricciuolo E, Minelli G, Cleri A, Cleri E et al., DoloMiNet: building a network of vertical farms in the heart of Italian Alps. *Acta Hort* **1298**:497–510 (2020).
 - 86 Bae JH, Park SY and Oh MM, Supplemental irradiation with far-red light-emitting diodes improves growth and phenolic contents in *Crepidiasrum denticulatum* in a plant factory with artificial lighting. *Hortic Environ Biotechnol* **58**:357–366 (2017).
 - 87 Park SY, Bae JH and Oh MM, Manipulating light quality to promote shoot growth and bioactive compound biosynthesis of *Crepidiasrum denticulatum* (Houtt) Pak & Kawano cultivated in plant factories. *J Appl Res Med Aromat Plants* **16**:100237 (2020).
 - 88 Bora KS and Sharma A, Phytochemical and pharmacological potential of *Medicago sativa*: a review. *Pharm Biol* **49**:211–220 (2011).
 - 89 Fiutak G, Michalczyk M, Filipczak-Fiutak M, Fiedor L and Surówka K, The impact of LED lighting on the yield, morphological structure and some bioactive components in alfalfa (*Medicago sativa* L.) sprouts. *Food Chem* **285**:53–58 (2019).
 - 90 Ajdanian L, Babaei M and Aroiee H, The growth and development of cress (*Lepidium sativum*) affected by blue and red light. *Heliyon* **5**:e02109 (2019).
 - 91 Verma SK, Gantait S, Jeong BR and Hwang SJ, Enhanced growth and cardenolides production in *Digitalis purpurea* under the influence of different LED exposures in the plant factory. *Sci Rep* **8**:18009 (2018).
 - 92 Scheibner H, Björkz L, Schulz U, Diettrich B and Luckner M, Influence of light on cardenolide accumulation in somatic embryos of *Digitalis lanata*. *J Plant Physiol* **130**:211–219 (1987).
 - 93 Zhang S, Ma J, Zou H, Zhang L, Li S and Wang Y, The combination of blue and red LED light improves growth and phenolic acid contents in *salvia miltiorrhiza* Bunge. *Ind Crops Prod* **158**:112959 (2020).
 - 94 Ghaffari Z, Rahimmalek M and Sabzalian MR, Variation in the primary and secondary metabolites derived from the isoprenoid pathway in the *Perovskia* species in response to different wavelengths generated by light emitting diodes (LEDs). *Ind Crops Prod* **140**:111592 (2019).
 - 95 Gharibi S, Tabatabaei BES, Saeidi G, Talebi M and Matkowski A, The effect of drought stress on polyphenolic compounds and expression of flavonoid biosynthesis related genes in *Achillea pachycephala* Rech.F. *Phytochemistry* **162**:90–98 (2019).
 - 96 Jeong SY, Moon MY, Ryu EK, Lee JS and Cha J, Identification of the phytochemical compounds and their type I procollagen induction in *Astragalus membranaceus* sprouts grown under different light conditions. *J Basic Appl Pharm Sci* **8**:1–7 (2018).
 - 97 Park WT, Yeo SK, Sathasivam R, Park JS, Kim JK and Park SU, Influence of light-emitting diodes on phenylpropanoid biosynthetic gene expression and phenylpropanoid accumulation in *Agastache rugosa*. *Appl Biol Chem* **63**:25 (2020).
 - 98 Lin KH, Huang MY, Huang WD, Hsu MH, Yang ZW and Yang CM, The effects of red, blue, and white light-emitting diodes on the growth, development, and edible quality of hydroponically grown lettuce (*Lactuca sativa* L. var. capitata). *Sci Hortic* **150**:86–91 (2013).
 - 99 Livadariu O, Raiciu D, Maximilian C and Căpitanu E, Studies regarding treatments of LED-s emitted light on sprouting hemp (*Cannabis sativa* L.). *Rom Biotechnol Lett* **24**:485–490 (2019).
 - 100 Namdar D, Charuvi D, Ajjampura V, Mazuz M, Ion A, Kamara I et al., LED lighting affects the composition and biological activity of *Cannabis sativa* secondary metabolites. *Ind Crops Prod* **132**:177–185 (2019).
 - 101 Christaki E, Bonos E, Giannenas I and Florou-Paneri P, Aromatic plants as a source of bioactive compounds. *Agriculture* **2**:228–243 (2012).
 - 102 Franz C, Baser KHC and Windisch W, Essential oils and aromatic plants in animal feeding - a European perspective. A review. *Flavour Fragr J* **25**:327–340 (2010).
 - 103 Zeng Z, Zhang S, Wang H and Piao X, Essential oil and aromatic plants as feed additives in non-ruminant nutrition: a review. *J Anim Sci Biotechnol* **6**:7 (2015).
 - 104 Sgherri C, Cecconami S, Pinzino C, Navari-Izzo F and Izzo R, Levels of antioxidants and nutraceuticals in basil grown in hydroponics and soil. *Food Chem* **123**:416–422 (2010).
 - 105 Lin KH, Huang MY and Hsu MH, Morphological and physiological response in green and purple basil plants (*Ocimum basilicum*) under different proportions of red, green, and blue LED lightings. *Sci Hortic* **275**:109677 (2021).
 - 106 Hosseini A, Mehrjerdi MZ and Aliniaefard S, Alteration of bioactive compounds in two varieties of basil (*Ocimum basilicum*) grown under different light spectra. *J Essent Oil Bearing Plants* **21**:913–923 (2018).
 - 107 Taulavuori K, Hyöky V, Oksanen J, Taulavuori E and Julkunen-Tiitto R, Species-specific differences in synthesis of flavonoids and phenolic acids under increasing periods of enhanced blue light. *Environ Exp Bot* **121**:145–150 (2016).
 - 108 Chang X, Alderson PG and Wright CJ, Solar irradiance level alters the growth of basil (*Ocimum basilicum* L.) and its content of volatile oils. *Environ Exp Bot* **63**:216–223 (2008).
 - 109 Samuolienė G, Brazaitytė A, Viršilė A, Jankauskienė J, Sakalauskienė S and Duchovskis P, Red light-dose or wavelength-dependent photo-response of antioxidants in herb microgreens. *PLoS One* **11**:e0163405 (2016).
 - 110 Pennisi G, Pistillo A, Orsini F, Cellini A, Spinelli F, Nicola S et al., Optimal light intensity for sustainable water and energy use in indoor cultivation of lettuce and basil under red and blue LEDs. *Sci Hortic* **272**:109508 (2020).
 - 111 Bantis F, Ouzounis T and Radoglou K, Artificial LED lighting enhances growth characteristics and total phenolic content of *Ocimum basilicum*, but variably affects transplant success. *Sci Hortic* **198**:277–283 (2016).
 - 112 Heo JW, Kang DH, Bang HS, Hong SG and Chun C, Early growth, pigmentation, protein content, and phenylalanine ammonia-lyase activity of red curled lettuces grown under different lighting conditions. *Korean J Hortic Sci Technol* **30**:6–12 (2012).

- 113 Naznin MT, Lefsrud M, Gravel V and Azad M, Blue light added with red LEDs enhance growth characteristics, pigments content, and antioxidant capacity in lettuce, spinach, kale, basil, and sweet pepper in a controlled environment. *Plan Theory* **8**:93 (2019).
- 114 Gao S, Liu X, Liu Y, Cao B, Chen Z and Xu K, Photosynthetic characteristics and chloroplast ultrastructure of welsh onion (*Allium fistulosum* L.) grown under different LED wavelengths. *BMC Plant Biol* **20**:1–12 (2020).
- 115 Raiciu AD and Livadariu O, Maximilian C and Crețu AD, the assessment of the effect induced by LED-s irradiation on garlic sprouts (*Allium sativum* L.). *Rom Biotechnol Lett* **23**:14187–14192 (2018).
- 116 Nguyen DTP, Kitayama M, Lu N and Takagaki M, Improving secondary metabolite accumulation, mineral content, and growth of coriander (*Coriandrum sativum* L.) by regulating light quality in a plant factory. *J Horticult Sci Biotech* **95**:356–363 (2020).
- 117 Ahmadi T, Shabani L and Sabzalian MR, Improvement in drought tolerance of lemon balm, *Melissa officinalis* L under the pre-treatment of LED lighting. *Plant Physiol Biochem* **139**:548–557 (2019).
- 118 Mittler R, Oxidative stress, antioxidants and stress tolerance. *Trends Plant Sci* **7**:405–410 (2002).
- 119 Frąszczak B, Gąsecka M, Golcz A and Zawirska-Wojtasiak R, The effect of radiation of LED modules on the growth of dill (*Anethum graveolens* L.). *Open Life Sci* **11**:61–70 (2016).
- 120 Tohidi B, Rahimmalek M, Arzani A and Sabzalian MR, Thymol, carvacrol, and antioxidant accumulation in *thymus* species in response to different light spectra emitted by light-emitting diodes. *Food Chem* **307**:125521 (2020).
- 121 Rihan HZ, Aldarkazali M, Mohame SJ, McMulki NB, Jbara MH and Fulle MP, A novel new light recipe significantly increases the growth and yield of sweet basil (*Ocimum basilicum*) grown in plant factory system. *Agronomy* **10**:934 (2020).
- 122 Nguyen TL and Saleh MA, Effect of exposure to light emitted diode (LED) lights on essential oil composition of sweet mint plants. *J Environ Sci Health A Tox Hazard Subst Environ Eng* **54**:435–440 (2019).
- 123 Takahashi JA, Rezende FAGG, Moura MAF, Dominguet LCB and Sande D, Edible flowers: bioactive profile and its potential to be used in food development. *Food Res Int* **129**:108868 (2020).
- 124 Batista DS, de Castro KM, da Silva AR, Teixeira ML, Sales TA and Soares LI, Graças Cardoso M das, de Oliveira Santos M, Viccini LF and Otoni WC, light quality affects in vitro growth and essential oil profile in *Lippia alba* (Verbenaceae). *Vitr Cell Dev Biol – Plant* **52**:276–282 (2016).
- 125 Billore V, Khatediya L and Jain M, Sink-source system of in vitro suspension culture of *Celastrus paniculatus* under regulation of monochromatic lights. *Plant Tissue Cult Biotechnol* **26**:175–185 (2016).
- 126 Dutta Gupta S and Karmakar A, Machine vision based evaluation of impact of light emitting diodes (LEDs) on shoot regeneration and the effect of spectral quality on phenolic content and antioxidant capacity in *Swertia chirata*. *J Photochem Photobiol B* **174**:162–172 (2017).
- 127 Dutta Gupta S, Kumar A and Agarwal A, Impact of light-emitting diodes (LEDs) on the growth and morphogenesis of encapsulated shoot buds of *Curculigo orchoides* Gaertn., an endangered medicinal herb. *Acta Physiol Plant* **41**:50 (2019).
- 128 De Carvalho AA, Bertolucci SKV, da Cunha HA, Rocha TT, Silva ST and Pinto JEBP, Influence of light spectra and elicitors on growth and ascaridole content using in vitro cultures of *Dysphania ambrosioides* L. *Plant Cell Tiss Org Cult* **143**:277–290 (2020).
- 129 Khurshid R, Ullah MA, Tungmunnithum D, Drouet S, Shah M, Zaeem A et al., Lights triggered differential accumulation of antioxidant and antidiabetic secondary metabolites in callus culture of *Eclipta alba* L. *PLoS One* **15**:e0233963 (2020).
- 130 Kubica P, Szopa A, Prokopiuk B, Komsta Ł, Pawłowska B and Ekiert H, The influence of light quality on the production of bioactive metabolites – verbascoside, isoverbascoside and phenolic acids and the content of photosynthetic pigments in biomass of *Verbena officinalis* L. cultured in vitro. *J Photochem Photobiol B Biol* **203**:111768 (2020).
- 131 Zárata-Salazar JR, de Souza LM, de Moraes MB, Palhares Neto L, Willadino L, Gouveia-Neto A et al., Light-emitting diodes and gas exchange facilitation minimize hyperhydricity in *Lippia grata*: physiological, biochemical and morpho anatomical aspects. *S Afr J Bot* **135**:164–171 (2020).
- 132 Zotov VS, Bolychevtseva YV, Khapchaeva SA, Terekhova IV, Shubin VV, Yurina NP et al., Effect of light quality on the biomass yield, photosystem 2 fluorescence, and the total essential oil content of *Ocimum basilicum*. *Appl Biochem Microbiol* **56**:336–343 (2020).