







FULL-LENGTH ORIGINAL RESEARCH

Toward a better definition of focal cortical dysplasia: An iterative histopathological and genetic agreement trial

Ingmar Blümcke^{1,2}  | Roland Coras¹ | Robyn M. Busch^{2,3,4}  | Marcia Morita-Sherman² | Dennis Lal^{2,4} | Richard Prayson⁵ | Fernando Cendes^{6,7}  | Iscia Lopes-Cendes^{7,8}  | Fabio Rogerio^{7,9} | Vanessa S. Almeida^{7,8} | Cristiane S. Rocha^{7,8} | Nam Suk Sim¹⁰ | Jeong Ho Lee^{10,11} | Se Hoon Kim¹² | Stephanie Baulac¹³  | Sara Baldassari¹³  | Homa Adle-Biassette^{14,15} | Christopher A. Walsh^{16,17} | Sara Bizzotto^{16,17} | Ryan N. Doan^{16,17} | Katherine S. Morillo^{16,17} | Eleonora Aronica^{18,19} | Angelika Mühlebner^{18,20}  | Albert Becker²¹  | Jesus Cienfuegos^{22,23} | Rita Garbelli²⁴  | Caterina Giannini^{25,26} | Mrinalini Honavar²⁷ | Thomas S. Jacques^{28,29}  | Maria Thom³⁰  | Anita Mahadevan³¹  | Hajime Miyata³²  | Pitt Niehusmann³³ | Harvey B. Sarnat^{34,35,36} | Figen Söylemezoglu³⁷ | Imad Najm^{2,3}

¹Department of Neuropathology, University Hospital, Erlangen, Germany

²Epilepsy Center, Cleveland Clinic, Cleveland, OH, USA

³Department of Neurology, Cleveland Clinic, Cleveland, OH, USA

⁴Genomic Medicine Institute, Cleveland Clinic, Cleveland, OH, USA

⁵Department of Anatomic Pathology, Cleveland Clinic, Cleveland, OH, USA

⁶Department of Neurology, University of Campinas, Sao Paulo, Brazil

⁷Brazilian Institute of Neuroscience and Neurotechnology, Sao Paulo, Brazil

⁸Department of Medical Genetics and Genomic Medicine, University of Campinas, Sao Paulo, Brazil

⁹Department of Pathology, University of Campinas, Sao Paulo, Brazil

¹⁰Graduate School of Medical Science and Engineering, Korea Advanced Institute of Science and Technology, Daejeon, South Korea

¹¹SoVarGen, Inc., Daejeon, Korea

¹²Department of Pathology, College of Medicine, Yonsei University, Seoul, South Korea

¹³Sorbonne Université, Institut du Cerveau - Paris Brain Institute - ICM, Inserm, CNRS, Paris, France

¹⁴Pathological Anatomy Service, Public Hospital Network of Paris, Paris, France

¹⁵NeuroDiderot, Inserm U1141, University of Paris, Paris, France

¹⁶Division of Genetics and Genomics, Manton Center for Orphan Disease Research, Department of Pediatrics, and Howard Hughes Medical Institute, Boston Children's Hospital, Boston, MA, USA

¹⁷Departments of Pediatrics and Neurology, Harvard Medical School, Boston, MA, USA

Ingmar Blümcke, Fernando Cendes, Jeong Ho Lee, Stephanie Baulac, Christopher A. Walsh, Eleonora Aronica, Rita Garbelli, Maria Thom, Hajime Miyata, Harvey B. Sarnat, and Imad Najm are members of the International League Against Epilepsy Task Force on Focal Cortical Dysplasia.

Dennis Lal, Iscia Lopes-Cendes, Vanessa S. Almeida, Cristiane S. Rocha, Nam Suk Sim, Jeong Ho Lee, Stephanie Baulac, Sara Baldassari, Christopher A. Walsh, Sara Bizzotto, Ryan N. Doan, and Katherine S. Morillo are members of the Focal Cortical Dysplasia Gene Panel Analysis Group.

Ingmar Blümcke, Roland Coras, Richard Prayson, Fabio Rogerio, Se Hoon Kim, Homa Adle-Biassette, Eleonora Aronica, Angelika Mühlebner, Albert Becker, Jesus Cienfuegos, Rita Garbelli, Caterina Giannini, Mrinalini Honavar, Thomas S. Jacques, Maria Thom, Anita Mahadevan, Hajime Miyata, Pitt Niehusmann, Harvey B. Sarnat, and Figen Söylemezoglu are members of the Histopathology Review Group.

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

© 2021 The Authors. *Epilepsia* published by Wiley Periodicals LLC on behalf of International League Against Epilepsy

- ¹⁸Department of (Neuro)Pathology, Amsterdam UMC, location Academic Medical Center, Amsterdam, the Netherlands
- ¹⁹Epilepsy Institutes of the Netherlands Foundation, Heemstede, the Netherlands
- ²⁰Department of Pathology, University Medical Center Utrecht, Utrecht, the Netherlands
- ²¹Department of Neuropathology, University of Bonn Medical Center, Bonn, Germany
- ²²Department of Anatomic Pathology, International Center for Epilepsy Surgery, Humanitas Medical Group Hospital, Mexico City, Mexico
- ²³Department of Anatomic Pathology, Angeles Clínica Londres Hospital, Mexico City, Mexico
- ²⁴Epilepsy Unit, Carlo Besta Neurological Institute, Scientific Institute for Research and Health Care Foundation, Milan, Italy
- ²⁵Anatomic Pathology, Mayo Clinic, Rochester, Minnesota, USA
- ²⁶Department of Biomedical and Neuromotor Science, Alma Mater Studiorum, University of Bologna, Bologna, Italy
- ²⁷Department of Anatomic Pathology, Pedro Hispano Hospital, Matosinhos, Portugal
- ²⁸Developmental Biology and Cancer Research and Teaching Programme, University College London Great Ormond Street Institute of Child Health, London, UK
- ²⁹Department of Histopathology, Great Ormond Street Hospital for Children, National Health Service Foundation Trust, London, UK
- ³⁰Department of Neuropathology, Institute of Neurology, University College London, London, UK
- ³¹Department of Neuropathology, National Institute of Mental Health and Neurosciences, Bangalore, India
- ³²Department of Neuropathology, Research Institute for Brain and Blood Vessels, Akita Cerebrospinal and Cardiovascular Center, Akita, Japan
- ³³Department of Neuro-/Pathology, Translational Neurodegeneration Research and Neuropathology Lab, University of Oslo and Oslo University Hospital, Oslo, Norway
- ³⁴Department of Paediatrics, University of Calgary Faculty of Medicine, Alberta Children's Hospital Research Institute, Calgary, AB, Canada
- ³⁵Department of Pathology (Neuropathology), University of Calgary Faculty of Medicine, Alberta Children's Hospital Research Institute, Calgary, AB, Canada
- ³⁶Department of Clinical Neurosciences, University of Calgary Faculty of Medicine, Alberta Children's Hospital Research Institute, Calgary, AB, Canada
- ³⁷Department of Pathology, Faculty of Medicine, Hacettepe University, Ankara, Turkey

Correspondence

Ingmar Blümcke, Department of Neuropathology, University Hospitals Erlangen, Schwabachanlage 6, D – 91054 Germany.
Email: bluemcke@uk-erlangen.de

Abstract

Objective: Focal cortical dysplasia (FCD) is a major cause of difficult-to-treat epilepsy in children and young adults, and the diagnosis is currently based on microscopic review of surgical brain tissue using the International League Against Epilepsy classification scheme of 2011. We developed an iterative histopathological agreement trial with genetic testing to identify areas of diagnostic challenges in this widely used classification scheme.

Methods: Four web-based digital pathology trials were completed by 20 neuropathologists from 15 countries using a consecutive series of 196 surgical tissue blocks obtained from 22 epilepsy patients at a single center. Five independent genetic laboratories performed screening or validation sequencing of FCD-relevant genes in paired brain and blood samples from the same 22 epilepsy patients.

Results: Histopathology agreement based solely on hematoxylin and eosin stainings was low in Round 1, and gradually increased by adding a panel of immunostainings in Round 2 and the Delphi consensus method in Round 3. Interobserver agreement was good in Round 4 ($\kappa = .65$), when the results of genetic tests were disclosed, namely, *MTOR*, *AKT3*, and *SLC35A2* brain somatic mutations in five cases and germline mutations in *DEPDC5* and *NPRL3* in two cases.

Significance: The diagnoses of FCD 1 and 3 subtypes remained most challenging and were often difficult to differentiate from a normal homotypic or heterotypic cortical architecture. Immunohistochemistry was helpful, however, to confirm the diagnosis of FCD or no lesion. We observed a genotype–phenotype association for brain somatic mutations in *SLC35A2* in two cases with mild malformation of cortical

development with oligodendroglial hyperplasia in epilepsy. Our results suggest that the current FCD classification should recognize a panel of immunohistochemical stainings for a better histopathological workup and definition of FCD subtypes. We also propose adding the level of genetic findings to obtain a comprehensive, reliable, and integrative genotype–phenotype diagnosis in the near future.

KEY WORDS

brain, classification, epilepsy, genes, neuropathology, seizure

Key Points

- We performed an iterative histopathological agreement trial with genetic testing to challenge the widely used ILAE classification scheme for FCD
- Twenty neuropathologists from 15 countries completed four digital pathology trials of 196 surgical tissue blocks obtained from 22 epilepsy patients
- Five independent laboratories performed genetic screening or validation sequencing of FCD-relevant genes in paired brain and blood samples from the same 22 patients
- Interobserver agreement increased when immunohistochemistry was available and results of genetic tests were disclosed
- We propose adding the level of genetic findings to the ILAE classification scheme to obtain an integrative genotype–phenotype diagnosis in the near future

1 | INTRODUCTION

Etiology matters in the clinical management of patients with difficult-to-treat focal epilepsies.^{1,2} Epilepsy surgery has been established as a reliable treatment option in many of these patients,^{3,4} and the spectrum of structural brain lesions amenable for surgical treatment is well described.⁵ Malformations of cortical development (MCDs) account for the most common surgical pathologies in children and the third most common in adults, with focal cortical dysplasia (FCD) representing almost 75% of all MCD cases.^{2,5} Postsurgical seizure outcomes vary considerably between various FCD subtypes. In a recent European multicenter study of 9147 epilepsy surgery patients, FCD Type 1 and mild MCD (mMCD) showed the poorest outcome, whereas Type 2 FCDs were associated with much higher rates of long-term seizure freedom on and off antiseizure medications up to 5 years after surgery.²

FCD was first described in 1971,⁶ but its clinicopathological classification scheme remains a topic of ongoing discussion.⁷ FCD originally contained only cytoarchitectural dysplasia including cytomegalic neurons and balloon cells in an architecturally abnormal neocortex, for example, FCD 2A or 2B in the current terminology. It was then extended to add a new category, frequently encountered in patients with epilepsy, that involves architectural abnormalities of the neocortex but lacking cytopathologic features,⁸ termed FCD Type 1 by the Palmini classification in 2004.⁹ Microscopic hallmarks of a reliable histopathological diagnosis of FCD

Type 1 remained suboptimal¹⁰ and were revised in the international FCD consensus classification scheme of 2011.¹¹

This widely used International League Against Epilepsy (ILAE) classification further introduced FCD Type 3, wherein architectural abnormalities were associated with and adjacent to another principal lesion, for example, hippocampal sclerosis (HS; FCD Type 3A), low-grade developmental brain tumors (FCD Type 3B), vascular malformations (FCD Type 3C), or any other lesion acquired during early life (FCD Type 3D). The foundation for this classification was laid by observations that patients with isolated (pure) FCD Type 1 have a more severe clinical phenotype^{12,13} and may require more extensive resection or hemispheric disconnection procedures.⁴ In contrast, clinical phenotypes and postsurgical seizure outcome were not significantly different between patients with HS or other principal brain lesions with or without accompanying FCD.¹² Microscopic intra- and interobserver agreement was tested for the 2011 ILAE classification addressing different levels of professional experience.¹⁴ Overall kappa values reached good levels, but the study also highlighted the need for continuous training and that FCD Type 1 and 3 subtypes required further attention.

Challenges in interobserver agreement with respect to histopathological diagnosis are long-standing and occur not only in the arena of epilepsy surgery.¹⁵ In 2016, the World Health Organization Classification of Tumors of the Central Nervous System proposed a multilayered diagnostic scheme to address this issue by integrating genetic testing with

TABLE 1 The 22 patients included in the study

Case	Sex	Age at seizure onset, years	Age at surgery, years	Localization ^a	MRI	RF	iEEG	Outcome, Engel ^b	FU, years
1	F	13	28	L Fr	pos	No	SDG, icEEG	IA	1.25
2	F	4	14	R T C	pos	DD	icEEG	IC	4.17
3	M	8	50	R Fr T	pos	FAM	SDG, icEEG	IA	4
4	F	13	25	L Fr	pos	TBI	SDG, icEEG	IA	1.17
5	M	9	19	R P	pos	FAM	icEEG	II	2.5
6	F	17	27	L T	neg	TBI	icEEG	IV	2.25
7	M	5	16	L Fr	pos	FAM	icEEG + ECoG	II	1.58
8	M	49	53	L T	neg	TBI	icEEG + ECoG	IA	1
9	M	36	49	L T	pos	Npl	No	III	1
10	F	22	44	L T	pos	No	No	IA	1.33
11	F	4	26	R T C	pos	TBI	No	IA	1.08
12	M	.8	8	L Fr	pos	DD	icEEG + ECoG	n.a.	.17
13	F	0	37	R P	pos	BIRTH	icEEG	III	4.75
14	F	2	39	L T	pos	DD	icEEG	IA	.25
15	F	23	34	R T C	neg	FS	No	IA	.5
16	F	6	28	L Fr	pos	No	icEEG	IA	.58
17	F	20	28	R Fr T	pos	TBI	icEEG	IA	.75
18	F	5	24	R P O	pos	No	SDG + ECoG	II	1
19	F	25	42	R T C	pos	FS	No	IA	1.08
20	F	5	17	R Fr T	pos	BIRTH	icEEG + ECoG	IB	.58
21	M	2.5	19	L Fr	pos	No	ECoG	IA	.42
22	F	.3	19	R T P O	pos	DD	SEEG	IV ^c	n.a. ^d

Note: Overview of 22 consecutive patients, who met the inclusion criteria.

Abbreviations: BIRTH, perinatal stroke; C, central; DD, developmental delay; ECoG, electrocorticography; EEG, electroencephalography; F, female; FAM, family history of seizures; Fr, frontal; FS, febrile seizure; FU, postsurgical follow-up period; icEEG, intracerebral EEG; iEEG, invasive EEG; L, left; M, male; MRI, magnetic resonance imaging; n.a., not available; neg, negative; Npl, neoplasia; O, occipital; P, parietal; pos, positive; R, right; RF, risk factors; SDG, subdural grids; SEEG, stereo-EEG; T, temporal; TBI, traumatic brain injury.

^aBrain localization of surgery.

^bPostsurgical outcome using Engel scale.

^cEngel Class IV outcome with elimination of drop attacks after previous surgery in outside hospital.

^dNo follow-up data available after last surgery at the Cleveland Clinic Epilepsy Center.

the microscopic assessment.¹⁶ The integrated genotype–phenotype classification of brain tumors significantly moved the field forward with more targeted therapies becoming available, despite the lack of resources for genetic testing in many regions of the world. A similar progress in FCD genetics emerged, revealing that somatic and/or germline mutations activating the mTOR–GATOR signaling pathway cause FCD Type 2.^{17–19} Also, somatic mutations in *SCL35A2* leading to aberrant N-glycosylation were reported in FCD 1 or mMCD,^{20,21} which has recently been confirmed and assigned histopathologically to mild malformation of cortical development with oligodendroglial hyperplasia in epilepsy (MOGHE)²² in a collaborative series of 26 cases.²³

To identify the challenges in clinical practice and improve the validity of FCD diagnoses we selected a single center (Cleveland Clinic, Cleveland, Ohio, USA) to access brain

samples surgically resected from a group of highly characterized patients with medically intractable focal epilepsies. These samples were histologically and immunohistochemically reviewed by a group of neuropathologists with experience in the diagnosis of epilepsy-related specimens through a web-based digital pathology platform. DNA extracted from resected brain tissue and peripheral blood from the same patients were genetically studied at five centers to assess the impact of a genotype–phenotype integration for the classification of FCD.^{24,25}

2 | MATERIALS AND METHODS

Twenty-two consecutive cases were selected from the Cleveland Clinic Epilepsy Center Biorepository according to

TABLE 2 Requested immunostains

Case	NeuN	GFAP	MAP2	NFL	SYN	VIM	αBC	CDs	O2	Other
1	x	x	x	x		x	x			CV-LFB
2	x		x	x		x	x			CV-LFB
3	x	x	x	x		x	x			CV-LFB
4	x		x	x		x	x			CV-LFB
5	x		x	x	x	x	x			CV-LFB
6 ^a	x	x	x	x						CV-LFB
7 ^a	x	x	x	x	x	x			x	Ki67
8	x	x	x	x			x	68		Ki67
9 ^a	x	x	x	x					x	IDH1
10 ^a	x	x	x	x				3		CV-LFB
11 ^a	x	x	x	x						CV-LFB
12	x	x	x	x	x	x			x	Ki67
13	x	x	x	x						CV-LFB
14	x		x	x		x				CV-LFB
15 ^a	x	x	x	x				3		CV-LFB
16	x	x	x	x		x	x			CV-LFB
17 ^a	x	x	x	x				3		Ki67
18	x		x	x		x	x			CV-LFB
19 ^a	x	x	x	x					x	CV-LFB
20	x	x	x	x	x	x		68		PB
21	x	x	x	x	x	x	x	68		Ki67
22	x	x	x	x	x	x	x	68		PB

Note: Case identification numbers are the same as used in Table 1. Specific stainings were prepared as indicated (x).

Abbreviations: CDs, T-cell marker CD3 (MRQ-39; Cell Marque); CV-LFB, cresyl violet–Luxol fast blue (Sigma); GFAP, glial fibrillary acidic protein (Dako Cytomation); IDH1, R132H point mutation-specific antibody (clone H09; Dianova); Ki67, proliferation marker (clone MIB-1; Cell Marque); MAP2, microtubule-associated protein 2 (clone HM2; Dako); NeuN, neuronal nuclei (clone A60; Millipore); NFL, nonphosphorylated neurofilament protein (clone SMI32; BioLegend); O2, oligodendroglial marker protein Olig2 (JP18953; IBL International); PB, Prussian blue (Merck); SYN, synaptophysin (SP11; Thermo Fisher Scientific); VIM, vimentin (SP20; Thermo Fisher Scientific); αBC, heat shock protein αB-crystallin (ADI-SPA-223; Enzo).

^aIn these eight cases, more than one formalin-fixed and paraffin-embedded block was requested for special stainings.

the following inclusion criteria: (1) presurgical evaluation and patient management conference protocols suggested a disease etiology within the spectrum of FCD; (2) formalin-fixed and paraffin-embedded (FFPE) surgical specimens were available for microscopic re-evaluation and immunohistochemistry, if requested (see below); (3) fresh frozen surgical brain samples and a matched blood sample would be available for DNA extraction and genetic testing; (4) the selection period included years 2018 and earlier to allow for postsurgical follow-up; and (5) all patients had provided informed consent for the scientific use of their biomaterials (Table 1). The use of biorepository tissues and the study protocol were approved by the Cleveland Clinic Institutional Review Board (IRB#18-1134), the institutional review boards of Boston Children's Hospital, the Committee for the Protection of Persons Ile de France II (No. ID-RCB/EUDRACT-2015-A00671-48) registered on ClinicalTrials.gov (No. NCT02890641), and the Korea Advanced Institute of Science and Technology Institutional Review Board and Committee on Human Research.

2.1 | First and second agreement trials (hematoxylin and eosin vs. immunohistochemistry)

Thirty-five neuropathologists from 18 countries were invited to review the series of 22 cases. A total of 196 hematoxylin and eosin (H&E)-stained FFPE tissue blocks and their H&E-stained sections were retrieved from the pathology archives of the Cleveland Clinic. Whole slide images (WSIs) were digitally scanned using Aperio ScanScope (Leica). All WSIs were uploaded into the open microscopy environment platform for secured web-based access (OMERO.web, University of Dundee). Each case included a short description of the patient's age at surgery, age at seizure onset, whether a lesion was visible by magnetic resonance imaging (MRI), and lobe localization for brain surgery. All reviewers were asked in an open text format to (1) identify the FFPE tissue block with best visible abnormalities; (2) indicate which, if any, additional

immunostainings they would need; and (3) submit a provisional diagnosis. Twenty-four neuropathologists submitted their reports within the given time limit of 6 weeks. None of the reviewers had access to the results of other reviewers. The OMERO.web administrator, who is a senior pathologist neither experienced in epilepsy surgery nor enrolled in this survey, retrieved and forwarded all answers pseudonymously to the study coordinator. Requested immunostainings (see Table 2) were prepared from blocks with most votes using a semiautomated immunostainer (Ventana Ultra), digitally scanned and uploaded onto the OMERO.web platform. After a 3-month interval, the same group of 24 participants was invited to rereview the case series including requested immunostainings. Twenty reviewers submitted their final diagnosis within a 6-week time period. Interobserver agreement was calculated by kappa coefficient analysis.²⁶ Kappa values are always equal to 1 or less. In our study, kappa values were interpreted as follows: <.2, slight; .2–<.4, fair; .4–<.6, moderate; .6–<.8, substantial/good; .8–<1.0, very good/perfect.²⁷

2.2 | Third and fourth agreement trials (Delphi consensus and gene panel analysis)

In a third agreement trial, we applied the Delphi method to build consensus.^{28,29} All answers of the first and second agreement trials were pseudonymously disclosed to the 20 colleagues who completed both previous rounds. Answers were assembled with (1) histopathological landmark images from selected FFPE blocks and requested immunostainings; and (2) a summary of clinical information including preoperative MRI findings, postsurgical outcome, information about invasive diagnostic evaluation, and potential epilepsy risk factors (Table 1). All reviewers were invited to answer an online survey evaluation form (the open access LimeSurvey application) and choose a final diagnosis from the list of all previously submitted diagnoses. The OMERO.web platform remained open for microscopy review during the entire period. All 20 collaborators submitted their Delphi consensus survey within 6 weeks, and interobserver agreement was calculated by kappa coefficient analysis. In December 2019, the Task Force convened at a group meeting to discuss all previous results. Histopathology patterns were reviewed for all cases, regardless of whether consensus agreement had been achieved. Commentaries were prepared to inform the reviewers when anatomical landmarks or clinical histories were perceived as important for the differential diagnosis (Figure S1). The same group of 20 participants was invited to a second, and final, Delphi consensus trial using a similar LimeSurvey questionnaire, which included all previous answers from the third round, assembled with histology

images and the glossary mentioned above. FCD gene panel results collected from four different laboratories became available at this time and were also disclosed to the reviewers. All reviewers submitted their final diagnosis within 8 weeks. Interobserver agreement was calculated by kappa coefficient analysis. Please see Figure 1 for an overview of the study design.

2.3 | Deep targeted sequencing analysis of FCD-relevant genes

DNA extracted from snap-frozen surgical brain tissue and matched blood samples of 22 patients were sent to five genetic laboratories. Each group performed an independent analysis with hybrid capture sequencing targeting their own set of FCD-relevant genes (Table S1). Genetic results from four of the five laboratories were available for the final round of reviews. In Lab A (Daejeon), technical replication of deep sequencing targeting of 13 genes was performed (SoVarGen). Analysis for replicated data was performed in accordance with a previous study.¹⁸ The RePlo variant caller was applied for somatic mutations.³⁰ After a filtering process, all candidate variants were validated using targeted amplicon sequencing (Miseq Dx sequencer; Illumina). Germline mutations were analyzed in a similar pipeline as Genome Analysis Toolkit (GATK)'s best practices. In Lab B (Paris Brain Institute), libraries were prepared and sequenced as previously described,¹⁹ using a targeted panel of 57 genes (Twist Bioscience). Brain somatic variants at low variant allele frequency were validated by droplet digital polymerase chain reaction (ddPCR) as previously described; ddPCR Mutation Detection Assays (FAM + HEX) were purchased from Bio-Rad Laboratories to detect the *MTOR* variant p.Thr1977Lys and *AKT3* variant p.Glu17Lys. For Lab C (Campinas), a hybrid capture panel was used. Four bioinformatics algorithms were used (Mutect2³¹ from GATK, Strelka2,³² VarScan,³³ and Somatic Sniper³⁴; see codes used for bioinformatic analysis in Appendix S1). In Lab D (Boston) a custom double-stranded molecular inversion probe (MIP) panel was used covering 18 FCD-relevant genes (see Table S1). For the MIP design, custom scripts incorporating the MipGen tool were generated that allowed for dense tiling of all exons and up to 60 bp of flanking intronic regions, and putative splicing regulators for all genes. All 33 183 unique MIPs were designed to include a custom backbone consisting of primer binding sites and a dual 8-nt unique molecular index (UMI). MIP were re-balanced in the pool based on the percent of GC content within the regions. MIP pool was amplified with low cycles (17 cycles), high-fidelity polymerase, and custom common primers, allowing for large quantities to be generated.




FIGURE 1 Summary of the four rounds of histopathology agreement. The color coding of pie charts was applied as follows (plain colors for cases with final agreement, shaded colors for cases without agreement; see Figure S1 for more details): gray, focal cortical dysplasia (FCD) 1A, 1B, 1C; green, FCD 2A, 2B; blue, associated FCD Type 3A, 3B, 3C, and 3D; red, no FCD and/or no lesion; yellow, lesions of the white matter (WM); mild malformation of cortical development [MCD] with oligodendroglial hyperplasia in epilepsy defined as oligodendroglial hyperplasia and mild MCD defined as excessive neurons in the WM); purple, MCD, including polymicrogyria, nodular heterotopia, tuberous sclerosis complex; brown, undetermined (undet.). Please note the increase of plain colors from Round 1 to Round 4. Only hematoxylin–eosin (HE) staining was available during Round 1. Immunohistochemistry (IHC) was available during Rounds 2–4. The Delphi consensus method, that is, informed decision based on anonymous disclosure of all previous results, was introduced in Round 3. Results of magnetic resonance imaging review and gene panel sequencing were made available for Round 4

Captured products were amplified using 15 cycles of PCR with dual 8-nt indexing primers and sequenced on the Illumina HiSeq platform with 1×150 -bp paired end reads. All regions were sequenced to a minimum of $\times 1000$ coverage, allowing for accurate assessment of somatic alleles down to .5% alternate allele frequency. Raw sequencing data were analyzed using existing in-house pipelines that, briefly, include quality assessment, mapping using BWA-mem, UMI-collapsing, additional trimming, and local realignment. Finally, variants were called using multiple calling algorithms including Mutect2 and CLC Genomics' Low-Frequency algorithm. Lab E (Amsterdam) carried out additional validation of somatic single nucleotide variants (SNVs) using a dedicated custom Ion AmpliSeq FCD next generation sequencing (NGS) panel and ddPCR for *MTOR*

and *AKT3* variants in Cases 3, 12, 14, 16, 21, and 22 (Table 3; results from individual centers are summarized in Tables S2–S6).

3 | RESULTS

Twenty reviewers completed the four histopathological agreement trials, and 1760 answers were collected for the analysis (Figure S2). In the first round, which was based on 196 H&E stainings, the interobserver agreement was low, with a kappa value of .16. Two cases of FCD 2B were agreed upon by more than 75% of reviewers ($2/22 = 9\%$). An additional 11 cases only reached a majority vote (defined as simple majority) for the diagnosis, as finally agreed upon after the fourth round.

TABLE 3 Summary of consensus genetic findings validated by independent laboratories

Case	Diagnosis ^a	Gene ^b	Amino acid change	Lab A (hybrid capture), brain ^c /blood ^d	Lab B, (hybrid capture), brain ^c /blood ^d	Lab C (hybrid capture), brain ^c /blood ^d	Lab D (MIP), brain ^c /blood ^d	Lab E (validation), brain ^c /blood ^d
3	FCD 2A	<i>DEPDC5</i>	p.Arg874*	46.6%/53.3%	48%/47%	51%/51.1%	19%/39.5%	49.9%/54%
14	FCD 2B	<i>MTOR</i>	p.Thr1977Lys	.3%/1.01%	.4%/0%	.4%/0%	2.8%/1.02%	.3%/nd
22	FCD 2A	<i>MTOR</i>	p.Ser2215Tyr	.6%/1.02%	1.1%/0%	1.1%/1.3%	nd	.6%/nd
21	FCD 2A	<i>AKT3</i>	p.Glu17Lys	1%/0%	.2%/0%	.5%/0%	nd	.9%/nd
16	PMG	<i>NPRL3</i>	p.Ala384fs	47.6%/50%	50%/50%	57.7%/42.8%	nd	47.2%/49.2%
12	MOGHE	<i>SLC35A2</i>	p.Cys210Tyr	51.4%/0%	37%/0%	nd	nd	40.1%/nd
17	MOGHE	<i>SLC35A2</i>	p.Pro15Thr	5.1% ^e /0%	nd	nd	3.3%/0%	nd/nd

Note: Case identification numbers are the same as used in Table 1.

Abbreviations: FCD, focal cortical dysplasia; MIP, molecular inversion probe; MOGHE, mild malformation of cortical development with oligodendroglial hyperplasia in epilepsy; nd, not detected due to low coverage or absence of the gene in the targeted panel; PMG, polymicrogyria.

^aFinal histopathology diagnosis after the fourth round.

^b*DEPDC5* transcript ID: NM_001242896, *MTOR* transcript ID: NM_004958, *AKT3* transcript ID: NM_005465, *NPRL3* transcript ID: NM_001077350, *SLC35A2* transcript ID: NM_005660.

^cVariant allele frequency detected in DNA from fresh frozen surgical brain sample.

^dVariant allele frequency detected in DNA from a peripheral blood draw.

^eMutation-negative in bulk frozen tissue, but mutation-positive in pathologically reviewed formalin-fixed and paraffin-embedded tissue samples.

We added 10 different antibody stainings to the second round (Table 2), which yielded >75% agreement in two additional cases of FCD 2B and two cases of FCD 2A (6/22 = 27%). This resulted in an improvement of the interobserver agreement to “fair” (kappa value of .35). The Delphi consensus method was applied in the third round and reached a moderate interobserver agreement, with a kappa value of .5. Diagnosis for nine cases was agreed upon by more than 75% of reviewers (41%), seven of which were FCD Type 2A or 2B. An agreement was reached by more than 75% of reviewers for 14 cases in the fourth round (64%). An agreement was reached for all subtypes of FCD 2, polymicrogyria, FCD 3 associated with perinatal stroke (FCD Type 3D), MOGHE, and a case with no lesion on MRI and histopathologically. The overall kappa value of .68 was “substantial.” Simple majority votes without consensus agreement were obtained for FCD Type 1A, an MRI-negative mMCD, and an MRI-negative nonlesional case, as well as four cases with a principal lesion and no associated FCD (three with HS and one with a tumor).

Eight cases finally achieved a consensus agreement for FCD Type 2A and 2B ($n = 4$ each); 86.7% of all FCD Type 2 diagnostic labels were already assigned to these eight cases in the first round, and this increased to 97.5% in the fourth round. Results differed in FCD Type 1. Only 20% of all FCD Type 1 labels were assigned in the first round to Case 5, which received the final majority vote for FCD Type 1A (Figure 2H). FCD Type 1 labels were scattered among all other cases and represented a common differential diagnosis for mMCD.

FCD Type 3 labels represented 14.3% of all answers in the first round, which increased to 17.5% in Round 2 and

20.5% in Round 3 (Figure 1, Figure S2). FCD 3A was the most frequent differential diagnosis in all cases with HS, representing 33% of all answers in the first, 50% in the second, and 56% in the third round. A Task Force group meeting was organized to clarify common patterns of FCD Type 3A recognized by the current ILAE classification scheme and provided as a commentary to the reviewers. In the fourth round, only 28% of answers selected FCD 3A in cases with HS and all cases reached a majority vote for “HS without FCD Type 3A” (Figure S2). Instead, 61.6% of all FCD 3 labels in Round 4 were assigned to both cases with the final diagnosis of FCD 3D (Figure 2J).

Only 4.7%–8.8% of all answers specified mMCD, which is defined as excessive (heterotopic) neurons in the white matter.⁹ Another lesion preferentially affecting the white matter was mMCD with oligodendroglial hyperplasia, for example, MOGHE, which also showed heterotopic neurons in the white matter,²² and collected 3.5%–7% of all answers. Based on consensus diagnosis, two cases reached the final agreement of MOGHE (Figure 2D). Interestingly, both cases revealed a somatic brain mutation for the *SLC35A2* gene (Table 3). The widespread distribution of diagnostic MOGHE labels observed during Rounds 1–3 collated to the two *SLC35A2* mutant cases after disclosure of the genetic information in the fourth round. The same holds true for one case with polymicrogyria, which was difficult to identify in the oblique cutting plane of the neocortical specimen. A germline *NPRL3* mutation was detected in blood and tissue and the diagnosis reached agreement status in the fourth round, as already suggested by a majority after the second round.




FIGURE 2 Illustrative cases of the patient series. (A) Patient 3 with focal cortical dysplasia (FCD) Type 2A and *DEPDC5* germline mutation. The arrow in the fluid-attenuated inversion recovery (FLAIR) image points to a lesion and site of surgical resection. (B) Surgical specimen A7 with an area of dysmorphic neurons (arrow) as further magnified in C (immunostain against nonphosphorylated neurofilaments with antibody SMI32) NCx = neocortex. (D) Patient 12 with mild malformation of cortical development with oligodendroglial hyperplasia in epilepsy and *SLC35A2* brain somatic mutation. The arrow in the FLAIR image points to a lesion and site of surgical resection. (E) Oligodendrocyte lineage transcription factor 2 (Olig2) immunohistochemistry revealed increased cell density at the gray–white matter border (arrow). (F) Oligodendroglial hyperplasia (arrow) visible at high-power magnification. Please compare the region of high cell density (arrow) with a region of deep white matter and normal oligodendroglial cell density (asterisk). (G) Patient 5 with FCD 1A. The arrow in the T2 image points to a lesion and site of surgical resection. (H) NeuN immunohistochemistry revealed a predominant microcolumnar architecture of the neocortex (arrow). (I) Patient 20 with FCD Type 3D. The arrow in the FLAIR image points to a lesion and site of surgical resection. (J) Nodules of neurons (arrow) were embedded in a glial scar as a hallmark of FCD Type 3D in this patient with perinatal stroke. Scale bar in C = 250µm, in F = 500µm and J = 2.5 mm.

All genetic laboratories independently performed genetic analysis of somatic and germline mutations in FCD-relevant genes (Table S1). To detect low-frequency somatic mutations, high-depth hybrid capture sequencing of FCD-relevant genes was performed (>×1000 read depth), followed by comprehensive bioinformatic analysis as described in the Materials and Methods section. All somatic and germline mutations were validated by alternative sequencing methods including target-specific amplicon sequencing, ddPCR, and Sanger sequencing. Most laboratories reproducibly found somatic brain SNVs in *MTOR*, *AKT3*, and *SLC35A2* in four patients and germline mutations in *DEPDC5* and *NPRL3* in two patients. Six panel-negative cases with a histopathologically verified lesion after the third histopathology review (Cases 1, 4, and 18 with FCD 2B, Case 2 with FCD 2A, Case 5 with FCD 1A, and Case 17 with MOGHE) were subjected

to a second round of genetic testing using DNA extracted from FFPE slides of a paraffin block containing the lesion. In this study, a new somatic brain mutation was identified in *SLC35A2* in Case 17 with MOGHE. As a result, genetic tests of FCD-relevant genes performed by independent laboratories identified genetic causes in seven of 22 (31.8%) patients including five of 22 (22.7%) patients with somatic brain mutations and two of 22 (9%) patients with germline mutations.

4 | DISCUSSION

Despite the ubiquitous use of the international consensus classification of FCD in research and clinical practice, the histopathological characterization of FCD subtypes remains a matter of ongoing debate.⁷ The presented results show

that the diagnosis of (1) FCD Type 2 subtypes can be improved using a selected protocol of immunostains, for example, NeuN, nonphosphorylated neurofilament, vimentin, and microtubule-associated protein 2 (Table 2) and genetic tests¹⁸; (2) FCD Type 1 and 3 subtypes remain difficult to differentiate from the normal variation in cortical architecture, particularly in the temporal lobe; (3) new clinicopathological and genetically defined subtypes can be recognized and should be added to the current FCD classification scheme.

Overall, there was 80% accuracy in identifying the eight FCD 2 cases of this series. However, an agreement was achieved in only two cases (25%) when the diagnosis was based solely on H&E sections. This number increased to six cases (75%) when a panel of immunohistochemical stainings was provided, as recommended by an ILAE Task Force in 2016.³⁵ NeuN staining was most helpful in deciphering the six-layered architecture of the neocortex. This highlighted the value of NeuN immunohistochemistry for routine application in order not to overlook focal regions of dysplasia in large resection specimens. NeuN immunoreactivity also helped to anticipate the regional variability of the laminar neuroanatomical organization (Figure S1). Antibodies directed against nonphosphorylated neurofilaments (SMI32³⁶) are sensitive markers to identify dysmorphic neurons in all FCD 2 subtypes, and dense accumulation of neurofilaments in large cell bodies of up to 50 µm diameter must be considered pathognomonic. Of note, SMI32 labeling remains weak in the normal neocortex of children and young adults but increases with age.³⁷ Vimentin or αB-crystallin were frequently requested to confirm the presence of balloon cells in FCD Type 2B. However, both antibodies also label reactive astrocytes. Their distinction from vimentin- or αB-crystallin-positive balloon cells can be challenging, therefore, whereas balloon cells do not exhibit cellular processes or end feet extending toward capillaries.

Consistent with previous genetic studies of FCD, pathogenic somatic and germline mutations in mTOR or GATOR pathway genes were identified only in FCD Type 2.^{17–19} Interestingly, we observed a pathogenic germline *DEPDC5* mutation in one case of FCD Type 2A. *DEPDC5* mutations have been assigned mostly to FCD Type 2A^{19,38–40} but were also reported in other FCD subtypes.⁴¹ In the case of *DEPDC5*, a repressor of the mTOR pathway, a second-hit somatic mutation has been shown to lead to the development of dysmorphic neurons in the dysplastic lesion.^{20,42} This second hit is yet to be identified in Cases 3 and 16, however. Regarding the laboratory requirements for the reliable detection of low-level somatic mutations implicated in FCD, our result suggested that FCD genetic tests require (1) hybrid capture and high-depth NGS sequencing (>×1000 read depth) of FCD-relevant genes; (2) the use of somatic mutation callers, for example, MuTect2, Replow, and Strelka2; and (3) the validation sequencing of candidate variations

using orthogonal technology, for example, ddPCR or target site-specific amplicon sequencing. Although genetic testing of somatic and germline mutations for FCD is not yet available in most pathology laboratories, it is important information for the genetic consultation whether FCD patients carry pathogenic somatic (not inherited, not transmissible) or germline (inherited and transmissible) variants. A hyperactivated mTOR kinase due to somatic or germline mutations in FCD Type 2 is also a promising candidate for targeted drug treatment of mTOR inhibitor (ClinicalTrials.gov identifier NCT03198949).⁴³ Thus, genetic diagnosis will be helpful and add an objective measure for an integrated diagnosis of FCD, thereby better stratifying the patient cohort, advancing our knowledge about the underlying disease pathomechanism, and promoting personalized medicine in epileptology.

MOGHE is a recently described new entity observed in young children with early onset and difficult-to-treat frontal lobe epilepsy, and histopathologically defined by oligodendroglial hyperplasia and heterotopic neurons in the shallow subcortical white matter.^{22,44} There was a majority vote from only 23% of reviewers in the first and second rounds for Cases 12 and 17. The Delphi process assisted most reviewers in recognizing this entity. Surprisingly, *SLC35A2* somatic variants were identified in both MOGHE cases. MOGHE neuropathology has only recently been associated with *SLC35A2* mutations in an unprecedented cohort of 26 cases²³ including those *SLC35A2* mutated cases previously published in the literature as mMCD or FCD Type 1.^{19–21,45} Both differentials were also commonly assigned to MOGHE cases in our study. We suggest, however, recognizing MOGHE diagnosis by the ILAE consensus classification scheme as a specific disease entity preferentially affecting the white matter (Table 4).

FCD 1 subtypes pose a major challenge in clinical practice.^{7,13,46} Their histopathological hallmarks comprise a large spectrum of horizontal or vertical abnormalities of the six-layered neocortex.¹¹ The human neocortex reveals different architectural patterns across the cortical ribbon (Figure S1), which are cytologically classified into 52 Brodmann areas.⁴⁷ However, it is almost impossible to recognize these Brodmann areas on a thin H&E-stained section obtained from an epilepsy surgery tissue sample. Clinical data, as well as knowledge of the cortical region resected, are essential when considering FCD Type 1 diagnoses. We finally agreed, therefore, that FCD Type 1 was unlikely in Patient 8, in whom seizures started after a motor vehicle accident at age 49 years (Table 1), and who received 15%–20% of all FCD Type 1 votes in Rounds 1, 2, and 3. In contrast, histopathologically well-documented patients with FCD Type 1A were reported to have early seizure onset, daily seizures, and a developmental delay.^{13,46}

The FCD Type 3 subgroup presented another diagnostic challenge during this agreement trial. The ILAE classification has specified several histopathology patterns and conditions

TABLE 4 Proposed amendments to the FCD classification scheme

Subtypes		
FCD Type 1	FCD 1A, vertical microcolumns	FCD 1B, abnormal layering
FCD Type 2	FCD 2A, dysmorphic neurons	FCD 1C, vertical and horizontal abnormalities
FCD Type 3	FCD 3A, cortical dyslamination associated with HS	FCD 2B, dysmorphic neurons and balloon cells
White matter lesions	mMCD ^a with excessive heterotopic neurons	FCD 3C, cortical dyslamination adjacent to vascular malformation
No FCD	When histopathological findings are not compatible with FCD Type 1, 2, or 3 ^b	

Abbreviations: FCD, focal cortical dysplasia; HS, hippocampal sclerosis; mMCD, mild malformation of cortical development; MOGHE, mMCD with oligodendroglial hyperplasia in epilepsy.

^amMCD not associated with any other principal lesion, such as hippocampal sclerosis, brain tumor, or vascular malformation; for reference values see.^{50–53}

^bNo FCD: a descriptive report is recommended to highlight anatomical ambiguities (if applicable).

for this FCD category. A pattern of temporal lobe sclerosis as defined by neuronal cell loss and astrogliosis in neocortical Layers 2 and 3 in patients with HS has been assigned to FCD Type 3A.¹¹ Lentiform heterotopias in the white matter of patients with HS⁴⁹ was also assigned to FCD 3A. It came as no surprise, therefore, that 65% of reviewers recognized these lentiform heterotopias in Patient 11 as FCD Type 3A (Figure S1). However, neuroanatomists would classify lentiform heterotopias as part of the ventral claustrum, fragmented by fibers of the fasciculus uncinatus,⁵⁴ which frequently extend into the whiter matter of the superior temporal gyrus, an area often included in the surgical resection for epilepsy. This information was disclosed to our reviewers in Round 4, and the final majority vote was HS, no FCD.

In addition, this agreement study highlighted cases where no FCD was reported despite focal epilepsy and subtle features on MRI; "no FCD" was used in 10% in Round 1, 13% in Round 2, and 12% in Round 3, and was finally assigned by 70% of the reviewers to Case 8 and 85% of the reviewers to Case 15. We propose that "no FCD" should be incorporated into the diagnostic framework of the international FCD classification scheme (Table 4). It needs to be recognized that even where dysplastic cortical lesions are suspected on MRI (or quantitative MRI) there may not be a confident FCD diagnosis possible within the current limitations of histology and genetic analysis. Acknowledging this option may reduce any tendency to overdiagnose, particularly the FCD 1 and FCD 3 subtypes.

In conclusion, the international FCD classification scheme is widely accepted in research and clinical practice and is helpful in stratifying clinicopathological disease conditions. Inclusion of newly recognized disease categories, a better definition of histopathological features, and genotype integration will help to improve the reliability of diagnosing FCD. This may also stimulate our research efforts into elucidating disease pathomechanisms and the development of targeted therapies.

ACKNOWLEDGMENTS

I.B. is funded by the German Research Council (DFG BI 421/4-1). M.M.-S. is funded by the National Institutes of Health (National Institute of Neurological Disorders and Stroke R01 NS097719). This work was funded by the European Research Council (No. 682345 to S.Bau.), the program "Investments of the Future" (ANR-10-IAIHU-06 and ANR-18-RHUS-0005 to S.Bau.), and the Cleveland Clinic Epilepsy Center. R.G. is funded by the Italian Ministry of Health. E.A. is funded by ZonMw, Programme Translational Research (project number 95105004). S.Bi. was supported by the Manton Center for Orphan Disease Research at Boston Children's Hospital. F.C. and I.L.-C. are funded by São Paulo Research Foundation (FAPESP) grant No. 2013/07559-3. T.S.J. is grateful for funding from the Brain Tumour Charity,

Children with Cancer UK, Great Ormond Street Hospital (GOSH) Children's Charity, Olivia Hodson Cancer Fund, Cancer Research UK, and the National Institute of Health Research (NIHR). All research at GOSH National Health Service Foundation Trust and University College London Great Ormond Street Institute of Child Health is made possible by the NIHR GOSH Biomedical Research Centre. The views expressed are those of the authors and not necessarily those of the National Health Service, the NIHR, or the Department of Health. We cordially thank Lisa Ferguson and Christina O'Connor from the Cleveland Clinic for their assistance with data collection and tissue preparation, respectively. Also, we would like to acknowledge Jennifer Neil Partlow, research coordinator/genetic counselor, for assistance at Boston Children's Hospital.

CONFLICT OF INTEREST

None of the authors has any conflict of interest to disclose, with the exception of J.H.L., who is a cofounder and chief technology officer of SoVarGen, which seeks to develop new diagnostics and therapeutics for brain disorders.

ORCID

Ingmar Blümcke  <https://orcid.org/0000-0001-8676-0788>

Robyn M. Busch  <https://orcid.org/0000-0002-5442-4912>

Fernando Cendes  <https://orcid.org/0000-0001-9336-9568>

Iscia Lopes-Cendes  <https://orcid.org/0000-0002-6221-6822>

Stephanie Baulac  <https://orcid.org/0000-0001-6430-4693>

Sara Baldassari  <https://orcid.org/0000-0003-4851-2796>

Angelika Mühlebner  <https://orcid.org/0000-0001-9102-7353>

Albert Becker  <https://orcid.org/0000-0003-2661-3705>

Rita Garbelli  <https://orcid.org/0000-0002-6475-4974>

Maria Thom  <https://orcid.org/0000-0001-7712-2629>

Anita Mahadevan  <https://orcid.org/0000-0002-4837-1463>

Hajime Miyata  <https://orcid.org/0000-0003-2551-5978>

REFERENCES

- Blumcke I, Russo GL, Najm I, Palmini A. Pathology-based approach to epilepsy surgery. *Acta Neuropathol.* 2014;128:1–3.
- Lamberink HJ, Otte WM, Blümcke I, Braun KPJ, Aichholzer M, Amorim I, et al. Seizure outcome and use of antiepileptic drugs after epilepsy surgery according to histopathological diagnosis: a retrospective multicentre cohort study. *Lancet Neurol.* 2020;19:748–57.
- Wiebe S, Blume WT, Girvin JP, Eliasziw M. A randomized, controlled trial of surgery for temporal-lobe epilepsy. *N Engl J Med.* 2001;345:311–8.
- Baud MO, Perneger T, Racz A, Pensel MC, Elger C, Rydenhag B, et al. European trends in epilepsy surgery. *Neurology.* 2018;91:e96–106.
- Blumcke I, Spreafico R, Haaker G, Coras R, Kobow K, Bien CG, et al. Histopathological findings in brain tissue obtained during epilepsy surgery. *N Engl J Med.* 2017;377:1648–56.
- Taylor DC, Falconer MA, Bruton CJ, Corsellis JA. Focal dysplasia of the cerebral cortex in epilepsy. *J Neurol Neurosurg Psychiatry.* 1971;34:369–87.
- Najm IM, Sarnat HB, Blümcke I. Review: the international consensus classification of focal cortical dysplasia—a critical update 2018. *Neuropathol Appl Neurobiol.* 2018;44:18–31.
- Tassi L, Colombo N, Garbelli R, Francione S, Lo Russo G, Mai R, et al. Focal cortical dysplasia: neuropathological subtypes, EEG, neuroimaging and surgical outcome. *Brain.* 2002;125:1719–32.
- Palmini A, Najm I, Avanzini G, Babb T, Guerrini R, Foldvary-Schaefer N, et al. Terminology and classification of the cortical dysplasias. *Neurology.* 2004;62:S2–8.
- Chamberlain WA, Cohen ML, Gyure KA, Kleinschmidt-DeMasters BK, Perry A, Powell SZ, et al. Interobserver and intraobserver reproducibility in focal cortical dysplasia (malformations of cortical development). *Epilepsia.* 2009;50:2593–8.
- Blumcke I, Thom M, Aronica E, Armstrong DD, Vinters HV, Palmini A, et al. The clinico-pathological spectrum of focal cortical dysplasias: a consensus classification proposed by an ad hoc Task Force of the ILAE Diagnostic Methods Commission. *Epilepsia.* 2011;52:158–74.
- Tassi L, Garbelli R, Colombo N, Bramerio M, Lo Russo G, Deleo F, et al. Type I focal cortical dysplasia: surgical outcome is related to histopathology. *Epileptic Disord.* 2010;12:181–91.
- Blumcke I, Pieper T, Pauli E, Hildebrandt M, Kudernatsch M, Winkler P, et al. A distinct variant of focal cortical dysplasia type I characterised by magnetic resonance imaging and neuropathological examination in children with severe epilepsies. *Epileptic Disord.* 2010;12:172–80.
- Coras R, de Boer OJ, Armstrong D, Becker A, Jacques TS, Miyata H, et al. Good interobserver and intraobserver agreement in the evaluation of the new ILAE classification of focal cortical dysplasias. *Epilepsia.* 2012;53:1341–8.
- Ramaswamy V, Taylor MD. Fall of the optical wall: freedom from the tyranny of the microscope improves glioma risk stratification. *Cancer Cell.* 2016;29:137–8.
- Louis DN, Perry A, Reifenberger G, von Deimling A, Figarella-Branger D, Cavenee WK, et al. The 2016 World Health Organization Classification of Tumors of the Central Nervous System: a summary. *Acta Neuropathol.* 2016;131:803–20.
- Lim JS, Kim WI, Kang HC, Kim SH, Park AH, Park EK, et al. Brain somatic mutations in MTOR cause focal cortical dysplasia type II leading to intractable epilepsy. *Nat Med.* 2015;21:395–400.
- Sim NS, Ko A, Kim WK, Kim SH, Kim JS, Shim KW, et al. Precise detection of low-level somatic mutation in resected epilepsy brain tissue. *Acta Neuropathol.* 2019;138:901–12.
- Baldassari S, Ribierre T, Marsan E, Adle-Biasette H, Ferrand-Sorbets S, Bulteau C, et al. Dissecting the genetic basis of focal cortical dysplasia: a large cohort study. *Acta Neuropathol.* 2019;138:885–900.
- Sim NS, Seo Y, Lim JS, Kim WK, Son H, Kim HD, et al. Brain somatic mutations in SLC35A2 cause intractable epilepsy with aberrant N-glycosylation. *Neurol Genet.* 2018;4:e294.
- Winawer MR, Griffin NG, Samanamud J, Baugh EH, Rathakrishnan D, Ramalingam S, et al. Somatic SLC35A2 variants in the brain are associated with intractable neocortical epilepsy. *Ann Neurol.* 2018;83:1133–46.
- Schurr J, Coras R, Rossler K, Pieper T, Kudernatsch M, Holthausen H, et al. Mild malformation of cortical development with oligodendroglial hyperplasia in frontal lobe epilepsy: a new clinicopathological entity. *Brain Pathol.* 2017;27:26–35.
- Bonduelle T, Hartlieb T, Baldassari S, Sim NS, Kim SH, Kang HC, et al. Frequent SLC35A2 brain mosaicism in mild malformation

- of cortical development with oligodendroglial hyperplasia in epilepsy (MOGHE). *Acta Neuropathol Commun*. 2021;9:3.
24. Benova B, Jacques TS. Genotype-phenotype correlations in focal malformations of cortical development: a pathway to integrated pathological diagnosis in epilepsy surgery. *Brain Pathol*. 2019;29:473–84.
 25. Blumcke I. It is time to move on: Commentary to: Genotype-phenotype correlations in focal malformations of cortical development: a pathway to integrated pathological diagnosis in epilepsy surgery. *Brain Pathol*. 2019;29:467–8.
 26. Sim J, Wright CC. The kappa statistic in reliability studies: use, interpretation, and sample size requirements. *Phys Ther*. 2005;85:257–68.
 27. Landis JR, Koch GG. The measurement of observer agreement for categorical data. *Biometrics*. 1977;33:159–74.
 28. Nair R, Aggarwal R, Khanna D. Methods of formal consensus in classification/diagnostic criteria and guideline development. *Semin Arthritis Rheum*. 2011;41:95–105.
 29. Sauro KM, Wiebe S, Macrodimitris S, Jetté N, EMU Quality Improvement Team. Quality indicators for the adult epilepsy monitoring unit. *Epilepsia*. 2016;57:1771–8.
 30. Kim J, Kim D, Lim JS, Maeng JH, Son H, Kang HC, et al. The use of technical replication for detection of low-level somatic mutations in next-generation sequencing. *Nat Commun*. 2019;10:1047.
 31. Cibulskis K, Lawrence MS, Carter SL, Sivachenko A, Jaffe D, Sougnez C, et al. Sensitive detection of somatic point mutations in impure and heterogeneous cancer samples. *Nat Biotechnol*. 2013;31:213–9.
 32. Kim S, Scheffler K, Halpern AL, Bekritsky MA, Noh E, Kallberg M, et al. Strelka2: fast and accurate calling of germline and somatic variants. *Nat Methods*. 2018;15:591–4.
 33. Koboldt DC, Zhang Q, Larson DE, Shen D, McLellan MD, Lin L, et al. VarScan 2: somatic mutation and copy number alteration discovery in cancer by exome sequencing. *Genome Res*. 2012;22:568–76.
 34. Larson DE, Harris CC, Chen K, Koboldt DC, Abbott TE, Dooling DJ, et al. SomaticSniper: identification of somatic point mutations in whole genome sequencing data. *Bioinformatics*. 2012;28:311–7.
 35. Blumcke I, Aronica E, Miyata H, Sarnat HB, Thom M, Roessler K, et al. International recommendation for a comprehensive neuropathologic workup of epilepsy surgery brain tissue: a consensus Task Force report from the ILAE Commission on Diagnostic Methods. *Epilepsia*. 2016;57:348–58.
 36. Sternberger LA, Sternberger NH. Monoclonal antibodies distinguish phosphorylated and nonphosphorylated forms of neurofilaments in situ. *Proc Natl Acad Sci U S A*. 1983;80:6126–30.
 37. Sternberger NH, Sternberger LA, Ulrich J. Aberrant neurofilament phosphorylation in Alzheimer disease. *Proc Natl Acad Sci U S A*. 1985;82:4274–6.
 38. Baulac S, Ishida S, Marsan E, Miquel C, Biraben A, Nguyen DK, et al. Familial focal epilepsy with focal cortical dysplasia due to DEPDC5 mutations. *Ann Neurol*. 2015;77:675–83.
 39. Scerri T, Riseley JR, Gillies G, Pope K, Burgess R, Mandelstam SA, et al. Familial cortical dysplasia type IIA caused by a germline mutation in DEPDC5. *Ann Clin Transl Neurol*. 2015;2:575–80.
 40. Carvill GL, Crompton DE, Regan BM, McMahon JM, Saykally J, Zemel M, et al. Epileptic spasms are a feature of DEPDC5 mTORopathy. *Neurol Genet*. 2015;1:e17.
 41. Marsan E, Review BS. Review: Mechanistic target of rapamycin (mTOR) pathway, focal cortical dysplasia and epilepsy. *Neuropathol Appl Neurobiol*. 2018;44(1):6–17.
 42. Ribierre T, Deleuze C, Bacq A, Baldassari S, Marsan E, Chipaux M, et al. Second-hit mosaic mutation in mTORC1 repressor DEPDC5 causes focal cortical dysplasia-associated epilepsy. *J Clin Invest*. 2018;128:2452–8.
 43. Krueger DA, Care MM, Holland K, Agricola K, Tudor C, Mangeskar P, et al. Everolimus for subependymal giant-cell astrocytomas in tuberous sclerosis. *N Engl J Med*. 2010;363:1801–11.
 44. Hartlieb T, Winkler P, Coras R, Pieper T, Holthausen H, Blumcke I, et al. Age-related MR characteristics in mild malformation of cortical development with oligodendroglial hyperplasia and epilepsy (MOGHE). *Epilepsy Behav*. 2018;91:68–74.
 45. Miller KE, Koboldt DC, Schieffer KM, Bedrosian TA, Crist E, Sheline A, et al. Somatic SLC35A2 mosaicism correlates with clinical findings in epilepsy brain tissue. *Neurol Genet*. 2020;6:e460.
 46. Hildebrandt M, Pieper T, Winkler P, Kolodziejczyk D, Holthausen H, Blumcke I. Neuropathological spectrum of cortical dysplasia in children with severe focal epilepsies. *Acta Neuropathol*. 2005;110:1–11.
 47. Brodmann K. Beiträge zur histologischen Lokalisierung der Grosshirnrinde. VI. Mitteilung: die Cortexgliederung des Menschen. *J Physiol Neurol*. 1908;10:231–46.
 48. Thom M, Eriksson S, Martinian L, Caboclo LO, McEvoy AW, Duncan JS, et al. Temporal lobe sclerosis associated with hippocampal sclerosis in temporal lobe epilepsy: neuropathological features. *J Neuropathol Exp Neurol*. 2009;68:928–38.
 49. Meroni A, Galli C, Bramerio M, Tassi L, Colombo N, Cossu M, et al. Nodular heterotopia: a neuropathological study of 24 patients undergoing surgery for drug-resistant epilepsy. *Epilepsia*. 2009;50:116–24.
 50. Emery JA, Roper SN, Rojiani AM. White matter neuronal heterotopia in temporal lobe epilepsy: a morphometric and immunohistochemical study. *J Neuropathol Exp Neurol*. 1997;56(12):1276–82. <https://doi.org/10.1097/00005072-199712000-00002>
 51. Mühlebner A, Coras R, Kobow K, Feucht M, Czech T, Stefan H, et al. Neuropathologic measurements in focal cortical dysplasias: validation of the ILAE 2011 classification system and diagnostic implications for MRI. *Acta Neuropathol*. 2012;123(2):259–72. <https://doi.org/10.1007/s00401-011-0920-1>
 52. Rojiani AM, Emery JA, Anderson KJ, Massey JK. Distribution of heterotopic neurons in normal hemispheric white matter: a morphometric analysis. *J Neuropathol Exp Neurol*. 1996;55(2):178–83. <https://doi.org/10.1097/00005072-199602000-00006>
 53. Thom M, Sisodiya S, Harkness W, Scaravilli F. Microdysgenesis in temporal lobe epilepsy. A quantitative and immunohistochemical study of white matter neurones. *Brain*. 2001;124(11):2299–309. <https://doi.org/10.1093/brain/124.11.2299>
 54. Rae AS. The form and structure of the human claustrum. *J Comp Neurol*. 1954;100:15–39.

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section.

How to cite this article: Blümcke I, Coras R, Busch RM, et al. Toward a better definition of focal cortical dysplasia: An iterative histopathological and genetic agreement trial. *Epilepsia*. 2021;62:1416–1428. <https://doi.org/10.1111/epi.16899>