



Systematic Review and Meta-Analysis

The performance of mid-upper arm circumference for identifying children and adolescents with overweight and obesity: a systematic review and meta-analysis

Binyam Girma Sisay^{1,*} , Hamid Yimam Hassen² , Beshada Rago Jima¹, Evan Atlantis^{3,4} and Seifu Hagos Gebreyesus¹

¹Department of Nutrition and Dietetics, School of Public Health, Addis Ababa University, Addis Ababa, Ethiopia:

²Department of Primary and Interdisciplinary Care, Faculty of Medicine and Health Sciences, University of Antwerp, Antwerp, Belgium: ³School of Health Sciences, Western Sydney University, Penrith, NSW, Australia: ⁴Discipline of Medicine, Nepean Clinical School, Faculty of Medicine and Health, The University of Sydney, Nepean, NSW, Australia

Submitted 20 August 2021: Final revision received 20 December 2021: Accepted 12 January 2022: First published online 17 January 2022

Abstract

Objective: This study aimed to synthesise the existing evidence on the performance of mid-upper arm circumference (MUAC) to identify children and adolescents with overweight and obesity.

Design: Systematic review and meta-analysis.

Setting: We searched PubMed, EMBASE, SCOPUS, Cochrane Library, Web of Science, CINAHL and Google scholar databases from their inception to December 10, 2021, for relevant studies. There were no restrictions regarding the language of publication. Studies reporting measures for the diagnostic performance of MUAC compared with a reference standard for diagnosing overweight and obesity in children and adolescents aged 2–19 years were included.

Participants: A total of 54 381 children and adolescents from twenty-one studies were reviewed; ten studies contributed to meta-analyses.

Results: In boys, MUAC showed a pooled AUC of 0.92 (95% CI 0.89, 0.94), sensitivity of 84.4 (95% CI 84.6, 90.8) and a specificity of 86.0 (95% CI 79.2, 90.8), when compared against BMI z-score, defined overweight and obesity. As for girls, MUAC showed a pooled AUC of 0.93 (95% CI 0.90, 0.95), sensitivity of 86.4 (95% CI 79.8, 91.0), specificity of 86.6 (95% CI 82.2, 90.1) when compared against overweight and obesity defined using BMI z-scores.

Conclusion: In comparison with BMI, MUAC has an excellent performance to identify overweight and obesity in children and adolescents. However, no sufficient evidence on the performance of MUAC compared with gold standard measures of adiposity. Future research should compare performance of MUAC to the 'golden standard' measure of excess adiposity.

Keywords

Mid-upper arm circumference
ROC curve
Overweight
Obesity
Children and adolescents

Overweight and obesity in childhood and adolescence is one of the greatest public health problems facing most countries in the world, which has increased dramatically in recent decades^(1,2). The global prevalence of obesity in children and adolescents was 5.6% in girls and 7.8% in boys, respectively, in 2016⁽²⁾. Childhood obesity can persist into adulthood, leading to an increased risk of chronic

non-communicable diseases and premature mortality^(3–6). Timely recognition and early diagnosis of overweight and obesity in young people are essential to mitigate the short- and long-term health risks associated with excess body weight, especially fat mass^(7,8). Health care professionals need to use an accurate and efficient screening tool to diagnose overweight and obesity for early intervention⁽⁹⁾.

*Corresponding author: Email binyamgirma3@gmail.com

© The Author(s), 2022. Published by Cambridge University Press on behalf of The Nutrition Society. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited.

According to the WHO, obesity is defined as an abnormal or excessive fat accumulation that may impair health⁽¹⁰⁾. Total body fat can be accurately measured using several methods such as hydrostatic weighing, air displacement plethysmography, deuterium oxide dilution or dual-energy X-ray absorptiometry, which estimates total body fat by measuring total body water and based on the absorption patterns of X-rays^(11,12). However, these methods are laboratory-based, expensive, time-consuming and not feasible for routine use⁽¹³⁾.

Alternatively, there are a range of anthropometric measurements that are simple and inexpensive options for overweight and obesity screening in children and adolescents⁽¹⁴⁾. However, the results of anthropometric measurements need to be interpreted using reference standards to define overweight and obesity in young people. While the BMI z-score is a widely utilised method to identify those with overweight and obesity in epidemiological studies but has limited applicability in routine clinical practice^(2,15). Simple and inexpensive alternatives to BMI z-scoring would be helpful to promote screening and early identification, especially in low- and middle-income countries with limited health care resources⁽¹⁶⁾.

The mid-upper arm circumference (MUAC) has been proposed as one such alternative to screen for overweight and obesity in children and adolescents^(17–20). It is a simple measure commonly used to screen for undernutrition in infants and children aged 6–59 months⁽²¹⁾ as well as thinness and severe thinness in adolescents^(22,23). The existing evidence of the usefulness of the MUAC against the BMI Z-score among children and adolescents is limited and unclear^(17–20,24–26). Therefore, this systematic review and meta-analysis aim to summarise the currently available evidence on the performance of MUAC to identify children and adolescents with overweight and obesity.

Methods

This systematic review and meta-analysis were conducted per the Preferred Reporting Items for Systematic Reviews and Meta-Analyses for Diagnostic Test (PRISMA-DTA) statement (see online supplementary material, Supplemental Table S1)⁽²⁷⁾. The protocol of this systematic review and meta-analysis was registered on PROSPERO (Registration number CRD42020183148) and published⁽²⁸⁾; minor deviations from the original protocol have also been explained (see online supplementary material, Supplemental Table S2).

Review question

This systematic review aims to generate an evidence summary to answer the following review question: what is the diagnostic performance of the MUAC assessment for diagnosing overweight and obesity in children and adolescents.

Search strategy and study selection

A systematic search was performed in PubMed, EMBASE, SCOPUS, Cochrane Library, Web of Science and CINAHL database of references for peer-reviewed articles. To retrieve grey literature, systematic search was performed using Google Scholar. The databases were systematically searched from their inception to December 10, 2021. A detailed search strategy is provided in the supplementary material (see online supplementary material, Supplemental Table S3).

All articles identified through the systematic search databases were imported into EndNote as a single library. Duplicate articles from the searches were verified and removed. The remaining articles were imported into rayyan.QCRI.org⁽²⁹⁾, a web-based tool that facilitates screening and collaboration among researchers.

Two independent reviewers (BGS and BRJ) conducted the title and abstract screening and included articles for the full-text review. Disagreement between the two reviewers was resolved by inviting the third reviewer (HYH) to make the final decision. The following inclusion criteria were used:

1. Population: children or adolescents aged 2–19 years.
2. Index test: studies that assessed the diagnostic performance of MUAC as an index test to identify children and adolescents with overweight/obesity.
3. Comparator: compared to reference standards such as BMI z-score, weight to height, waist circumference, skinfold thickness, dual-energy X-ray absorptiometry, air-displacement plethysmography, bioelectrical impedance and hydro densitometry.
4. Outcome: overweight and obesity.
5. Study design: observational studies including cross-sectional, cohort and case-control were included.
6. Language: studies published in any language were included.
7. Year of publication: no restriction was made based on the year of publication.

Studies that fulfill any of the following criteria were excluded:

1. Articles available only in abstract form, letters, reviews, commentaries, editorials, case series.
2. Duplicate publication of the same study

Data extraction

Two independent reviewers (BGS and BRJ) have extracted the following information from included studies, using pilot-tested data collection form: first author's name, year of publication, country or region, funding source, study design, total sample size, number of males and females, response rate, age of study participants, MUAC cut-off values, reference standard, diagnostic criteria of

overweight and obesity (reference standard), sensitivity, specificity, AUC, positive and negative likelihood ratio, prevalence of overweight and obesity, positive and negative predictive values. The extracted data by independent reviewers were compared and any discrepancy was resolved by consensus. When relevant information was missing from the article, we contacted the primary authors twice via e-mail.

Risk of bias and certainty of evidence assessment

Two independent reviewers (BGS and BRJ) assessed the risk of bias and applicability of the included studies using Quality Assessment of Diagnostic Accuracy Studies (QUADAS-2)⁽³⁰⁾. We have also assessed the certainty of evidence for relevant outcomes using the Grading of Recommendations Assessment, Development and Evaluation (GRADE) approach for diagnostic tests⁽³¹⁾. Discrepancies among reviewers on individual items were resolved by discussion and consensus.

Statistical analysis

The extracted data were exported to STATA/SE Version 16 for further processing and analysis. We have summarised the diagnostic test accuracy by creating a 2 × 2 table for each study. We have performed a graphical descriptive analysis of the included studies. We have reported coupled forest plots (sensitivity and specificity separately, along with the 95 % CI), and we provided a graphical representation of studies in the summary receiver operating characteristic (SROC) curve (sensitivity against 1 – specificity).

Among the included studies, those that report the number of true positive, true negative, false positive, false negative or values that are required to calculate them are included in the meta-analysis.

We have used the hierarchical summary receiver-operating characteristic curve model to produce SROC curves⁽³²⁾. The AUC – SROC curve values were used to describe test accuracy. We evaluated the discriminatory power by the AUC-SROC using values proposed by Swets⁽³³⁾, with ≤ 0.5 considered to have no discriminatory power, > 0.5 and ≤ 0.7 to have low discriminatory power, > 0.7 and ≤ 0.9 to have good discriminatory power and 1 to be a perfect test. In addition, we have estimated the pooled sensitivity, specificity, positive likelihood ratio and negative likelihood ratio to complement the findings of SROC. Subgroup analysis was performed for boys, girls, children (2–9 years), adolescents (10–19 years), with overweight and obesity.

We assessed the heterogeneity of diagnostic test parameters by visual inspection of the paired forest plots and SROC plots. One of the major sources of heterogeneity in diagnostic accuracy systematic review and meta-analysis is the use of different thresholds. We have assessed the presence of a threshold effect using spearman's correlation coefficient between the logit of sensitivity and the logit of 1 – specificity⁽³⁴⁾. We have explored potential sources

of bias including sex, cut-off point, age group and weight status (overweight, obesity) variables. To further investigate heterogeneity, we performed subgroup analysis for boys, girls, children (2–9 years), adolescents (10–19 years), children and adolescents with overweight, children and adolescents with obesity. To assess possible publication bias, we used Deeks' funnel plot, with Deeks' asymmetry test, where $P < 0.05$ was considered as significant asymmetry⁽³⁵⁾.

Results

Selection of studies

The search strategy results in a total of 3039 references. Of these, 2041 were duplicates, resulting in 998 articles. After screening titles and abstracts, 963 studies were excluded. Thirty-five papers were retained for review after full-text evaluation. Of these, fourteen articles were excluded with reasons; did not report measures of diagnostic performance; different target condition; was conducted on age group outside of the scope of this review. Therefore, twenty-one studies met the inclusion criteria and were included in the systematic review, however, ten studies with sufficient information are included in the meta-analysis. These eleven articles reported neither the required parameters for meta-analyses (true positive, true negative, false negative or false positive) nor other parameters to calculate them (prevalence, positive predictive value or negative predictive value). Figure 1 shows the detailed description of the article screening process.

Characteristics of studies

Table 1 shows characteristics of studies included in the systematic review. All of the studies were cross-sectional and were published from 2013 to 2021. They were conducted in twenty-one countries including Brazil⁽³⁶⁾, China⁽³⁷⁾, Netherlands⁽³⁸⁾, Ethiopia⁽²⁰⁾, India^(39–41), Indonesia⁽⁴²⁾, Nigeria^(43,44), Pakistan^(25,45), Sri Lanka⁽²⁴⁾, South Africa^(46,47), Thailand⁽⁴⁸⁾, Turkey^(49,50), Trinidad and Tobago⁽⁵¹⁾ and twelve countries⁽¹⁷⁾ (Australia, Brazil, Canada, China, Colombia, Finland, India, Kenya, Portugal, South Africa, the UK and the USA) ($n = 1$). The number of participants varied substantially between studies (range from 211 to 31 471), with a pooled population of 54 381 children and adolescents.

Studies used different reference methods: three studies used bioelectrical impedance^(18,44); two used waist circumferences^(49,52); the rest used BMI. Most studies used an 85th percentile (Z score $> 1 + SD$) cut-off of BMI growth curves for overweight and a 95th percentile cut-off (Z score $> 2 + SD$) of BMI curve for obesity. Both studies that use bioelectrical impedance classify participants with 85th percentile as overweight^(18,44). The cut-off points of MUAC values for defining overweight and obesity varied




Fig. 1 Study selection flow chart

substantially between studies (see online supplementary material, Supplemental Table S4).

Meta-analysis

A meta-analysis was conducted to assess the performance of MUAC for identifying overweight and obesity in children and adolescents. The meta-analysis showed that a pooled AUC of 0.92 (95% CI 0.89, 0.94), sensitivity of 85.2 (95% CI 77.3, 90.6) and a pooled specificity of 85.6 (95% CI 81.0, 98.2) for boys and girls, respectively. In boys, MUAC showed a pooled AUC of 0.92 (0.89, 0.94), sensitivity of 84.4 (95% CI 84.6, 90.8) and a pooled specificity of 86.0 (95% CI 79.2, 90.8) (Fig. 2 and Table 2). As for girls, MUAC showed a pooled AUC of 0.93 (95% CI 0.90, 0.95), sensitivity of 86.4 (95% CI 79.8, 91.0) and pooled specificity of 86.6 (95% CI 82.2, 90.1) (Fig. 3 and Table 2).

We have further explored the diagnostic performance of the MUAC by weight status. The highest discriminatory

ability was observed among children and adolescents with obesity, resulting in a pooled AUC of 0.95 (95% CI 0.92, 0.96), sensitivity of 89.4 (95% CI 80.2, 94.6) and a pooled specificity of 88.9 (95% CI 83.4, 92.8).

Moreover, we have analysed the diagnostic performance of MUAC by the age of participants 2–9 years and 10–19 years. The MUAC showed higher performance in adolescents between the age of 10–19 compared to those aged 2–9 years with an AUC of 0.96 (0.93, 0.97) *v.* 0.88 (95% CI 0.85, 0.90).

Risk of bias and publication bias

The full results of the risk of bias and applicability were assessed using QUADAS-2 is shown in supplementary material (see online supplementary material, Supplemental Table S5). The study design and procedure were homogeneous and almost all met all QUADAS-2 domains. One study was classified as ‘high risk’ in the



Table 1 Characteristics of included studies

Study author	Location	Study design	Sample size	Boys/girls	Sex	Age	Reference standard	Outcome threshold	Cut-off	Sensitivity	Specificity
Lu <i>et al.</i> ⁽²⁶⁾ (2013)	China	Cross-sectional	2847	1475/1372	Boys and girls	7–12	BMI	≥ 85th percentiles	18.9–23.4	83–95 %	82–96 %
Rerksuppaphol & Rerksuppaphol ⁽⁵⁷⁾ (2017)	Thailand	Cross-sectional	3618		Boys and girls	7–12	BMI	BMI Z score > 2 + SD	19.8–25.5	82–96 %	89.2–97 %
Asif <i>et al.</i> ⁽²⁵⁾ (2018)	Pakistan	Cross-sectional	7921	4021/3900	Boys and girls	8		BMI Z score > 1 + SD	18.0–23.2	93 %	94 %
Craig <i>et al.</i> ⁽¹⁸⁾ (2014)	South Africa	Cross-sectional	978		Boys and girls	5–14	BMI	BMI Z score > +1SD	16.38–22.73	60–90 %	59–90 %
Chaput <i>et al.</i> ⁽¹⁷⁾ (2016)	12 countries	Cross-sectional	7337	3408/3929	Boys and girls	5–14	BIA	BMI Z score > +1SD		64–97 %	76–93 %
Jaiswal <i>et al.</i> ⁽³⁹⁾ (2017)	India	Cross-sectional	875	436/439	Boys and girls	9–11	BMI	> 85th percentile		26–100 %	73–95 %
Talma <i>et al.</i> ⁽¹⁹⁾ (2018)	Dutch	Cross-sectional	6167		Boys and girls	5–14	BMI	BMI Z score > +2SD	24.6–25.2	94–95 %	90–92 %
Mazicioğlu <i>et al.</i> ⁽⁵⁸⁾ (2010)	Turkey	Cross-sectional	5358	2621/2737	Boys and girls	2–18	BMI	≥ 85th percentiles	18.8–23.3	91–100 %	95.1–100 %
Ayu <i>et al.</i> ⁽⁴²⁾ (2017)	Indonesia	Cross-sectional	2258		Boys and girls	6–17	BMI	≥ 85th percentiles	>1.3 SDS	52–95 %	71–94 %
Shinsugi <i>et al.</i> ⁽²⁴⁾ (2020)	Serilanka	Cross-sectional	528		Boys and girls	6–17	WC	≥ 90th percentile	18.1–24.9	50–95 %	76–98 %
Dumith <i>et al.</i> ⁽³⁶⁾ (2018)	Brazil	Cross-sectional	1075	512/563	Boys and girls	6–7	BMI	≥ 85th percentile	17.9–25.7	53–100 %	63–92 %
Otitoola <i>et al.</i> ⁽⁵⁹⁾ (2020)	South Africa	Cross-sectional	211	102/109	Boys and girls	6–7		≥ 85th percentile	18.5	88–89 %	78 %
Orimadegun ⁽⁴³⁾ (2019)	Nigeria	Cross-sectional	920	403/517	Boys and girls	6–7		≥ 95th percentile	19.5	85–93 %	86–87 %
Asif <i>et al.</i> ⁽⁴⁵⁾ (2018)	Pakistan	Cross-sectional	4962		Boys and girls	6–7		≥ 85–< 95 percentile	18.5	73–83 %	78 %
de Almeida <i>et al.</i> ⁽⁶⁰⁾ (2003)	Brazil	Cross-sectional	1090		Boys and girls	5–10		BMI Z score > +1 SD	19.05–21.8	95–100 %	85–96 %
Oriaifo <i>et al.</i> ⁽⁴⁴⁾ (2019)	Nigeria	Cross-sectional	1067	538/529	Boys and girls	13–19	BMI	BMI Z score > +1SD		58–84 %	57–89 %
Sisay <i>et al.</i> ⁽²⁰⁾ (2020)	Ethiopia	Cross-sectional	851	456/395	Boys and girls	6–19		BMI Z score > +1SD			
Khiamniungan & Mondal ⁽⁴⁰⁾ (2019)	India	Cross-sectional	960	400	Boys and girls	5–18		BMI Z score > +2SD			
Ozturk <i>et al.</i> ⁽⁵²⁾ (2015)	Turkey	Cross-sectional	5358	968	Boys	12–18	BMI	≥ 85th percentile	19.43–24.76	58–94 %	20–88 %
Ramcharitar-Bourne <i>et al.</i> ⁽⁵¹⁾ (2021)	Trinidad and Tobago	Cross-sectional	595	301/295	Boys and girls	12–18	BMI	0.7 (z score)	0.6–0.7 (z score)	77–79 %	78 %
Nitika ⁽⁴¹⁾ (2021)	India	Cross-sectional	31 471	16 158/15 313	10–19	BMI	BMI	≥ 85th percentile	18.75–30.0	60–100 %	53–100 %
								BMI Z score > +1 SD	27.75–27.95	90–94 %	89–91 %
								WC ≥ 90th percentile	16.5–21.6	50–95 %	65–80 %
								≥ 85th percentile	MUAC z-score	43–92 %	95–98 %
								BMI Z score > +1SD	21.2–29.8	74.5–90 %	74.5–90 %

BIA, bioelectrical impedance analysis; WC, waist circumference.




Fig. 2 Summary receiver operating characteristic (SROC) curve for boys

question of applicability domain (Index test) because the reference population they use to identify children with overweight and obesity was inappropriate. All included studies had a ‘high risk’ of bias domain (reference standard) because they used BMI as their reference standard, which is not a golden standard to measure excess adiposity. Even though BMI is highly correlated with excess adiposity, it misclassifies a significant number of children and adolescents⁽⁹⁾. Except for one⁽²⁰⁾, none of the included studies reported the time interval between performing the index test and reference standard. However, it is unlikely that any time delay between conducting the index test and the reference standard would introduce bias. The description of the index tests and reference standards was adequately reported. Evidence of publication bias was not observed by Deek’s test ($P = 0.71$) (see online supplementary material, Supplemental Fig. S1).

Investigation of heterogeneity

Visual inspection of the paired forest plot and SROC curve revealed substantial heterogeneity. (Fig. 4) A Spearman’s rank correlation test showed the presence of a threshold effect $r = 0.63$. We have examined the influence of covariate sex, age group and weight status. We have observed that MUAC cut-off (P -value = 0.00), age group (P -value = 0.00) and weight status (P -value = 0.05) significantly contribute to the heterogeneity of the study. However, sex of participants did not contribute to the

Table 2 Diagnostic performance of mid-upper arm circumference

Sex	Pooled AUC	95 % CI	Pooled sensitivity	95 % CI	Pooled specificity	95 % CI	Positive likelihood ratio	95 % CI	Negative likelihood ratio	95 % CI
Boys	0.92	0.89, 0.94	84.4	84.6, 90.8	86.0	79.0, 90.8	6.0	3.7, 9.7	0.12	0.10, 0.31
Girls	0.93	0.90, 0.95	86.4	79.8, 91.0	86.6	82.2, 90.1	6.5	4.6, 9.1	0.16	0.11, 0.24
Boys and girls	0.92	0.89, 0.94	85.2	77.3, 90.6	85.6	81.0, 89.2	5.9	4.0, 8.9	0.17	0.11, 0.28
2–9	0.88	0.85, 0.90	83.6	67.9, 92.1	78.8	66.1, 87.7	3.9	2.2, 7.2	0.21	0.01, 0.47
10–19	0.96	0.93, 0.97	91.5	87.1, 94.5	88.9	85.5, 91.6	8.3	6.4, 10.7	0.09	0.06, 0.14
Overweight	0.91	0.88, 0.93	84.1	75.3, 90.1	84.3	79.4, 88.3	5.4	3.8, 7.5	0.18	0.11, 0.31
Obesity	0.95	0.92, 0.96	89.4	80.2, 94.6	88.9	83.4, 92.8	8.1	5.0, 16.9	0.12	0.06, 0.24




Fig. 3 Summary receiver operating characteristic (SROC) curve for girls

heterogeneity (P -value = 0.10). To further explore sources of heterogeneity, we performed subgroup analyses by sex, age group (children, adolescents) and weight status (overweight, obese). The results suggest that weight status might contribute to the heterogeneity among studies. Since pooling sensitivity and specificity are more reliable in the absence of a threshold effect, the findings should be primarily judged based on the SROC curve.

Certainty of evidence assessment

We rated the certainty of evidence of the pooled studies and considered it as moderate for all pooled measures of diagnostic accuracy. The reasons for downgrading the certainty of evidence included the marked heterogeneity observed⁽³¹⁾. The result of certainty of evidence assessment is available on supplementary material (see online supplementary material, Supplemental Table S6).

Discussion

This meta-analysis showed that when assessing the diagnostic performance of MUAC compared with BMI defined overweight and obesity. MUAC has high sensitivity and specificity, correctly identifying about 86.4% of children and adolescents with BMI defined overweight and obese. In addition, MUAC has a greater discriminatory ability in identifying children and adolescents with BMI-defined obesity than those with BMI-defined overweight.




Fig. 4 Paired forest plot of sensitivity and specificity with 95% CI



However, no sufficient evidence on the performance of MUAC compared with gold standard measures of adiposity.

The MUAC was initially developed to screen and diagnose under-five children with moderate and severe acute malnutrition⁽²¹⁾. Moreover, MUAC has also been used to identify adolescents and women of reproductive age with thinness and severe thinness^(22,23). In recent years, MUAC has been explored as an alternative screening tool for children and adolescents with overweight and obesity^(17–20,24–26). MUAC has attractive characteristics that make it desirable as a simple screening tool.

To the best of our knowledge, this is the first systematic review and meta-analysis to examine the discriminatory performance of MUAC to identify children and adolescents with overweight and obesity considering age group, sex and weight status. The previous systematic review had included MUAC as one of the uncommonly used measures of overweight and obesity. However, this systematic review has not assessed the discriminatory ability of MUAC in identifying children and adolescents with overweight and obesity⁽⁵³⁾.

This meta-analysis showed that MUAC has a high discriminatory ability to identify both boys and girls with BMI defined overweight and obesity with a comparable value of sensitivity and specificity. However, concerning weight status, MUAC has superior discriminatory ability among children and adolescents with obesity than those who are BMI defined overweight. Similar findings have been reported by a meta-analysis conducted on the performance of neck circumference⁽⁵⁴⁾. This might be due to the fact that sensitivity and specificity of screening tools depend on the spectrum of a condition, in this case, the degree of adiposity; those who are obese are more likely to be easily identified by screening tools than those who are overweight.

An ideal anthropometric measurement to identify adolescents with overweight and obesity should be easy to use, accurate and reliable⁽⁵⁵⁾. MUAC fulfil almost all characteristics; MUAC is simple to use since its measurement requires only a non-stretchable MUAC tape; MUAC measurement is easy to interpret since it does not require the use of an additional reference chart. Moreover, MUAC can easily be used by an illiterate person if it is colour-coded. Traffic light colours of red (obese), amber (overweight) and green (normal weight) may also be considered by non-numerate field workers in developing countries to facilitate screening⁽¹⁷⁾. The other important characteristic of ideal measurement is its accuracy in identifying overweight and obesity among children and adolescents. As we have observed in this meta-analysis, MUAC has high accuracy in identifying overweight and obesity. However, in this meta-analysis, all included articles compare the discriminatory ability of MUAC against BMI defined overweight and obesity. Even though BMI is highly correlated with percent body fat, it does not differentiate between lean mass and fat mass⁽⁵⁶⁾. Only two studies have

compared the discriminatory ability of MUAC with total body fat which was measured with bioelectrical impedance. A high level of discriminatory ability of MUAC has been reported by both studies^(18,44). None of the studies have assessed the discriminatory performance of MUAC with the four-compartment model, the known golden standard to measure total body fat. There is a need for future researches to explore the performance of MUAC to identify overweight and obesity, ideally as defined by the 'golden standard' measure of total body fat.

The major limitation of this meta-analysis was the presence of marked heterogeneity. First, a diagnostic threshold bias was identified as a cause of heterogeneity in the pooled results. In this meta-analysis, there was no consistent cut-off value. To overcome this limitation, we have used a hierarchical SROC curve that accounts for the threshold effect. Furthermore, we have conducted a subgroup analysis to reduce heterogeneity, we calculated measures of diagnostic performance according to age group, sex or weight status. Second, we have attempted to reach the corresponding authors of articles with insufficient data (prevalence, sample size by category) to be included in the meta-analysis, but without success. The main strength of this study was the rigorous statistical methods used to pool data across diagnostic accuracy studies. The other strength of this study is the comprehensive search strategy in several electronic databases.

Conclusions

In conclusion, MUAC has high sensitivity and specificity compared with BMI defined overweight and obesity. However, there is no sufficient evidence on the performance of MUAC compared to gold standard measures of adiposity. There is a need for future studies to evaluate the diagnostic performance compared with the 'golden standard' measure of total body fat and evaluate the predictive value of the MUAC measurement for developing weight-related complications.

Acknowledgements

Acknowledgements: We would like to express our deepest gratitude to authors that respond to our request for additional information and explanations. *Financial support:* This research received no specific grant from any funding agency, commercial or not-for-profit sectors. *Conflict of interest:* There are no conflicts of interest. *Authorship:* B.G.S. was the lead author of the manuscript including study conception, design, conduct, statistical analysis and writing of the first draft. S.H.G., H.Y.H., B.R.J. and E.A. were involved in formulating a research question and critical revision of the manuscript. B.G.S. and E.A. performed



systematic searches of bibliographic databases. H.Y.H. and B.R.J. were also involved in the screening of potentially relevant articles for review, risk of bias assessment, data extraction and data analysis. H.Y.H., B.R.J. and E.A. have all critically reviewed and edited the manuscript. All authors read and approved the final manuscript. None of the authors reported a conflict of interest related to the study. *Ethics of human subject participation:* Not applicable since this study is systematic review and meta-analysis.

Supplementary material

For supplementary material/s referred to in this article, please visit <https://doi.org/10.1017/S1368980022000143>

Reference

- World Health Organization (2021) Obesity and Overweight. <https://www.who.int/news-room/fact-sheets/detail/obesity-and-overweight> (accessed February 2021).
- Abarca-Gómez L, Abdeen ZA, Hamid ZA *et al.* (2017) Worldwide trends in body-mass index, underweight, overweight, and obesity from 1975 to 2016: a pooled analysis of 2416 population-based measurement studies in 128.9 million children, adolescents, and adults. *Lancet* **390**, 2627–2642.
- Avgerinos KI, Spyrou N, Mantzoros CS *et al.* (2019) Obesity and cancer risk: emerging biological mechanisms and perspectives. *Metabolism* **92**, 121–135.
- Shields M, Tremblay MS, Connor Gorber S *et al.* (2012) Abdominal obesity and cardiovascular disease risk factors within body mass index categories. *Health Rep* **23**, 7–15.
- Gepstein V & Weiss R (2019) Obesity as the main risk factor for metabolic syndrome in children. *Front Endocrinol* **10**, 568–568.
- Han TS & Lean ME (2016) A clinical perspective of obesity, metabolic syndrome and cardiovascular disease. *JRSM Cardiovasc Dis* **5**, 2048004016633371.
- Simmonds M, Llewellyn A, Owen CG *et al.* (2016) Predicting adult obesity from childhood obesity: a systematic review and meta-analysis. *Obes Rev* **17**, 95–107.
- Llewellyn A, Simmonds M, Owen CG *et al.* (2016) Childhood obesity as a predictor of morbidity in adulthood: a systematic review and meta-analysis. *Obes Rev* **17**, 56–67.
- Javed A, Jumean M, Murad MH *et al.* (2015) Diagnostic performance of body mass index to identify obesity as defined by body adiposity in children and adolescents: a systematic review and meta-analysis. *Pediatr Obes* **10**, 234–244.
- WHO (2011) World Health Organisation Factsheet No. 311. <http://www.who.int/mediacentre/factsheets/fs311/en/> (accessed June 2021).
- Wells JC & Fewtrell MS (2006) Measuring body composition. *Arch Dis Child* **91**, 612–617.
- Cornier MA, Després JP, Davis N *et al.* (2011) Assessing adiposity: a scientific statement from the American Heart Association. *Circulation* **124**, 1996–2019.
- Simmonds M, Llewellyn A, Owen CG *et al.* (2016) Simple tests for the diagnosis of childhood obesity: a systematic review and meta-analysis. *Obes Rev* **17**, 1301–1315.
- WHO (1995) Physical status: the use and interpretation of anthropometry. Report of a WHO Expert Committee. *World Health Organ Tech Rep Ser* **854**, 1–452.
- Pischon T, Boeing H, Hoffmann K *et al.* (2008) General and abdominal adiposity and risk of death in Europe. *N Engl J Med* **359**, 2105–2120.
- Prentice AM (2006) The emerging epidemic of obesity in developing countries. *Int J Epidemiol* **35**, 93–99.
- Chaput JP, Katzmarzyk PT, Barnes JD *et al.* (2017) Mid-upper arm circumference as a screening tool for identifying children with obesity: a 12-country study. *Pediatr Obes* **12**, 439–445.
- Craig E, Bland R, Ndirangu J *et al.* (2014) Use of mid-upper arm circumference for determining overweight and overfatness in children and adolescents. *Arch Dis Child* **99**, 763–766.
- Talma H, van Dommelen P, Schweizer JJ *et al.* (2019) Is mid-upper arm circumference in Dutch children useful in identifying obesity? *Arch Dis Child* **104**, 159–165.
- Sisay BG, Haile D, Hassen HY *et al.* (2019) Performance of mid-upper arm circumference as a screening tool for identifying adolescents with overweight and obesity. *PLoS One* **15**, e0235063.
- Myatt M, Khara T & Collins S (2006) A review of methods to detect cases of severely malnourished children in the community for their admission into community-based therapeutic care programs. *Food Nutr Bull* **27**, S7–S23.
- Sethi V, Gupta N, Pedgaonkar S *et al.* (2019) Mid-upper arm circumference cut-offs for screening thinness and severe thinness in Indian adolescent girls aged 10–19 years in field settings. *Public Health Nutr* **22**, 2189–2199.
- Sisay BG, Haile D, Hassen HY *et al.* (2021) Mid-upper arm circumference as a screening tool for identifying adolescents with thinness. *Public Health Nutr* **24**, 457–466.
- Shinsugi C, Gunasekara D & Takimoto H (2020) Use of Mid-Upper Arm Circumference (MUAC) to predict malnutrition among Sri Lankan schoolchildren. *Nutrients* **12**, 168.
- Asif M, Aslam M & Altaf S (2018) Mid-upper-arm circumference as a screening measure for identifying children with elevated body mass index: a study for Pakistan. *Korean J Pediatr* **61**, 6–11.
- Lu Q, Wang R, Lou DH *et al.* (2014) Mid-upper-arm circumference and arm-to-height ratio in evaluation of overweight and obesity in Han children. *Pediatr Neonatol* **55**, 14–19.
- McInnes MDF, Moher D, Thoms BD *et al.* (2018) Preferred reporting items for a systematic review and meta-analysis of diagnostic test accuracy studies: the PRISMA-DTA statement. *JAMA* **319**, 388–396.
- Sisay BG, Hassen HY & Gebreyesus SH (2021) Diagnostic performance of mid-upper arm circumference to identify overweight and obesity in children and adolescents: a protocol for a systematic review and meta-analysis. *BMJ Open* **11**, e044624.
- Ouzzani M, Hammady H, Fedorowicz Z *et al.* (2016) Rayyan – a web and mobile app for systematic reviews. *Syst Rev* **5**, 210.
- Whiting PF, Rutjes AW, Westwood ME *et al.* (2011) QUADAS-2: a revised tool for the quality assessment of diagnostic accuracy studies. *Ann Intern Med* **155**, 529–536.
- Schünemann HJ, Oxman AD, Brozek J *et al.* (2008) Grading quality of evidence and strength of recommendations for diagnostic tests and strategies. *BMJ* **336**, 1106–1110.
- Rutter CM & Gatsonis CA (2001) A hierarchical regression approach to meta-analysis of diagnostic test accuracy evaluations. *Stat Med* **20**, 2865–2884.
- Swets JA (1988) Measuring the accuracy of diagnostic systems. *Science* **240**, 1285–1293.
- Moses LE, Shapiro D & Littenberg B (1993) Combining independent studies of a diagnostic test into a summary ROC curve: data-analytic approaches and some additional considerations. *Stat Med* **12**, 1293–1316.

35. Deeks JJ, Macaskill P & Irwig L (2005) The performance of tests of publication bias and other sample size effects in systematic reviews of diagnostic test accuracy was assessed. *J Clin Epidemiol* **58**, 882–893.
36. Dumith SC, Muraro MFR, Monteiro AR *et al.* (2018) Diagnostic properties and cutoff points for overweight prediction through anthropometric indicators in adolescents from Caracol, Piauí, Brazil, 2011. *Epidemiol Serv Saude* **27**, e201715013.
37. Lu Q, Wang R, Lou DH *et al.* (2014) Mid-upper-arm circumference and arm-to-height ratio in evaluation of overweight and obesity in han children. *Pediatr Neonatol* **55**, 14–19.
38. Talma H, Van Dommelen P, Schweizer JJ *et al.* (2019) Is mid-upper arm circumference in Dutch children useful in identifying obesity? *Arch Dis Childhood* **104**, 159–165.
39. Jaiswal M, Bansal R & Agarwal A (2017) Role of mid-upper arm circumference for determining overweight and obesity in children and adolescents. *J Clin Diagn Res* **11**, SC05–SC08.
40. Khiamniungan KS & Mondal PR (2019) Mid-upper arm circumference: an alternative to bmi for screening overweight and obesity among the Khiamniungan tribal children and adolescents of Nagaland, Northeast India. *Online J Health Allied Sci* **18**, 1–4.
41. Nitika N (2021) Discriminatory ability of mid-upper arm circumference in identifying overweight and obese adolescents: findings from the comprehensive national nutrition survey, India. *na J Public Health* **65**, 269–274.
42. Ayu DR, Aditiawati A, Anzar J *et al.* (2017) Upper arm circumference measurement for detecting overweight and obesity in children aged 6–7 years. *Paediatr Indones* **57**, 23–29.
43. Orimadegun A (2019) Accuracy of mid upper arm circumference in detection of obesity among school children in Yenagoa City, South-south region of Nigeria. *Niger J Paediatr* **46**, 48–54.
44. Oriafio S, Abiodun P, Atimati A *et al.* (2019) Determination of overnutrition using mid-upper arm circumference in comparison with bioelectrical impedance analysis in children and adolescents in Benin, Nigeria. *J Health Res* **34**, 68–78.
45. Craig E, Bland R, Ndirangu J *et al.* (2014) Use of mid-upper arm circumference for determining overweight and overfatness in children and adolescents. *Arch Dis Child* **99**, 763–766.
46. Asif M, Aslam M & Altaf S (2018) Use of mid-upper arm circumference in evaluation of overweight and obesity in the Pakistani children and adolescent, aged 12–18 years. *Pak Paediatr J* **42**, 43–48.
47. Otitoola O, Oldewage-Theron W & Egal A (2021) Prevalence of overweight and obesity among selected schoolchildren and adolescents in Cofimvaba, South Africa. *S Afr J Clin Nutr* **34**, 97–102.
48. Rerksuppaphol S & Rerksuppaphol L (2017) Mid-upper-arm circumference and arm-to-height ratio to identify obesity in school-age children. *Clin Med Res* **15**, 53–58.
49. Mazicioğlu MM, Hatipoğlu N, Öztürk A *et al.* (2010) Waist circumference and mid-upper arm circumference in evaluation of obesity in children aged between 6 and 17 years. *JCRPE J Clin Res Pediatr Endocrinol* **2**, 144–150.
50. Ozturk A, Cicek B, Mazicioğlu MM *et al.* (2015) Determining abdominal obesity cut-offs and relevant risk factors for anthropometric indices in Turkish children and adolescents. *J Pediatr Endocrinol Metab* **28**, 525–532.
51. Ramcharitar-Bourne A, Nichols S & Badrie N (2021) Predictive utility of anthropometric based cut-offs in assessing excess adiposity among preschool children in a multiethnic population. *West Indian Med J* **69**, 114.
52. Ozturk A, Cicek B, Mazicioğlu MM *et al.* (2015) Determining abdominal obesity cut-offs and relevant risk factors for anthropometric indices in Turkish children and adolescents. *J Pediatr Endocrinol Metab* **28**, 525–532.
53. Ranasinghe P, Jayawardena R, Gamage N *et al.* (2021) The range of non-traditional anthropometric parameters to define obesity and obesity-related disease in children: a systematic review. *Eur J Clin Nutr* **75**, 373–384.
54. Kroll C, Mastroeni S, Czarnobay SA *et al.* (2017) The accuracy of neck circumference for assessing overweight and obesity: a systematic review and meta-analysis. *Ann Hum Biol* **44**, 667–677.
55. Roper HP (2003) Child and adolescent obesity: causes and consequences, prevention and management. *J R Soc Med* **96**, 312–313.
56. Freedman DS, Horlick M & Berenson GS (2013) A comparison of the Slaughter skinfold-thickness equations and BMI in predicting body fatness and cardiovascular disease risk factor levels in children. *Am J Clin Nutr* **98**, 1417–1424.
57. Rerksuppaphol S & Rerksuppaphol L (2017) Mid-upper-arm circumference and arm-to-height ratio to identify obesity in school-age children. *Clin Med Res* **15**, 53–58.
58. Mazicioğlu MM, Hatipoğlu N, Öztürk A *et al.* (2010) Waist circumference and mid-upper arm circumference in evaluation of obesity in children aged between 6 and 17 years. *J Clin Res Pediatr Endocrinol* **2**, 144–150.
59. Otitoola O, Oldewage-Theron W & Egal A (2020) Prevalence of overweight and obesity among selected schoolchildren and adolescents in Cofimvaba, South Africa. *South Afr J Clin Nutr* **34**, 1–6.
60. de Almeida CA, Del Ciampo LA, Ricco RG *et al.* (2003) Assessment of mid-upper arm circumference as a method for obesity screening in preschool children. *J Pediatr* **79**, 455–460.