



Influence of 5-fluorouracil-loaded microsphere formulation on efficient rat glioma radiosensitization

Submitted by Laurent Lemaire on Mon, 12/01/2014 - 16:34

Titre	Influence of 5-fluorouracil-loaded microsphere formulation on efficient rat glioma radiosensitization
Type de publication	Article de revue
Auteur	Roullin, Valérie-Gaëlle [1], Mege, Martine [2], Lemaire, Laurent [3], Cueyssac, Jean-Pierre [4], Venier-Julienne, Marie-Claire [5], Menei, Philippe [6], Gamelin, Erik [7], Benoît, Jean-Pierre [8]
Editeur	American Association of Pharmaceutical Scientists
Type	Article scientifique dans une revue à comité de lecture
Année	2004
Langue	Anglais
Date	2004
Pagination	1558-63
Volume	21
Titre de la revue	Pharmaceutical Research
ISSN	0724-8741
Mots-clés	Animals [9], Brain [10], Brain Neoplasms [11], Drug Evaluation, Preclinical [12], Fluorouracil [13], Glioma [14], Magnetic Resonance Imaging [15], Microspheres [16], Radiation-Sensitizing Agents [17], Rats [18], Rats, Sprague-Dawley [19] PURPOSE: To determine (i) the efficiency of radiosensitizing 5-FU-loaded microspheres and (ii) the impact of microparticle formulation on response to treatment. METHODS: C6 tumor-bearing rats were stereotactically implanted with microspheres and/or allocated to: control groups (untreated) or treatment (only radiotherapy; fast-release 5-FU microspheres + radiotherapy; slow-release 5-FU microspheres + radiotherapy). The next day, fractionated radiotherapy, limited to the hemibrain, was initiated in all treated animals. The irradiation cycle included 36 Gy, given in 9 sessions for 3 consecutive weeks. Tumor development was assessed by T2-weighted MRI. RESULTS: 5-FU microspheres associated with radiotherapy caused a 47% complete remission rate (9/19) as opposed to the 8% rate (1/12) when radiotherapy alone or 0% in control animals. Drug delivery for 3 weeks produced better survival results (57%) compared to one-week sustained release (41%). MR images showed exponentially increasing tumor volumes during the first half of the radiotherapy cycle, followed by a decrease, and the disappearance of the tumor if survival exceeded 120 days. CONCLUSIONS: 5-FU controlled delivery is a promising strategy for radiosensitizing gliomas. Drug delivery system formulation is unambiguously implicated in both the response to treatment and the limitation of toxic side effects.
Résumé en anglais	

URL de la notice	http://okina.univ-angers.fr/publications/ua5699 [20]
Autre titre	Pharm. Res.
Identifiant (ID) PubMed	15497679 [21]

Liens

- [1] [http://okina.univ-angers.fr/publications?f\[author\]=7096](http://okina.univ-angers.fr/publications?f[author]=7096)
- [2] [http://okina.univ-angers.fr/publications?f\[author\]=9582](http://okina.univ-angers.fr/publications?f[author]=9582)
- [3] <http://okina.univ-angers.fr/l.lemaire/publications>
- [4] [http://okina.univ-angers.fr/publications?f\[author\]=9583](http://okina.univ-angers.fr/publications?f[author]=9583)
- [5] <http://okina.univ-angers.fr/ma.venier/publications>
- [6] <http://okina.univ-angers.fr/ph.menei/publications>
- [7] [http://okina.univ-angers.fr/publications?f\[author\]=9584](http://okina.univ-angers.fr/publications?f[author]=9584)
- [8] <http://okina.univ-angers.fr/j.benoit/publications>
- [9] [http://okina.univ-angers.fr/publications?f\[keyword\]=964](http://okina.univ-angers.fr/publications?f[keyword]=964)
- [10] [http://okina.univ-angers.fr/publications?f\[keyword\]=1866](http://okina.univ-angers.fr/publications?f[keyword]=1866)
- [11] [http://okina.univ-angers.fr/publications?f\[keyword\]=8329](http://okina.univ-angers.fr/publications?f[keyword]=8329)
- [12] [http://okina.univ-angers.fr/publications?f\[keyword\]=10301](http://okina.univ-angers.fr/publications?f[keyword]=10301)
- [13] [http://okina.univ-angers.fr/publications?f\[keyword\]=8410](http://okina.univ-angers.fr/publications?f[keyword]=8410)
- [14] [http://okina.univ-angers.fr/publications?f\[keyword\]=8435](http://okina.univ-angers.fr/publications?f[keyword]=8435)
- [15] [http://okina.univ-angers.fr/publications?f\[keyword\]=6040](http://okina.univ-angers.fr/publications?f[keyword]=6040)
- [16] [http://okina.univ-angers.fr/publications?f\[keyword\]=7833](http://okina.univ-angers.fr/publications?f[keyword]=7833)
- [17] [http://okina.univ-angers.fr/publications?f\[keyword\]=10302](http://okina.univ-angers.fr/publications?f[keyword]=10302)
- [18] [http://okina.univ-angers.fr/publications?f\[keyword\]=975](http://okina.univ-angers.fr/publications?f[keyword]=975)
- [19] [http://okina.univ-angers.fr/publications?f\[keyword\]=1300](http://okina.univ-angers.fr/publications?f[keyword]=1300)
- [20] <http://okina.univ-angers.fr/publications/ua5699>
- [21] <http://www.ncbi.nlm.nih.gov/pubmed/15497679?dopt=Abstract>

Publié sur *Okina* (<http://okina.univ-angers.fr>)