



Correlation between structural studies and third order NLO properties of selected new quinolinium semi-organic compounds

Submitted by Emmanuel Lemoine on Tue, 02/04/2014 - 16:12

Titre	Correlation between structural studies and third order NLO properties of selected new quinolinium semi-organic compounds
Type de publication	Article de revue
Auteur	Bouchouit, K. [1], Bendeif, E.-E. [2], El Ouazzani, Hasnaa [3], Dahaoui, S. [4], Lecomte, C. [5], Benali-Cherif, N. [6], Sahraoui, Bouchta [7]
Type	Article scientifique dans une revue à comité de lecture
Année	2010
Langue	Anglais
Date	2010/09/14
Numéro	1
Pagination	1 - 7
Volume	375
Titre de la revue	Chemical Physics
ISSN	0301-0104
Mots-clés	DFWM [8], Nonlinear [9], Semi-organic [10], Single [11], Structural [12], THG [13], Thin [14] New quinolinium semi-organic compounds of formula (C ₉ H ₈ N) ₂ ⁺ · SO ₄ ²⁻ , H ₂ O (I) (bis-quinolinium sulphate monohydrate) and (C ₉ H ₈ N) ₂ ⁺ · NO ₃ ⁻ (II) (quinolinium nitrate) have been synthesized and characterized by UV-Vis absorption spectroscopy, nonlinear optical (NLO) measurements and by single crystal X-ray diffraction. The third order nonlinear optical properties of (I) and (II) were investigated using two methods: the degenerate four wave mixing technique (DFWM) performed in solution at $\lambda = 532$ nm and the third-harmonic generation (THG) measurements carried out on thin films at $\lambda = 1064$ nm. The NLO measurements showed that compound (I) presents better nonlinear optical properties compared to compound (II). To understand further the optical behaviour of (I) and (II), the crystal structures of both compounds were determined from accurate single crystal X-ray diffraction measurements performed at 100 K. The crystallographic studies revealed the key role of the intermolecular interactions and the molecular arrangements in the enhancement of the NLO properties.
Résumé en anglais	New quinolinium semi-organic compounds of formula (C ₉ H ₈ N) ₂ ⁺ · SO ₄ ²⁻ , H ₂ O (I) (bis-quinolinium sulphate monohydrate) and (C ₉ H ₈ N) ₂ ⁺ · NO ₃ ⁻ (II) (quinolinium nitrate) have been synthesized and characterized by UV-Vis absorption spectroscopy, nonlinear optical (NLO) measurements and by single crystal X-ray diffraction. The third order nonlinear optical properties of (I) and (II) were investigated using two methods: the degenerate four wave mixing technique (DFWM) performed in solution at $\lambda = 532$ nm and the third-harmonic generation (THG) measurements carried out on thin films at $\lambda = 1064$ nm. The NLO measurements showed that compound (I) presents better nonlinear optical properties compared to compound (II). To understand further the optical behaviour of (I) and (II), the crystal structures of both compounds were determined from accurate single crystal X-ray diffraction measurements performed at 100 K. The crystallographic studies revealed the key role of the intermolecular interactions and the molecular arrangements in the enhancement of the NLO properties.
URL de la notice	http://okina.univ-angers.fr/publications/ua1954 [15]
DOI	10.1016/j.chemphys.2010.07.007 [16]
Lien vers le document	http://dx.doi.org/10.1016/j.chemphys.2010.07.007 [16]

Liens

- [1] [http://okina.univ-angers.fr/publications?f\[author\]=21962](http://okina.univ-angers.fr/publications?f[author]=21962)
- [2] [http://okina.univ-angers.fr/publications?f\[author\]=2623](http://okina.univ-angers.fr/publications?f[author]=2623)
- [3] [http://okina.univ-angers.fr/publications?f\[author\]=3112](http://okina.univ-angers.fr/publications?f[author]=3112)
- [4] [http://okina.univ-angers.fr/publications?f\[author\]=2625](http://okina.univ-angers.fr/publications?f[author]=2625)
- [5] [http://okina.univ-angers.fr/publications?f\[author\]=2626](http://okina.univ-angers.fr/publications?f[author]=2626)
- [6] [http://okina.univ-angers.fr/publications?f\[author\]=2627](http://okina.univ-angers.fr/publications?f[author]=2627)
- [7] <http://okina.univ-angers.fr/bouchta.sahraoui/publications>
- [8] [http://okina.univ-angers.fr/publications?f\[keyword\]=4871](http://okina.univ-angers.fr/publications?f[keyword]=4871)
- [9] [http://okina.univ-angers.fr/publications?f\[keyword\]=1919](http://okina.univ-angers.fr/publications?f[keyword]=1919)
- [10] [http://okina.univ-angers.fr/publications?f\[keyword\]=5771](http://okina.univ-angers.fr/publications?f[keyword]=5771)
- [11] [http://okina.univ-angers.fr/publications?f\[keyword\]=5756](http://okina.univ-angers.fr/publications?f[keyword]=5756)
- [12] [http://okina.univ-angers.fr/publications?f\[keyword\]=4380](http://okina.univ-angers.fr/publications?f[keyword]=4380)
- [13] [http://okina.univ-angers.fr/publications?f\[keyword\]=4820](http://okina.univ-angers.fr/publications?f[keyword]=4820)
- [14] [http://okina.univ-angers.fr/publications?f\[keyword\]=5770](http://okina.univ-angers.fr/publications?f[keyword]=5770)
- [15] <http://okina.univ-angers.fr/publications/ua1954>
- [16] <http://dx.doi.org/10.1016/j.chemphys.2010.07.007>

Publié sur *Okina* (<http://okina.univ-angers.fr>)