

## Evaluation of Sacubitril/Valsartan Prescribing Patterns Within an Inpatient Hospital Setting

Megan Taylor PharmD  
*Lehigh Valley Health Network*, [megan.taylor@lvhn.org](mailto:megan.taylor@lvhn.org)

Katie Nesbitt PharmD, BCPS  
*Lehigh Valley Health Network*, [katie.nesbitt@lvhn.org](mailto:katie.nesbitt@lvhn.org)

Follow this and additional works at: <https://scholarlyworks.lvhn.org/pharmacy>



Part of the [Pharmacy and Pharmaceutical Sciences Commons](#)

---

### Published In/Presented At

Taylor, M. & Nesbitt, K. (2021). *Evaluation of sacubitril/valsartan prescribing patterns within an inpatient hospital setting*. Presented at Lehigh Valley Health Network, Allentown, PA.

This Poster is brought to you for free and open access by LVHN Scholarly Works. It has been accepted for inclusion in LVHN Scholarly Works by an authorized administrator. For more information, please contact [LibraryServices@lvhn.org](mailto:LibraryServices@lvhn.org).



# Evaluation of Sacubitril/Valsartan Prescribing Patterns Within an Inpatient Hospital Setting

Megan Taylor, PharmD, MSCR, Katie Nesbitt, PharmD, BCPS  
Lehigh Valley Health Network, Allentown, Pa.

## Background

- The American College of Cardiology and American Heart Association (ACC/AHA) revised heart failure consensus guidance preferentially recommending angiotensin receptor-neprilysin inhibitor (ARNI) therapy over angiotensin-converting enzyme (ACE) inhibitor or angiotensin receptor blocker (ARB) therapy in heart failure (preserved and reduced ejection fraction)
- Guideline Directed Medication Therapy (GDMT) for heart failure with reduced ejection fraction (HFrEF) include ARNI/ACE/ARB, beta blockers, mineralocorticoid receptor antagonists, and sodium glucose transporter 2 inhibitors
- The purpose of this study was to assess prescribing patterns of sacubitril/valsartan in patients with heart failure admitted to a Lehigh Valley Health Network (LVHN) hospital

## Methods

- Retrospective chart review of all adult patients with heart failure who received ≥1 dose of sacubitril/valsartan at a LVHN hospital from January 1, 2021 to March 31, 2021

### DISCLOSURE STATEMENTS

Authors of this presentation have nothing to disclose concerning possible financial or personal relationships with commercial entities that may have a direct or indirect interest in the subject matter of this presentation.

Patient Characteristics (n=218)	Mean (SD)
<b>Age, years</b>	<b>69.1 (13.3)</b>
<b>Length of Stay, days</b>	<b>5.5 (6.8)</b>
	<b>N (%)</b>
Female	78 (35.8)
<b>Race</b>	
White	182 (83.5)
Black	12 (5.5)
Asian	2 (0.9)
Other	22 (10.1)
<b>Ethnicity</b>	
Hispanic	23 (10.6)
Non-Hispanic	189 (86.7)
Unknown	6 (2.7)
<b>Heart Failure Category</b>	
HFrEF (EF ≤40%)	174 (79.8)
HFmrEF (EF 41-49%)	15 (6.9)
HFpEF (EF ≥50%)	27 (12.4)
Unknown	2 (0.9)
<b>NYHA Classification</b>	
1	4 (1.8)
2	46 (21.1)
3	36 (16.5)
4	13 (6.0)
Unknown	119 (54.6)

Table 1




Figure 1




Figure 2




Figure 3a




Figure 3b

## Results

- 11 of 42 patients transitioned from ACE inhibitor/ARB therapy initiated on not recommended doses, 9 of which initiated on a more conservative dosing regimen
- 22 patients required a washout from an ACE inhibitor, 18 patients (81.8%) received the recommended 36-hour washout period
- Most common adverse event was hypotension, requiring held doses or discontinuation of therapy in 25% of patients

## Conclusions

- 80% patients prescribed sacubitril/valsartan in the in-patient setting received maximally optimized care
- 88% of patients discharged on sacubitril/valsartan, substantially higher than national estimates of 3.8% reported in 2016
- Areas for development:
  - Appropriate initial dosing of sacubitril/valsartan
  - Initiation of other heart failure GDMT as tolerated
- Majority of patients prescribed sacubitril/valsartan in a LVHN hospital were managed according to guideline and manufacturer recommendations