

Seasonal Variation in Incidence of Stroke in a General Population of 1.4 Million Japanese : The Shiga Stroke Registry.

著者	FUJII Takako, ARIMA Hisatomi, TAKASHIMA Naoyuki, KITA Yoshikuni, MIYAMATSU Naomi, TANAKA-MIZUNO Sachiko, SHITARA Satoshi, URUSHITANI Makoto, MIURA Katsuyuki, NOZAKI Kazuhiko
journal or publication title	Cerebrovascular diseases (Basel, Switzerland)
volume	51
number	1
page range	75-81
year	2022-02
URL	http://hdl.handle.net/10422/00013219

doi: 10.1159/000518370(<https://doi.org/10.1159/000518370>)

Seasonal Variation in Incidence of Stroke in a General Population of 1.4 Million Japanese: The Shiga Stroke Registry

Takako Fujii^{a, b} Hisatomi Arima^{b, c} Naoyuki Takashima^{c, d} Yoshikuni Kita^{c, e}
Naomi Miyamatsu^f Sachiko Tanaka-Mizuno^g Satoshi Shitara^a
Makoto Urushitani^h Katsuyuki Miura^{c, i} Kazuhiko Nozaki^a

^aDepartment of Neurosurgery, Shiga University of Medical Science, Otsu, Japan; ^bDepartment of Preventive Medicine and Public Health, Faculty of Medicine, Fukuoka University, Fukuoka, Japan; ^cDepartment of Public Health, Shiga University of Medical Science, Otsu, Japan; ^dDepartment of Public Health, Kindai University Faculty of Medicine, Osaka, Japan; ^eTsuruga Nursing University, Fukui, Japan; ^fDepartment of Clinical Nursing, Shiga University of Medical Science, Otsu, Japan; ^gDepartment of Digital Health and Epidemiology, Kyoto University, Kyoto, Japan; ^hDepartment of Neurology, Shiga University of Medical Science, Otsu, Japan; ⁱNCD Epidemiology Research Center, Shiga University of Medical Science, Otsu, Japan

Keywords

Stroke · Seasonal variation · Incidence · Community based · Epidemiology of stroke

Abstract

Introduction: The purpose of this study was to investigate seasonal variation in stroke incidence using data from a large-scale stroke registry of general population in current Japan. **Methods:** Shiga Stroke Registry (SSR) is an ongoing population-based registry of stroke that occurred in the Shiga Prefecture in central Honshu, Japan. A total 6,688 cases of first-ever stroke, with onset dates ranging from 1 January 2011 to 31 December in 2013 were included in this study. Incidence rates of first-ever stroke in each season were estimated using the person-year approach and adjusted for age and sex using the Poisson regression models. **Results:** From 2011 to 2013, we identified a total of 6,688 stroke cases (3,570 men, 3,118 women), of which 4,480 cases had ischemic stroke (2,518 men, 1,962 women), 1,588 had intracerebral hemorrhage (857 men, 731 women) and 563 had subarachnoid hemorrhage (166 men, 397 women). Age- and

sex-adjusted incidence rates of total stroke were 151 (95% confidence interval [CI] 144–160, $p = <0.001$ vs. summer) in spring, 130 (95% CI 122–137) in summer, 141 (95% CI 133–149, $p = 0.020$ vs. summer) in autumn and 170 (95% CI 161–179, $p = <0.001$ vs. summer) in winter. Seasonal variation was more pronounced in intracerebral hemorrhage than in ischemic stroke. **Conclusion:** In the present large-scale stroke registry of general population, incidence rates of stroke were highest in winter and lowest in summer in current Japan.

© 2021 The Author(s).
Published by S. Karger AG, Basel

Introduction

Lifetime risk of stroke is estimated to be 25% [1], and stroke affects 12 million people worldwide [2]. Stroke is also leading causes of premature death affecting 6.2 million people worldwide annually [3]. In addition, many stroke survivors suffer from physical, cognitive, and communication deficits, placing burden on family and community [4–7]. Effective prevention of stroke requires strategies based on up-to-date knowledge of stroke risk factors [8].

A number of epidemiological studies reported seasonal variation in incidence of stroke [9–16]. However, current evidence is mainly derived from Western populations [9, 10]. There are some epidemiological studies which showed seasonal variation in stroke incidence among Japanese, but they were based on population-based data obtained during 1960s–2000s [11, 12] or based on hospital-based populations [13–16]. Therefore, it is not quite clear whether current evidence is applicable to general population in current Japan. The purpose of this study was to investigate seasonal variation in stroke incidence using data from a large-scale stroke registry of general population (Shiga Stroke Registry (SSR) as a part of the Shiga Stroke and Heart attack Registry Study [SSHR]) in current Japan [17–22].

Methods

Study Design

SSHR is an ongoing population-based registry of stroke and coronary artery disease that occurred in the Shiga Prefecture in central Japan. As a part of SSHR, we have established a population-based registry study of stroke (SSR) designed to build a complete information system regarding acute ischemic and non-traumatic hemorrhagic stroke management in Shiga Prefecture, Japan. The design of SSR has been described elsewhere in detail. In brief, the SSR uses central and local coordination and monitoring, combined with remote data collection and quality control systems, to create an integrated surveillance system involving the registration of cases among a network of all acute-care hospitals with neurology/neurosurgery facilities and smaller hospitals with rehabilitation facilities in Shiga Prefecture. Information on the death certificates of all deceased residents is also collected to help detect all cases of stroke, including those leading to rapid death outside hospital, with the approval of the Ministry of Health, Labour and Welfare [17–22].

Definition of Stroke

A total of 6,688 cases of first-ever stroke, with onset dates ranging from January 1, 2011 to 31 December in 2013 were included in this study. Diagnosis of stroke was defined as a sudden onset of focal neurological deficits persisting for >24 h, according to the Monitoring Trends and Determinants in Cardiovascular Disease (WHO-MONICA) Project [23]. Stroke was classified into ischemic stroke, intracerebral hemorrhage, subarachnoid hemorrhage, and other/unclassified stroke. The present study was approved by the Institutional Review Board of Shiga University of Medical Science (R2011-186).

Temperature Data

Information on average, maximum, and minimum temperature at the Hikone Meteorological Observatory, which located in the center of the Shiga Prefecture, in each month during the study period (2011–2013) was obtained from website of the Japan Meteorological Agency [24].

Statistical Analysis

Incidence rates of first-ever stroke in each month during the study period (2011–2013) were estimated using the person-year approach. The number of residents in the Shiga Prefecture was obtained by sex and 10-year age groupings from the national population estimates data on 30th June 2012 [25]. Age- and sex-adjusted incidence rates were estimated and compared using the Poisson regression models. Age- and sex-adjusted incidence rates in spring (March–May), summer (June–August), autumn (September–November), and winter (December–February) were also estimated and compared using the Poisson regression models. A p value <0.05 was considered statistically significant. All analyses were performed using SAS 9.4 (SAS Institute, Cary, NC, USA).

Results

Temperature characteristics of the Shiga Prefecture was shown in Table 1. Average temperature was lowest (<5°C) in January/February and highest (>25°C) in July/August.

From 2011 to 2013, we identified a total of 6,688 stroke cases (3,570 men, 3,118 women), of which 4,480 cases had ischemic stroke (2,518 men, 1,962 women) (1,042 lacunar infarction, 1,333 large artery infarction and 1,235 cardioembolic infarction), 1,588 had intracerebral hemorrhage (857 men, 731 women), and 563 had subarachnoid hemorrhage (166 men, 397 women). Figure 1 shows monthly variation in average temperature (top part of Fig. 1) and age- and sex-adjusted incidence rates of total stroke in the Shiga Prefecture (Fig. 1a). Incidence rates of total stroke tended to be low in summer (June–August) high in winter and (December–February). When age- and sex-adjusted incidence rates of total stroke were estimated by season with pooling of data from 2011 to 2013, incidence (per 100,000 person-years) was lowest in summer (130, 95% confidence interval [CI] 122–137) (Table 2). Compared with summer, age- and sex-adjusted incidence rates of total stroke were significantly higher in winter (170, 95% CI 161–179, $p < 0.001$), in autumn (141 95% CI 133–149,

Fig. 1. Monthly variation in average temperature and age- and sex-adjusted incidence rates of total stroke (a), ischemic stroke (b), intracerebral hemorrhage (c), and subarachnoid hemorrhage: SSR 2011–2013 (d). The top part of the figure shows average temperature at the Hikone Meteorological Observatory, which is located in the center of the Shiga Prefecture, in each month from 2011 to 2013. The lower parts of the figure show incidence rates of total stroke (a), ischemic stroke (b), intracerebral hemorrhage (c), and subarachnoid hemorrhage (d) in each month from 2011 to 2013, which was estimated using the person-year approach and standardized for age and sex using the Poisson regression models. SSR, Shiga Stroke Registry. (For figure see next page.)




Table 1. Temperature characteristics of the Shiga Prefecture from 2011 to 2013

2011												
Month	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Average, °C	2.0	4.8	5.5	10.8	17.4	23	27.3	28.1	24.3	17.8	13.1	6.0
Maximum, °C	8.3	15.8	15.6	26.6	27.2	35.2	35.2	35.9	33	25.4	23.1	16.2
Minimum, °C	-3.9	-2.9	-1.1	0.7	8.4	15	20.8	22.2	13.7	9.4	3.9	-0.3
2012												
Month	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Average, °C	3.4	3.1	6.8	12.4	16.9	21.3	26.5	28.5	25.3	17.9	10.9	4.9
Maximum, °C	9.7	10.5	19.4	26.2	27.6	29.9	35.8	35.5	33.5	27.4	20.2	12.1
Minimum, °C	-1.9	-3.8	-1.6	1.4	8.6	15.3	19.3	22.6	16.6	8.8	1.8	-2.0
2013												
Month	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Average, °C	3.3	3.4	7.9	11.7	17.2	22.7	27.1	28.3	23.5	19.3	11.2	6.0
Maximum, °C	10.9	13.1	19.8	24.0	28.7	31.8	36.2	35.6	32.2	32.1	20.5	14.7
Minimum, °C	-3.7	-2.4	-1.9	2.4	4.7	15.9	19.3	20.5	13.1	7.1	2.1	-1.2

Table shows average, maximum, and minimum temperature at the Hikone Meteorological Observatory, which located in the center of the Shiga Prefecture, in each month from 2011 to 2013.

$p = 0.020$), and in spring (151 95% CI 144–160, $p < 0.001$). Stratified analysis by age and sex demonstrated similar seasonal patterns in 4 groups (men aged <70 years, men aged ≥ 70 years, women aged <70 years and women aged ≥ 70 years) (see online suppl. Table 1; see www.karger.com/doi/10.1159/000518370 for all online suppl. material).

Similar patterns were observed for ischemic stroke (Fig. 1b), intracerebral hemorrhage (Fig. 1c) and subarachnoid hemorrhage (Fig. 1d) but seasonal variation was more evident for intracerebral hemorrhage than ischemic stroke. Age- and sex-adjusted incidence rates of ischemic stroke were 88 (95% CI 82–94, $p = 0.004$ vs. summer) in spring, 78 (95% CI 72–83) in summer, 80 (95% CI 74–86, $p = 0.549$ vs. summer) in autumn and 90 (95% CI 84–97, $p < 0.001$ vs. summer) in winter (Table 2). When ischemic stroke was classified into subtypes, there were no clear seasonal variations in incidence rates of lacunar infarction (Table 2). In contrast, incidence rates of large artery infarction were highest in winter (26 per 100,000 person-years) ($p = 0.008$ vs. summer) (Table 2). Incidence rates of cardioembolic infarction were also highest in winter (18 per 100,000 person-years) and spring (18 per 100,000 person-years) ($p < 0.001$ for both winter and spring vs. summer) (Table 2). Age- and sex-adjusted incidence rates of intracerebral hemorrhage were 39 (95% CI 35–43, $p < 0.001$ vs. summer) in spring, 30 (95% CI 27–34) in summer, 38 (95% CI 34–42, $p = 0.002$ vs. summer) in autumn and 52 (95% CI 47–57, $p < 0.001$ vs. summer) in winter (Table 2). Age- and sex-

adjusted incidence rates of subarachnoid hemorrhage were 16 (95% CI 13–19, $p = 0.170$ vs. summer) in spring, 13 (95% CI 11–16) in summer, 16 (95% CI 14–19, $p = 0.129$ vs. summer) in autumn, and 20 (95% CI 17–23 $p = 0.002$ vs. summer) in winter (Table 2).

Discussion

In the present large-scale stroke registry of general population, incidence rates of stroke were high in winter and low in summer in current Japan. Seasonal variation was more evident for intracerebral hemorrhage than for ischemic stroke.

Some prior epidemiological studies have investigated seasonal variation in stroke incidence in Japan [11–16]. A hospital-based stroke registry in Akita investigated 2,168 stroke patients from 1983 to 1985 and demonstrated higher risks of total stroke and intracerebral hemorrhage in winter than in summer, but did not report clear seasonal variation for ischemic stroke [13]. Hospital-based stroke registry in Kyoto investigated 13,788 stroke patients from 1999 to 2009 and demonstrated increased incidence rates of total stroke, intracerebral hemorrhage and subarachnoid hemorrhage in winter and spring compared with those in summer, but no clear seasonal variation was observed for ischemic stroke [15]. Another nation-wide hospital-based stroke registry (Japanese Standard Stroke Registry) investigated 35,631 stroke patients admitted to 163 acute stroke hospitals from 1999 to 2007

Table 2. Seasonal variation in age- and sex-adjusted incidence rates of stroke and its subtypes: SSR 2011–2013

	Age- and sex-adjusted incidence rate, per 100,000 person-years (95% confidence interval)		<i>p</i> value (vs. summer)
Total stroke			
Spring	151	(144–160)	<0.001
Summer	130	(122–137)	Reference
Autumn	141	(133–149)	0.020
Winter	170	(161–179)	<0.001
Ischemic stroke			
Spring	88	(82–94)	0.004
Summer	78	(72–83)	Reference
Autumn	80	(74–86)	0.549
Winter	90	(84–97)	<0.001
Lacunar infarction			
Spring	20	(17–23)	0.478
Summer	21	(19–25)	Reference
Autumn	23	(20–27)	0.325
Winter	21	(18–24)	0.815
Large artery infarction			
Spring	25	(22–28)	0.031
Summer	21	(18–24)	Reference
Autumn	20	(18–23)	0.661
Winter	26	(23–30)	0.008
Cardioembolic infarction			
Spring	18	(12–18)	<0.001
Summer	13	(9–13)	Reference
Autumn	14	(9–14)	0.517
Winter	18	(12–18)	<0.001
Intracerebral hemorrhage			
Spring	39	(35–43)	<0.001
Summer	30	(27–34)	Reference
Autumn	38	(34–42)	0.002
Winter	52	(47–57)	<0.0001
Subarachnoid hemorrhage			
Spring	16	(13–19)	0.170
Summer	13	(11–16)	Reference
Autumn	16	(14–19)	0.129
Winter	20	(17–23)	0.002

Incidence rates in spring (March–May in 2011–2013), summer (June–August in 2011–2013), autumn (September–November in 2011–2013), and winter (December–February in 2011–2013) were estimated using the person-year approach and standardized for age and sex using the Poisson regression models. Age- and sex-adjusted incidence rates were compared to the reference group of summer using the Poisson regression models. SSR, Shiga Stroke Registry.

and demonstrated increased risks of hemorrhagic stroke and cardioembolic infarction in winter, while the risks of non-cardioembolic infarction were higher in summer than winter [14]. A recent hospital-based observational study of 2,965 acute ischemic stroke patients admitted to

an acute stroke hospital from 2011 to 2015 demonstrated no clear seasonal variation in incidence rates of ischemic stroke, while the risks of cardioembolic infarction were highest in winter [16]. However, findings from hospital-based studies might be affected by seasonal variation of stroke patients' intention to visit acute-care hospitals or by capacity of each hospital. With regard to population-based study, the Hisayama Study investigated 311 stroke patients during follow-up duration from 1961 to 1985 and showed higher incidence of intracerebral hemorrhage and ischemic stroke in winter than in summer [11]. A population-based stroke registry (Takashima Stroke Registry) investigated 1,665 stroke patients from 1988 to 2001 and reported highest risks of total stroke, intracerebral hemorrhage, and ischemic stroke in spring and lowest risks in summer [12]. The results of the present large-scale population-based study of 6,688 stroke patients were comparable with those of some prior studies and this study clearly demonstrated that incidence rates of total stroke and its subtypes including ischemic stroke were higher in winter than in summer in current Japan.

Exposure to the cold provokes a range of physiological responses, predominantly as a consequence of activation of the sympathetic nervous system, including elevation in blood pressure and heart rate [26, 27]. In fact, several observational studies reported seasonal variation of BP levels (i.e., higher BP levels in winter than those in summer) [28–34]. Cold weather was also reported to be associated with increased risks of arrhythmia including atrial fibrillation [35–38]. Framingham study also reported decreased endothelial function in the cold weather [39]. Other possible mechanisms involve seasonal variation in serum lipids, fibrinogen, and viscosity [40, 41], which are reported to be risk factors of ischemic stroke [42–45]. Influenza and influenza-like symptoms, which are more likely to strike in winter than other seasons, have also been shown to be associated with stroke [46, 47]. Exacerbation of air pollution including gaseous pollutants (such as ozone) and particulate matter (such as PM_{2.5}) in winter and spring in East Asia [48] might also be attributable to seasonal variation of stroke, as exposure to air pollution has been reported to be associated with admissions to hospitals for stroke [49]. These mechanisms might be in part attributable to seasonal variation in incidence rates of stroke.

Although this is a large-scale, comprehensive, population-based registry in current Japan, several limitations should be discussed. First, we could not examine the association between seasonal variations in incidence of transient ischemic attack due to no information on the

patients with transient ischemic attack. Another limitation is that we do not have information on environmental temperature of patients at the time of stroke onset. Third, our findings might not be generalizable to other regions of Japan where climate conditions and cardiovascular risk factors of residents are different from those in the Shiga Prefecture.

Conclusion

In the present large-scale stroke registry of general population, incidence rates of stroke were high in winter and low in summer in current Japan.

Acknowledgments

We are deeply grateful to all investigators, participating hospitals, study staff at the Shiga Stroke Data Center, Shiga Medical Association, Shiga Prefecture Hospital Association, Biwako Brain and Heart attack Consortium (BIWA-BHAC), Department of Public Health Care and Welfare, Shiga prefecture government.

Statement of Ethics

The present study was approved by the Institutional Review Board of Shiga University of Medical Science (R2011-186) and the president of Shiga University of Medical Science gave the waiver for written informed consent.

References

- 1 GBD 2016 Lifetime Risk of Stroke Collaborators; Feigin VL, Nguyen G, Cercy K, Johnson CO, Alam T, et al. GBD 2016 lifetime risk of stroke collaborators. Global, regional, and country-specific lifetime risks of stroke, 1990 and 2016. *N Engl J Med* 2018;379:2429–37.
- 2 Krishnamurthi RV, Ikeda T, Feigin VL. Global, regional and country-specific burden of ischaemic stroke, intracerebral haemorrhage and subarachnoid haemorrhage: a systematic analysis of the global burden of disease study 2017. *Neuroepidemiology*. 2020;54:171–9.
- 3 GBD 2017 Causes of Death Collaborators. Global, regional, and national age-sex-specific mortality for 282 causes of death in 195 countries and territories, 1980–2017: a systematic analysis for the global burden of disease study 2017. *Lancet*. 2018;392:1736–88.
- 4 Feigin VL, Norrving B, Mensah GA. Global burden of stroke. *Circ Res*. 2017;120:439–48.
- 5 GBD 2017 DALYs and HALE Collaborators. Global, regional, and national disability-adjusted life-years (DALYs) for 359 diseases and injuries and healthy life expectancy (HALE) for 195 countries and territories, 1990–2017: a systematic analysis for the global burden of disease study 2017. *Lancet*. 2018;392:1859–922.
- 6 Katan M, Luft A. Global burden of stroke. *Semin Neurol*. 2018;38(2):208–11.
- 7 GBD 2016 Stroke Collaborators. Global, regional, and national burden of stroke, 1990–2016: a systematic analysis for the global burden of disease study 2016. *Lancet Neurol*. 2019;18:439–58.
- 8 Pandian JD, Gall SL, Kate MP, Silva GS, Akinyemi RO, Ovbiagele BI, et al. Prevention of stroke: a global perspective. *Lancet*. 2018;392:1269–78.
- 9 Rothwell PM, Wroe SJ, Slattery J, Warlow CP. Is stroke incidence related to season or temperature? The Oxfordshire community stroke project. *Lancet*. 1996;347:934–6.
- 10 Sipilä JO, Ruuskanen JO, Kauko T, Rautava P, Kytö V. Seasonality of stroke in Finland. *Ann Med*. 2017;49:310–8.
- 11 Shinkawa A, Ueda K, Hasuo Y, Kiyohara Y, Fujishima M. Seasonal variation in stroke incidence in Hisayama, Japan. *Stroke*. 1990;21:1262–7.
- 12 Turin TC, Kita Y, Murakami Y, Rumana N, Sugihara H, Morita Y, et al. Higher stroke incidence in the spring season regardless of conventional risk factors: Takashima stroke registry. *Stroke*. 1988;39:745–52.
- 13 Suzuki K, Kutsuzawa T, Takita K, Ito M, Sakamoto T, Hirayama A, et al. Clinico-epidemiologic study of stroke in Akita, Japan. *Stroke*. 1987;18:402–6.
- 14 Takizawa S, Shibata T, Takagi S, Kobayashi S. Japan standard stroke registry study group. Seasonal variation of stroke incidence in Japan for 35,631 stroke patients in the Japanese standard stroke registry, 1998–2007. *J Stroke Cerebrovasc Dis*. 2013;22:36–41.

Conflict of Interest Statement

Arima H. is an associate editor of the Journal *Cerebrovascular Disease*. The other authors have no conflicts of interest to declare.

Funding Sources

Shiga Stroke and Heart Registry is supported by Shiga Prefecture and the Japan Agency for Medical Research and Development (Grant No. 17ek0210090).

Author Contributions

Nozaki K., Kita Y., and Takashima N. contributed to the conception of this study and Nozaki K., Kita Y., Takashima N., Miyamatsu N., and Miura K. contributed to the design of this study. Fujii T., Takashima N., Kita Y., Tanaka-Mizuno S., Shitara S., Miyamatsu N., Miura K., and Nozaki K. contributed to the acquisition of data. Fujii T. and Arima H. contributed to analysis of this study, and Fujii T., Takashima N., Kita Y., Tanaka-Mizuno S., Urushitani M., Miura K., and Nozaki K. contributed to interpretation of data. Fujii T. drafted the manuscript for the work and Arima H. supervised the writing of the manuscript. All authors revised the manuscript, approved the manuscript to be published, and agree to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Data Availability Statement

Research data are not publicly available on ethical grounds.

- 15 Shigematsu K, Watanabe Y, Nakano H. Kyoto stroke registry committee. Higher ratio of ischemic stroke to hemorrhagic stroke in summer. *Acta Neurol Scand*. 2015;132:423–9.
- 16 Toyoda K, Koga M, Yamagami H, Yokota C, Sato S, Inoue M, et al. Seasonal variations in neurological severity and outcomes of ischemic stroke: 5-year single-center observational study. *Circ J*. 2018;82:1443–50.
- 17 Takashima N, Arima H, Kita Y, Fujii T, Miyamatsu N, Komori M, et al. Incidence, management and short-term outcome of stroke in a general population of 1.4 million Japanese: Shiga stroke registry. *Circ J*. 2017;81:1636–46.
- 18 Widhi Nugroho A, Arima H, Miyazawa I, Fujii T, Miyamatsu N, Sugimoto Y, et al. The association between glomerular filtration rate estimated on admission and acute stroke outcome: the Shiga stroke registry. *J Atheroscler Thromb*. 2018;25:570–9.
- 19 Widhi Nugroho A, Arima H, Takashima N, Fujii T, Shitara S, Miyamatsu N, et al. The JAGUAR score predicts 1-month disability/death in ischemic stroke patient ineligible for recanalization therapy. *J Stroke Cerebrovasc Dis*. 2018;27:2579–86.
- 20 Takashima N, Arima H, Kita Y, Fujii T, Miyamatsu N, Komori M, et al. Two-year survival after first-ever stroke in a general population of 1.4 million Japanese: Shiga stroke registry. *Circ J*. 2018;82:2549–56.
- 21 Takashima N, Arima H, Kita Y, Fujii T, Tanaka-Mizuno S, Shitara S, et al. Shiga stroke and heart attack registry group. Two-year recurrence after first-ever stroke in a general population of 1.4 million Japanese patients- the Shiga stroke and heart attack registry study. *Circ J*. 2020;84:943–8.
- 22 Takashima N, Arima H, Kita Y, Fujii T, Tanaka-Mizuno S, Shitara S, et al. Long-term survival after stroke in 1.4 million Japanese population: shiga stroke and heart attack registry. *J Stroke*. 2020;22:336–44.
- 23 Thorvaldsen P, Kuulasmaa K, Rajakangas A-M, Rastenyte D, Sarti C, Wilhelmsen L. Stroke trends in the WHO MONICA project. *Stroke*. 1997;28(3):500.
- 24 Japan Meteorological Agency. [Weather data in the past \(in Japanese\)](https://www.data.jma.go.jp/obd/stats/etrn/index.php?prec_no=60&block_no=47761&year=&month=&day=&view=); 2021. Available from: https://www.data.jma.go.jp/obd/stats/etrn/index.php?prec_no=60&block_no=47761&year=&month=&day=&view= Accessed 2021 Mar 3.
- 25 Shiga Prefecture. [Demographic statistics in the Shiga Prefecture in 2012](https://www.pref.shiga.lg.jp/kensei/tokei/tokeisyu/305448.html); 2012. Available from: <https://www.pref.shiga.lg.jp/kensei/tokei/tokeisyu/305448.html> Accessed 2021 March 3.
- 26 Okada M, Kakehashi M. Effects of outdoor temperature on changes in physiological variables before and after lunch in healthy women. *Int J Biometeorol*. 2014;58:1973–81.
- 27 Cui J, Muller MD, Blaha C, Kunselman AR, Sinoway LI. Seasonal variation in muscle sympathetic nerve activity. *Physiol Rep*. 2015;3:e12492.
- 28 Brennan PJ, Greenberg G, Miall WE, Thompson SG. Seasonal variation in arterial blood pressure. *Br Med J*. 1982;285:919–23.
- 29 Sega R, Cesana G, Bombelli M, Grassi G, Stella ML, Zanchetti A, et al. Seasonal variations in home and ambulatory blood pressure in the PAMELA population. *Pressione Arteriose Monitorate E Loro Associazioni*. *J Hypertens*. 1998;16:1585–92.
- 30 Madsen C, Nafstad P. Associations between environmental exposure and blood pressure among participants in the Oslo health study (HUBRO). *Eur J Epidemiol*. 2006;21:485–91.
- 31 Alperovitch A, Lacombe JM, Hanon O, Dartigues JF, Ritchie K, Ducimetière P, et al. Relationship between blood pressure and outdoor temperature in a large sample of elderly individuals: the three-city study. *Arch Intern Med*. 2009;169:75–80.
- 32 Lewington S, Li L, Sherkliker P, Guo Y, Millwood I, Bian Z, et al. Seasonal variation in blood pressure and its relationship with outdoor temperature in 10 diverse regions of China: the China Kadoorie Biobank. *J Hypertens*. 2012;30:1383–91.
- 33 Modesti PA, Morabito M, Massetti L, Rapi S, Orlandini S, Mancia G, et al. Seasonal blood pressure changes: an independent relationship with temperature and daylight hours. *Hypertension*. 2013;61:908–14.
- 34 Iwahori T, Miura K, Obayashi K, Ohkubo T, Nakajima H, Shiga T, et al. Seasonal variation in home blood pressure: findings from nationwide web-based monitoring in Japan. *BMJ Open*. 2018;8:e017351.
- 35 Frost L, Johnsen SP, Pedersen L, Husted S, Engholm G, Sørensen HT, et al. Seasonal variation in hospital discharge diagnosis of atrial fibrillation: a population-based study. *Epidemiology*. 2002;13:211–5.
- 36 Watanabe E, Kuno Y, Takasuga H, Tong M, Sobue Y, Uchiyama T, et al. Seasonal variation in paroxysmal atrial fibrillation documented by 24-h Holter electrocardiogram. *Heart Rhythm*. 2007;4:27–31.
- 37 Fustinoni O, Saposnik G, Esnaola y Rojas MM, Lakkis SG, Sposato LA; ReNACer Investigators. Higher frequency of atrial fibrillation linked to colder seasons and air temperature on the day of ischemic stroke onset. *J Stroke Cerebrovasc Dis*. 2013;22:476–81.
- 38 Loomba RS. Seasonal variation in paroxysmal atrial fibrillation: a systematic review. *J Atr Fibrillation*. 2015 Feb;7:1201.
- 39 Widlansky ME, Vita JA, Keyes MJ, Larson MG, Hamburg NM, Levy D, et al. Relation of season and temperature to endothelium-dependent flow-mediated vasodilation in subjects without clinical evidence of cardiovascular disease (from the Framingham Heart Study). *Am J Cardiol*. 2007;100:518–23.
- 40 Ockene IS, Chiriboga DE, Stanek EJ 3rd, Harmatz MG, Nicolosi R, Saperia G, et al. Seasonal variation in serum cholesterol levels: treatment implications and possible mechanisms. *Arch Intern Med*. 2004;164:863–70.
- 41 Hermida RC, Calvo C, Ayala DE, López JE, Fernández JR, Mojón A, et al. Seasonal variation of fibrinogen in dipper and nondipper hypertensive patients. *Circulation*. 2003;108:1101–6.
- 42 Zhang X, Patel A, Horibe H, Wu Z, Barzi F, Rodgers A, et al. Cholesterol, coronary heart disease, and stroke in the Asia Pacific region. *Int J Epidemiol*. 2003;32:563–72.
- 43 Miller M, Stone NJ, Ballantyne C, Bittner V, Criqui MH, Ginsberg HN, et al. American heart association clinical lipidology, thrombosis, and prevention committee of the council on nutrition, physical activity, and metabolism; council on arteriosclerosis, thrombosis and vascular biology; council on cardiovascular nursing; council on the kidney in cardiovascular disease. triglycerides and cardiovascular disease: a scientific statement from the American heart association. *Circulation*. 2011;123:2292–333.
- 44 Baeten KM, Akassoglou K. Extracellular matrix and matrix receptors in blood-brain barrier formation and stroke. *Dev Neurobiol*. 2011;71:1018–39.
- 45 Song SH, Kim JH, Lee JH, Yun YM, Choi DH, Kim HY. Elevated blood viscosity is associated with cerebral small vessel disease in patients with acute ischemic stroke. *BMC Neurol*. 2017;17:20.
- 46 Kwok CS, Aslam S, Kontopantelis E, Myint PK, Zaman MJ, Buchan I, et al. Influenza, influenza-like symptoms and their association with cardiovascular risks: a systematic review and meta-analysis of observational studies. *Int J Clin Pract*. 2015;69:928–37.
- 47 Elkind MSV, Boehme AK, Smith CJ, Meisel A, Buckwalter MS. Infection as a Stroke risk factor and determinant of outcome after stroke. *Stroke*. 2020;51:3156–68.
- 48 Turnock ST, Allen RJ, Andrews M, Bauer SE, Deushi M, Emmons L, et al. Historical and future changes in air pollutants from CMIP6 models. *Atmos Chem Phys*. 2020;20(23):14547–79.
- 49 Shah AS, Lee KK, McAllister DA, Hunter A, Nair H, Whiteley W, et al. Short term exposure to air pollution and stroke: systematic review and meta-analysis. *BMJ*. 2015;350:h1295.