

A numerical property of hilbert functions and lex segment ideals

*Original*

A numerical property of hilbert functions and lex segment ideals / Favacchio, G.. - In: JOURNAL OF THE KOREAN MATHEMATICAL SOCIETY. - ISSN 0304-9914. - STAMPA. - 57:3(2020), pp. 777-792. [10.4134/JKMS.j190380]

*Availability:*

This version is available at: 11583/2859978 since: 2021-01-12T16:20:33Z

*Publisher:*

Korean Mathematical Society

*Published*

DOI:10.4134/JKMS.j190380

*Terms of use:*

openAccess

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

*Publisher copyright*

Nature --&gt; vedi Generico

[DA NON USARE] ex default\_article\_draft

(Article begins on next page)

# A NUMERICAL PROPERTY OF HILBERT FUNCTIONS AND LEX SEGMENT IDEALS

GIUSEPPE FAVACCHIO

ABSTRACT. We introduce the *fractal expansions*, sequences of integers associated to a number. We show that these sequences characterize the  $O$ -sequences and encode some information about lex segment ideals. Moreover, we introduce a numerical functions called *fractal functions*, and we use them to solve the open problem of the classification of the Hilbert functions of any bigraded algebra.

## 1. INTRODUCTION

In Commutative Algebra (and other fields of pure mathematics) it often happens that easy numerical conditions describe some deeper algebraic results. A significant example are the  $O$ -sequences.

Let  $S := k[x_1, \dots, x_n]$  be the standard graded polynomial ring and let  $I \subseteq S$  be a homogeneous ideal. The quotient ring  $S/I$  is called a *standard graded  $k$ -algebra*. The Hilbert function of  $S/I$  is defined as  $H_{S/I} : \mathbb{N} \rightarrow \mathbb{N}$  such that

$$H_{S/I}(t) := \dim_k (S/I)_t = \dim_k S_t - \dim_k I_t.$$

A famous theorem, due to Macaulay (cf. [14]) and pointed out by Stanley (cf. [16]), characterizes the numerical functions that are Hilbert functions of a standard graded  $k$ -algebra, i.e. the functions  $H$  such that  $H = H_{S/I}$  for some homogeneous ideal  $I \subseteq S$ . To introduce this fundamental result we need some preparatory material.

Let  $h, i > 0$  be integers, we can uniquely write  $h$  as

$$(1) \quad h = \binom{m_i}{i} + \binom{m_{i-1}}{i-1} + \dots + \binom{m_j}{j}$$

where  $m_i > m_{i-1} > \dots > m_j \geq j \geq 1$ . This expression is called the  *$i$ -binomial expansion* of the integer  $h$ .

If  $h > 0$  has  $i$ -binomial expansion as in 1, then we set

$$h^{(i)} = \binom{m_i + 1}{i + 1} + \binom{m_i}{i} + \dots + \binom{m_j + 1}{j + 1}.$$

We use the convention that  $0^{(i)} = 0$ .

For example, since  $7 = \binom{4}{3} + \binom{3}{2}$ , the 3-binomial expansion of 7 is and  $7^{(3)} = \binom{5}{4} + \binom{4}{3} = 9$ .

**Definition 1.1.** A sequence of non-negative integers  $(h_0, h_1, h_2 \dots)$  is called an  $O$ -sequence if

- i)  $h_0 = 1$ ;
- ii)  $h_{i+1} \leq h_i^{(i)}$  for all  $i > 0$ .

---

2000 *Mathematics Subject Classification.* 13F20, 13A15, 13D40.

*Key words and phrases.* Hilbert function, lex segment ideal, bigraded algebra .

Version: May 30, 2019.

**e-mail address:** favacchio@dmi.unict.it .

An  $O$ -sequence is said to have maximal growth from degree  $i$  to degree  $i + 1$  if  $h_{i+1} = h_i^{\binom{i}{i}}$

We are now ready to enunciate the Macaulay's Theorem. It characterizes the Hilbert function of standard graded  $k$ -algebras bounding the growth from any degree to the next. The proof of this theorem and more details about  $O$ -sequences are also discussed in [3] Chapter 4. We represent  $H_{S/I}$  as a sequence of integers  $(h_0, h_1, h_2, \dots)$  where  $h_t := H_{S/I}(t)$ .

**Theorem 1.2** (Macaulay, [14]). *Let  $H := (h_0, h_1, h_2, \dots)$  be a sequence of integers, then the following are equivalent:*

- 1)  $H$  is the Hilbert function of a standard graded  $k$ -algebra;
- 2)  $H$  is an  $O$ -sequence.

It is interesting to find an extension of the above theorem to the multi-graded case. Multi-graded Hilbert functions and resolutions arise in many contexts, see for instance [7, 9, 11, 17, 18], and properties related to the Hilbert function of multi-graded algebras are currently studied, see [6, 8, 10, 15].

The generalization of Macaulay's theorem to multi-graded rings is an open problem. A first answer was given by the author in [5] where the Hilbert functions of a bigraded algebra in  $k[x_1, x_2, y_1, y_2]$  are classified.

In this paper, see Section 4 Theorem 4.12, we generalize the Macaulay's theorem to any bigraded algebra. First, in Section 2, we introduce  $\Phi(n)$  a list of finite sequences called fractal expansion of  $n$ . Then we define a *coherent truncation* of these vectors and we show that these objects are strictly related to the  $O$ -sequences. Indeed, in Section 3 we show that they also characterize the Hilbert function of standard graded  $k$ -algebras. Furthermore, we show that these sequences can be used to compute the Betti numbers of a lex ideal.

The computer program CoCoA [1] was indispensable for all the computations.

## 2. THE EXPANSION OF A FRACTAL SEQUENCE

In this section we introduce a sequence of tuples, called *coherent fractal growth*, and we study its properties. In the main result of this section, Theorem 2.14, we prove that these sequences have the same behavior of the  $O$ -sequences.

Roughly speaking a numerical sequence  $\sigma$  is called fractal if once we delete the first occurrence of each number it remains identical to the original. Such property thus implies we can repeat this process indefinitely, and  $\sigma$  contains infinitely many copy of itself. It has something like a *fractal* behavior. See [13] for a formal definition and further properties.

For instance, one can show that the sequence

$$\sigma = (1, 1, 2, 1, 2, 3, 1, 2, 3, 4, 1, 2, 3, 4, 5, \dots)$$

is fractal. Indeed, after removing the first occurrence of each number we get a sequence that is the same as the starting one

$$(\cancel{1}, 1, \cancel{2}, 1, 2, \cancel{3}, 1, 2, 3, \cancel{4}, 1, 2, 3, 4, \cancel{5}, \dots) = \sigma.$$

We introduce some notation. Given a positive integer  $a \in \mathbb{N}$  we denote by  $[a] := (1, 2, \dots, a) \in \mathbb{N}^a$  the tuple of length  $a$  consisting of all the positive integers less than or equal to  $a$  written in increasing order.

Given a (finite or infinite) sequence  $\sigma$  of positive integers we construct a new sequence, named the expansion of  $\sigma$ , denoted by  $[\sigma]$ . If  $\sigma := (s_1, s_2, s_3, \dots)$  we set  $[\sigma] := [s_1] \parallel [s_2], \parallel [s_3] \parallel \dots$ , where the symbol " $\parallel$ " denotes the associative operation of concatenation of two vectors. E.g.  $(3, 5, 4) \parallel (2, 3) = (3, 5, 4, 2, 3)$ .

This construction can be recursively applied. We denote by  $[\sigma]^d := [[\sigma]^{d-1}]$  where we set  $[\sigma]^0 := \sigma$ . For a positive integer  $a$ , we also denote by  $[a]^d := [[a]^{d-1}]$ , where  $[a]^0 := (a)$ .

For instance we have

$$[3] = (1, 2, 3), \quad [3]^2 = (1, 1, 2, 1, 2, 3), \quad [3]^3 = (1, 1, 1, 2, 1, 1, 2, 1, 2, 3).$$

**Lemma 2.1.** *Let  $\sigma := (s_1, s_2, s_3, \dots)$  be a sequence of positive integers. Then  $[\sigma]^d = [s_1]^d \parallel [s_2]^d \parallel [s_3]^d \parallel \dots$ .*

*Proof.* If  $d = 0$  the statement is true. Assume  $d > 0$ . By definition  $[\sigma]^d = [[\sigma]^{d-1}]$  then by the inductive hypothesis we have

$$[\sigma]^d = [[s_1]^{d-1}] \parallel [[s_2]^{d-1}] \parallel [[s_3]^{d-1}] \parallel \dots = [s_1]^d \parallel [s_2]^d \parallel [s_3]^d \parallel \dots.$$

□

**Corollary 2.2.** *Let  $a \in \mathbb{N}$  be a positive integer. Then*

$$[a]^d = [1]^{d-1} \parallel [2]^{d-1} \parallel \dots \parallel [a]^{d-1}.$$

*Proof.* The statement follows by Lemma 2.1 since  $[a]^d = [(1, 2, \dots, a)]^{d-1}$ . □

**Remark 2.3.** The sequence  $[\mathbb{N}] := [1] \parallel [2] \parallel \dots \parallel [n] \parallel \dots$  is a fractal sequence.

Given a sequence  $\sigma$ , the symbols  $\sigma_i$  and  $\sigma(i)$  both denote the  $i$ -th entry of  $\sigma$ . If  $\sigma$  is finite then  $|\sigma|$  denotes the number of entries and  $\sum \sigma$  their sum. We use the convention that these values are " $\infty$ " for infinite sequences of positive integers. For instance,  $[3]_4^3 = 2$ ,  $|[3]^3| = 10$  and  $\sum [3]^3 = 15$ .

Throughout this paper we use the convention that, for a finite sequence  $\sigma$ , the notation  $\sigma_a$  or  $\sigma(a)$  implies  $a \leq |\sigma|$ .

**Remark 2.4.** Note that, for a finite sequence of positive integers  $\sigma$ , the definition of  $[\sigma]$  easily implies the equality  $\sum \sigma = |[\sigma]|$ .

Given a positive integer  $n$ , we define the fractal expansion of  $n$  as the list

$$\Phi(n) := ([n]^0, [n]^1, \dots, [n]^d, \dots).$$

Each element in  $\Phi(n)$  is a finite sequence of positive integers. In the following lemma we compute the number of their entries.

**Lemma 2.5.** *Let  $n$  be a positive integer. Then  $|[n]^d| = \binom{d+n-1}{d}$  and  $\sum [n]^d = \binom{d+n}{d+1}$ .*

*Proof.* By definition we have  $|[n]^0| = \binom{n-1}{0} = 1$  and  $|[n]^1| = \binom{n}{1} = n$ . By Corollary 2.2 we have  $[n]^d = [1]^{d-1} \parallel [2]^{d-1} \parallel \dots \parallel [n]^{d-1}$ , therefore

$$|[n]^d| = \sum_{j=1}^n |[j]^{d-1}| = \sum_{j=1}^n \binom{d+j-2}{n-1} = \binom{d+n-1}{d}.$$

Moreover, by Remark 2.4 we have  $\sum [n]^d = |[n]^{d+1}| = \binom{d+n}{d+1}$ . □

Next lemma introduces a way to decompose a number as sum of binomial coefficients that is slight different to the Macaulay decomposition. We use the convention that  $\binom{a}{b} = 0$  whenever  $a < b$ .

**Lemma 2.6.** *Let  $d$  be a positive integer. Any  $a \in \mathbb{N}$  can be written uniquely in the form*

$$(2) \quad a = \binom{k_d}{d} + \binom{k_{d-1}}{d-1} + \dots + \binom{k_2}{2} + \binom{k_1}{1}$$

where  $k_d > k_{d-1} > \dots > k_2 \geq k_1 \geq 1$

*Proof.* In order to prove the existence, we choose  $k_d$  maximal such that  $\binom{k_d}{d} < a$ . If  $d = 1$  the statement is trivial. If  $d = 2$  then let  $k_2$  be the maximal integer such that  $a - \binom{k_2}{2} > 0$ . Then set  $k_1 := a - \binom{k_2}{2}$ . Since  $a \leq \binom{k_2+1}{2}$  we have  $k_1 = a - \binom{k_2}{2} \leq \binom{k_2+1}{2} - \binom{k_2}{2} = k_2$ . Assume  $d > 2$ , and let  $k_d$  be the maximum integer such that  $a - \binom{k_d}{d} > 0$ . By induction,  $a - \binom{k_d}{d}$  can be written as  $a - \binom{k_d}{d} = \sum_{i=1}^{d-1} \binom{k_i}{i}$ , where  $k_{d-1} > \dots > k_2 \geq k_1 \geq 1$ . Since  $k_1 \geq 1$  we have  $a - \binom{k_d}{d} > \binom{k_{d-1}}{d-1}$ . Moreover, since  $\binom{k_{d+1}}{d} \geq a$ , it follows that

$$\binom{k_d}{d-1} = \binom{k_d+1}{d} - \binom{k_d}{d} \geq a - \binom{k_d}{d} > \binom{k_{d-1}}{d-1}.$$

Hence  $k_d > k_{d-1}$ . The uniqueness follows by induction on  $d$ . If  $d = 1$  it is trivial. Now let  $d > 1$  and assume for each  $b \in \mathbb{N}$  the uniqueness of the decomposition

$$(3) \quad b = \binom{k'_{d-1}}{d-1} + \dots + \binom{k'_2}{2} + \binom{k'_1}{1}$$

where  $k'_{d-1} > \dots > k'_2 \geq k'_1 \geq 1$ . Let  $a \in \mathbb{N}$  and let

$$(4) \quad a = \binom{k_d}{d} + \binom{k_{d-1}}{d-1} + \dots + \binom{k_2}{2} + \binom{k_1}{1}$$

be a decomposition of  $a$ . Then we claim that  $k_d$  is the maximal integer such that  $\binom{k_d}{d} < a$ . Indeed, if  $\binom{k_{d+1}}{d} < a$  we get  $a > \binom{k_{d+1}}{d} \geq \binom{k_d}{d} + \binom{k_{d-1}}{d-1} + \dots + \binom{k_2}{2} + \binom{k_1}{1} = a$ . Consider now the number  $a - \binom{k_d}{d}$ . From equation 4 we have  $a - \binom{k_d}{d} = \sum_{i=1}^{d-1} \binom{k_i}{i}$  where  $k_{d-1} > \dots > k_2 \geq k_1 \geq 1$ . The uniqueness of this decomposition follows from inductive hypothesis. This also implies the uniqueness of equation 4.  $\square$

**Remark 2.7.** The decomposition in Lemma 2.6 is different from the Macaulay decomposition since it is always required that  $k_1 \geq 1$ . Moreover, for any  $j \geq 2$  we only have that  $k_j \geq j - 1$ , thus some binomial coefficient could be zero. For instance, we have  $1 = \binom{d-1}{d} + \binom{d-2}{d-1} + \dots + \binom{1}{2} + \binom{1}{1}$  where the first  $d - 1$  binomial coefficients in the sum are equal to zero.

**Definition 2.8.** We refer to equation 2 as the  $d$ -fractal decomposition of  $a$ . We denote by

$$[a]^{(d)} := (k_d - d + 2, k_{d-1} - d + 3, \dots, k_4 - 2, k_3 - 1, k_2, k_1) \in \mathbb{N}^d.$$

We call these numbers the  $d$ -fractal coefficients of  $a$ .

$[a]^{(d)}$  is a not increasing sequence of positive integers. Indeed,  $[a]_d^{(d)} > 0$  and by construction  $k_1 \leq k_2$ . Moreover, for  $j > 1$ , since  $k_j > k_{j-1}$ , we have  $k_j - j + 2 \geq k_{j-1} - j + 1$ .

Next result explains the name " $d$ -fractal decomposition". We show that  $k_1$  is the  $a$ -th entry in  $[n]^d$  i.e. in  $[\mathbb{N}]^{d-1}$ . We use the convention that  $[0]^d$  is the empty sequence and  $|[0]^d| = 0$ .

**Theorem 2.9.**  $[\mathbb{N}]_a^{d-1} = [a]_a^{(d)}$ .

*Proof.* Assume  $n$  is large enough, we will show that  $[n]_a^d = [a]_a^{(d)}$ . If  $d = 1$  then  $[n]_a^1 = a$  and  $a = \binom{a}{1}$  thus  $[a]_a^{(1)} = a = [n]_a^1$ . We now assume  $d > 1$ . Let  $a = \binom{k_d}{d} + \binom{k_{d-1}}{d-1} + \dots + \binom{k_2}{2} + \binom{k_1}{1}$  be the  $d$ -fractal decomposition of  $a$ . Note that, since  $\binom{k_{d-1}}{d-1} + \dots + \binom{k_2}{2} + \binom{k_1}{1}$  is the  $(d-1)$  fractal decomposition of  $a - \binom{k_d}{d}$ , by the inductive hypothesis, we have  $k_1 = [n]_{a - \binom{k_d}{d}}^{d-1}$ . Therefore, we only have to show that  $[n]_a^d = [n]_{a - \binom{k_d}{d}}^{d-1}$ . Since  $a \leq \binom{k_{d+1}}{d}$ , by Lemma 2.5, we have  $|[k_d - d + 2]^d| \geq a > |[k_d - d + 1]^d|$ . Since from Corollary 2.2 we have the equivalence  $[n]^d = [1]^{d-1} \parallel [2]^{d-1} \parallel \dots \parallel [k_d - d +$

$1]^{d-1} \parallel [k_d - d + 2]^{d-1} \parallel \dots$ , it follows that the  $a$ -th entry in  $[n]^d$  is the  $\binom{a - \binom{k_d}{d}}{d}$ -th in  $[k_d - d + 2]^{d-1}$ , i.e.,  $[n]_a^d = [k_d - d + 2]_{a - \binom{k_d}{d}}^{d-1} = [n]_{a - \binom{k_d}{d}}^{d-1}$ .  $\square$

We introduce the lexicographic order for elements in  $\mathbb{N}^d$ . Given  $\alpha, \beta \in \mathbb{N}^d$  then  $\alpha <_{lex} \beta$  if and only if for some  $i \leq d$  we have  $\alpha_j = \beta_j$  for  $j < i$  and  $\alpha_i < \beta_i$ . The following lemma is crucial for our intent. We prove that the  $d$ -fractal coefficients have a good behavior with respect the lex order.

**Lemma 2.10.**  $[a]^{(d)} <_{lex} [b]^{(d)}$  iff  $a < b$ .

*Proof.* If  $d = 1$  the assertion is trivial. Let  $d > 1$  and let  $a, b$  be two integers with fractal decomposition

$$a = \binom{a_d}{d} + \binom{a_{d-1}}{d-1} + \dots + \binom{a_2}{2} + \binom{a_1}{1}, \quad a_d > \dots > a_2 \geq a_1 \geq 1$$

and

$$b = \binom{b_d}{d} + \binom{b_{d-1}}{d-1} + \dots + \binom{b_2}{2} + \binom{b_1}{1}, \quad b_d > \dots > b_2 \geq b_1 \geq 1.$$

If  $[a]^{(d)} <_{lex} [b]^{(d)}$  then there is an index  $j$  such that  $[a]_i^{(d)} = [b]_i^{(d)}$  for any  $i < j$  and  $[a]_j^{(d)} < [b]_j^{(d)}$ . Hence,  $\binom{a_i}{i} = \binom{b_i}{i}$  for any  $i < j$  and  $\binom{a_j}{j} < \binom{b_j}{j}$ . If  $j = d$  then easily  $b > a$ , otherwise we have

$$b > \binom{b_d}{d} + \binom{b_{d-1}}{d-1} + \dots + \binom{b_j}{j} \geq \binom{a_d}{d} + \binom{a_{d-1}}{d-1} + \dots + \binom{a_i + 1}{j} \geq a.$$

Vice versa let  $b > a$ . We claim that  $b_d \geq a_d$ . Indeed, if  $b_d < a_d$  we get  $a > \binom{a_d}{d} \geq \binom{b_d + 1}{d} \geq b$  contradicting  $b > a$ . So if  $b_d > a_d$  we are done otherwise the statement follows by induction.  $\square$

Given two sequences  $\tau$  and  $\sigma$  we say that  $\tau$  is a truncation of  $\sigma$  if  $|\tau| \leq |\sigma|$  and  $\tau(j) = \sigma(j)$  for any  $j \leq |\tau|$ . For instance  $(1, 1, 2, 1, 2)$  is a truncation of  $[3]^2$ .

Next definition introduces the main tool of the paper. The *coherent fractal growths* are suitable truncations of the elements in the fractal expansion of  $n$ .

**Definition 2.11.** We say that  $T := (\tau_0, \tau_1, \tau_2, \dots)$  is a coherent fractal growth if  $\tau_0 := (n)$  and  $\tau_j$  is a truncation of  $[\tau_{j-1}]$  for each  $j \geq 1$ .

For instance  $((3), (1, 2, 3), (1, 1, 2, 1, 2), (1, 1, 1, 2, 1, 1), (1, 1, 1, 1))$  is a coherent fractal growth. Indeed one can check that each elements is truncation of the expansion of the previous one. On the other hand, for instance,  $((3), (1, 2, 3), (1, 1, 2), (1, 1, 1, 2, 1))$  is not a coherent fractal growth. Indeed,  $(1, 1, 1, 2, 1)$  is not a truncation of  $[(1, 1, 2)] = (1, 1, 1, 2)$ .

**Remark 2.12.** Note that, in a coherent fractal growth,  $\tau_d$  consists of the first  $|\tau_d|$  elements in  $[n]^d$ . Moreover, the length of the elements in  $T := (\tau_0, \tau_1, \tau_2, \dots)$ , a coherent fractal growth, is bounded for any  $d$ . Indeed, by Remark 2.4 we have

$$(5) \quad |\tau_d| \leq \sum \tau_{d-1}$$

for each  $d \geq 1$ .

In the last part of this section we prove that the bound in Remark 2.12 is equivalent to the binomial expansion for a  $O$ -sequence. In order to relate the coherent fractal growth with  $O$ -sequences we need the following lemma.

**Lemma 2.13.** Let  $[a]^{(d)} = (c_d, c_{d-1}, \dots, c_1)$  be the  $d$ -fractal coefficients of  $a$ . Then the  $(d+1)$ -fractal coefficients of  $a^{(d)}$  are

$$[a^{(d)}]^{(d+1)} = (c_d, c_{d-1}, \dots, c_2, c_1, c_1).$$

*Proof.* The  $d$ -fractal decomposition of  $a$  is, by definition,

$$a = \binom{c_d + d - 2}{d} + \binom{c_{d-1} + d - 3}{d-1} + \cdots + \binom{c_3 + 1}{3} + \binom{c_2}{2} + \binom{c_1}{1}.$$

If  $c_1 < c_2$ , we get the Macaulay decomposition of  $a$  by removing the binomials  $\binom{j}{i}$  equal to 0. Since  $\binom{j}{i} = 0$  implies  $\binom{j+1}{i+1} = 0$  we have

$$a^{(d)} = \binom{c_d + d - 1}{d+1} + \binom{c_d + d - 2}{d} + \cdots + \binom{c_3 + 2}{4} + \binom{c_2 + 1}{3} + \binom{c_1 + 1}{2}.$$

Since  $\binom{c_1+1}{2} = \binom{c_1}{2} + \binom{c_1}{1}$  we are done. Now we consider the case  $c_1 = c_2$ . Then we get the following decomposition of  $a$

$$a = \binom{c_d + d - 2}{d} + \binom{c_{d-1} + d - 3}{d-1} + \cdots + \binom{c_3 + 1}{3} + \binom{c_2 + 1}{2}.$$

Thus, if  $c_3 > c_2$ , this representation is the Macaulay decomposition of  $a$  once we remove the binomials  $\binom{j}{i}$  equal to 0.

$$\begin{aligned} a^{(d)} &= \binom{c_d + d - 1}{d+1} + \binom{c_{d-1} + d - 2}{d} + \cdots + \binom{c_3 + 2}{4} + \binom{c_2 + 2}{3} = \\ &= \binom{c_d + d - 1}{d+1} + \binom{c_{d-1} + d - 2}{d} + \cdots + \binom{c_3 + 2}{4} + \binom{c_2 + 1}{3} + \binom{c_1 + 1}{2} = \\ &= \binom{c_d + d - 1}{d+1} + \binom{c_{d-1} + d - 2}{d} + \cdots + \binom{c_3 + 2}{4} + \binom{c_2 + 1}{3} + \binom{c_1}{2} + \binom{c_1}{1}. \end{aligned}$$

The proof follows in a finite number of steps by iterating this argument.  $\square$

The following theorem is the main result of this section. We show that the length of the elements in a coherent fractal growth is an  $O$ -sequence.

**Theorem 2.14.** *Let  $T = (\tau_0, \tau_1, \tau_2 \dots)$  be a list of truncations of  $\Phi(n) = ((n), [n], [n]^2, \dots)$ . Then the following are equivalent*

- i)  $T = (\tau_0, \tau_1, \tau_2 \dots)$  is a coherent fractal growth;
- ii)  $(|\tau_0|, |\tau_1|, |\tau_2|, \dots)$  is an  $O$ -sequence.

*Proof.* In order to prove (i)  $\Rightarrow$  (ii) we need to show  $|\tau_{d+1}| \leq |\tau_d|^{(d)}$ , for each  $d \geq 0$ . Set  $a = |\tau_d|$  and take the  $d$ -fractal decomposition of  $a$

$$a = \binom{k_d}{d} + \binom{k_{d-1}}{d-1} + \cdots + \binom{k_3}{3} + \binom{k_2}{2} + \binom{k_1}{1}.$$

If  $a = \binom{n+d-1}{d} = |[n]^d|$  the statement follows by Lemma 2.5. Assume now  $a < \binom{n+d-1}{d}$ . Since  $\tau_d$  is a truncation of  $[n]^d = [1]^{d-1} \parallel [2]^{d-1} \parallel \cdots \parallel [n]^{d-1}$  and, by Lemma 2.5,

$$|[k_d - d + 1]^d| = \binom{k_d}{d} < a \leq \binom{k_d + 1}{d} = |[k_d - d + 2]^d|$$

we get (denoted by  $[0]$  the empty sequence)

$$\tau_d = [k_d - d + 1]^d \parallel \tau'_d$$

where  $\tau'_d$  is a truncation of  $[k_d - d + 2]^{d-1}$ . Therefore, iterating this argument, we have

$$\tau_d = [k_d - d + 1]^d \parallel [k_{d-1} - d + 2]^{d-1} \parallel \cdots \parallel [k_3 - 1]^3 \parallel [k_2 - 1]^2 \parallel [k_1].$$

By equation 5 in Remark 2.12, we have  $|\tau_{d+1}| \leq \sum \tau_d = \sum_{i=1}^d \binom{k_i+1}{d+1} =$

$$= \binom{k_d + 1}{d+1} + \binom{k_{d-1} + 1}{d} + \cdots + \binom{k_2 + 1}{3} + \binom{k_1}{2} + \binom{k_1}{1}.$$

This sum, by Lemma 2.13, is equal to  $a^{(d)}$ .

Vice versa, to prove  $(ii) \Rightarrow (i)$ , we have to show that, for each  $d \geq 0$ , the sequence  $\tau_{d+1}$  is a truncation of  $[\tau_d]$  i.e.  $|\tau_{d+1}| \leq |[\tau_d]|$ . By using the same argument as above, we get  $|\tau_d|^{(d)} = \sum \tau_d$ . Then, the statement follows since  $|\tau_{d+1}| \leq |\tau_d|^{(d)}$ , by hypothesis, and  $|[\tau_d]| = \sum \tau_d$ , by Remark 2.12.  $\square$

Let's check, for instance, that  $H := (1, 3, 3, 4)$  is an  $O$ -sequence. We write a sequence of truncations of  $\Phi(3)$  of length 1, 3, 3, 4 respectively. We get

$$T := ((3), (1, 2, 3), (1, 1, 2), (1, 1, 1, 2)).$$

It is a coherent fractal growth. Indeed, by definition each sequence is a truncation of the bracket of the previous one, e.g.  $[(1, 2, 3)] = (1, 1, 2) \parallel (1, 2, 3)$  and  $[(1, 1, 2)] = (1, 1, 1, 2)$ . Now, we check that  $H := (1, 3, 5, 8)$  is not an  $O$ -sequence. Indeed, take a coherent fractal growth

$$((3), (1, 2, 3), (1, 1, 2, 1, 2), \tau_3)$$

where the first three sequences are truncations of length 1, 3, 5 of the elements in  $\Phi(3)$ . Then  $\tau_3$  should be a truncation of  $[(1, 1, 2, 1, 2)] = (1, 1, 1, 2, 1, 1, 2)$  that has length 7. Thus  $|\tau_3| \leq 7$  and  $(1, 3, 5, 8)$  is not an  $O$ -sequence.

### 3. FRACTAL EXPANSIONS AND HOMOLOGICAL INVARIANTS

In Section 2 we introduced a novel approach to describe the  $O$ -sequences. In this section we show the "algebraic" meaning of a coherent fractal growth. We directly relate these sequences to lex segment ideals and their homological invariants. In particular, the Eliahou-Kervaire formula is naturally applied to our case. Therefore, the fractal expansion of  $n$  is used in Proposition 3.4 to compute the Betti numbers of a lex algebra. See Section 2.1.2 in [12] for a complete discussion on monomial orders, including the lexicographic order.

Let  $a$  and  $d$  be positive integers. Let  $[a]^{(d)} := (c_d, c_{d-1}, \dots, c_2, c_1)$  be the  $d$ -fractal coefficient of  $a$ , see Lemma 2.6 and Definition 2.8. We associate to  $a$  and  $d$  a monomial  $X_a^{(d)}$  of degree  $d$  in the variables  $X := \{x_1, \dots, x_n\}$  in such a way

$$X_a^{(d)} := x_{c_1} x_{c_2} \cdots x_{c_{d-1}} x_{c_d}.$$

Vice versa, a monomial  $T = x_{c_1} x_{c_2} \cdots x_{c_{d-1}} x_{c_d}$  of degree  $d$  in the variables in  $X$  identifies a  $d$ -upla  $(c_d, c_{d-1}, \dots, c_1)$  such that  $c_i \geq c_{i+1}$  for each  $i$ .

**Remark 3.1.** An immediate consequence of Lemma 2.10 is that  $X_a^{(d)} >_{lex} X_b^{(d)}$  if and only if  $a > b$ , with respect the lexicographic order induced by  $x_1 < x_2 < \cdots < x_n$ . Therefore, with respect the same order,  $X_a^{(d)}$  is the  $a$ -th greatest monomial (in the variables in  $X$ ) of degree  $d$ .

Let  $S := k[X] = k[x_1, \dots, x_n]$  be the standard graded polynomial ring.

We set

$$\mathcal{G}(X)_{\leq t}^{(d)} := \{X_a^{(d)} \mid a \leq t\}$$

and

$$\mathcal{G}(X)_{> t}^{(d)} := \{X_a^{(d)} \mid a > t\}.$$

$S_d$  is spanned by the monomials in  $\mathcal{G}(X)_{\leq t}^{(d)} \cup \mathcal{G}(X)_{> t}^{(d)}$ .

By Remark 3.1  $\mathcal{G}(X)_{> t}^{(d)}$  is a lex set of monomials of degree  $d$  with respect the degree lexicographic order  $x_1 < x_2 < \cdots < x_n$ .

Given  $T := \{\tau_0, \tau_1, \tau_2, \dots\}$  a coherent fractal growth, we set  $I(T)_d := \langle \mathcal{G}(X)_{> |\tau_d|}^{(d)} \rangle_k$  the  $k$ -vector space spanned by the monomial in  $\mathcal{G}(X)_{> |\tau_d|}^{(d)}$ . Then, by Theorem, 2.14 and Theorem 1.2, the following result holds.

**Proposition 3.2.**  $I(T) := \bigoplus_d I_d(T) \subseteq R$  is a lex segment ideal and  $H_{R/I}(d) = |\tau_d|$ .

Given a minimal free resolution of a lex segment ideal

$$0 \rightarrow \bigoplus_j S(-j)^{\beta_{p,p+j}} \rightarrow \dots \rightarrow \bigoplus_j S(-j)^{\beta_{1,1+j}} \rightarrow S \rightarrow S/I \rightarrow 0$$

the Betti numbers can be computed by the Eliahou-Kervaire formula, see [4]. See also [12] equation 7.7 of Section 7.

**Theorem 3.3** (Eliahou-Kervaire formula). *Let  $I$  be a lex segment ideal. For  $u \in \mathcal{G}(I)$ , a monomial minimal generator of  $I$ , let  $m(u)$  denotes the largest index  $j$  such that  $x_j$  divides  $u$ . Let  $m_{kj}$  be the number of monomials  $u \in \mathcal{G}(I)$  with  $m(u) = k$ . Then*

$$\beta_{i,i+j}(S/I) = \sum_{u \in \mathcal{G}(I)_j} \binom{m(u) - 1}{i} = \sum_{i=1}^n \binom{k-1}{i-1} m_{kj}.$$

This result can be written in terms of coherent fractal growth.

**Proposition 3.4.** *Given  $T := \{\tau_0, \tau_1, \tau_2, \dots\}$  a coherent fractal growth. Then*

$$\beta_{i,i+j}(S/I(T)) = \sum_{a=|\tau_j|+1}^{|\tau_{i-1}|} \binom{[a]_j^{(j)} - 1}{i} = \sum_{i=1}^n \binom{k-1}{i-1} w_{kj}$$

where  $w_{kj}$  is the number of occurrence of " $k$ " in  $\tau'_i$  with  $[\tau_{i-1}] = \tau_i \parallel \tau'_i$ .

*Proof.* It is an immediate consequence of Theorem 3.3 and Theorem 2.9.  $\square$

#### 4. THE HILBERT FUNCTION OF A BIGRADED ALGEBRA AND THE FRACTAL FUNCTIONS

Let  $k$  be an infinite field, and let  $R := k[x_1, \dots, x_n, y_1, \dots, y_m]$  be the polynomial ring in  $n+m$  indeterminates with the grading defined by  $\deg x_i = (1, 0)$  and  $\deg y_j = (0, 1)$ . Then  $R = \bigoplus_{(i,j) \in \mathbb{N}^2} R_{(i,j)}$  where  $R_{(i,j)}$  denotes the set of all homogeneous elements in  $R$  of degree  $(i, j)$ . Moreover,  $R_{(i,j)}$  is generated, as a  $k$ -vector space, by the monomials  $x_1^{i_1} \dots x_n^{i_n} y_1^{j_1} \dots y_m^{j_m}$  such that  $i_1 + \dots + i_n = i$  and  $j_1 + \dots + j_m = j$ . An ideal  $I \subseteq R$  is called a *bigraded* ideal if it is generated by homogeneous elements with respect to this grading. A bigraded algebra  $R/I$  is the quotient of  $R$  with a bigraded ideal  $I$ . The Hilbert function of a bigraded algebra  $R/I$  is defined such that  $H_{R/I} : \mathbb{N}^2 \rightarrow \mathbb{N}$  and  $H_{R/I}(i, j) := \dim_k (R/I)_{(i,j)} = \dim_k R_{(i,j)} - \dim_k I_{(i,j)}$  where  $I_{(i,j)} = I \cap R_{(i,j)}$  is the set of the bihomogeneous elements of degree  $(i, j)$  in  $I$ .

From now on, we will work with the degree lexicographical order on  $R$  induced by  $x_n > \dots > x_1 > y_m > \dots > y_1$ . With this ordering, we recall the definition of blex ideal, introduced and studied in [2]. We refer to [2] for all preliminaries and for further results on blex ideals.

**Definition 4.1** ([2], Definition 4.4). A set of monomials  $L \subseteq R_{(i,j)}$  is called *blex* if for every monomial  $uv \in L$ , where  $u \in R_{(i,0)}$  and  $v \in R_{(0,j)}$ , the following conditions are satisfied:

- if  $u' \in R_{(i,0)}$  and  $u' > u$ , then  $u'v \in L$ ;
- if  $v' \in R_{(0,j)}$  and  $v' > v$ , then  $uv' \in L$ .

A monomial ideal  $I \subseteq R$  is called a *blex ideal* if  $I_{(i,j)}$  is generated as  $k$ -vector space by a blex set of monomials, for every  $i, j \geq 0$ .

Blex ideals play a crucial role in the study of the Hilbert function of bigraded algebras.

**Theorem 4.2** ([2], Theorem 4.14). *Let  $J \subseteq R$  be a bigraded ideal. Then there exists a bilex ideal  $I$  such that  $H_{R/I} = H_{R/J}$ .*

In [5] was solved the question of characterize the Hilbert functions of bigraded algebras of  $k[x_1, x_2, y_1, y_2]$  by introducing the Ferrers functions. In this section we generalize these functions by introducing the fractal functions, see Definition 4.6. We prove, Theorem 4.12, that these classify the Hilbert functions of bigraded algebras.

We need some preparatory material. We denote by  $\mathcal{M}^{a \times b}(U)$  the set of all the matrices with size  $(a, b)$  -  $a$  rows and  $b$  columns - and entries in a set  $U \subseteq \mathbb{N}$ . Given a matrix  $M = (m_{ij}) \in \mathcal{M}^{a \times b}(U)$  we denote by

$$\sum M := \sum_{i \leq a} \sum_{j \leq b} m_{i,j}.$$

Next definition introduces the objects we need in this section. The  $\leq$  symbol denote the partial order on  $\mathbb{N}^2$  given by componentwise comparison.

**Definition 4.3.** A *Ferrers matrix* of size  $(a, b)$  is a matrix  $M = (m_{ij}) \in \mathcal{M}(\{0, 1\})^{a \times b}$  such that

$$\text{if } m_{ij} = 1 \text{ then } m_{i'j'} = 1 \text{ for any } (i', j') \leq (i, j).$$

We set by  $\mathcal{F}^{a \times b}$  the family of all the Ferrers matrices of size  $(a, b)$ .

In the following definition we introduce expansions of a matrix.

**Definition 4.4.** Let  $M \in \mathcal{M}(U)^{a \times b}$  be a matrix of size  $(a, b)$  and let  $\mathbf{v} := (v_1, \dots, v_a) \in \mathbb{N}^a$  and  $\mathbf{w} := (w_1, \dots, w_b) \in \mathbb{N}^b$  be vectors of non negative integers. We denote by  $M^{(\mathbf{v}, \bullet)}$  an element in  $M(U)^{\sum \mathbf{v} \times b}$  constructed by

$$\text{repeating the } i\text{-th row of } M \text{ } v_i \text{ times, for } i = 1, \dots, a.$$

We denote by  $M^{(\bullet, \mathbf{w})}$  an element in  $M(U)^{a \times \sum \mathbf{w}}$  constructed by

$$\text{repeating the } j\text{-th column of } M \text{ } w_j \text{ times, for } j = 1, \dots, b.$$

**Remark 4.5.** The expansions of a Ferrers matrix are also Ferrers matrices. Take, for instance,

$$M = \begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 0 \\ 1 & 1 & 0 \\ 1 & 0 & 0 \end{pmatrix} \in \mathcal{F}^{(4,3)}.$$

Set  $\mathbf{v} := (2, 1, 0, 3)$  and  $\mathbf{w} := (3, 1, 3)$ . Then

$$M^{(\mathbf{v}, \bullet)} = \begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 0 \\ 1 & 0 & 0 \\ 1 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix} \in \mathcal{F}^{(6,3)} \quad M^{(\bullet, \mathbf{w})} = \begin{pmatrix} 1 & 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 & 0 & 0 & 0 \\ 1 & 1 & 1 & 1 & 0 & 0 & 0 \\ 1 & 1 & 1 & 0 & 0 & 0 & 0 \end{pmatrix} \in \mathcal{F}^{(4,7)}.$$

Given  $M, N \in \mathcal{F}^{a \times b}$  we say that  $M \leq N$  if and only if  $m_{ij} \leq n_{ij}$  for any  $i, j$ .

We are ready to introduce the fractal functions.

**Definition 4.6.** Let  $H : \mathbb{N}^2 \rightarrow \mathbb{N}$  be a numerical function. We say that  $H$  is a *fractal function* if  $H(0, 0) = 1$  and, for any  $(i, j) \in \mathbb{N}^2$ , there exists a matrix of  $M_{ij} \in \mathcal{F}^{\binom{i-1+n}{n-1} \times \binom{j-1+m}{m-1}}$  with  $\sum M_{ij} = H(i, j)$  and such that all the matrices satisfy the condition

$$\begin{cases} M_{ij} \leq M_{i-1, j}^{\langle [n]^{i-1}, \bullet \rangle} & \text{if } i > 0 \\ M_{ij} \leq M_{i, j-1}^{\langle \bullet, [m]^{j-1} \rangle} & \text{if } j > 0 \end{cases}$$

**Remark 4.7.** Let  $H : \mathbb{N}^2 \rightarrow \mathbb{N}$  be the numerical function

$$H(i, j) = \binom{i-1+n}{n-1} \binom{j-1+m}{m-1}.$$

For any  $i, j \in \mathbb{N}$ , there is only one element in  $M_{ij} \in \mathcal{F}^{\binom{i-1+n}{n-1} \times \binom{j-1+m}{m-1}}$  satisfying the condition in Definition 4.6 that is the matrix with all "1" entries. Therefore  $H$  is a fractal function.

**Remark 4.8.** If  $n = m = 2$ , the definition of fractal functions agrees with Definition 3.3. in [5]. Indeed it is enough to write each partition  $\alpha_{ij} = (a_1, a_2, \dots)$  as a matrix  $M_{ij} = (m_{hk}) \in \mathcal{F}^{(i+1) \times (j+1)}$  where  $m_{hk} = 1$  iff  $k \leq a_k$  otherwise  $m_{hk} = 0$ . In this case the expansions are given by the elements in  $\Phi(2) := ((2), (1, 2), (1, 1, 2), (1, 1, 1, 2), \dots)$ .

In general the matrices  $M_{ij}$  are not uniquely determined by the conditions of Definition 4.6, even in "small" cases.

**Example 4.9.** Set  $n = m = 2$  and let  $H : \mathbb{N}^2 \rightarrow \mathbb{N}$  be the numerical function

$$H := \begin{array}{c|cccc} & 0 & 1 & 2 & 3 & \dots \\ \hline 0 & 1 & 2 & 3 & 0 & \dots \\ 1 & 2 & 4 & 3 & 0 & \dots \\ 2 & 3 & 3 & 3 & 0 & \dots \\ 3 & 0 & 0 & 0 & 0 & \dots \end{array}$$

The only possibility for  $M_{11}$  is

$$M_{11} = \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix} \in \mathcal{F}^{(2,2)}.$$

Thus, we have no restriction on  $M_{21}$  and  $M_{12}$  but the number of non zero entries.

$$M_{12} \in \left\{ \begin{pmatrix} 1 & 1 & 1 \\ 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 1 & 1 & 0 \\ 1 & 0 & 0 \end{pmatrix} \right\} \subseteq \mathcal{F}^{(2,3)}.$$

$$M_{21} \in \left\{ \begin{pmatrix} 1 & 0 \\ 1 & 0 \\ 1 & 0 \end{pmatrix}, \begin{pmatrix} 1 & 1 \\ 1 & 0 \\ 0 & 0 \end{pmatrix} \right\} \subseteq \mathcal{F}^{(3,2)}.$$

Now we check if all these choices are allowed. We have to look at the conditions on  $M_{22}$ . We have  $M_{22} \leq M_{1,2}^{((1,2), \bullet)}$  and  $M_{22} \leq M_{2,1}^{(\bullet, (1,2))}$ . In the next table we collect all the possibilities for  $M_{22}$  depending on  $M_{1,2}$  and  $M_{2,1}$ .

$$\begin{array}{c|c|c} M_{22} \leq & \begin{pmatrix} 1 & 1 & 1 \\ 0 & 0 & 0 \end{pmatrix} & \begin{pmatrix} 1 & 1 & 0 \\ 1 & 0 & 0 \end{pmatrix} \\ \hline \begin{pmatrix} 1 & 0 \\ 1 & 0 \\ 1 & 0 \end{pmatrix} & \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} & \begin{pmatrix} 1 & 0 & 0 \\ 1 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix} \\ \hline \begin{pmatrix} 1 & 1 \\ 1 & 0 \\ 0 & 0 \end{pmatrix} & \begin{pmatrix} 1 & 1 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} & \begin{pmatrix} 1 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \end{array}$$

Since  $H(2, 2) = 3$ , we note that

- only one of the above four matrices in the table must be excluded;
- in the other three cases  $M_{22}$  is uniquely determined by  $M_{12}$  and  $M_{21}$ .

Therefore,  $H$  is a fractal function and three different set of matrices satisfy the conditions in Definition 4.6.

In the following we denote by  $X := \{x_1, \dots, x_n\}$  and  $Y := \{y_1, \dots, y_m\}$  the set of the variables of degree  $(1, 0)$  and  $(0, 1)$  respectively.

Next lemma is useful for our purpose. It is an immediate consequence of Lemma 2.13.

**Lemma 4.10.**  $x_1 \cdot X_a^{(d)} = X_{a^{(d)}}^{(d+1)}$ .

To shorten the notation we set  $\alpha_i := \binom{n+i-1}{n-1}$  and  $\beta_j := \binom{m+j-1}{m-1}$ . In order to relate fractal functions and Hilbert functions of bigraded algebras we need to introduce a correspondence between Ferrers matrices and monomials.

Let  $M = (m_{ab}) \in \mathcal{F}^{\alpha_i \times \beta_j}$ . We denote by  $\lambda(M)_{(ij)}$  the set of the monomials

$$\lambda(M) := \{X_a^{(i)} \cdot Y_b^{(j)} \mid m_{ab} = 0\}.$$

Let  $L \subseteq R_{(i,j)}$  be a bilex set of monomials of bidegree  $(i, j)$ . We denote by  $\mu(L) \in \mathcal{M}(\{0, 1\})^{\alpha_i \times \beta_j}$  the matrix  $(m_{ab})$  such that  $m_{ab} = 1$  iff  $X_a^{(i)} \cdot Y_b^{(j)} \notin L$  otherwise  $m_{ab} = 0$ .

**Proposition 4.11.** *There is a one to one correspondence between bilex sets of monomials of degree  $(i, j)$  and elements in  $\mathcal{F}^{\alpha_i \times \beta_j}$ .*

*Proof.* Let  $M = (m_{ab}) \in \mathcal{F}^{\alpha_i \times \beta_j}$ , we claim that

$$\lambda(M) \text{ is a bilex set of monomials of bidegree } (i, j).$$

To prove the claim we use Lemma 2.10 and Remark 3.1. Let  $X_a^{(i)} \cdot Y_b^{(j)}$  be an element in  $\lambda(M)$  and  $X_u^{(i)} > X_a^{(i)}$ . Since  $(u, b) > (a, b)$  and  $m_{ab} = 0$  we get  $m_{ub} = 0$  i.e.  $X_u^{(i)} \cdot Y_b^{(j)} \in \lambda(M)$ . In a similar way it follows that  $X_a^{(i)} \cdot Y_v^{(j)} \in \lambda(M)$  for  $v > b$ .

Let  $L \subseteq R_{(i,j)}$  be a bilex set of monomials of bidegree  $(i, j)$ . We claim that

$$\mu(L) \in \mathcal{F}^{\alpha_i \times \beta_j}.$$

The claim follows by using Lemma 2.10 and Remark 3.1. Indeed, say  $m_{ab} = 0$  for an entry of  $\mu(L)$ . This implies  $X_a^{(i)} \cdot Y_b^{(j)} \in L$ . Thus, for  $u > a$ , we have  $X_u^{(i)} \cdot Y_b^{(j)} \in L$  i.e.  $m_{ub} = 0$ . Analogously, we see that  $m_{av} = 0$  for  $v > b$ .  $\square$

We are ready to prove the main result of this paper.

**Theorem 4.12.** *Let  $H : \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$  be a numerical function. Then the following are equivalent*

- 1)  $H$  is a fractal function;
- 2)  $H = H_{R/I}$  for some bilex ideal  $I \subseteq R = k[x_1, \dots, x_n, y_1, \dots, y_m]$ .

*Proof.* (1)  $\Rightarrow$  (2) Let  $H$  be a fractal function. For each  $(i, j) \in \mathbb{N}^2$ , let  $I_{(i,j)}$  be the  $k$ -vector space spanned by the elements in  $\lambda(M_{ij})$ . We claim that  $I := \bigoplus_{(i,j) \in \mathbb{N}^2} I_{(i,j)}$  is an ideal of  $R$ . To prove the claim, it is enough to show that if  $X_a^{(i)} \cdot Y_b^{(j)} \in I_{(i,j)}$  then  $x_u \cdot X_a^{(i)} \cdot Y_b^{(j)} \in I_{(i+1,j)}$  for any  $x_u \in X$  and  $y_v \cdot X_a^{(i)} \cdot Y_b^{(j)} \in I_{(i,j+1)}$  for any  $y_v \in Y$ . We have, see Lemma 4.10,

$$x_u \cdot X_a^{(i)} \cdot Y_b^{(j)} \geq x_1 \cdot X_a^{(i)} \cdot Y_b^{(j)} = X_{a^{(i)}}^{(i+1)} \cdot Y_b^{(j)}.$$

Then, by Definition 4.6 and Theorem 2.14, the entry  $(a^{(i)}, b)$  in the matrix  $M_{i+1,j}$  is 0. Thus  $X_{a^{(i)}}^{(i+1)} \cdot Y_b^{(j)} \in I_{(i+1,j)}$  and furthermore  $x_u \cdot X_a^{(i)} \cdot Y_b^{(j)} \in I_{(i+1,j)}$ . In a similar way it follows that  $y_v \cdot X_a^{(i)} \cdot Y_b^{(j)} \in I_{(i,j+1)}$ .

(2)  $\Rightarrow$  (1) Let  $I \subseteq R$  be a bilex ideal such that  $H_{R/I} = H$ . Set  $M_{ij} := \mu(I_{ij})$ , we claim that the  $M_{ij}$ s satisfy the condition in Definition 4.6. By Theorem 2.14 it is enough to show that if  $M_{ij}(a, b) = 0$  (the entry  $(a, b)$  in  $M_{ij}$  is 0), then also

$M_{i+1,j}(a^{(i)}, b) = 0$  (the entry  $(a^{(i)}, b)$  in  $M_{i+1,j}$  is 0). Set  $J := (X_u^{(h)} \mid M_{hj}(u, b) = 0)$ , then the claim is an immediate consequence of the fact that  $J$  is a lex ideal of  $k[X]$ .  $\square$

**Remark 4.13.** Note that since the  $M_{ij}$  are not uniquely determined, as shown in Example 4.9, then there could be several blex ideals having the same Hilbert function.

The following question is motivated by the arguments in Section 3.

**Question 4.14.** Can the bigraded Betti numbers of a blex ideal  $I = \bigoplus I_{(i,j)}$  be computed from the matrices  $\mu(I_{(i,j)})$ ?

#### REFERENCES

- [1] Abbott J, Bigatti AM, Robbiano L. *CoCoA: a system for doing Computations in Commutative Algebra*. Available at <http://cocoa.dima.unige.it>
- [2] Aramova A, Crona K, De Negri E. *Bigeneric initial ideals, diagonal subalgebras and bigraded Hilbert functions*. Journal of Pure and Applied Algebra. **150**(3):215–35 (2000).
- [3] Bruns W, Herzog HJ. *Cohen-macaulay rings*. Cambridge University Press; 1998.
- [4] Eliahou S., Kervaire M., *Minimal resolutions of some monomial ideals*. J. Algebra **129**:1–25 (1990).
- [5] Favacchio G. *The Hilbert Function of Bigraded Algebras in  $k[\mathbb{P}^1 \times \mathbb{P}^1]$* . To appear in Journal of Commutative Algebra. (2019). <https://projecteuclid.org/euclid.jca/1523433691>
- [6] Favacchio G., Guardo E. *The minimal free resolution of fat almost complete intersections in  $\mathbb{P}^1 \times \mathbb{P}^1$* , Canadian Journal of Mathematics 69(2017), no. 6, 1274-1291.
- [7] Favacchio G., Guardo E. *On the Betti numbers of three fat points in  $\mathbb{P}^1 \times \mathbb{P}^1$*  Journal of the Korean Mathematical Society 56 (3), 751-766
- [8] Favacchio G., Guardo E., Migliore J. *On the arithmetically Cohen-Macaulay property for sets of points in multiprojective spaces*. Proc. Amer. Math. Soc. 146 (2018), pp. 2811-2825.
- [9] Favacchio G., Guardo E., Picone B *Special arrangements of lines: Codimension 2 ACM varieties in  $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$*  Journal of Algebra and Its Applications Vol. 18, No. 04, 1950073 (2019) 7
- [10] G. Favacchio, J. Migliore *Multiprojective spaces and the arithmetically Cohen-Macaulay property*. Mathematical Proceedings of the Cambridge Philosophical Society Volume 166, Issue 3 May 2019 , pp. 583-597
- [11] Guardo E, Van Tuyl A. *Arithmetically Cohen-Macaulay sets of points in  $\mathbb{P}^1 \times \mathbb{P}^1$* . Springer-Briefs in Mathematics, Springer 2015.
- [12] Herzog J, Hibi T. *Monomial ideals*. Springer. 2011.
- [13] Kimberling C. *Fractal sequences and interspersions*. Ars Combinatoria. **45**:157-68 (1997).
- [14] Macaulay FS. *Some properties of enumeration in the theory of modular systems*. Proceedings of the London Mathematical Society. **2**(1):531-55(1927).
- [15] Peeva I, Stillman M. *Open problems on syzygies and Hilbert functions*. Journal of Commutative Algebra. **1**(1):159-95 (2009).
- [16] Stanley RP. *Hilbert functions of graded algebras*. Advances in Mathematics. **28**(1):57-83 (1978).
- [17] Van Tuyl A. *The border of the Hilbert function of a set of points in  $\mathbb{P}^{n_1} \times \mathbb{P}^{n_k}$* . Journal of Pure and Applied Algebra. 2002 Dec 24;176(2-3):223-47.
- [18] Van Tuyl A. *The Hilbert functions of ACM sets of points in  $\mathbb{P}^{n_1} \times \mathbb{P}^{n_k}$* . Journal of Algebra. 2003 Jun 15;264(2):420-41.

DIPARTIMENTO DI MATEMATICA E INFORMATICA, VIALE A. DORIA, 6 - 95100 - CATANIA, ITALY

E-mail address: [favacchio@dmi.unict.it](mailto:favacchio@dmi.unict.it)

URL: [www.dmi.unict.it/~gfavacchio](http://www.dmi.unict.it/~gfavacchio)