

Effect of the good solvent nature in flash nano-precipitation via population balance modelling and computational fluid dynamics coupling approach

*Original*

Effect of the good solvent nature in flash nano-precipitation via population balance modelling and computational fluid dynamics coupling approach / Ferrari, M.; Lavino, A. D.; Barresi, A.; Marchisio, D.; Matar, O. K.. - ELETTRONICO. - (2021). ((Intervento presentato al convegno 74th Annual Meeting of the APS Division of Fluid Dynamics tenutosi a Phoenix (AZ), U.S.A. nel 21-23 November 2021).

*Availability:*

This version is available at: 11583/2953972 since: 2022-01-28T12:17:51Z

*Publisher:*

APS

*Published*

DOI:

*Terms of use:*

openAccess

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

*Publisher copyright*

(Article begins on next page)

# 74th Annual Meeting of the APS Division of Fluid Dynamics Sunday–Tuesday, November 21–23, 2021; Phoenix Convention Center, Phoenix, Arizona

## Session Q08: Particle-laden Flows: Simulations

8:00 AM–9:31 AM, Tuesday, November 23, 2021  
Room: North 123

Chair: JESSE CAPECELATRO, University of Michigan

### **Abstract: Q08.00006 :**

## **Effect of the good solvent nature in flash nano-precipitation via population balance modelling and computational fluid dynamics coupling approach**

9:05 AM–9:18 AM

### **Presenter:**

Marco Ferrari

(Department of Applied Science and Technology, Institute of Chemical Engineering, Politecnico di Torino, 10129 Torino, Italy)

### **Authors:**

Marco Ferrari

(Department of Applied Science and Technology, Institute of Chemical Engineering, Politecnico di Torino, 10129 Torino, Italy)

Alessio D Lavino

(Department of Chemical Engineering, Imperial College London, South Kensington Campus, London SW7 2AZ, UK)

Antonello Barresi

(Department of Applied Science and Technology, Institute of Chemical Engineering, Politecnico di Torino, 10129 Torino, Italy)

Daniele Marchisio

(Department of Applied Science and Technology, Institute of Chemical Engineering, Politecnico di Torino, 10129 Torino, Italy)

Omar K Matar

(Department of Chemical Engineering, Imperial College London, South Kensington Campus, London SW7 2AZ, UK)

The effect of the good solvent nature on polymer nano-particles (NP) formation in flash nano-precipitation is here investigated through a combined population balance model-computational fluid dynamics approach (PBM-CFD). Four good solvents are considered: acetone, acetonitrile, tetrahydrofuran and tert-butanol and the different resulting mean NP size is predicted in terms of mean radius of gyration via the Flory law of real polymers. Good solvents effects are here modelled in terms of solute–solvent interactions, using the Flory–Huggins theory and the Hansen solubility parameters. In this way, kinetics and thermodynamics are intertwined in a unique modelling tool. Our results show that the proposed methodology is able to predict the role played by the different good solvents, analysing single factors at the time. More specifically, the dynamics of mixing is decoupled from the dynamics of aggregation achieving a deeper insight into the fundamental fluid properties which affect the final NP size, pointing out the main mechanisms involved and showing a good agreement with experimental data.