

RESEARCH PAPER

## Effects of insect cadavers infected by *Heterorhabditis bacteriophora* and *Steinernema diaprepesi* on *Meloidogyne incognita* parasitism in pepper and summer squash plants

Eleodoro E. Del Valle<sup>1,3</sup>, Paola Lax<sup>2,3</sup>, Juan Rondán Dueñas<sup>4</sup>, and Marcelo E. Doucet<sup>2,3</sup>

<sup>1</sup>Facultad de Ciencias Agrarias, Universidad Nacional del Litoral, Kreder 2805, C.P. 3080HOF, Esperanza, Santa Fe, Argentina.

<sup>2</sup>Laboratorio de Nematología, Centro de Zoología Aplicada, Jardín Zoológico, Rondeau 798, C.P. X5000AVP, Córdoba, Argentina.

<sup>3</sup>Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Av. Rivadavia 1917, C.P. C1033AAJ, Ciudad Autónoma de Buenos Aires, Argentina.

<sup>4</sup>Laboratorio de Biología Molecular, Pabellón CEPROCOR, Santa María de Punilla, X5164 Córdoba, Argentina.

### Abstract

**E. E. Del Valle, P. Lax, J. Rondán Dueñas, and M. E. Doucet. 2013. Effects of insect cadavers infected by *Heterorhabditis bacteriophora* and *Steinernema diaprepesi* on *Meloidogyne incognita* parasitism in pepper and summer squash plants. Cien. Inv. Agr. 40(1):109-118.**

The effects of insect cadavers infected with three isolates of *Heterorhabditis bacteriophora* and one isolate of *Steinernema diaprepesi* on a population of *Meloidogyne incognita* in pepper (*Capsicum annuum*) and summer squash (*Cucurbita maxima*) were evaluated in greenhouse experiments carried out in Santa Fe (Argentina). Insect cadavers were obtained for the experiments from last instar larvae of *Galleria mellonella* and *Tenebrio molitor* that had been infected with entomopathogenic nematodes. Two six-day-old insect cadavers per pot were placed below the soil surface, and the soil was inoculated with 100 second-stage juveniles of *M. incognita*. Sixty days after inoculation, the following parameters were recorded for each plant: number of leaves; dry weight of aerial parts; numbers of galls, egg masses and eggs; and numbers of galls, egg masses and eggs g<sup>-1</sup> of root fresh matter. In pepper, the only variable affected by the infected cadavers with respect to control was the number of eggs in the treatment involving *T. molitor* cadavers infected with the *H. bacteriophora* isolate Rama Caída. In summer squash, several treatments using infected cadavers resulted in a decrease in the numbers of galls and egg masses. Only the treatment involving *G. mellonella* cadavers infected with the *H. bacteriophora* isolate Rama Caída proved to be efficient in reducing the number of *M. incognita* eggs. Our results indicated that the application of insect cadavers infected with the entomopathogenic nematodes studied might reduce *M. incognita* damage in pepper and summer squash plants.

**Key words:** Biological control, entomopathogenic nematodes, insect host cadavers.

### Introduction

Root-knot nematodes (*Meloidogyne* spp.) are plant parasites that cause serious losses to economically

important crops (Koenning *et al.*, 1999). The major strategy for the control of these organisms in horticultural crops in recent decades has been the use of chemical products, especially methyl bromide (Halbrendt and LaMondia, 2004). However, the use of methyl bromide was banned

Received July 7, 2012. Accepted January 29, 2013.

Corresponding author: edelvalle@fca.unl.edu.ar

and/or restricted in several countries due to the environmental damage that it produces and its impact on human health, thus generating the need for new, environmentally friendly alternatives (Schafer, 1999). Accordingly, treatment with entomopathogenic nematodes (EPNs) is regarded as an alternative for the management of plant-parasitic nematodes (Grewal *et al.*, 2005).

Insect-parasitic (entomopathogenic) nematodes belong to the genera *Heterorhabditis* and *Steinernema* (Nematoda: Heterorhabditidae and Steinernematidae). Infective juveniles (IJs) search for a host, penetrate it through natural openings (mouth, anus or spiracles) or through the cuticle (Poinar, 1990). Once inside the host hemocoel, IJs release a symbiotic bacterium from their digestive tract. *Steinernema* species are associated with bacteria of the genus *Xenorhabdus*, whereas *Heterorhabditis* species are associated with bacteria of the genus *Photorhabdus* (Boemare *et al.*, 1993; Thomas and Poinar, 1979). Infected insects usually die of septicemia approximately 48 h after infection (Akhurst and Boemare, 1990; Burman, 1982). Nematodes feed on the bacteria and digested tissues and complete one to three generations within the host until IJs emerge into the soil (Poinar, 1990).

The interaction between EPNs and plant-parasitic nematodes has been documented previously (Lewis and Grewal, 2005). Previous studies using EPNs did not find any definitive results in terms of their action against nematodes of the genus *Meloidogyne* (Fallon *et al.*, 2002; Molina *et al.*, 2007; Shapiro-Ilan *et al.*, 2006a). However, several cases of suppressive action on *M. incognita* have been reported. Lewis *et al.* (2001) demonstrated that the use of IJs of *Steinernema feltiae* (Filipjev) significantly reduced the number of galls and eggs in tomato plants, even affecting egg hatching. In experiments conducted in growth chambers and a greenhouse, Pérez and Lewis (2002) found that IJs of *H. bacteriophora*, *Steinernema riobrave*

(Cabanillas, Poinar and Raulston) and *S. feltiae* reduced parasitism of *M. incognita* in tomato plants. Moreover, *M. incognita* was suppressed by IJs of *Steinernema glaseri* (Steiner) in an experiment evaluating application doses under greenhouse conditions.

Most of the works conducted until now have been based on the application of IJs of EPNs in aqueous suspension (Jagdale *et al.*, 2002; Molina *et al.*, 2007; Perry *et al.*, 1998; Smitley *et al.*, 1992). The use of this method has some disadvantages, such as IJs formulation, decrease of infectivity, survival of IJs during storage, transportation difficulties, and the need for adequate irrigation equipment (Grewal, 2002).

The use of EPN-infected insect cadavers provides an option for the control of root-knot nematodes because they lack some of the disadvantages of aqueous suspensions (Bruck *et al.*, 2005; Creighton and Fassuliotis, 1985; Shapiro-Ilan *et al.*, 2010; 2012). Infected insect cadavers are placed below the soil surface, and IJs emerge in the same environment where the target organisms are found. In laboratory assays, the use of this mode of EPN application has been shown to result in greater infectivity to insect hosts and increased survival and dispersal capacity than aqueous suspensions (Pérez *et al.*, 2003; Shapiro and Glazer, 1996; Shapiro-Ilan and Lewis, 1999). Similar results were obtained in greenhouse experiments. Infective juveniles that had emerged from the cadavers of *Tenebrio molitor* infected with *Heterorhabditis indica* (Poinar, Karunaka and David) Hom1 caused a greater decrease in the survival of *Diaprepes abbreviatus* (Linnaeus) than IJs applied in aqueous suspensions (Shapiro-Ilan *et al.*, 2003). In the same work, the authors found similar results regarding the control of *Otiorynchus sulcatus* (Fabricius) 7 days after the application of *T. molitor* (Linnaeus) cadavers infected with *Heterorhabditis bacteriophora* (Poinar) Oswego.

Suppression of plant-parasitic nematodes using infected cadavers has been reported by Jagdale and Grewal (2008), who studied the efficacy of preventive and curative applications of infective cadavers of *S. carpocapsae* (Weiser) for the management of *Aphelenchoides fragariae* (Ritzema Bos) in plants of *Hosta* spp.; the authors attributed the observed suppression to toxins produced by the entomopathogens and/or their symbiotic bacteria. The secondary metabolites 3,5-dihydroxy-4-isopropylstilbene (DST) and indole, which are obtained from filtrates of *P. luminescens*, exhibit nematocidal properties. Second-stage juveniles of *M. incognita* were affected by indole, and both components inhibited egg hatching in that species (Hu *et al.*, 1999). Hatching inhibition is most likely related to the fact that anion transporters act as target sites for blockers and DST in nematodes, thereby altering physiological processes through voltage-gated chloride channels, resulting in nematode and egg mortality (Boina *et al.*, 2008). In addition, ammonia, which is produced by symbiotic bacteria in the host, exhibits nematocidal activity (Grewal *et al.*, 1999).

Variable and/or confusing results about the EPNs-*Meloidogyne* spp. interaction have also been reported in studies involving the application of IJs (Fallon *et al.*, 2002; 2004; Molina *et al.*, 2007; Pérez and Lewis, 2004). Hence, knowledge of the suppressive effect of local EPN isolates applied as insect cadavers on *M. incognita* is of great importance. The aim of the present work was to determine the effect of insect cadavers infected with three local isolates of *H. bacteriophora* and one of *Steinernema diaprepesi* (Nguyen and Duncan) on a population of *M. incognita* in pepper (*Capsicum annuum* Linnaeus) and summer squash (*Cucurbita maxima* Duch.). We hypothesize that the application of cadavers infected with local isolates of *H. bacteriophora* and *S. diaprepesi* would reduce *M. incognita* parasitism in pepper and summer squash.

## Materials and methods

### *Nematodes and infected cadavers*

Three Argentinean isolates of *H. bacteriophora* Rama Caída (Rama Caída, Mendoza), Jn (Recreo, Santa Fe) and Mo (Ángel Gallardo, Santa Fe) and one isolate of *S. diaprepesi* (Santa Rosa de Calchines, Santa Fe) were used as EPNs. Nematodes were multiplied on larvae of the greater wax moth, *Galleria mellonella* (Lepidoptera: Pyralidae) at 25 °C, following the procedure described by Woodring and Kaya (1988). Third-stage juveniles were harvested from modified White traps (Kaya and Stock, 1997) and stored at 16 °C until use. *G. mellonella* larvae were reared on the diet formulated by Eischen and Dietz (1990), and larvae of *T. molitor* (Coleoptera: Tenebrionidae) were fed with wheat bran.

Insect hosts were infected in Petri dishes (90 mm in diameter) lined with filter paper to obtain insect cadavers. Last-instar *G. mellonella* or *T. molitor* larvae were exposed to 100 IJs of each isolate. The Petri dishes were incubated at 25 °C for 4 days, and infected cadavers were subsequently transferred to new Petri dishes lined with dry filter paper for a further two days of incubation to allow the development of typical signs of EPN infection (the cadavers become flaccid and exhibit a color change) (Woodring and Kaya, 1988).

The population of *M. incognita* was obtained from pepper and summer squash maintained under greenhouse conditions and inoculated with nematodes from the locality of Santa Rosa de Calchines. These plant species were chosen because they are severely parasitized by *M. incognita* and of regional economic importance. Second-stage juveniles were obtained by incubation of egg masses of *M. incognita* in a wet chamber under laboratory conditions.

### Greenhouse experiments

One week after germination, seedlings of pepper, cultivar K. Resistant Giant (Nasco Seed Co.), and of summer squash (*Cucurbita maxima* var. zapallito (Carr.) Millán), cultivar Nacional (CAPS), were transplanted into pots (10 cm in diameter) containing 700 cm<sup>3</sup> of sterile soil (1.1% organic matter; pH: 7.1; clay: 6%, silt: 8% and sand: 87%). Two days after transplanting, EPN-infected cadavers were applied. The treatments were as follows: (i) control; (ii) carbofuran 25 mg L<sup>-1</sup> as a chemical control; (iii) *T. molitor* cadavers infected with *H. bacteriophora* Rama Caída (HbRCT); (iv) *T. molitor* cadavers infected with *H. bacteriophora* Jn (HbJnT); (v) *T. molitor* cadavers infected with *H. bacteriophora* Mo (HbMoT); (vi) *T. molitor* cadavers infected with *S. diaprepesi* (SdT); (vii) *G. mellonella* cadavers infected with *H. bacteriophora* Rama Caída (HbRCG); (viii) *G. mellonella* cadavers infected with *H. bacteriophora* Jn (HbJnG); (ix) *G. mellonella* cadavers infected with *H. bacteriophora* Mo (HbMoG); and (x) *G. mellonella* cadavers infected with *S. diaprepesi* (SdG).

In treatments involving the use of infected cadavers, two cadavers were added per pot. The cadavers were buried 2 cm below the soil surface and 2.5 cm from the stem, diametrically opposite each other. The insect cadavers used were infected 6 days before application (Del Valle *et al.*, 2008). Five days after the introduction of the cadavers into the pots, the plants were inoculated by adding an aqueous suspension containing 100 J2s of *M. incognita* with a micropipette. The suspension was poured into holes made in the soil around the plant stem. The plants were watered as needed. During the experimental period, the mean temperature was maintained at 18.1 °C to ensure normal development and multiplication of nematodes inside the hosts (Grewal *et al.*, 2006).

The results were evaluated 60 days after *M. incognita* inoculation. The following variables were recorded for each plant: number of leaves,

dry weight of aerial parts, number of galls, number of egg masses, number of eggs, number of galls/g of root fresh matter, number of egg masses g<sup>-1</sup> of root fresh matter, and number of eggs g<sup>-1</sup> of root fresh matter. Total numbers of galls, egg masses and eggs were counted under a stereoscopic microscope. The eggs were removed from the roots using the procedure described by Hussey and Barker (1973).

The experiment was conducted using a completely randomized design with six replications. The entire experiment was performed twice. No significant interactions between trials ( $\alpha=0.05$ ) were detected; therefore, pooled data were subjected to analysis of variance (ANOVA). The averages of experiments are shown in Tables 1 and 2. Means were compared among treatments using Tukey's HDS test at  $P\leq 0.05$  probability.

## Results

### Experiments using pepper plants

The results observed in pepper plants are presented in Table 1. The use of cadavers infected with EPN isolates did not show significant differences from control treatment with respect to the number of leaves per plant or the dry weight of aerial parts ( $F_{9,110}=4.54$ ,  $P<0.0001$  and  $F_{9,110}=4.06$ ,  $P<0.0002$ , respectively). The treatment with carbofuran was the only treatment to exhibit statistically significant differences from the control with respect to the number of galls. However, treatments HbRCT, HbRCG, HbMoG and SdG did not differ from the chemical treatment ( $F_{9,110}=5.97$ ,  $P\leq 0.0001$ ). Carbofuran application caused a significant reduction in the number of egg masses in pepper plants, without showing significant differences from HbRCT, HbRCG, HbMoG or SdG ( $F_{9,110}=5.47$ ;  $P\leq 0.0001$ ). The number of *M. incognita* eggs produced by nematodes in the root systems of plants treated with carbofuran and HbRCT was significantly lower than that observed for the control ( $F_{9,110}=7.17$ ,  $P\leq 0.0001$ ).

### Experiments using summer squash

The treatments evaluated did not significantly affect the number of leaves per plant ( $F_{9,110} = 0.56$ ,  $P \leq 0.8242$ ). Plants treated with HbRCT showed the highest value of dry weight of aerial parts among treatments ( $F_{9,110} = 3.82$ ,  $P \leq 0.0003$ ). Infected cadavers significantly decreased the numbers of galls and egg masses ( $F_{9,110} = 3.88$ ,  $P \leq 0.0003$  and  $F_{9,110} = 3.88$ ,  $P \leq 0.0003$ , respectively). HbJnG did not affect the variables mentioned, and HbRCT did not differ from the control in terms of the number of egg masses per plant. HbRCG was the only treatment that differed from control in reducing the total number of eggs per plant ( $F_{9,110} = 3.89$ ,  $P \leq 0.0003$ ). There were no statistically significant differences among treatments with respect to the number of galls/g of root fresh weight and the number of egg masses/g of root fresh weight ( $F_{9,110} = 4.45$ ,  $P \leq 0.0001$  and  $F_{9,110} = 3.09$ ,  $P \leq 0.0024$ , respectively). The negative control was statistically similar to treatments HbRCT, HbJnT, HbMoT, SdT and HbRCG in terms of the number of eggs/g of root fresh weight ( $F_{9,110} = 6.49$ ,  $P \leq 0.0001$ ). The results of this experiment are presented in Table 2.

### Discussion

The use of EPN-infected cadavers of *G. mellonella* and *T. molitor* had different effects on pepper and summer squash. In pepper, the only variable affected by infected cadavers with respect to the control was the number of eggs in the treatment involving *T. molitor* cadavers infected with *H. bacteriophora* Rama Caída. In summer squash, several treatments using infected cadavers reduced the number of galls and egg masses. Only the treatment including *G. mellonella* cadavers infected with *H. bacteriophora* Rama Caída proved efficient in reducing the number of *M. incognita* eggs.

The lack of efficacy of EPN aqueous suspensions for the control of *M. incognita* has also been reported. For example, *Heterorhabditis megidis*

(Poinar, Jackson and Klein) did not produce any effects on tomato, even at high doses (Pérez and Lewis, 2004). In the only work that studied plants of the family Cucurbitaceae, the species *S. riobrave* and *H. bacteriophora* were not effective in the control of zucchini squash (*Cucurbita pepo* Linnaeus) (Riegel *et al.*, 1998). The results presented here demonstrate that the action of EPNs may reduce damage caused by *M. incognita* in summer squash plants.

Determining the activity of metabolites and toxins produced by the symbiotic bacteria studied was beyond the scope of the present work; however, such activity may have affected our results. The action of metabolites and toxins might explain, at least in part, the suppression observed in the treatments including *H. bacteriophora* isolate Rama Caída. In addition, Grewal *et al.* (1999) confirmed that *G. mellonella* cadavers infected with *H. bacteriophora* had repelling effects on juveniles of *M. incognita* and that allelochemicals produced by symbiotic bacteria are plant-parasitic nematode antagonists.

In our research, two cadavers per pot were applied, similar to a greenhouse trial conducted by Shapiro-Ilan *et al.* (2006b), who demonstrated that the application of two *T. molitor* cadavers infected with *S. riobrave* reduced the number of egg masses of *Meloidogyne partityla* (Kleynhans) in pecan seedlings. Larvae of *G. mellonella* and *T. molitor* were used as insect hosts in the present work due to the great commercial potential of this application method (Deol *et al.*, 2011; Shapiro-Ilan *et al.*, 2001, 2010). The host species did not show a marked influence on the suppression of *M. incognita* with the EPNs used. Both insect hosts might be employed in EPN application. *T. molitor* cadavers have a more rigid cuticle than those of *G. mellonella* and therefore exhibit greater resistance to physical damage during manipulation and transportation (Shapiro-Ilan *et al.*, 2010). However, the formulation of *G. mellonella* cadavers is an alternative that facilitates handling (Del Valle *et al.*, 2009; Shapiro-Ilan *et al.*, 2001).

**Table 1.** Application of cadavers of *Tenebrio molitor* (T) and *Galleria mellonella* (G) infected with the entomopathogenic nematodes *Heterorhabditis bacteriophora* (Hb, isolates RC, Mo and Jn) and *Steinernema diaprepesi* (Sd) in pepper plants inoculated with *Meloidogyne incognita*. Means followed by different letters are statistically significant according to Tukey's test (differences were significant at 5%).

Treatment	Leaf number	Dry weight of aerial part (g)	Number of galls	Number of egg masses	Number of eggs	Galls g root <sup>-1</sup>	Egg masses g root <sup>-1</sup>	Eggs g root <sup>-1</sup>
Control	12.42 ± 0.63 abc	0.75 ± 0.11 a	58.67 ± 9.79 bcd	30.25 ± 5.28 bc	14858 ± 2973 bc	9.71 ± 1.51 bc	5.00 ± 0.94 ab	2368 ± 481 ab
Carbofuran	9.00 ± 0.74 a	0.61 ± 0.18 a	8.17 ± 3.16 a	4.50 ± 2.04 a	2608 ± 960 a	3.04 ± 1.33 a	1.61 ± 0.72 a	774 ± 283 a
HbRCT	10.92 ± 0.54 ab	0.60 ± 0.07 a	25.00 ± 5.96 ab	14.58 ± 3.63 ab	4121 ± 1231 a	6.50 ± 0.80 ab	3.75 ± 0.51 ab	1065 ± 160 a
HbJnT	12.67 ± 0.57 abc	0.84 ± 0.06 a	77.00 ± 9.75 d	43.83 ± 7.04 c	18021 ± 3251 bc	14.24 ± 2.32 c	8.46 ± 1.92 b	3326 ± 494 bc
HbMoT	12.83 ± 0.61 bc	0.75 ± 0.06 a	57.50 ± 9.14 bcd	39.08 ± 6.72 c	18913 ± 2443 c	11.63 ± 1.44 bc	8.03 ± 1.25 b	4131 ± 534 c
SdT	13.50 ± 0.87 bc	0.88 ± 0.09 a	56.17 ± 5.37 bcd	36.00 ± 3.16 bc	16021 ± 1852 bc	11.99 ± 1.26 bc	7.83 ± 0.87 b	3402 ± 458 bc
HbRCG	13.83 ± 1.45 bc	0.98 ± 0.15 a	41.75 ± 9.67abcd	23.83 ± 4.91 abc	9142 ± 1891 ab	7.69 ± 1.31 ab	4.31 ± 0.65 ab	1493 ± 207 a
HbJnG	14.25 ± 0.83bc	1.09 ± 0.12 a	65.25 ± 11.16 cd	37.67 ± 5.61 bc	15404 ± 2231 bc	8.99 ± 1.34 abc	5.30 ± 0.83 ab	2242 ± 376 ab
HbMoG	15.00 ± 1.03 c	0.91 ± 0.09 a	36.58 ± 7.48 abc	23.75 ± 5.72 abc	11283 ± 1788 abc	6.50 ± 1.18 ab	4.18 ± 0.94 ab	2013 ± 257 ab
SdG	13.67 ± 0.40 cb	0.89 ± 0.08 a	43.67 ± 8.17abcd	27.17 ± 5.16 abc	8883 ± 1319 ab	8.31 ± 1.30 abc	5.39 ± 0.96 ab	1610 ± 145 a

**Table 2.** Application of cadavers of *Tenebrio molitor* (T) and *Galleria mellonella* (G) infected with the entomopathogenic nematodes *Heterorhabditis bacteriophora* (Hb, isolates RC, Mo and Jn) and *Steinernema diaprepesi* (Sd) in summer squash inoculated with *Meloidogyne incognita*. Means followed by different letters are statistically significant according to Tukey's test (differences were significant at 5%).

Treatment	Leaf number	Dry weight of aerial part (g)	Number of galls	Number of egg masses	Number of eggs	Galls g root <sup>-1</sup>	Egg masses g root <sup>-1</sup>	Eggs g root <sup>-1</sup>
Control	7.42 ± 0.42 a	1.84 ± 0.07 ab	80.50 ± 15.13 b	30.67 ± 2.89 b	9875 ± 1596 bc	5.00 ± 1.22 ab	1.84 ± 0.26 ab	542 ± 83 ab
Carbofuran	7.50 ± 0.26 a	1.92 ± 0.10 ab	39.67 ± 4.65 a	12.25 ± 3.84 a	4600 ± 663 ab	2.37 ± 0.31 a	0.76 ± 0.25 ab	296 ± 54 a
HbRCT	7.75 ± 0.39 a	2.28 ± 0.21 c	43.67 ± 3.78 a	19.00 ± 2.61 ab	4550 ± 1296 ab	2.28 ± 0.21 a	0.98 ± 0.12 ab	243 ± 68 a
HbJnT	7.42 ± 0.43 a	1.68 ± 0.09 a	47.5 ± 5.26 a	16.58 ± 2.65 a	6925 ± 1111 abc	4.11 ± 0.73 ab	1.40 ± 0.30 ab	611 ± 122 ab
HbMoT	7.83 ± 0.41 a	1.86 ± 0.08 ab	45.92 ± 5.26 a	17.33 ± 3.23 a	6575 ± 1371 abc	5.18 ± 0.66 b	1.89 ± 0.35 ab	740 ± 151 abc
SdT	7.58 ± 0.38 a	1.79 ± 0.10 ab	44.58 ± 4.19 a	15.75 ± 2.61 a	5875 ± 1164 abc	5.41 ± 0.57 b	1.89 ± 0.31 ab	718 ± 145 abc
HbRCG	7.92 ± 0.29 a	1.77 ± 0.07 a	38.67 ± 2.26 a	12.17 ± 2.09 a	3450 ± 663 a	3.71 ± 0.37 ab	1.16 ± 0.21 ab	338 ± 73 a
HbJnG	7.58 ± 0.40 a	1.90 ± 0.11 ab	57.00 ± 4.82 ab	22.00 ± 3.21 ab	8900 ± 979 bc	5.77 ± 0.23 b	2.10 ± 0.18 b	977 ± 133 bc
HbMoG	6.92 ± 0.70 a	1.44 ± 0.09 a	40.83 ± 4.37 a	13.58 ± 2.56 a	7925 ± 1354 abc	5.18 ± 0.59 b	1.64 ± 0.27 ab	1045 ± 213 bc
SdG	7.08 ± 0.36 a	1.91 ± 0.08 ab	45.42 ± 3.44 a	15.17 ± 1.93 a	10050 ± 1123 c	5.40 ± 0.45 b	1.81 ± 0.23 ab	1243 ± 182 c

Inconsistent results were also reported by Shapiro-Ilan *et al.* (2006a) in an assessment of the effect of the application of *T. molitor* cadavers infected with *S. feltiae* and *S. riobrave* on *M. partityla*. Further research is necessary to elucidate the reasons why the efficacy observed in laboratory assays is frequently not found in greenhouse or field experiments.

In the present work, we observed a marked difference in the suppression of *M. incognita* among the three isolates of *H. bacteriophora* evaluated. The performance of each isolate may be associated with metabolites produced by their symbiotic bacteria, which differ in quality and quantity among isolates (Webster *et al.*, 2002). *Heterorhabditis bacteriophora* Rama Caida produced the greatest reduction in the parasitism of *M. incognita* in pepper and summer squash and will require field investigations for commercial application.

The species *S. diaprepesi* did not exhibit potential for the suppression of *M. incognita* in pepper and summer squash, although IJs of this species exhibited increased survival in the soil than *H. bacteriophora* IJs (Shapiro-Ilan *et al.*, 2006b). Pérez and Lewis (2004) suggested that *Steinernema* species might be more effective in the suppression

of root-knot nematodes than *H. bacteriophora* because of their greater capacity to penetrate roots and to release symbiotic bacteria inside them. Within roots, the bacteria would release allelochemicals that are toxic and repellent to *Meloidogyne* spp. (Grewal *et al.*, 1999). *Steinernema diaprepesi* IJs remained near the roots in pepper and summer squash plants. The greatest efficacy of steinernematids suggested by Pérez and Lewis (2004) was not evident in our experiments. Additional studies are necessary to determine the behavior of *S. diaprepesi* IJs and the action of their allelochemicals against root-knot nematodes parasitizing pepper and summer squash.

Our results indicated that the application of insect cadavers infected with the EPNs studied might reduce *M. incognita* damage in pepper and summer squash plants. However, further studies are required to determine their efficacy in field applications.

### Acknowledgments

The present work was funded by the program Curso de Acción para la Investigación y el Desarrollo, CAI+D 2009 of the Universidad Nacional del Litoral.

### Resumen

**E.E. Del Valle, P. Lax, J. Rondán Dueñas y M.E. Doucet. 2013. Efecto de insectos cadáveres infectados por *Heterorhabditis bacteriophora* y *Steinernema diaprepesi* sobre el parasitismo de *Meloidogyne incognita* en plantas de pimiento y zapallito redondo de tronco. Cien. Inv. Agr. 40(1):109-118.** El efecto de cadáveres de insectos infectados con tres aislados de *Heterorhabditis bacteriophora* y un aislado de *Steinernema diaprepesi* sobre una población de *Meloidogyne incognita* en plantas de pimiento (*Capsicum annuum*) y zapallito redondo de tronco (*Cucurbita maxima*) fue evaluado en experiencias de invernadero conducidas en Santa Fe (Argentina). Los cadáveres de insectos necesarios para la experiencia se obtuvieron infectando larvas de último estadio de *Galleria mellonella* y *Tenebrio monitor* con nematodos entomopatógenos. Dos cadáveres de seis días de infección se colocaron debajo de la superficie del suelo de macetas que fueron inoculadas con 100 juveniles de segundo estadio de *M. incognita*. A los sesenta días se registraron los siguientes parámetros en cada planta: número de hojas, peso seco de la parte aérea, número de agallas, masas de huevos

y huevos, y número de agallas, masas de huevos y huevos g<sup>1</sup> de materia fresca radical. En pimiento, la única variable afectada por los cadáveres en relación al testigo fue el número de huevos de *M. incognita* en el tratamiento de cadáveres de *T. molitor* infectados con *H. bacteriophora* aislado Rama Caída. En zapallito redondo de tronco, varios tratamientos provocaron una disminución en el número de agallas y masas de huevos. Sólo la aplicación de cadáveres de *G. mellonella* infectados con *H. bacteriophora* aislado Rama Caída demostró ser eficiente en reducir el número de huevos de *M. incognita*. Nuestros resultados indicaron que la aplicación de cadáveres de insectos infectados con los nematodos entomopatógenos estudiados podría reducir los daños causados por *M. incognita* en plantas de pimiento y zapallito redondo de tronco.

**Palabras clave:** Cadáveres de insectos infectados, control biológico, nematodos entomopatógenos.

## References

- Akhurst R., and N.E. Boemare. 1990. Biology and taxonomy of *Xenorhabdus*. In: Gaugler R., and Kaya H.K. (eds.). Entomopathogenic nematodes in biological control. Boca Raton, FL, CRC Press. p. 75-87.
- Boemare N.E., Akhurst R.J., and R.G. Mourant. 1993. Deoxyribonucleic acid relatedness between *Xenorhabdus* spp. (Enterobacteriaceae), symbiotic bacteria of entomopathogenic nematodes, with a proposal to transfer *Xenorhabdus luminescens* to a new genus *Photorhabdus* gen. nov. International Journal of Systematic Bacteriology 43:249-255.
- Boina D.R., Lewis E.E., and J.R. Bloomquist. 2008. Nematicidal activity of anion transport blockers against *Meloidogyne incognita*, *Caenorhabditis elegans*, and *Heterorhabditis bacteriophora*. Pest Management Science 64:646-653.
- Bruck D.J., Shapiro-Ilan D.I., and E.E. Lewis. 2005. Evaluation of application technologies of entomopathogenic nematodes for control of the black vine weevil, *Otiorynchus sulcatus*. Journal of Economic Entomology 98:1884-1889.
- Burman M. 1982. *Neoplectana carpocapsae*: a toxin production by axenic insect parasitic nematode. Nematologica 28:62-70.
- Creighton C.S., and G. Fassuliotis. 1985. *Heterorhabditis* sp. (Nematoda: Heterorhabditidae): A nematode parasite isolated from the banded cucumber beetle *Diabrotica balteata*. Journal of Nematology 17:150-153.
- Del Valle E.E., Dolinski C., Barreto E.L.S., and R.M. Souza. 2009. Effect of cadaver coatings on emergence and infectivity of the entomopathogenic nematode *Heterorhabditis baujardi* LPP7 (Rhabditida: Heterorhabditidae) and the removal of cadavers by ants. Biological Control 50:21-24.
- Del Valle E.E., Dolinski C., Barreto E.L.S., and R.M. Souza. 2008. Efficacy of *Heterorhabditis baujardi* LPP7 (Nematoda: Rhabditida) applied in *Galleria mellonella* (Lepidoptera: Pyralidae) insect cadavers to *Conotrachelus psidii*, (Coleoptera: Curculionidae) larvae. Biocontrol Science and Technology 18:33-41.
- Deol Y.S., Jagdale G.B., Cañas L., and P.S. Grewal. 2011. Delivery of entomopathogenic nematodes directly through commercial growing medium via the inclusion of infected host cadavers: A novel approach. Biological Control 58:60-67.
- Eischen F.A., and A. Dietz. 1990. Improved culture techniques for mass rearing *Galleria mellonella*. Entomological News 101:123-128.
- Fallon D.J., Kaya H.K., Gaugler R., and B.S. Sipes. 2002. Effects of entomopathogenic nematodes on *Meloidogyne javanica* on tomatoes and soybeans. Journal of Nematology 34:239-245.
- Fallon D.J., Kaya H.K., Gaugler R., and B.S. Sipes. 2004. Effect of *Steinernema feltiae*-*Xenorhabdus bovienii* insect pathogen complex on *Meloidogyne javanica*. Nematology 6:671-680.
- Grewal P.S. 2002. Formulation and application technology. In: Gaugler R. (ed.). Entomopathogenic nematology. CABI, Wallingford, UK. p. 265-287.



- Grewal P.S., Bornstein-Forst S., Burnell A.M., Glazer I., and G.B. Jagdale. 2006. Physiological, genetic, and molecular mechanisms of chemoreception, thermobiosis, and anhydrobiosis in entomopathogenic nematodes. *Biological Control* 38:54-65.
- Grewal P.S., Ehlers R.-U., and D.I. Shapiro-Ilan. 2005. *Nematodes As Biocontrol Agents*. CABI, U.K. 505 pp.
- Grewal P.S., Lewis E.E., and S. Venkatachari. 1999. Allelopathy: A possible mechanism of suppression of plant-parasitic nematodes by entomopathogenic nematodes. *Nematology* 1:735-743.
- Halbrendt J.M., and J.A. LaMondia. 2004. Crop Rotation And Other Cultural Practices. *In: Chen, Z.X., Chen S.Y., and D.W. Dickson (eds.). Nematology, Advances and Perspectives Vol 2: Nematode Management and Utilization*. CABI, Cambridge. p. 909-930.
- Hu K., Li J., and J.M. Webster. 1999. Nematicidal metabolites produced by *Photorhabdus luminescens* (Enterobacteriaceae), bacterial symbiont of entomopathogenic nematodes. *Nematology* 1:457-469.
- Hussey R.S., and K.R. Barker. 1973. A comparison of methods of collecting inocula of *Meloidogyne* spp. including a new technique. *Plant Disease Report* 57:1025-1028.
- Jagdale G.B., and P.S. Grewal. 2008. Influence of the entomopathogenic nematode *Steinernema carpocapsae* infected host cadavers or their extracts on the foliar nematode *Aphelenchoides fragariae* on *Hosta* in the greenhouse and laboratory. *Biological Control* 44:13-23.
- Jagdale G.B., Somasekhar N., Grewal P.S., and M.G. Klein. 2002. Suppression of plant-parasitic nematodes by application of live and dead infective juveniles of an entomopathogenic nematode, *Steinernema carpocapsae*, on boxwood (*Buxus* spp.). *Biological Control* 24:42-49.
- Kaya H.K., and S.P. Stock. 1997. Techniques In Insect Nematology. *In: Lacey, L.A. (ed.). Manual of techniques in insect pathology Biological Techniques Series*. London, Academic Press. p. 281-324.
- Koenning S.R., Overstreet C., Noling J.W., Donald P.A., Becker J.O., and B.A. Fortnum. 1999. Survey of crop losses in response to phytoparasitic nematodes in the United States for 1994. *Journal of Nematology* 31:587-618.
- Lewis E.E., Grewal P.S., and S. Sardanelli. 2001. Interactions between the *Steinernema feltiae*-*Xenorhabdus bovienii* insect pathogen complex and the root-knot nematode *Meloidogyne incognita*. *Biological Control* 21:55-62.
- Lewis E.E., and P.S. Grewal. 2005. Interactions With Plant-Parasitic Nematodes. *In: Grewal P.S., Ehlers R.-U., and D.I. Shapiro-Ilan (eds.). Nematodes as Biocontrol Agents*. CABI, UK. p. 349-361.
- Molina J.P., Dolinski C., Souza R.M., and E.E. Lewis. 2007. Effect of entomopathogenic nematodes (Rhabditida: Steinernematidae and Heterorhabditidae) on *Meloidogyne mayaguensis* Rammah and Hirschmann (Tylenchida: Meloidoginidae) infection in tomato plants. *Journal of Nematology* 39:338-342.
- Pérez E.E., and E.E. Lewis. 2002. Use of entomopathogenic nematodes to suppress *Meloidogyne incognita* on greenhouse tomatoes. *Journal of Nematology* 34:171-174.
- Pérez E.E., Lewis E.E., and D.I. Shapiro-Ilan. 2003. Impact of host cadaver on survival and infectivity of entomopathogenic nematodes (Rhabditida: Steinernematidae and Heterorhabditidae) under desiccating conditions. *Journal of Invertebrate Pathology* 82:111-118.
- Pérez E.E., and E.E. Lewis. 2004. Suppression of *Meloidogyne incognita* and *Meloidogyne hapla* with entomopathogenic nematodes on greenhouse peanuts and tomatoes. *Biological Control* 30:336-341.
- Perry R.N., Hominick W.M., Beane J., and B. Briscoe. 1998. Effects of the entomopathogenic nematodes, *Steinernema feltiae* and *S. carpocapsae*, on the potato cyst nematode, *Globodera rostochiensis*, in pot trials. *Biocontrol Science and Technology* 8: 175-180.
- Poinar G.O. 1990. Biology And Taxonomy Of *Steinernematidae*. *In: Gaugler R., and H.K. Kaya (eds.). Entomopathogenic Nematodes in Biological Control*, CRC Press. FL. p. 23-61.

- Riegel C., Dickson D.W., Nguyen K.B., and G.C. Smart. 1998. Management of root-knot nematodes with entomopathogenic nematodes. *Journal of Nematology Supplement* 24:637-641.
- Schafer K.S. 1999. Methyl Bromide Phase-out Strategies: A Global Compilation of Laws and Regulations, OzonAction Programme of the United Nations Environment Programme Division of Technology, Industry and Economics (UNEP TIE), Chatelaine, Geneva. 149 pp.
- Shapiro D.I., and I. Glazer. 1996. Comparison of entomopathogenic nematode dispersal from infected hosts versus aqueous suspension. *Environmental Entomology* 25:1455-1461.
- Shapiro-Ilan D.I., Nyczepir A.P., and E.E. Lewis. 2006a. Entomopathogenic nematodes and bacteria applications for control of the pecan root-knot nematode, *Meloidogyne partityla*, in the greenhouse. *Journal of Nematology* 38:449-454.
- Shapiro-Ilan D.I., Stuart R., and C.W. McCoy. 2006b. A comparison of entomopathogenic nematode longevity in soil under laboratory conditions. *Journal of Nematology* 38:119-129.
- Shapiro-Ilan D.I., and E.E. Lewis. 1999. Comparison of entomopathogenic nematode infectivity from infected hosts versus aqueous suspension. *Environmental Entomology* 28:907-911.
- Shapiro-Ilan D.I., Han, R., and C. Dolinski. 2012. Entomopathogenic nematode production and application technology. *Journal of Nematology* 44:206-217.
- Shapiro-Ilan D.I., Lewis E.E., and W.L. Tedders. 2003. Superior efficacy observed in entomopathogenic nematodes applied in infected-host cadavers compared with application in aqueous suspension. *Journal of Invertebrate Pathology* 83:270-272.
- Shapiro-Ilan D.I., Lewis E.E., Behle R.W., and M.R. McGuire. 2001. Formulation of entomopathogenic nematode-infected-cadavers. *Journal of Invertebrate Pathology* 78:17-23.
- Shapiro-Ilan D.I., Morales-Ramos J.A., Rojas M.G., and W.L. Tedders. 2010. Effects of a novel entomopathogenic nematode-infected host formulation on cadaver integrity, nematode yield, and suppression of *Diaprepes abbreviatus* and *Aethina tumida*. *Journal of Invertebrate Pathology* 103:103-108.
- Smitley D.R., Warner F.W., and G.W. Bird 1992. Influence of irrigation and *Heterorhabditis bacteriophora* on plant-parasitic nematodes in turf. *Journal of Nematology* 24:637- 641.
- Thomas G.M., and G.O. Poinar Jr. 1979. *Xenorhabdus* gen. nov., a genus of entomopathogenic and nematophilic bacteria of the family Enterobacteriaceae. *International Journal of Systematic Bacteriology* 29:352-360.
- Webster J.M., Chen G., Hu K., and J. Li. 2002. Bacterial metabolites. In: Gaugler R. (ed.). *Entomopathogenic nematology* CABI, Wallingford, UK. p. 99-114.
- Woodring J.L., and H.K. Kaya. 1988. Steinemematid and heterorhabditid nematodes: A handbook of techniques, Arkansas Agricultural Experiment Station, Fayetteville, AK. Series Bulletin 331.