

Accepted Manuscript

Note

Synthetic tools for the characterization of galactofuranosyl transferases. Glycosylations via acylated glycosyl iodides

Luciana Baldoni, Carla Marino

PII: S0008-6215(13)00128-6

DOI: <http://dx.doi.org/10.1016/j.carres.2013.03.032>

Reference: CAR 6443

To appear in: *Carbohydrate Research*

Received Date: 18 December 2012

Revised Date: 26 March 2013

Accepted Date: 28 March 2013

Please cite this article as: Baldoni, L., Marino, C., Synthetic tools for the characterization of galactofuranosyl transferases. Glycosylations via acylated glycosyl iodides, *Carbohydrate Research* (2013), doi: <http://dx.doi.org/10.1016/j.carres.2013.03.032>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



1
2
3
4 **Synthetic tools for the characterization of galactofuranosyl**
5
6
7 **transferases. Glycosylations via acylated glycosyl iodides**
8
9

10
11
12
13 Luciana Baldoni, Carla Marino*

14
15
16
17
18 *CIHIDECAR-CONICET, Departamento de Química Orgánica, Facultad de Ciencias Exactas*
19 *y Naturales, Universidad de Buenos Aires, Buenos Aires (1428), Argentina.*
20 *cmarino@qo.fcen.uba.ar*
21

22
23 **Abstract**
24

25
26
27
28 With the aim to develop synthetic tools for the characterization of
29 galactofuranosyltransferases, the synthesis of 9-decenyl glycosides of D-Man_p, D-Galf and β-
30 D-Galf-(1→3)-D-Man_p was targeted. The interest in the alkenyl aglycone arises from its
31 potential conjugation reactions, once the terminal double bond has been conveniently
32 functionalized. The glycosylation of β-D-Galf-(1→3)-D-Man_p was attempted by two different
33 approaches: the trichloroacetimidate method and the glycosylation via the glycosyl iodide.
34 The conditions for the latter were established on the basis of glycosylation assays of per-*O*-
35 acetylmannose. On the other hand, the study of glycosylation reactions *via* per-*O*-benzoylated
36 galactofuranosyl iodide confirms the versatility of glycosyl iodides as donors.
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52

53 **Keywords:** Mannopyranosyl iodide/ per-*O*-Benzoylated-galactofuranosyl iodide / per-*O*-*tert*-
54 Butyldimethylsilyl-β-D-galactofuranose / Galactofuranosyl transferases.
55
56

57 *Corresponding author.
58 Tel/Fax: 54-11-45763352
59 E-mail address: cmarino@qo.fcen.uba.ar
60
61
62
63
64
65

Among the numerous structures of pathogenic microorganisms in which D-galactofuranose occurs,¹ the disaccharide β -D-Galf-(1 \rightarrow 3)-D-Manp (**1**, Figure 1) is found in glycoconjugates of protozoa (*Trypanosoma cruzi* and *Leishmania* spp.) and in fungi, as *Aspergillus fumigatus*^{2,3}. In *T. cruzi*, motif **1** is present as non-reducing terminal units of the glycoinositolphospholipids (GIPLs),⁴⁻⁶ which are important for the interaction with the intestine of the insect vector.⁷ In *Leishmania*, disaccharide **1** is present as an internal unit in the lipophosphoglycan (LPG),⁸ which was also shown to play a critical role in the attachment of *Leishmania* promastigotes to the fly midgut.⁹

For elucidating the biosynthesis of D-Galf containing glycoconjugates, synthetic substrates of the involved enzymes are required.¹⁰⁻¹² Oligosaccharides containing the β -D-Galf-(1 \rightarrow 3)-D-Manp (**1**) motif have been synthesized,^{13,14} as well as some derivatives of **1**, which were afforded by different approaches.¹⁵ We have described the synthesis of free disaccharide **1** using the glycosyl-aldonolactone approach, and we have shown that is hydrolyzed by the *exo* β -D-galactofuranosidase of *P. fellutanum*, our non-pathogenic model to evaluate the synthetic tools developed for studying the D-Galf related enzymes.¹⁶





Figure 1. Synthetic targets.


With the aim of obtaining a derivative of **1** for the characterization of the galactofuranosyltransferases of *P. fellutanum* and *T. cruzi*, we have now targeted the synthesis of 9-decenyl β -D-Galf-(1 \rightarrow 3)-D-Manp (**2**, Figure 1), which is expected to be an acceptor of D-Galf units and as precursor of other derivatives designed to study the immunogenic activity of **1**. For the synthesis of **2** it was necessary to introduce the alkenyl moiety with a glycosylation method that would preserve the Galf-(1 \rightarrow 3)-Manp linkage. We decided to carry on this synthesis by two alternative approaches: one based on the trichloroacetimidate method, which required transformations that we have previously applied,^{17,18} and the other involving a glycosyl iodide donor (Scheme 1). In this case, the conditions for the glycosylation of the acetylated mannose unit must be established. For the glycobiological studies, we have also decided to synthesize acceptors **3** and **4** (Figure 1). The optimized conditions for the synthesis of **3** would be useful for the glycosylation of **1**. On the other hand, as continuation of our studies on the scope of glycosylations *via* galactofuranosyl iodides,¹⁸⁻²⁰ this reaction was investigated for the synthesis of compound **4** from acylated precursors of D-Galf. Thus, we report here the studies of glycosylations of per-*O*-acetylmannose *via* glycosyl iodides, the exploration of the galactofuranosylation *via* iodides prepared from per-*O*-acylated precursors, and the synthesis of glycosyl disaccharide **2**.



Scheme 1. Retrosynthetic approaches for 9-decenyl β -D-Galf-(1 \rightarrow 3)-D-Manp (**2**)

For the synthesis of mannopyranoside **3** we used the glycosyl iodide method starting from penta-*O*-acetyl- α,β -D-mannopyranose (**5**). In first instance, we established the reaction conditions for the formation of the mannopyranosyl iodide and its subsequent glycosylation, in order to apply similar conditions to the synthesis of disaccharide **2**. According to the reported conditions for similar substrates,^{21,22} compound **5** was treated with TMSI (2.4 equiv) at room temperature and after 2 h, the medium was neutralized with EtN(*i*Pr)₂, and 9-decen-1-ol was added as acceptor (Scheme 2). Under this conditions, starting compound **5** was not completely consumed, and the reaction proceeded slowly towards the formation of a main product, but on the basis of the ¹³C NMR spectrum, this compound was identified as the orthoester **7** (Table 1, entry 1).^{23,24}

Attempts to rearrange the orthoester **7** with TMSOTf^{25,26} were not satisfactory as, although the NMR spectra of the crude product showed the formation of **8**, a complex mixture of products has been obtained as result of partial *O*-deacetylation.



Scheme 2. Synthesis of 9-decanyl α -D-mannopyranoside (**3**)

It has been reported that ZnI₂ accelerates the formation of peracetylated glycosyl iodides and prevents the formation of the orthoester.^{21,22} The ZnI₂ also acted as a iodide source and in

Table 1. Reaction conditions assayed for glycosylation of penta-*O*-acetyl-D-mannopyranose (**5**) via mannosyl iodide.

Entry	Iodide 6 formation			Glycosylation			
	TMSI (equiv)	ZnI ₂ (equiv)	Conditions	Base/promoter (equiv)	MS 4Å	Conditions	Products/Observations
1	2.4		25 °C, 2 h	EtN(<i>i</i> Pr) ₂ (2.4)	-	25 °C, 144 h	7 (50 %)
2	1.5	0.4	45 °C, 0.5 h	ZnI ₂ (1.0)	yes	45 °C, 3.5 h	8 (46%) (80% after reacetylation)

this way the halogen exchange in the anomeric carbon, that would occur with other halogenated Lewis acids, was avoided. Hence, ZnI₂ was added in a substoichiometric amount and iodide **6** was formed in just 0.5 h (Table 1, entry 2). As the glycosylation of acylated iodides generally requires a promoter,^{23,24} after the complete transformation of **5** into **6**, an additional amount of ZnI₂ was added together with the 4Å powdered molecular sieves. They were not used in the first step because it has been reported that they retard the iodide formation.^{23,24} Under these conditions the 9-decenyl glycoside **8** was obtained as major product (46 %), along with an important amount of partially de-*O*-acylated products. Therefore, after reacetylation, compound **8** was obtained in 80 % combined yield (Table 1, entry 2). The NMR spectra of **8** showed that the glycosylation occurred with complete 1,2-*trans* stereoselectivity, due to participation of the neighboring acetyl group on O-2. *O*-Deacetylation of crude compound **8** afforded **3** in 80 % overall yield from **5**.

The synthesis of **8** and **3** as precursors of oligosaccharides present in the antigenic lipophosphoglycan of *Leishmania donovani*, had been previously accomplished by the Koenigs-Knorr method from acetobromomannose.²⁷ The glycosyl iodide method here

1
2
3
4 described, besides avoiding the use of mercuric salts, affords **8** in higher yield, and the
5
6 complete NMR spectroscopic characterization of both **8** and **3** is now provided.
7


8
9 Previously, we have described the synthesis of per-*O*-TBS- β -D-galactofuranose (**9**) and
10
11 its glycosylation by *in situ* activation with TMSI as the galactofuranosyl iodide **12** (Scheme
12
13 3). Compound **12** was effectively glycosylated to afford *O*-,¹⁸ *S*-, *C*-galactofuranosides,²⁰ and
14
15 some nitrogenated derivatives,¹⁹ under mild conditions compatible with labile acceptors.
16
17 Recently, the lipoteichoic acid from *Streptococcus* sp. DSM 8747 has been synthesized by
18
19 glycosylation of **9** via the galactofuranosyl iodide, and the method showed to be significantly
20
21 more efficient than those using traditional glycosyl donors.²⁸ Condensation of persilylated **9**
22
23 with 9-decen-1-ol under the conditions previously described,¹⁸ afforded glycoside **14** in 83 %
24
25 yield as an anomeric mixture β/α in a 3:1 ratio. A similar diastereoselectivity was observed
26
27 with simple acceptors, which was increased when bulky acceptors were used.¹⁸ *O*-Desilylation
28
29 of **14** with TBAF afforded the free galactofuranoside **4** in 66 % yield.
30
31
32
33
34
35

36 Although β -D-galactofuranosides can be stereoselectively obtained by neighboring-
37
38 group participation from acylated precursors using SnCl₄ or other Lewis acids³ as promoters,
39
40 the glycosyl acceptors are limited to acid stable derivatives. We aimed to investigate the scope
41
42 of the galactofuranosyl iodide glycosylations from the easily available peracylated Galf
43
44 derivatives **10**²⁹ and **11**,³⁰ which are expected to give glycosides with higher
45
46 diastereoselectivity than **9**, due to the anchimeric effect. In order to optimize the reaction
47
48 conditions, *n*BuOH was employed as a model acceptor. As peracylated precursors are less
49
50 reactive than persilylated,³¹⁻³³ more drastic conditions than those employed for **9** would be
51
52 required. The assayed conditions involving variations in the amounts of TMSI, temperature
53
54 and reaction time are summarized in Table 2. The effect of molecular sieves and promoters
55
56
57
58
59
60
61
62
63
64
65

was also examined. As expected, **11** was more reactive than **10**, but less than **9** (Table 2, entries 1-3). The best condition for the preparation of iodide **13**, in the absence of a catalyst or a promoter, was the treatment of **11** with 3 equiv of TMSI (Table 2, entry 5). Compound **10** required 4.5 equiv of TMSI to complete the reaction (Table 2, entry 4). Iodide **13** was not stable enough to be isolated.

The addition of powdered molecular sieves during the second step of the reaction avoided the formation of TMSOAc and TMSOBz^{23,34} and the subsequent reaction with **13**, which would afford **10** and **11** as recombination products.

The effect of the addition of ZnI₂ during the iodide formation was also studied (Table 2, entries 6-8). In the presence of 0.6 equiv of ZnI₂, compound **13** was formed at room temperature with only 1.2 equiv of TMSI and in 0.5 h (Table 2, entry 8). When the amount of ZnI₂ was reduced, the consumption of acetate **11** was not complete, although the formed iodide was consumed in 1 h (Table 2, entries 7 and 8). The best results were obtained conducting the reactions at room temperature, without the need of heating to 45 °C, as in the case of the mannopyranosyl iodide **6**.



Scheme 3. Synthesis of *O*-galactofuranosides *via* in situ formed galactofuranosyl iodides.

Table 2. Reaction conditions assayed for *O*-glycosylations via in situ formed galactofuranosyl iodides.


Entry	Precursor	TMSI (equiv)	ZnI ₂ (equiv)	Conditions	MS 4Å	Conversion to iodide ^a	Products/ Observations
1	9	1.2	-	0 °C, 0.5 h	yes	100 %	14 β:α 3:1
2	10	1.2	-	0 °C, 0.5 h	yes	0 %	-
3	11	1.2	-	0 °C, 0.5 h	yes	20 %	15 β
4	10	4.5	-	0→25 °C, 1 h	yes	100 %	15 β
5	11	3.0	-	0→25 °C, 1.5 h	yes	100 %	15 β
6	11	1.2	0.6	0→25 °C, 0.5 h	yes	100 %	15 β in 1 h (80 %)
7	11	1.2	0.4	0→25 °C, 1-2 h	yes	80 %	15 β in 1 h
8	11	1.2	0.2	0→25 °C, 1-2 h	yes	60 %	15 β in 1 h

^aEstimated by TLC, ^bIsolated by column chromatography

Under the optimized conditions established for the synthesis of **15** (Table 2, entry 8), the analogous decenyl glycoside **16** was obtained in 66 % yield, mainly in the β-configuration (9:1). *O*-Debenzylation of **16** with NaOMe/MeOH in CH₂Cl₂ afforded **4** in almost quantitative yield. The β-configuration of the major component of **16** and **4** was confirmed on the basis of the ¹³C NMR spectra, which showed characteristic resonances for C-1 (105.6 and 109.4 ppm, respectively) and signals corresponding to C-2 and C-4 above 80 ppm, also characteristic of the β-D-Galf configuration.

Despite the convenience of the use of iodide **13** to achieve stereoselectively β-D-galactofuranosides, the disarmed character of this benzoylated iodide was evidenced when allylTMS, (TMS)₂S or 2,4,6-tri-*O*-benzoyl-D-manono-1,4-lactone were used as acceptors.

While these compounds were effectively coupled with **9**,^{18,20} the glycosylation of **13** failed.



Scheme 4. Synthesis of 9-decanyl β -D-Galf-(1 \rightarrow 3)-D-Manp (**2**).

Both strategies designed to accomplish the synthesis of the decenyl glycoside **2**, required per-*O*-acetylated disaccharide **17**, which was afforded as an anomeric mixture in 75 % yield by treatment of **1**¹⁶ with $\text{Ac}_2\text{O}/\text{py}$ (Scheme 4). The approach involving a trichloroacetimidate donor required the selective anomeric *O*-deacetylation of **17**. Hence, **17** was treated with ethylenediamine and acetic acid to afford the hemiacetal **18** in 84 % yield (Scheme 4), exclusively in the α -configuration as indicated by the ^1H NMR spectrum. Treatment of **18** with trichloroacetonitrile and DBU afforded the trichloroacetimidate **20** (83 %). Glycosylation of **20** with 1.5 equiv of 9-decen-1-ol in CH_2Cl_2 using TMSOTf as catalyst gave **21** in 72 % yield. The NMR spectra showed that, despite the anchimeric assistance from

1
2
3
4 the participating acetyl group at O-2, compound **21** was obtained as an inseparable mixture of
5
6 β/α anomers, in a 3:2 ratio. The ^{13}C NMR spectrum showed resonances at 102.7 and 102.5
7
8 ppm corresponding to C-1' of the β - and the α -anomers, respectively, and signals at δ 99.6
9
10 (C-1 β) and 99.2 (C-1 α) due to the mannopyranosyl unit. The ^1H NMR spectrum showed
11
12 singlets at 5.19 and 5.13 ppm for H-1' of the β - and the α -anomers, respectively, and doublets
13
14 at 4.90 (H-1 α) and 4.52 ppm (H-1 β) for the mannopyranosyl unit.
15
16
17
18

19
20 On the other hand, peracetylated compound **17** was treated with TMSI/ZnI₂, according
21
22 to the conditions optimized for the formation and glycosylation of iodide **6** (Table 1, entry 2),
23
24 although a greater amount of ZnI₂ (0.7 equiv) was necessary to obtain iodide **19**. Then, 9-
25
26 decen-1-ol and 4Å powdered molecular sieves were added (Scheme 4). Compound **21** (78 %)
27
28 was obtained, along with a small amount of **18**.
29
30

31
32 In the ^1H NMR spectrum of **21** obtained in this way, it was observed that an anomeric
33
34 mixture in a β/α ratio of 3:2 was actually obtained. This ratio was almost equal to that
35
36 obtained in the glycosylation *via* the trichloroacetimidate **20**, suggesting that the
37
38 stereoselectivity depends on the substrate itself rather than the glycosylation method used.
39
40 Probably, the β -D-Galp unit as substituent on the O-3 of D-Manp would be responsible for a
41
42 distortion in the intermediate bicyclic 1,2-acyloxonium ion, making the anchimeric
43
44 participation less efficient.
45
46
47
48

49 Finally, de-*O*-acetylation of **21** with NaOMe/MeOH in CH₂Cl₂ afforded **2** in
50
51 quantitative yield (Scheme 4). The ^1H NMR spectrum of **2**, showed signals corresponding to
52
53 H-1' (δ 5.04 and 5.00) of both anomers, which correlated with signals at 104.3 and 105.4 ppm
54
55 in the HSQC experiment. The broad singlets at δ 4.74 (H-1 α) and 4.47 (H-1 β) corresponding
56
57 to the Manp unit, correlated with signals at 99.9 (C-1 β) and 99.7 (C-1 α) ppm. The assignment
58
59
60
61
62
63
64
65

1
2
3
4 of the anomeric configuration in the Man_p moiety was confirmed by a 2D NOESY
5
6 experiment which showed cross peaks between H-1/ H-3 and H-1/H-5 for the β-anomer.
7
8

9
10 Since its development, the trichloroacetimidate glycosylation method has been widely
11 used as it has the advantage of being mild enough to preserve other glycosidic linkages
12 present in the acceptor or in the donor.³⁵ This aspect is particularly critic in the case of
13 furanosyl units, due to their lability. Glycosyl iodides have long been underused as they were
14 considered too reactive to be of synthetic utility. However, their use as glycosyl donors has
15 been revalued over the past 15 years mainly due to the development of new methods of
16 preparation.³⁶ Our studies on the glycosylation *via* the benzoylated galactofuranosyl iodide **13**
17 confirms once more the versatility of glycosyl iodides as donors.
18
19
20
21
22
23
24
25
26
27

28
29 For the synthesis of **2** by the trichloroacetimidate approach compound **21** was obtained
30 from **17** in three steps with the corresponding column chromatography purifications in 50 %
31 overall yield. The synthesis of **21** from **17** by means of the glycosyl iodide strategy involved
32 two reaction steps and two column chromatography purifications, in 78 % overall yield.
33 Beyond the yield, the advantage of the iodide approach was that the sequence was shorter and
34 the reaction times were significantly reduced. On the other hand, in the synthesis of **2** *via* a
35 mannosyl iodide, we demonstrated that a β-D-Galf unit in the glycosyl donor resists the
36 glycosylation, without degradation.
37
38
39
40
41
42
43
44
45
46
47
48
49
50

51 **1. Experimental section**

52 **1.1. General synthetic methods**

53
54
55
56
57
58
59
60
61
62
63
64
65

Analytical thin layer chromatography (TLC) was performed on Silica Gel 60 F254 (Merck) aluminum supported plates (layer thickness 0.2 mm) with solvent systems given in the text. Visualization of the spots was effected by exposure to UV light and charring with a solution of 10 % (v/v) sulphuric acid in EtOH, containing 0.5 % p-anisaldehyde. Column chromatography was carried out with Silica Gel 60 (230-400 mesh, Merck). Optical rotations were measured with a Perkin-Elmer 343 digital polarimeter. Nuclear magnetic resonance (NMR) spectra were recorded with a Bruker AMX 500 spectrometer. Assignments of ^1H and ^{13}C were assisted by 2D ^1H -COSY and HSQC experiments. High resolution mass spectra (HRMS ESI⁺) were recorded in a Bruker micrOTOF-Q II spectrometer.

1.2. 9-Decenyl 2,3,4,6-tetra-*O*-acetyl- α -D-mannopyranoside (**8**)

A suspension of **5** (0.1 g, 0.25 mmol) and ZnI_2 (0.4 equiv, 0.032 g, 0.1 mmol) in anhydrous CH_2Cl_2 (10.0 mL) was stirred under argon atmosphere at 0 °C for 15 min. TMSI (1.5 equiv, 50.0 μL , 0.375 mmol) was slowly added and the stirring was continued for another 15 min. The suspension was allowed to reach room temperature and then heated at 45 °C. After 30 min of stirring TLC analysis showed total consumption of the starting material ($R_f = 0.36$, 1:1 hexane/EtOAc) and a single spot of $R_f = 0.52$ (1:1 hexane/EtOAc). Powdered molecular sieves 4Å and 9-decen-1-ol (0.14 mL, 0.75 mmol, 3.0 equiv) were added. After 3 h of stirring at 45 °C and 18 h at room temperature, the solution was diluted with CH_2Cl_2 (250 mL), washed with NaHCO_3 (ss) (2 x 140 mL) and water (3 x 100 mL), dried (Na_2SO_4) and concentrated. The residue was purified by column chromatography (3:1→3:2 hexane/EtOAc) affording syrupy compound **8** (0.056 g, 46 %), $R_f = 0.70$ (1:1 hexane/EtOAc), $[\alpha]_D +38.4$ (c 0.9, CHCl_3). Lit:²⁷ $[\alpha]_D +40$ (c 1, CHCl_3). Fractions of $R_f = 0.34$ (1:1 hexane/EtOAc) were

1
2
3
4 reacetylated affording **195** in 80 % overall yield. ^1H NMR (CDCl_3 , 500 MHz) δ 5.80 (m, 1H,
5
6 $\text{CH}=\text{CH}_2$), 5.34 (dd, $J = 3.4, 10.0$ Hz, 1H, H-3), 5.26 (at, $J = 10.0$ Hz, 1H, H-4), 5.22 (dd, $J =$
7
8 1.7, 3.4 Hz, 1H, H-2), 4.96 (m, 1H, $\text{CH}=\text{CH}_a\text{H}$), 4.92 (m, 1H, $\text{CH}=\text{CH}_b\text{H}$), 4.79 (d, $J = 1.7$ Hz,
9
10 1H, H-1), 4.27 (dd, $J = 5.3, 12.2$ Hz, 1H, H-6), 4.09 (dd, $J = 2.3, 12.2$ Hz, 1H, H-6'), 3.97
11
12 (ddd, $J = 2.3, 5.2, 10.0$ Hz, 1H, H-5), 3.67 (m, 1H, OCH_aH), 3.43 (dt, $J = 6.6, 9.6$ Hz, 1H,
13
14 OCH_bH), 2.14, 2.09, 2.03, 1.98, (4s, COCH_3), 1.58 (CH_2), 1.28 (CH_2). ^{13}C NMR (CDCl_3 ,
15
16 125,8 MHz) δ 170.6, 170.1, 169.9, 169.7 (COCH_3), 139.1 ($\text{CH}=\text{CH}_2$), 114.1 ($\text{CH}=\text{CH}_2$), 97.5
17
18 (C-1), 69.7 (C-2), 69.1 (C-3), 68.5 (O CH_2), 68.3 (C-5), 66.2 (C-4), 63.0 (OCH_2), 62.5 (C-6),
19
20 33.7, 32.7, 29.3, 29.2, 29.0, 28.8, 26.0 (CH_2), 20.9, 20.72, 20.67 x 2 (COCH_3). ^1H NMR data
21
22 matches data reported in the literature.²⁷ HRMS (ESI) m/z calcd. for $\text{C}_{24}\text{H}_{38}\text{NaO}_{10}$ $[\text{M}+\text{Na}]^+$:
23
24 509.2357. Found: 509.2376.
25
26
27
28
29
30
31
32
33

34 **1.3. 9-Decenyl α -D-mannopyranoside (3)**

35
36 To a solution of **8** (0.05 g, 0.1 mmol) in anhydrous 2:1 $\text{CH}_2\text{Cl}_2/\text{MeOH}$ (10 mL) at 0 °C, 1.3 M
37
38 NaOMe/MeOH (0.5 mL) was added. After 1 h of stirring at 0 °C, the mixture was
39
40 concentrated to 3 mL and deionized by elution with MeOH through a column of strongly
41
42 acidic cation exchange resin (H^+). The eluate was evaporated under reduced pressure to afford
43
44 compound **3** (0.032 g, 99 %) as a syrup, $R_f = 0.65$ (7:1:2 $n\text{PrOH}/\text{NH}_3/\text{H}_2\text{O}$), $[\alpha]_D +50$ (c 0.9,
45
46 MeOH). Lit.:²⁷ $[\alpha]_D +56$ (c 0.5, MeOH). ^1H NMR (CD_3OD , 500 MHz) δ 5.81 (ddt, $J = 6.8,$
47
48 10.2, 13.9 Hz, 1H, $\text{CH}=\text{CH}_2$), 4.98 (ddt, $J = 1.6, 2.2, 17.1$ Hz, 1H, $\text{CH}=\text{CH}_a\text{H}$), 4.91 (ddt, $J =$
49
50 1.2, 2.3, 10.2 Hz, 1H, $\text{CH}=\text{CH}_b\text{H}$), 4.73 (d, $J = 1.6$ Hz, 1H, H-1), 3.82 (dd, $J = 2.4, 11.8$ Hz,
51
52 1H, H-6), 3.78 (dd, $J = 1.7, 3.4$ Hz, 1H, H-2), 3.73 (dt, $J = 6.7, 9.6$ Hz, 1H, OCH_aH partially
53
54 overlapped with H-6'), 3.71 (dd, $J = 5.8, 11.8$ Hz, 1H, H-6'), 3.69 (dd, $J = 3.4, 9.2$ Hz, 1H, H-
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 3), 3.61 (at, $J = 9.5$ Hz, 1H, H-4), 3.52 (ddd, $J = 2.4, 5.8, 9.6$ Hz, 1H, H-5), 3.41 (dt, $J = 6.3,$
5
6 9.7 Hz, 1H, OCHH_b), 2.08–1.27 (CH₂). ¹³C NMR (CD₃OD, 125.8 MHz) δ 140.1 (CH=CH₂),
7
8 114.7 (CH=CH₂), 101.5 (C-1), 74.5 (C-5), 72.7 (C-3), 72.3 (C-2), 68.63 (OCH₂), 68.56 (C-4),
9
10 62.9 (C-6), 34.9, 30.6, 30.53, 30.50, 30.2, 30.1, 27.3 (CH₂). ¹³C NMR data matches data
11
12 reported in the literature.²⁷ HRMS (ESI) m/z calcd. for C₁₆H₃₀NaO₆ [M+Na]⁺: 341.19346.
13
14 Found: 341.19476.
15
16
17
18
19
20

21 **1.4. 9-Decenyl 2,3,5,6-tetra-*O*-*tert*-butyldimethylsilyl- α,β -D-galactofuranoside (14)**

22
23
24 A solution of **9** (0.20 g, 0.26 mmol) in anhydrous CH₂Cl₂ (10.0 mL) containing dry 4 Å
25
26 powdered molecular sieves was cooled to 0 °C and stirred during 10 min under Ar. Then,
27
28 TMSI (1.2 equiv, 0.042 mL, 0.32 mmol) was added and the solution was stirred at 0 °C until
29
30 TLC monitoring showed complete transformation of **9** into two lower moving products, the 1-
31
32 iodo intermediate **12** ($R_f = 0.70$, 10:1 hexane-EtOAc) and some 2,3,5,6-tetra-*O*-TBS- α,β -D-
33
34 galactofuranose ($R_f = 0.54$), formed as a result of the hydrolysis of **12** on the silica gel plate.¹⁶
35
36 9-Decen-1-ol (1.3 equiv, 0.34 mmol, 0.061 mL) and EtN(*i*Pr)₂ (0.054 mL, 0.32 mmol), were
37
38 added by syringe. After stirring at room temperature during 2 h the solution was diluted with
39
40 CH₂Cl₂ (250 mL), washed with NaHCO₃ (ss) (2 x 140 mL) and water (3 x 100 mL), dried
41
42 (Na₂SO₄) and concentrated. The syrup obtained was purified by column chromatography
43
44 (99.7:0.3→99.5:0.5 hexane/EtOAc) affording syrupy compound **14** (0.167 g, 83 %) as an
45
46 inseparable β/α mixture in a 3:1 ratio, which gave $R_f = 0.40$ (7:0.1 hexane/EtOAc twice
47
48 developed), $[\alpha]_D -11.7$ (c 1, CHCl₃). ¹H NMR (CDCl₃, 500 MHz) δ 5.81 (m, 1.29H, CH=CH₂
49
50 α,β), 4.99 (m, 1.29H, CH=CH_aH α,β), 4.92 (m, 1.29H, CH=CHH_b α,β), 4.84 (d, $J = 4.2$ Hz,
51
52 0.36H, H-1 α), 4.79 (d, $J = 2.6$ Hz, 1H, H-1 β), 4.20 (apparent t, $J = 5.0$ Hz, 0.36 H, H-3 α),
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 4.13 (dd, $J = 3.6, 6.0$ Hz, 1H, H-3 β), 3.98 (dd, $J = 2.5, 3.5$ Hz, 1H, H-2 β), 3.94 (dd, $J = 2.5,$
5
6 6.0 Hz, 1H, H-4 β), 3.90 (dd, $J = 4.0, 5.2$ Hz, 0.36H, H-2 α), 3.75 (m, 2.3H, H-5 β , H-4 α , H-5 α ,
7
8 $OCH_aH\alpha$), 3.67 (m, 2.6H, H-6 α,β , $OCH_aH\beta$), 3.57 (m, 1.43H, H-6' α,β), 3.35 (dt, $J = 6.7, 9.6$
9
10 Hz, 1H, $OCHH_b\beta$), 3.28 (m, 0.32H, $OCHH_b\alpha$), 2.04–1.28 (CH₂ α,β), 0.91–0.87 (SiC(CH₃)₃),
11
12 0.11–0.05 (Si(CH₃)₂). ¹³C NMR (CDCl₃, 125.8 MHz) δ 139.2 (2C, CH=CH₂ α,β), 114.1 (2C,
13
14 CH=CH₂ α,β), 108.0 (C-1 β), 102.2 (C-1 α), 84.7 (C-2 β), 83.8 (C-4 β), 79.5 (C-3 β), 78.8 (C-
15
16 2 α), 76.4 (C-3 α), 73.5 (C-5 α), 73.3 (C-5 β), 68.7 (OCH₂ α), 68.0 (OCH₂ β), 65.2 (C-6 α), 64.5
17
18 (C-6 β), 33.8, 29.7, 29.6, 29.5, 29.44, 29.42, 29.1, 29.08, 28.94, 28.92 (CH₂), 26.2–25.7
19
20 (SiC(CH₃)₃), 18.4–17.8 (SiC(CH₃)₃), –3.5–(–5.4) (Si(CH₃)₂). HRMS (ESI) m/z calcd for
21
22 C₄₀H₈₆NaO₆Si₄ [M+Na]⁺: 797.53937. Found: 797.54188.
23
24
25
26
27
28
29
30
31

32 **1.5. 9-Decenyl 2,3,5,6-tetra-*O*-benzoyl- β -D-galactofuranoside (16)**

33
34 A suspension of **11** (0.20 g, 0.31 mmol) in anhydrous CH₂Cl₂ (10.0 mL) containing dry 4 Å
35
36 powdered molecular sieves cooled to 0 °C and stirred during 10 min under Ar. TMSI (1.2
37
38 equiv, 0.048 mL, 0.37 mmol) and ZnI₂ (0.6 eq, 0.059 g, 0.18 mmol) were added and the
39
40 stirring was continued at 0 °C for 15 min and then the suspension was allowed to reach room
41
42 temperature. After 0.5 h TLC monitoring showed complete transformation of **11** ($R_f = 0.61,$
43
44 9:1 toluene-EtOAc) into a lower moving product ($R_f = 0.27$), presumable 2,3,5,6-tetra-*O*-
45
46 benzoyl-D-Galf. 9-Decen-1-ol (1.3 equiv, 0.40 mmol, 0.072 mL) was added and the stirring
47
48 was continued during 1 h. Then, the suspension was filtered and the filtrate was diluted with
49
50 CH₂Cl₂ (250 mL), washed with NaHCO₃ (ss) (2 x 140 mL) and water (3 x 100 mL), dried
51
52 (Na₂SO₄) and concentrated. After purification by column chromatography (95:5 toluene-
53
54 EtOAc) fractions of $R_f = 0.67$ (9:1 toluene-EtOAc) afforded syrupy compound **16** (0.15 g, 66
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

%), $[\alpha]_D +10.3$ (*c* 1.2, CHCl₃). For the β anomer: ¹H NMR (CDCl₃, 500 MHz) δ 8.16–7.21 (aromatic), 6.08 (m, 1H, H-5), 5.81 (m, 1H, CH=CH₂), 5.63 (d, *J* = 5.2 Hz, 1H, H-3), 5.47 (s, 1H, H-2), 5.30 (s, 1H, H-1), 5.01–4.90 (m, 2H, CH=CH₂), 4.79–4.72 (m, 2H, H-6,6'), 4.64 (m, 1H, H-4), 3.75 (m, 1H, OCH_aH), 3.54 (m, 1H, OCHH_b), 1.72–1.57 (CH₂), 1.51–1.47 (CH₂), 1.41–1.22 (CH₂). ¹³C NMR (CDCl₃, 125.8 MHz) δ 166.1, 165.7, 165.6, 165.4 (COPh), 139.1 (CH=CH₂), 133.42, 133.40, 133.29, 133.28, 133.2, 133.04, 133.03 (C-aromatic), 114.1 (CH=CH₂), 105.6 (C-1), 82.0 (C-2), 81.2 (C-4), 77.6 (C-3), 70.3 (C-5), 67.6 (OCH₂), 63.5 (C-6), 40.3, 33.7, 29.3, 29.04, 29.02, 28.9, 26.6 (CH₂). HRMS (ESI) *m/z* calcd for C₄₄H₄₆NaO₁₀ [M+Na]⁺: 757.2983. Found: 757.2999.

1.6. 9-Decenyl α,β -D-galactofuranoside (4)

1.6.1. From 14. To a solution of compound 14 (0.077 g, 0.1 mmol) in freshly distilled THF (10.0 mL) cooled at 0 °C, TBAF (0.209 g, 0.8 mmol) was added.¹⁸ The stirring was continued for 10 min at 0 °C and then at room temperature for 1 h. The solution was evaporated and the residue was purified by column chromatography (EtOAc). Fractions of *R_f* = 0.85 (7:1:2 *n*PrOH/NH₃/H₂O) gave compound 4 (0.028 g, 87 %) as β/α mixture in a 3:1 ratio, $[\alpha]_D -34.3$ (*c* 0.8, CH₃OH). ¹H NMR (CD₃OD, 500 MHz) δ 5.81 (ddt, *J* = 6.7, 10.3, 17.1 Hz, 1.23H, CH=CH₂ α,β), 4.98 (m, 2H, CH=CH₂ β), 4.91 (m, 0.89H, CH=CH₂ α), 4.85–4.83 (m, 1.44H, H-1 α,β), 4.08 (at, *J* = 7.3 Hz, 0.44H, H-3 α), 4.00 (dd, *J* = 4.0, 6.7 Hz, 1H, H-3 β), 3.94 (m, 0.44H, H-2 α), 3.93 (dd, *J* = 2.0, 4.0 Hz, 1H, H-2 β), 3.91 (dd, *J* = 3.3, 6.7 Hz, 1H, H-4 β), 3.80 (dt, *J* = 6.9, 9.6 Hz, 0.44H, OCH_aH α), 3.74–3.67 (m, 2.44H, H-4 α , H-5 β , OCH_aH β), 3.65–3.58 (m, 2.88H, H-5 α , H-6 α , H-6 β , H-6' β), 3.55 (m, 0.44H, H-6' α), 3.46 (dt, *J* = 6.7, 9.4 Hz, 0.44H, OCHH_b α), 3.41 (dt, *J* = 6.6, 9.6 Hz, 1H, OCHH_b β), 2.08–1.27 (7 CH₂). ¹³C

1
2
3
4 NMR (CD₃OD, 125.8 MHz) δ 140.1 (CH=CH₂), 114.7 (CH=CH₂), 109.4 (C-1 β), 102.8 (C-
5
6 1 α), 84.1 (C-4 β), 83.5 (C-4 α), 83.4 (C-2 β), 78.9 (C-2 α), 78.7 (C-3 β), 76.4 (C-3 α), 74.5 (C-
7
8 5 α), 72.4 (C-5 β), 69.7 (OCH₂ α), 68.9 (OCH₂ β), 64.6 (C-6 β), 64.2 (C-6 α), 34.9, 30.7, 30.6,
9
10 30.55, 30.49, 30.2, 30.1, 27.2 (CH₂). HRMS (ESI) m/z calcd for C₁₆H₃₀NaO₆ [M+Na]⁺:
11
12 341.19346. Found: 341.19470.
13
14
15
16
17
18

19 **1.6.2. From 16.** To a solution of compound **16** (0.073 g, 0.1 mmol) in anhydrous 3:2
20 CH₂Cl₂/MeOH (10 mL) at 0 °C, 1.3 M NaOMe/MeOH (0.4 mL) was added. After 1 h of
21 stirring at 0 °C, the mixture was concentrated to 4 mL and deionized by elution with MeOH
22 through a column of strongly acidic cation exchange resin (H⁺). The eluate was evaporated
23 under reduced pressure to afford compound **4** (0.030 g, 94 %) as a syrup, R_f = 0.9 (7:1:2
24 *n*PrOH/NH₃/H₂O), [α]_D -25.6 (c 1.1, MeOH). The NMR spectra were showed that compound
25
26
27
28
29
30
31
32
33
34 **4** was a mixture of anomers in 9:1 ratio.
35
36
37
38

39 **1.7. 1,2,4,6-Tetra-*O*-acetyl-3-*O*-(2,3,5,6-tetra-*O*-acetyl- β -D-galactofuranosyl)- α,β -D-**
40 **mannopyranose (17).**– To a solution of **1** (1.43 g, 4.19 mmol)¹⁹ in dry pyridine (10 mL)
41 cooled at 0°C, Ac₂O (4.74 mL, 50.26 mmol) was added dropwise, and the mixture was stirred
42 overnight at 5 °C. After cooling to 0 °C, the reaction was quenched by slow addition of water
43 (0.5 mL) and the stirring continued for 30 min at room temperature. The solution was diluted
44 with CH₂Cl₂ (250 mL) and then successively washed with HCl 10% (150 mL), NaHCO₃ ss
45 (150 mL) and water (3 × 150 mL). The organic layer was dried (Na₂SO₄), filtered and then
46 concentrated under reduced pressure. Purification of the crude mixture by column
47 chromatography (8:1 →1:1 hexane/EtOAc) afforded compound **17** (2.13 g, 75 %) as an
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 anomeric mixture in 2:1 α/β ratio, $R_f = 0.55$ (1:3 hexane/EtOAc), $[\alpha]_D -20.1$ (c 0.9, CHCl_3).
5
6
7 ^1H NMR (CDCl_3 , 500 MHz) δ 6.09 (d, 1H, $J = 2.0$ Hz, H-1 α), 5.80 (d, 0.4H, $J = 1.2$ Hz, H-
8
9 1 β), 5.51 (dd, 0.4H, $J = 1.2, 3.7$ Hz, H-2 β), 5.40–5.33 (m, 1.4H, H-5' α,β), 5.29 (m, 1H, H-
10
11 2 α), 5.27–5.17 (m, 1.4H, H-4 α,β), 5.14 (s, 1H, H-1' α), 5.11 (s, 0.4H, H-1' β), 4.97–4.95 (m,
12
13 2.8H, H-2' α,β , H-3' α,β), 4.39 (dd, 1H, $J = 4.5, 11.6$ Hz, H-6' α), 4.37–4.25 (m, 1.8H, H-
14
15 6 α,β , H-6' α), 4.20–4.12 (m, 4.2H, H-4' α,β , H-6' β , H-3 α), 4.09 (dd, 1H, $J = 2.5,$
16
17 12.2 Hz, H-6 α), 4.03–3.97 (m, 1.4H, H-3 β , H-5 α), 3.77–3.73 (m, 0.4H, H-5 β), 2.17–2.05
18
19 (CH_3CO). ^{13}C NMR (CDCl_3 , 125.8 MHz) δ 170.7, 170.4, 170.3, 169.96, 169.9, 169.2, 169.1,
20
21 169.06, 169.03 (CH_3CO), 102.5 (C-1' α), 102.2 (C-1' β), 90.9 (C-1 α), 90.8 (C-1 β), 80.8, 80.7
22
23 (C-2' α,β), 80.6 (2C, C-4' α,β), 76.4 (2C, C-3' α,β), 73.4 (C-5 β), 72.3 (C-3 β), 70.6 (C-5 α),
24
25 70.5 (C-3 α), 69.3 (2C, C-5' α,β), 66.0 (C-2 α), 65.9 (2C, C-4 α,β), 65.8 (C-2 β), 62.5 (2C, C-
26
27 6' α,β), 62.2 (2C, C-6 α,β), 20.9, 20.8, 20.79, 20.77, 20.73, 20.71, 20.67, 20.66, 20.5 (CH_3CO).
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

HRMS (ESI) calcd for $\text{C}_{28}\text{H}_{38}\text{NaO}_{19}$ $[\text{M} + \text{Na}]^+$: 701.1900. Found 701.1897.

1.8. 2,4,6-Tri-*O*-acetyl-3-*O*-(2,3,5,6-tetra-*O*-acetyl- β -D-galactofuranosyl)- α -D-mannopyranose (18).

To a stirred solution of ethylenediamine (0.025 mL, 0.38 mmol) in THF (5 ml) cooled to 0 °C glacial acetic acid (0.025 mL, 0.46 mmol) was added dropwise. Immediately, this mixture was transferred to a flask containing compound **17** (0.23 g, 0.34 mmol) and the solution was stirred for 21 h at room temperature. The mixture was diluted with CH_2Cl_2 (50 ml), washed with 5% HCl (30 ml), NaHCO_3 ss (30 ml) and water (3x 30 ml). The organic layer was dried (Na_2SO_4), filtered and evaporated under reduced pressure. Purification by column chromatography (2:3 hexane/EtOAc) gave compound **18** in 84 % yield (0.17 g), $R_f = 0.28$ (1:3

1
2
3
4 hexane/EtOAc), $[\alpha]_D -12.7$ (*c* 1, CHCl₃). ¹H NMR (CDCl₃, 500 MHz) δ 5.35 (dt, 1H, *J* = 2.9,
5
6 4.1 Hz, H-5'), 5.30 (dd, 1H, *J* = 1.9, 3.5 Hz, H-2), 5.23 (s, 1H, H-1), 5.21 (at, *J* = 9.9 Hz, 1H,
7
8 H-4), 5.12 (s, 1H, H-1'), 4.96–4.93 (m, 2H, H-2', H-3'), 4.33 (dd, 1H, *J* = 4.2, 11.9 Hz, H-
9
10 6'a), 4.25–4.08 (m, 6H, H-3, H-6a, H-6'b, H-4', H-5, H-6b). ¹³C NMR (CDCl₃, 125.8 MHz) δ
11
12 170.8, 170.5, 170.3, 169.9, 169.3, 169.28, 169.25 (CH₃CO), 102.2 (C-1'), 92.4 (C-1), 80.8 (C-
13
14 2'), 80.5 (C-4'), 76.5 (C-3'), 70.3 (C-3), 69.3 (C-5'), 68.5 (C-5), 67.5 (C-2), 66.6 (C-4), 62.6,
15
16 62.5 (C-6, C-6'), 20.9, 20.79, 20.76, 20.75, 20.7, 20.5 (CH₃CO). HRMS (ESI) calcd for
17
18 C₂₆H₃₆NaO₁₈ [M + Na] 659.1794, found 659.1776.
19
20
21
22
23
24
25
26

27 **1.9. 9-Decenyl 2,4,6-tri-*O*-acetyl-3-*O*-(2,3,5,6-tetra-*O*-acetyl- β -D-galactofuranosyl)- α,β -D-**
28 **mannopyranoside (21)**
29
30

31
32 **1.9.1. Trichloroacetimidate method.** To a stirred solution of **18** (0.16 g, 0.25 mmol) and
33
34 trichloroacetonitrile (0.174 mL, 1.75 mmol) in anhydrous CH₂Cl₂ (15 mL) cooled to 0°C,
35
36 DBU (15.5 μ L, 0.1 mmol) was slowly added. After 1 h, the solution was carefully
37
38 concentrated under reduced pressure, and the residue was purified by column chromatography
39
40 (2:3 hexane/EtOAc) to give 0.163 g (83.4 %) of the trichloroacetimidate of **20** as a syrup, *R_f* =
41
42 0.63 (1:3 hexane/EtOAc). A stirred suspension of **20** (163 mg, 0.208 mmol), 9-decen-1-ol (55
43
44 μ L, 0.313 mmol), and 4 Å powdered molecular sieves (0.5 g) in anhydrous CH₂Cl₂ (15 mL)
45
46 was cooled to –78°C, and TMSOTf (11.3 μ L, 0.062 mmol) was slowly added. After 48 h of
47
48 stirring at room temperature, the mixture was quenched by addition of NaHCO₃ ss(10 mL)
49
50 and then extracted with CH₂Cl₂. Purification by column chromatography (2:1→1:1
51
52 hexane/EtOAc), afforded syrupy **21** (0.11 g, 72 %) as an anomeric mixture in 2:3 α/β ratio, *R_f*
53
54 = 0.58 (1:3 hexane/EtOAc), $[\alpha]_D -31.9$ (*c* 1.2, CHCl₃). ¹H NMR (CDCl₃, 500 MHz) δ 5.79
55
56
57
58
59
60
61
62
63
64
65

(m, 2.5H, CH=CH₂α,β), 5.38–5.34 (m, 2.5H, H-5'α,β), 5.30 (at, *J* = 8.5 Hz, 1.5H, H-4β), 5.28 (dd, *J* = 9.0, 10.0 Hz, 1H, H-4α), 5.19 (s, 1.5H, H-1'β), 5.13 (s, 1H, H-1'α), 5.06 (dd, *J* = 2.1, 6.1 Hz, 1H, H-3'α), 5.02–4.99 (m, 4.5H, H-2'β, CH=CH_aHβ, H-3'β), 4.97 (m, 1H, CH=CH_aHα), 4.94 (dd, *J* = 0.6, 2.3 Hz, 1H, H-2'α), 4.93 (m, 1H, CH=CH_bHα), 4.92–4.90 (m, 2.5H, CH=CH_bHβ, H-1α), 4.52 (d, *J* = 1.3 Hz, 1.5H, H-1β), 4.37–4.15 (m, 11.5H, H-6'α,β, H-6α,β, H-4'α,β, H-6'β, H-6β), 4.11 (dd, *J* = 1.4, 3.1 Hz, 1.5H, H-2β), 4.08 (dd, *J* = 2.4, 12.3 Hz, 1H, H-6β), 4.04–4.00 (m, 2H, H-2α, H-3α), 3.91 (dt, *J* = 6.8, 9.5 Hz, 1.5H, OCH_aHβ), 3.87 (ddd, *J* = 2.8, 5.3, 10.3 Hz, 1H, H-5α), 3.82 (dd, *J* = 3.1, 9.0 Hz, 1.5H, H-3β), 3.65 (dt, *J* = 6.8, 9.8 Hz, 1H, OCH_aHα), 3.59 (m, 1.5H, H-5β), 3.50 (dt, *J* = 6.8, 9.5 Hz, 1.5H, OCH_bHβ), 3.42 (dt, *J* = 6.8, 9.8 Hz, 1H, OCH_bHα), 2.13–2.06 (CH₃CO, CH₂), 1.64–1.54 (CH₂), 1.40–1.27 (CH₂). ¹³C NMR (CDCl₃, 125.8 MHz) δ 170.88, 170.84, 170.5, 170.4, 170.0, 169.99, 169.97, 169.7, 169.4, 169.1 (CH₃CO), 139.2 (2C, CH=CH₂α,β), 114.2, 114.1 (CH=CH₂α,β), 102.7 (C-1'β), 102.5 (C-1'α), 99.6 (C-1β), 99.2 (C-1α), 82.4 (C-2'α), 81.6 (C-2'β), 80.5 (C-4'β), 80.1 (C-4'α), 76.1 (C-3'β), 75.7 (C-3'α), 75.0 (C-3β), 74.5 (C-3α), 72.2 (C-5β), 70.1 (OCH₂), 69.28, 69.27 (C-5'α,β), 68.3 (C-5α), 68.1 (OCH₂), 67.3 (C-2α), 66.9 (C-4β), 66.8 (C-2β), 66.4 (C-4α), 62.8 (C-6'α), 62.7 (C-6α), 62.6 (C-6β), 62.4 (C-6'β), 33.8–25.9 (CH₂), 20.9–20.6 (CH₃CO).

1.9.2. Glycosyl iodide method. A suspension of **17** (0.050 g, 0.074 mmol) and ZnI₂ (0.038 g, 0.12 mmol) in anhydrous CH₂Cl₂ (10.0 mL) was stirred at 0 °C under argon atmosphere. After 15 min TMSI (35 μL, 0.26 mmol) was slowly added and the reaction was allowed to reach room temperature. After 30 min of stirring at 45 °C TLC analysis showed total consumption

of starting material ($R_f = 0.51$, 1:3 hexane/EtOAc) and a new compound of $R_f = 0.64$ (1:3 hexane/EtOAc), presumably **19**. Powdered molecular sieves 4Å (0.5 g) and 9-decen-1-ol (0.04 mL, 0.22 mmol) were then added. After 17 h of stirring at room temperature the reaction mixture was diluted with CH_2Cl_2 (250 mL), washed with NaHCO_3 ss (2 x 140 mL) and H_2O (3 x 100 mL), dried (Na_2SO_4) and concentrated under reduced pressure. The syrup was purified by silica gel column chromatography (2:1→1:1 hexane/EtOAc) and fractions of $R_f = 0.59$ (1:3 hexane/EtOAc) afforded compound **21** (0.034 g, 61 %). By reacetylation of the partial deprotected products formed during the purification by column chromatography the yield was improved (78 %).

1.10. 9-Decenyl 3-O-(β -D-galactofuranosyl)- α,β -D-mannopyranoside (**2**)

To a solution of **21** (0.05g, 0.064 mmol) in 2:1 anhydrous $\text{CH}_2\text{Cl}_2/\text{MeOH}$ (6 mL) stirred at 0°C , 1.3 M NaOMe/MeOH (0.75 mL) was added. After 1 h the solution was deionized by elution with MeOH through a column of strongly acidic cation exchange resin (H^+). The eluate was evaporated and the residue was dissolved in water and further purified through a RP18 cartridge. Compound **2** (0.031 g, 100%) was obtained as a 2:3 α/β anomeric mixture, $R_f = 0.62$ (7:1:2 $n\text{PrOH}/\text{NH}_3/\text{H}_2\text{O}$), $[\alpha]_D -12.7$ (c 1, MeOH). $^1\text{H NMR}$ (D_2O , 500 MHz) δ 5.68 (m, 2.5H, $\text{CH}=\text{CH}_2\alpha,\beta$), 5.04 (s, 1.5H, $\text{H}-1'\beta$), 5.00 (s, 1H, $\text{H}-1'\alpha$), 4.89 (dd, 2.5H, $J = 6.5$, 16.7 Hz, $\text{CH}=\text{CH}_a\text{H } \alpha,\beta$), 4.82 (m, 2.5H, $\text{CH}=\text{CHH}_b \alpha,\beta$), 4.74 (s, 1H, $\text{H}-1\alpha$), 4.47 (s, 1.5H, $\text{H}-1\beta$), 4.076 (m, 1.5H, $\text{H}-2'\beta$), 4.05 (m, 2.5H, $\text{H}-2'\alpha$, $\text{H}-2\beta$), 4.03–3.96 (m, 6H, $\text{H}-3'\alpha,\beta$, $\text{H}-4'\alpha,\beta$, $\text{H}-2\alpha$, 3.81–3.66 (m, 11H, $\text{OCH}_a\text{H } \beta$, $\text{H}-5'\alpha,\beta$, $\text{H}-6a \alpha,\beta$, $\text{H}-6b \alpha,\beta$, $\text{H}-3\alpha$, $\text{H}-4\alpha$), 3.64–3.51 (m, 9H, $\text{OCH}_a\text{H}\alpha$, $\text{H}-6'a \alpha,\beta$, $\text{H}-6'b \alpha,\beta$, $\text{H}-3\beta$, $\text{H}-4\beta$), 3.50–3.43 (m, 2.5H, $\text{OCHH}_b\beta$, $\text{H}-5\alpha$), 3.33 (m, 1H, $\text{OCHH}_b\alpha$), 3.24 (m, 1.5H, $\text{H}-5\beta$), 1.98–1.88, 1.57–1.45,

1
2
3
4 1.34–1.13 (7 CH₂). ¹³C NMR (D₂O, 125.8 MHz) δ 139.0, 138.8 (CH=CH₂), 114.16, 114.15
5
6 (CH=CH₂), 105.4 (C-1'α), 104.3 (C-1'β), 99.9(C-1β), 99.7 (C-1α), 83.3 (C-4'α), 83.1 (C-
7
8 4'β), 81.3 (C-2'β), 81.1 (C-2'α), 77.5 (C-3β), 77.1 (2C, C-3'α,β), 76.9 (C-3α), 76.1 (C-5β),
9
10 72.5 (C-5α), 70.8, 70.7 (2C, C-5'α,β), 69.8 (OCH₂β), 67.7 (OCH₂α), 67.6 (C-2α), 67.3 (C-
11
12 2β), 64.8 (C-4β), 64.6 (C-4α), 62.8 (2C, C-6'α,β), 60.9 (2C, C-6α,β), 33.7, 33.6
13
14 (CH₂CH=CH₂α,β), 29.4–25.7 (CH₂). HRMS (ESI) calcd for C₂₂H₄₁O₁₁ [M + H]⁺ 481.26434.
15
16 Found, 481.26331.
17
18
19
20
21
22
23
24

25 Acknowledgements

26
27 The authors are indebted to the University of Buenos Aires, Agencia Nacional de
28 Promoción Científica y Tecnológica (ANPCyT) and the National Research Council of
29 Argentina (CONICET). C.M. is Research Members of CONICET, and L.B. was supported by
30 a fellowship from CONICET.
31
32
33
34
35
36
37
38

39 Supplementary data

40
41 Supplementary data associated with this article (¹H and ¹³C NMR spectra for
42 compounds **2-4**, **8**, **14-18** and **21**) can be found online version at doi:
43
44
45
46
47
48

49 References

- 50
51 1. (a) Peltier, P.; Euzen, R.; Daniellou, R.; Nugier-Chauvin, C.; Ferrières, V. *Carbohydr. Res.*
52 **2008**, *343*, 1987-1923. (b) Richards, M. R.; Lowary, T. L. *ChemBioChem.* **2009**, *10*, 1920-
53 1938. (c) Imamura, A.; Lowary, T. *Trends Glycosci. Glycotechnol.* **2011**, *23*, 134-152.
54
55
56
57
58
59
60
61
62
63
64
65

- 1
2
3
4 2. (a) Lederkremer R. M.; Colli, W. *Glycobiology* **1995**, *5*, 547-552. (b) Pedersen, L. L.;
5
6 Turco, S. J. *Cell. Mol. Life Sci.* **2003**, *60*, 259-266.
7
8
9 3. Marino, C.; Gallo-Rodriguez, C.; Lederkremer, R. M. In *Glycans: Biochemistry,*
10
11 *Characterization and Applications*, Mora-Montes, H. M. Ed. Nova Science Publisher: New
12
13 York, **2012**; pp 207-268.
14
15
16 4. Lederkremer, R. M.; Agusti, R. *Adv. Carbohydr. Chem. Biochem.* **2009**, *62*, 311-366.
17
18
19 5. Previato, J. O.; Gorin, P. A.; Mazurek, M.; Xavier, M. T.; Fournet, B.; Wieruszczk, J. M.;
20
21 Mendonça-Previato, L. *J. Biol. Chem.* **1990**, *265*, 2518-2526.
22
23
24 6. Lederkremer, R. M.; Lima, C.; Ramirez, M. I.; Ferguson, M. A. J.; Homans, S. W.;
25
26 Thomas-Oates, J. *J. Biol. Chem.* **1991**, *266*, 23670-23675.
27
28
29 7. Nogueira, N. F.; Gonzalez, M. S.; Gomes, J. E.; de Souza, W.; Garcia, E. S.; Azambuja, P.;
30
31 Nohara, L. L.; Almeida, I. C.; Zingales, B.; Colli, W. *Exp. Parasitol.* **2007**, *116*, 120-128.
32
33
34 8. (a) Turco, S. J.; Descoteaux, A. *Annu. Rev. Microbiol.* **1992**, *46*, 65-94; (b) McConville, M.
35
36 J.; Collidge, T. A.; Ferguson, M. A.; Schneider, P. *J. Biol. Chem.* **1993**, *268*, 15595-15604;
37
38 (c) McConville, M. J.; Ferguson, M. A. *J. Biochem. J.* **1993**, *294*, 305-324.
39
40
41 9. (a) Soares, R. P.; Margonari, C.; Secundino, N. C.; Macêdo, M. E.; da Costa, S. M.; Rangel,
42
43 E. F.; Pimenta, P. F.; Turco, S. J. *J. Biomed. Biotechnol.* **2010**, doi: 10.1155/2010/439174;
44
45 (b) Wilson, R.; Bates, M. D.; Dostalova, A.; Jecna, L.; Dillon, R. J.; Volf, P.; Bates, P. A.
46
47 *PLoS Negl. Trop. Dis.* **2010**, *4*(9): e816. doi: 10.1371/journal.pntd.0000816.
48
49
50 10. Kremer, L.; Dover, L. G.; Morehouse, C.; Hitchin, P.; Everett, M.; Morris, H. R.; Dell, A.;
51
52 Brennan, P. J.; McNeil, M. R.; Flaherty, C.; Duncan, K.; Besra, G. S., *J. Biol. Chem.*, **2001**,
53
54 *276*, 26430-26440.
55
56
57 11. Completo, G. C.; Lowary, T. L. *J. Org. Chem.*, **2008**, *73*, 4513-4525.
58
59
60
61
62
63
64
65

- 1
2
3
4 12. (a) Levengood, M. R.; Splain, R. A.; Kiessling, L. L. *J. Am. Chem. Soc.*, **2011**, *133*,
5 12758-12766; (b) May, J. F.; Splain, R. A.; Brotschi, C.; Kiessling, L. L. *Proc. Natl. Acad.*
6
7 *Sci. USA* **2009**, *106*, 11851-11856.
8
9
10
11 13. Gandolfi-Donadio, L.; Gallo-Rodriguez, C.; Lederkremer R. M. *J. Org. Chem.* **2002**, *67*,
12 4430-4435.
13
14
15
16 14. Ruda, K.; Lindberg, J.; Garegg, P. J.; Oscarson, S.; Konradsson, P. *J. Am. Chem. Soc.*
17 **2000**, *122*, 11067-11072.
18
19
20
21 15. (a) Gorin P.A.J.; Barreto-Bergter E. M.; Da Cruz F. S. *Carbohydr. Res.* **1981**, *88*, 177-188;
22
23 (b) Tsui, D.S.; Gorin P. A. J. *Carbohydr. Res.* **1986**, *156*, 1-8; (c) Mc Auliffe J. C.;
24
25 Hindsgaul, O. *J. Org. Chem.* **1997**, *62*, 1234-1239; (d) Johnston, B. D.; Pinto, B. M.
26
27 *Carbohydr. Res.* **1997**, *305*, 289-292.
28
29
30
31 16. Marino, C.; Chioconi, A.; Varela, O.; Lederkremer, R. M. *Carbohydr. Res.* **1998**,
32 *311*, 183-189.
33
34
35
36 17. Marino, C.; Lima, C.; Mariño, K. ; Lederkremer, R. M. *Beilstein J. Org. Chem.* **2012**, *8*,
37 2142-2148
38
39
40
41 18. Baldoni, L.; Marino, C. *J. Org. Chem.* **2009**, *74*, 1994-2003.
42
43
44 19. Baldoni, L.; Stortz, C. A.; Marino, C. *Carbohydr. Res.* **2011**, *346*, 191-196.
45
46
47 20. Baldoni, L.; Marino, C. *Carbohydr. Res.* **2012**, *362*, 70-78.
48
49
50 21. Thiem, J.; Meyer, B. *Chem. Ber.* **1980**, *113*, 3075-3085.
51
52 22. Gervay, J.; Hadd, M. J. *J. Org. Chem.*, **1997**, *62*, 6961-6967.
53
54 23. Murakami, T.; Sato, Y.; Shibakami, M. *Carbohydr. Res.*, **2008**, *343*, 1297-1308.
55
56 24. Fanzuo, K. *Carbohydr. Res.* **2007**, *342*, 345-373.
57
58 25. Ogawa, Y.; Beppu, K.; Nakabashi, S. *Carbohydr. Res.* **1981**, *93*, C6-C9.
59
60 26. Wang, W.; Kong, F. *J. Org. Chem.* **1998**, *63*, 5744-5745.
61
62
63
64
65

- 1
2
3
4 27. Nikolaev, A. V.; Rutherford, T. J.; Ferguson, M. A. J.; Brimacombe, J. S. *J. Chem. Soc.*
5
6 *Perkin Trans. 1*, **1995**, 1977-1987.
7
8
9 28. Sauvageau, J.; Foster, A. J.; Khan, A. A.; Chee, S. H.; Sims, I. M.; Timmer, M. S.;
10
11 Stocker, B. L. *ChemBioChem* **2012**, *13*, 2416-2424.
12
13
14 29. Repetto, E.; Marino, C.; Uhrig, M. L.; Varela, O. *Bioorg. Med. Chem.* **2009**, *17*, 2703-
15
16 2711.
17
18
19 30. Marino, C.; Gandolfi-Donadío, L.; Gallo-Rodriguez, C.; Bai, Y.; Lederkremer, R. M.
20
21 (2011). One-step syntheses of 1,2,3,5,6-penta-*O*-benzoyl- α,β -D-galactofuranose and
22
23 1,2,3,5-tetra-*O*-benzoyl- α,β -D-arabinofuranose. *Carbohydrate Chemistry: Proven Methods*.
24
25 Paul Kovac, Ed. CRC Press. Vol. 1, 231-238.
26
27
28 31. Mukhopadhyay, B.; Kartha, K. P. R.; Russell, D. A.; Field, R. A. *J. Org. Chem.* **2004**, *69*,
29
30 7758-7760.
31
32
33 32. Caputo, R.; Kunz, H.; Mastroianni, D.; Palumbo, G.; Pedatella, S.; Sola, F. *Eur. J. Org.*
34
35 *Chem.* **1999**, 3147-3150.
36
37
38 33. Bickley, J.; Cottrell, J. A.; Ferguson, J. R.; Field, R. A.; Harding, J. R.; Hughes, D. L.;
39
40 Kartha, K. P. R.; Law, J. L.; Scheinmann, F.; Stachulski, A. V. *Chem. Commun.* **2003**,
41
42 1266-1267.
43
44
45 34. van Well, R. M.; Kartha, K. P. R.; Field, R. A. *J. Carbohydr. Chem.* **2005**, *24*, 463-474.
46
47
48 35. Schmidt, R. R. *Angew. Chem. Int. Ed. En.* **1986**, *25*, 212-235.
49
50
51 36. (a) Gervay, J. Glycosyl Iodides in Organic Chemistry. In *Organic synthesis: Theory and*
52
53 *Applications*; JAI Press monographs series, **1998**; Vol. 4, pp 121-153. (b) Meloncelli, P. J.;
54
55 Martin, A. D.; Lowary, T. L. *Carbohydr. Res.* **2009**, *344*, 1110-1122.
56
57
58
59
60
61
62
63
64
65







Figure 1. Synthetic targets.




Scheme 1. Retrosynthetic approaches for 9-decanyl β -D-Galp-(1 \rightarrow 3)-D-Manp (**2**)



Scheme 2. Synthesis of 9-decyl α -D-mannopyranoside (**3**)




Scheme 3. Synthesis of *O*-galactofuranosides *via* in situ formed galactofuranosyl iodides.



Scheme 4. Synthesis of 9-decanyl β -D-Galf-(1 \rightarrow 3)-D-Manp (**2**).

Graphical Abstract**Synthetic tools for the characterization of galactofuranosyl transferases. Glycosylations via acylated glycosyl iodides**

Luciana Baldoni and Carla Marino*



- 9-Decenyl glycosides of D-Man_p, D-Galf and β-D-Galf-(1→3)-D-Man_p were synthesized.
- Conditions for glycosylation per-*O*-acetyl-Man_p iodide were revised and optimized.
- The established conditions were used for the glycosylation of β-D-Galf-(1→3)-D-Man_p
- Galactofuranosylation *via* per-*O*-benzoyl-β-D-Galf iodide was investigated.

ACCEPTED MANUSCRIPT