

## Accepted Manuscript



Title: Fungal root colonization of *Puccinellia frigida* (Phil.) Johnston, a dominant grass species inhabiting the margins of high-altitude hypersaline Andean wetlands

Author: <ce:author id="aut0005"> Silvani Vanesa<ce:author id="aut0010"> Rothen Carolina<ce:author id="aut0015"> Rodríguez María Alejandra<ce:author id="aut0020"> Cisneros Gabriela<ce:author id="aut0025"> Godeas Alicia<ce:author id="aut0030"> Aranda-Rickert Adriana<ce:author id="aut0035"> Fracchia Sebastián

PII: S0304-3770(13)00039-9  
DOI: <http://dx.doi.org/doi:10.1016/j.aquabot.2013.03.001>  
Reference: AQBOT 2554

To appear in: *Aquatic Botany*

Received date: 29-9-2011  
Revised date: 2-3-2013  
Accepted date: 5-3-2013

Please cite this article as: Vanesa, S., Carolina, R., Alejandra, R.M., Gabriela, C., Alicia, G., Adriana, A.-R., Sebastián, F., Fungal root colonization of *Puccinellia frigida* (Phil.) Johnston, a dominant grass species inhabiting the margins of high-altitude hypersaline Andean wetlands, *Aquatic Botany* (2013), <http://dx.doi.org/10.1016/j.aquabot.2013.03.001>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Highlights**

1-Diverse root fungal endophytes colonize the grass *Puccinellia frigida* in high Andean wetlands

2-Septate fungi and chytrids were observed in all root samples under extreme soil conditions

3-No variation in AM colonization was recorded between sampling seasons in two sites

4-In highly stressful sites AM fungi colonization was not detected

5-Soil salinity and organic matter affect septate fungi and chytrids root colonization

Accepted Manuscript

**Fungal root colonization of *Puccinellia frigida* (Phil.) Johnston, a dominant grass species inhabiting the margins of high-altitude hypersaline Andean wetlands**

Silvani Vanesa<sup>a</sup>, Rothen Carolina<sup>a</sup>, Rodríguez María Alejandra<sup>a</sup>, Cisneros Gabriela<sup>a</sup>,  
Godeas Alicia<sup>a</sup>, Aranda-Rickert Adriana<sup>b</sup>, Fracchia Sebastián<sup>b\*</sup>

<sup>a</sup>Departamento de Biodiversidad y Biología Experimental. Facultad de Ciencias Exactas y Naturales (FCEyN). Universidad de Buenos Aires (UBA). Ciudad Universitaria, 4to piso Pabellón 2. 1428. Buenos Aires. Argentina.

<sup>b</sup>Centro Regional de Investigaciones Científicas y Transferencia Tecnológica, CRILAR-CONICET. Entre Ríos y Mendoza, 5301 Anillaco, La Rioja, Argentina

\*Corresponding author:

Tel.: +54-03-827-494251

Fax: +54-03-827-494231

E-mail address: sebafrac@yahoo.com.ar

**Abstract**

1  
2  
3  
4  
5 High-altitude hypersaline Andean wetlands are considered stressful environments by the  
6  
7 prevalence of extreme abiotic conditions affecting both plant host and fungal root  
8  
9 endophytes. *Puccinellia frigida* (Phil.) Johnston, a dominant plant species inhabiting the  
10  
11 margins of these wetlands over 4000 m a.s.l. in Northwest Argentina, is frequently  
12  
13 colonized by fungal root endophytes. Here we examined the nature and dynamics of  
14  
15 fungal root colonization on this plant species in three different wetlands (six sampling  
16  
17 sites) and two growing seasons. Morphologically diverse septate fungi and chytrids  
18  
19 were observed in root samples of *P. frigida* in all sampling sites, whereas arbuscular  
20  
21 mycorrhizal fungi were found only in two sites. The level of colonization of chytrids  
22  
23 and septate fungi differed significantly both between sites and seasons. Soil organic  
24  
25 matter and electrical conductivity were the only habitat parameters significantly related  
26  
27 to the observed differences. The widespread occurrence of non-destructive fungal  
28  
29 associations suggest an important role for plant survival in extreme environments.  
30  
31  
32  
33  
34  
35  
36  
37  
38

39 **Key words:** mycorrhiza, stressful environment, salinity, Andean wetlands, *Puccinellia*  
40  
41 *frigida*  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## 1. Introduction

1  
2  
3  
4  
5 In the high Andean mountains of west and northwest Argentina, several hypersaline  
6  
7 wetlands exist over 4000 m.a.s.l.; an elevation near the limit for plant life in the cold  
8  
9 and arid central Andes (Arroyo et al., 2004; Squeo et al., 2006). These environments  
10  
11 exhibit multiple extreme variables that limit the establishment and development of plant  
12  
13 species (Zack and Wildman, 2004). These factors include high water and soil salinity,  
14  
15 high UV irradiance, soil oligotrophy, high heavy metal content, hypoxia and low  
16  
17 temperatures (Fernández Zenoff et al., 2006; Dorador et al., 2008; Dib et al., 2009).

18  
19  
20  
21  
22 Plants in extreme environments are specifically challenged to live on limited amounts of  
23  
24 nutrients and water and to counteract soil toxicity problems. High salinity and low  
25  
26 temperatures have a strong inhibitory effect on inorganic nitrogen and phosphorus  
27  
28 uptake and availability (Clarkson et al., 1992; Navarro et al., 2001; Henry and Jefferies,  
29  
30 2003). Excessive  $\text{Na}^+$  and  $\text{Cl}^-$  ions have toxic effects, including structural disruption of  
31  
32 enzymes and other macromolecules, damage to cell organelles and plasma membrane,  
33  
34 and disruption of photosynthesis, respiration and protein synthesis (Juniper and Abbott,  
35  
36 1993). Also, high heavy metal content in soil induces oxidative stress and can severely  
37  
38 damage plant cell membranes (Barceló and Poschenrieder, 1990; Polle and Rennenberg,  
39  
40 1993), whereas low soil water activity is a key limiting factor for normal plant growth  
41  
42 in hypersaline and cold environments (Morgan, 1984). Therefore, plant species that  
43  
44 inhabit these environments must have developed efficient mechanisms and strategies in  
45  
46 order to grow and survive under such conditions.  
47  
48  
49  
50  
51

52  
53 Plant symbiotic fungi are widespread and occur naturally in diverse ecosystems, from  
54  
55 deserts to rainforests, as well as in boreal and austral regions just in the limit for plant  
56  
57 growth (Brundrett, 1991; Fracchia et al., 2009). It is well known that symbiotic fungi  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

are important to the structure, function and health of plant communities, and may be responsible for the acclimation of plants to environmental stresses (Rodríguez and Redman 1997; Brundrett 2006; Porrás-Alfaro et al., 2008). Specifically, it has been demonstrated that root-fungal associations can be determinant to alleviate diverse biotic and abiotic stresses for plant growth (Rodríguez et al., 2004). Understanding the nature and dynamics of these associations could be a first approach for the management and conservation of native plants living in these extreme habitats as well as to understand their ecological interactions (Barrow et al., 1997).

In preliminary studies, we found patches of the grass *Puccinellia frigida* growing as a dominant species in the margins of several high-altitude hypersaline Andean wetlands of Northwest Argentina. We observed this species in a wide range of soil conditions, being the only plant species able to grow between salt crusts but also growing close to freshwater springs in association with other macrophytes (Nicora, 1999, Fracchia, pers. obs)

The objective of this work is to describe the nature and extent of the fungal root colonization on *P. frigida* in three high-altitude Andean wetlands of Northwest Argentina, exploring differences in the level of colonization between six sampling sites and two growing seasons. Additionally, we analysed the main soil chemical characteristics of the six sites and set up the hypothesis that the colonization level of each fungal type is influenced by the soil chemical features at each site.

## 2. Material and Methods

### 2.1. Site description

1 The three wetlands studied (Laguna Brava, Laguna Mulas Muertas and Laguna  
2 Veladero) belong to the Central Andes mountains of La Rioja Province, Northwest  
3 Argentina, and are located within the protected area of the Reserva Provincial Laguna  
4 Brava (Fig. 1). The region comprises a network of hypersaline shallow lagoons  
5 associated with small communities of *bofedales* (wet-marshy meadows; Carrizo et al.,  
6 1997). These lands correspond to the Altoandina phytogeographic region, which is  
7 characterized by altitudes over 4000 m.a.s.l. (Cabrera, 1971). The wetlands are included  
8 in the List of Wetlands of International Importance (RAMSAR). Climatic conditions are  
9 similar for the three water systems (Table 1): precipitation is scarce (annual average 200  
10 mm), falling mostly as snow during the winter (Squeo et al, 2006), and air temperatures  
11 can reach -25°C in winter and 15°C in summer (annual average 0.8°C). Strong western  
12 winds with snow storms occur along the year, but mostly during fall and winter.  
13 Maximal UV-B irradiance for a nearby Andean wetland (Laguna Azul, Catamarca  
14 Province, 4400 m.a.s.l.) has been found to reach 10.8 Wm<sup>-2</sup> for the 300 to 325 nm  
15 range, measured at noon in the austral summer (Dib et al., 2009). Soils are sandy and  
16 rocky, with low organic matter content (Table 2), and are classified as entisols (Soil  
17 Survey Staff. 2010. Keys to Soil Taxonomy, 11th ed. USDA-Natural Resources  
18 Conservation Service, Washington, DC).

19 Recent analyzes have detected high heavy metal concentrations (As:3.5, Cd:6, Cu:20,  
20 Pb:9, Cr:35, Ni:25 µg/g) in the margins soils from Laguna Brava and Mulas Muertas  
21 wetlands (Fracchia, pers. com.).

## 22 **2.2. Field sampling and plant identification**

23 Six different sampling sites were selected on the margins of the three wetlands (Fig. 1,  
24 Table 1). Sites A, B and C belong to Laguna Brava, site A has salt crusts on the surface

1 and *P. frigida* grow there in the form of compact patches (Fig. 2A); site B has *P. frigida*  
2 as the only growing plant species and site C is a small *bofedal* with several  
3 macrophytes. Site D is on the margin of the small shallow hypersaline lagoon Veladero  
4 and has only small patches of *P. frigida*. Sites E and F belong to Mulas Muertas, site E  
5 is close to a freshwater spring and *P. frigida* is found there associated with a few  
6 macrophytes whereas site F has salt crusts on the surface and very few and  
7 inconspicuous individuals of *P. frigida*. At each sampling site, six to ten *P. frigida*  
8 individuals of similar size (8-10 cm tall; 2 cm wide at the base) were randomly collected  
9 at the beginning (November 2009) and at the end (March 2010) of the growing season.  
10 Around each plant, a hole of 20 cm diameter and 15 cm deep was dug and the entire  
11 plant and soil surrounding it was collected, in order to obtain their root system as  
12 complete as possible (Fig. 2B). Individual plants were sealed in polyethylene bags and  
13 kept in a cool chamber until their analysis in the laboratory.  
14 The taxonomical identification of the plant species was based on published regional  
15 Floras (Kiesling, 2009, Zuloaga et al., 1994; Zuloaga and Morrone, 1996). Voucher  
16 specimens were deposited in the Darwinion Botanical Institute, Buenos Aires.  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40

### 41 **2.3. Root preparation and staining**

42 Root samples of each *P. frigida* individual were carefully washed in running tap water  
43 to remove soil debris, and observed with a stereo binocular microscope (Leica MZ12) to  
44 detect external hyphal cover and extraradical mycelium. About 20 to 30 healthy root  
45 segments 1.5 cm long of each plant sample were processed according to the method of  
46 Barrow (2003) for dual staining (sudan IV and trypan blue). With this method, chitin  
47 stained dark blue with trypan blue and fungal and plant lipids stained bright red with  
48 sudan IV.  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65



#### 2.4. Root colonization

About 20 to 30 stained root segments per plant individual were screened for the presence/absence of the different endophytic fungal types with a binocular microscope (Leica DMLB) at a x1000 magnification. Arbuscular micorrhiza (AM) fungi, septate fungi and Chytridiomycetes were analyzed in each root fragment. Colonization level of target fungi was obtained with the colonization frequency formula (Ai-Rong and Kai-Yun, 2007). For AM fungi, the colonization type (Paris, Arum or intermediate) was determined as defined by Dickson (2004). Root samples were placed in lactic acid and deposited at the CRILAR (Centro Regional de Investigaciones La Rioja) in Anillaco, La Rioja, Argentina.

#### 2.5. Soil chemical analysis

For soil chemical analysis, five subsamples were taken from the top 20 cm at each sampling site and bulked into a composite pooled sample of about 1 kg. The soil was analyzed as follows: % humidity was determined for 10 g soil samples by subtracting the weight of the soil dried overnight at 105°C from its fresh weight; electric conductivity (EC) and pH were determined in 1:2.5 suspension of soil in water; organic matter (OM) content followed the Walkley and Black method (Walkley and Black, 1934); available P followed the Kurtz and Bray method (Kurtz and Bray, 1945) and soluble Na<sup>+</sup> content was determined by the direct flame photometer method (Halstead and Chairen, 1950). All measurements were carried out in triplicate.

#### 2.6. Statistical analysis

1 A two-way Analysis of Covariance (ANCOVA) was carried out separately for each  
2 fungal type to test for differences in the percent of fungal colonization (dependent  
3 variable) between sites and seasons (fixed factors), and the influence of environmental  
4 soil parameters (treated as covariates) on the dependent variable. When differences were  
5 significant, means were compared with a *post hoc* Tukey's test. Simple linear regression  
6 analyses were used to determine correlations between those significant covariates and  
7 the fungal colonization percentage. Percentage values were arcsine transformed prior  
8 the analyses to normalize variability. The p-value for hypothesis tests was set at 0.05.  
9 All the analyses were carried out with the STATISTICA program.  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23

### 24 **3. Results**

#### 25 **Root colonization morphology**

26 All *P. frigida* individuals sampled in the three wetlands were highly colonized by  
27 diverse root fungal endophytes. Plant defense-related symptoms (i.e. cell wall  
28 thickening and gall formation, cell death) were observed associated with fungal  
29 structures only in a few root samples. Before root staining, solitary external hyphae  
30 were scarcely observed on the root surface of individuals from sites A, B, D and F,  
31 whereas an inconspicuous mycelial network attached to sand particles and soil  
32 aggregates was observed in the roots of individuals from sites C and E.  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45

46 The co-occurrence of septate fungi and chytrids was observed in all root samples from  
47 the six sampling sites (Table 3), , whereas AM fungal structures were only present in  
48 the roots of *P. frigida* from site C (Laguna Brava) and site E (Mulas Muertas); in both  
49 cases with low colonization levels (Table 3). In site C, two AM fungal colonization  
50 types were distinguished: 1) typical arum colonization (Fig. 3 A-B) with arbuscules,  
51 vesicles and intra- and intercellular hyphae in the cortical tissue, and 2) a fine endophyte  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 type (Fig. 3C), with a few small and round vesicles, very thin hyphae and arbuscule-like  
2 structures inside the cortical cells.  
3

4 All *P. frigida* individuals from the six sites and both sampling seasons were colonized  
5 by septate fungi of variable morphology. Septate fungi were characterized by  
6 microsclerotia of variable appearance and structure inside the root tissue (Fig. 3 D-F)  
7 and diverse hyaline and melanized hyphae (Fig. 3 G-L). Some septate hyaline hyphae  
8 colonized the cortex, intra- and intercellularly, and in some root sections, hyaline fungal  
9 structures inside the vascular tissue were observed. In this case, septate fungi were  
10 characterized by the presence of profuse lipid stained bodies and a very thin wall that  
11 was not stained with trypan blue (Fig. 3G). Also, intracellular hyaline hyphae with  
12 negative reaction to the dual staining method used were scarcely visible (Fig. 3H), but  
13 observed in the root system from almost all sampled specimens from all sampling sites.  
14 A continuum between melanized and hyaline septate hyphae, as well as between  
15 globose and thin hyaline hyphae, was commonly observed (Fig. 3I). The colonization of  
16 some septate endophyte fungi seemed to spread from one cortical cell to another by  
17 narrow penetrating hyphae (Fig. 3J). Globose and septate hyaline mycelia that stained  
18 positively with sudan IV but not with trypan blue (Fig. 3K) were also visible in  
19 sampling roots from sites A, B and C. Finally, a septate mycelium that stained partially  
20 pale blue with trypan blue and colonized the cortical cells was also observed in *P.*  
21 *frigida* individuals from all sampling sites of Brava and Mulas Muertas wetlands (Fig.  
22 3L).  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49

50 Chytridiomycetes were observed in all sampling sites and were characterized by resting  
51 sporangia of variable size and structure inside root cortical cells of *P. frigida* (Fig. 3 M-  
52 O). No evidence of tissue disruption or necrosis was observed in the roots colonized by  
53 this fungal type.  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## Root colonization dynamics

Root colonization levels by chytrids and septate fungi differed significantly between sites and seasons (Table 4). Individuals from sites C and E were more heavily colonized by both fungal types compared to the other sites (post hoc Tukey's test  $p < 0.0001$ ) with no significant differences between these both sites. During March the colonization level of both fungal types was higher compared to September. The statistical analysis showed a significant interaction term between sites and season: during March colonization was higher at sites D and E for chytrids and at sites E and F for septate fungi (Table 3).

Also for chytrids and septate fungi, soil organic matter and electrical conductivity were the only significant soil parameters that were correlated to the observed differences in their level of colonization (Table 4). Simple linear regressions showed that colonization was positively correlated with the organic matter (chytridio:  $F = 108.38$ ,  $p < 0.0001$ ,  $r = 0.87$ , septate fungi:  $F = 65.28$ ,  $p < 0.0001$ ,  $r = 0.81$ ) and negatively correlated with the electrical conductivity (chytrids:  $F = 85.87$ ,  $p < 0.0001$ ,  $r = -0.84$ , septate fungi:  $F = 41.66$ ,  $p < 0.0001$ ,  $r = -0.74$ ). For Glomeromycetes, differences between sites were significant, but not between seasons or their interaction term. None of the tested habitat parameters had a significant influence on the differences in colonization percentage for this fungal type.

## 4. Discussion

The results of our study show that *P. frigida*, a plant species that grows in the margins of high-altitude hypersaline Andean wetlands, presents widespread fungal-root colonization by septate and chytridiomycetes endophytes. In contrast, we observed low values of AM colonization: AM fungi were recorded only in two sampling sites near

1 freshwater springs, where the plants were associated with other macrophytes. In sites  
2 where *P. frigida* grew as the only plant species and under very stressful conditions (i.e.  
3 high EC and Na<sup>+</sup> content), AM fungi were not observed. It is commonly accepted that  
4 AM fungi are prevalent in mesic conditions but nearly absent in stressful environments  
5 (Barrow and Aaltonen, 2001). It is interesting to note that we observed active AM  
6 colonization with arbuscules, vesicles and hyphal coils in *P. frigida* roots, even at EC  
7 values over 50 (dS/m) (site E). AM colonization was recorded at this salinity content  
8 only in a few other surveys (Juniper and Abbott, 1993; Evelin et al., 2009). Given that is  
9 well known that the establishment of specific mycorrhizal symbiosis allows plant  
10 species to protect or alleviate soil stress conditions (Schulz, 2006), the isolation,  
11 identification and characterization of these AM fungal strains could be useful to  
12 evaluate their protective role under salinity stress.  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27

28 No fluctuations in AM colonization were recorded between sampling seasons in both  
29 sites where AM fungi were observed. Seasonal fluctuation in AM colonization has been  
30 reported to be correlated mainly with the soil moisture/dry conditions, nutrient  
31 availability and seasonal climatic differences (Schulz et al., 1999; Barrow et al., 2001;  
32 Ruotsalainen et al., 2002); all variables that did not show significant seasonal changes in  
33 the high-altitude Andean wetlands of this study.  
34  
35  
36  
37  
38  
39  
40  
41  
42

43 Diverse septate hyphae morphotypes were observed in the roots of all sampling sites.  
44 Septate fungal endophytes could have a wider adaptability to the soil conditions than  
45 AM fungi. However, colonization fluctuations were observed between sites; particularly  
46 soil salinity and organic matter seem to be determinant soil variables that condition the  
47 root colonization by these fungi. It has been suggested that, in the absence of arbuscular  
48 mycorrhizas, septate fungi might act as surrogate root mutualists in cold-stressed soils  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 (Bledsoe et al., 1990; Urcelay et al., 2011), as well as in soils with high EC or Na<sup>+</sup>  
2 content (Mandyam and Jumpponen, 2005).  
3

4 Although some of the fungal morphologies were recognizable and clearly different from  
5 each other, we could not discriminate in separate categories, mainly due to the difficult  
6 to discern the nature of some structures belonging to a particular fungal morphotype.  
7 The observed continuum of melanized and hyaline hyphae, as well as between hyaline  
8 globose and thin hyphae, and the overlap of different fungal hyphae in the same root  
9 section and even inside the same cell, were the main causes that did not allow us to  
10 discriminate between different morphotypes. It is clear that a consortium of septate  
11 fungal taxonomic identities colonizes the plant species *P. frigida*, and these different  
12 strains might be less or better adapted to specific soil conditions. Using molecular tools,  
13 Porras-Alfaro et al. (2008) identified several orders of fungal taxa colonizing the roots  
14 of the grass species *Bouteloua gracilis* in semiarid grasslands. The orders included  
15 Hypocreales, Xylariales, Sordariales, Agaricales, Pleosporales and Pezizales, all of  
16 which are taxa with septate hyphae species.  
17

18 The extraradical mycelium of root endophytic fungi regularly explores the soil  
19 surrounding the root system (Brundrett, 2006). In our study, although we observed high  
20 root fungal colonization levels, we did not detect a well-developed extraradical  
21 mycelium network. The reduction of this mycelial network was notorious at sites where  
22 the most stressful soil conditions prevail (sites A, B, D, and F). These observations  
23 suggest that soil stressful conditions could be an obstacle for the development of the  
24 extraradical phase of endophytic fungi. However, obligate fungal species are known to  
25 inhabit the roots and not to extend outside the root tissue in their whole life cycle  
26 (Lucero et al., 2006). Thus, we can not exclude that some septate fungi with these  
27 characteristics are also present in the roots of *P. frigida*.  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

Chytrid sporangia were observed in all root samples of *P. frigida* and in sites with very high soil salinity levels. In all cases there was no evidence of harm to the host plant, even in roots that were highly colonized. True symbiotic interactions between plants and chytrids are not known, and species of some chytrid genera such as *Synchytrium* and *Physoderma* have been described as plant pathogens (Laidlaw, 1985). Nonetheless, it has been suggested that these fungi could regulate fungal colonization and nutrient uptake by the host (Barrow et al, 1997; Gleason et al, 2010). The regular, non-symptomatic colonization of *P. frigida* roots by chytrids suggest that they could play a significant but unknown function in the life cycle of *P. frigida* at least in wetland ecosystems.

Future research should be directed towards the isolation and identification of fungal strains associated with the roots of *P. frigida* of natural ecosystems. Some of these isolates could be of crucial importance for the establishment and survival of the plant species growing in these extreme environments. Furthermore, considering that extreme environments are a good source of microorganisms with exceptional phenotypic and genotypic characteristics (Dib et al., 2009), the detection of beneficial fungi for plant stress alleviation could be very interesting not only for basic research purposes, but also as valuable tools for biotechnological applications.

### **Acknowledgements**

Authors are grateful to The Environmental Agency of the La Rioja Province (Secretaría de Ambiente de La Rioja) for providing the sampling permission in the Reserva Provincial Laguna Brava. This research was financed by the CONICET (Concejo Nacional de Investigaciones Científicas y Técnicas- PIP 114201001 00022)

**References**

- 1  
2 Ai-Rong L., Kai-Yun G., 2007. Mycorrhizal and dark septate endophytic fungi of  
3  
4 Pedicularis species from northwest of Yunnan Province, China. *Mycorrhiza* 17, 103–  
5  
6 109.  
7  
8  
9  
10  
11 Arroyo, M., Squeo, F., Cavieres, L., Marticorena, C., 2004. Chilenische Anden. In:  
12  
13 Burga, C.A., Kötzli, F., Grabherr, G. (Eds.), *Gebirge der Erde: Landschaft, Klima,*  
14  
15 *Pflanzenwelt.* Eugen Ulmer Gmb and Co., Stuttgart, Germany, pp. 210-219.  
16  
17  
18  
19  
20  
21 Barceló, J., Poschenrieder, C., 1990. Plant water relations as affected by heavy metal  
22  
23 stress: A review. *J. Plant Nutr.* 13, 1-37.  
24  
25  
26  
27  
28  
29 Barrow, J.R., Havstad, K.M., McCaslin, B.D., 1997. Fungal root endophytes in  
30  
31 fourwing saltbush, *Atriplex canescens*, on arid rangelands of southwestern USA. *Arid*  
32  
33 *Soil Res. Rehabil.* 11, 177-185.  
34  
35  
36  
37  
38  
39 Barrow, J.R. Aaltonen, R.E., 2001. Evaluation of the internal colonization of *Atriplex*  
40  
41 *canescens* (Pursh) Nutt. roots by dark septate fungi and the influence of host  
42  
43 physiological activity. *Mycorrhiza* 11, 199-205.  
44  
45  
46  
47  
48  
49 Barrow, J.R., 2003. Atypical morphology of dark septate fungal root endophytes of  
50  
51 *Bouteloua* in arid southwestern USA rangelands. *Mycorrhiza* 13, 239-247.  
52  
53  
54  
55  
56 Bledsoe, C., Klein, P., Bliss, L.C., 1990. A survey of mycorrhizal plants on Truelove  
57  
58 Lowland, Devon Island, N.W.T., Canada. *Can. J. Bot.* 68, 1848-1856.  
59  
60  
61  
62  
63  
64  
65



1  
2 Brundrett, M.C., 1991. Mycorrhizas in natural ecosystems. In: Macfayden, A., Begon,  
3  
4 M., Fitter, A.H. (Eds.), Advances in ecological research. Academic Press, London, pp.  
5  
6 171-313.  
7

8  
9  
10  
11 Brundrett, M.C., 2006. Understanding the roles of multifunctional mycorrhizal and  
12  
13 endophytic fungi. In: Schulz, C., Boyle, T., Sieber, B. (Eds.), Microbial root  
14  
15 endophytes. Springer, Berlin, pp. 281-298.  
16  
17

18  
19  
20  
21 Cabrera, A.L., 1971. Fitogeografía de la República Argentina. Bol. Soc. Argent. Bot.  
22  
23 14, 1-42.  
24

25  
26  
27  
28 Carrizo, R., Baldoni, A., Cavallaro, S., Dzenoletas, M.A., 1997. Estudio preliminar de  
29  
30 las características geoambientales del área de reserva Laguna Brava, provincia de La  
31  
32 Rioja. Contribuciones Científicas 9, 269-281.  
33  
34

35  
36  
37  
38 Clarkson, D.T., Jones, L.H.P., Purves, J.V., 1992. Absorption of nitrate and ammonium  
39  
40 ions by *Lolium perenne* from flowing solution cultures at low root temperatures. Plant  
41  
42 Cell Environ. 15, 99-106.  
43  
44

45  
46  
47  
48 Dib, J.R., Weiss, A., Neumann, A., Ordoñez, O., Estévez, M.C., Farías, M.E., 2009.  
49  
50 Isolation of bacteria from remote high-altitude Andean lakes able to grow in the  
51  
52 presence of antibiotics. Recent Pat. Antiinfect. Drug Discov. 4, 66-76.  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 Dickson, S. 2004. The Arum–Paris continuum of mycorrhizal symbioses. *New Phytol.*  
2 163, 187-200.  
3

4  
5  
6  
7 Dorador, C., Busekow, A., Vila, I., Imhoff, J.F., Witzel, K.P., 2008. Molecular analysis  
8 of enrichment cultures of ammonia oxidizers from the Salar de Huasco, a high altitude  
9 saline wetland in northern Chile. *Extremophiles* 12, 405-414.  
10  
11  
12

13  
14  
15  
16  
17 Evelin, H., Kapoor, R., Giri, B., 2009. Arbuscular mycorrhizal fungi in alleviation of  
18 salt stress: a review. *Ann. Bot.* 104, 1263-1280.  
19  
20  
21

22  
23  
24 Fernández Zenoff, V., Heredia, J., Ferrero, M.A., Siñeriz, F., Farías, M.E., 2006.  
25 Isolation of UV resistant bacteria from high altitude Andean Lake bacterial  
26 community. *Curr. Microbiol.* 52, 359-362.  
27  
28  
29

30  
31  
32  
33  
34 Fracchia, S., Aranda-Rickert, A., Gopar, A., Silvani, V., Fernández, L., Godeas, A.,  
35 2009. Mycorrhizal status of plant species in the Chaco Serrano Woodland from  
36 central Argentina. *Mycorrhiza* 19, 205-214.  
37  
38  
39

40  
41  
42  
43 Gleason, F.H., Schmidt, S.K., Marano, A.V., 2010. Can zoosporic true fungi grow or  
44 survive in extreme or stressful environments?. *Extremophiles* 14, 417-425.  
45  
46  
47

48  
49  
50  
51 Halstead, W. J., Chairen, B., 1950. Flame-photometer determination of sodium and  
52 potassium in soils and other siliceous materials. *Pub. Rds.* 26, 99-104.  
53  
54  
55

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
- Henry, H.A.L., Jefferies, R.L., 2003. Interactions in the uptake of amino acids, ammonium and nitrate ions in the arctic salt-marsh grass, *Puccinellia phryganodes*. *Plant Cell Environ.* 26, 419-428.
- Juniper, S., Abbott, L.K., 1993. Vesicular-arbuscular mycorrhizas and soil salinity. *Mycorrhiza* 4, 45-57.
- Kiesling, R., 2009. Flora de San Juan IV. Monocotiledóneas. Editorial Fundación Universidad Nacional de San Juan, San Juan, Argentina.
- Kurtz, L.T., Bray, R.H., 1945. Determination of total, organic and available forms of phosphorus in soils. *Soil Sci.* 59, 39-45.
- Laidlaw, W.M.R., 1985. A method for detection of the resting sporangia of potato wart disease (*Synchytrium endobioticum*) in the soil of old outbreak sites. *Potato Res.* 28, 223-232.
- Lucero, M.E., Barrow, J.R., Osuna, P., Reyes, I., 2006. Plant–fungal interactions in arid and semi-arid ecosystems: Large-scale impacts from microscale processes. *J. Arid Environ.* 65, 276-284.
- Mandyam, K., Jumpponen, A., 2005. Seeking the elusive function of the root-colonising dark septate endophytic fungi. *Stud. Mycol.* 53, 173-189.



- 1  
2  
3  
4  
5  
6  
7 Ruotsalainen, A.L., Väre, H., Vestberg, M., 2002. Seasonality of root fungal  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
- colonization in low-alpine herbs. *Mycorrhiza* 12, 29-36.
- Schulz, B., Ak, R., Dammann, U., Aust, H.J., Strack, D., 1999. Fungal symbiosis: from mutualism to parasitism, who controls the outcome, host or invader?. *New Phytol.* 151, 705-716.
- Schulz, B., 2006. Mutualistic interactions with fungal root endophytes. In: *Microbial Root Endophytes*. Springer Verlag, Berlin, pp. 261-279.
- Squeo, F., Warner, B.R., Aravena, D., Espinoza, D., 2006. Bofedales: high altitude peatlands of the central Andes. *Rev. Soc. Chil. Hist. Nat.* 79, 245–255.
- Urcelay, C., Acho, A., Joffre, R., 2011. Fungal root symbionts and their relationship with fine root proportion in native plants from the Bolivian Andean highlands above 3,700 m elevation. *Mycorrhiza* 21, 323-330.
- Walkley, A., Black, I.A., 1934. An examination of Degtjareff method for determining soil organic matter and a proposed modification of the chromic acid titration method. *Soil Sci.* 37, 29-37.
- Zack, J.C., Wildman, H.G., 2004. Fungi in stressful environments. In: Mueller, G.M., Bills, G.F., Foster, M.S. (Eds.), *Biodiversity of fungi, inventory and monitoring methods*. Elsevier Academic Press, Amsterdam, pp. 303–315.

1 Zuloaga, F.O., Nicora, E.G., Rúgolo de Agrasar, Z.E., Morrone, O., Pensiero, J.,  
2 Cialdella, A.M., 1994. Catálogo de la familia Poaceae en la República Argentina.  
3 Missouri Botanical Garden, Missouri, USA.  
4  
5  
6  
7  
8

9 Zuloaga, F.O., Morrone, O., 1996. Catálogo de plantas vasculares de la Republica  
10 Argentina. I. Pteridophyta. Gymnospermae y Angiospermae (Monocotyledonea).  
11 Missouri Botanical Garden, Missouri, USA.  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25

## 26 **Figure legends**


27  
28  
29 **Fig. 1.** Locations of the three high-altitude Andean wetlands and the six sampling sites  
30 (A-F)  
31  
32  
33  
34  
35

36 **Fig. 2** A, view of Laguna Brava at sampling site A with patches of *P. frigida* (arrow); B,  
37 whole sample of *P. frigida* from site C  
38  
39  
40  
41  
42

43 **Fig 3.** *P. frigida* root fungal structures observed under binocular microscope after dual  
44 staining with tripan blue and sudan IV. A-C Arbuscular mycorrhizal structures in root  
45 samples from site C (A: arum type colonization with arbuscules and hyphae, B:  
46 arbuscule, C: hyphae of a fine endophyte); D-F Microsclerotia of different size and  
47 morphology (D: laxus and melanized microsclerotium structure, E: compact and  
48 melanized microsclerotium, F: non melanized microsclerotium with lipid content); G-L  
49 Septate fungi hyphae (G: hyaline hyphae inside the vascular tissue with positive stained  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65


1 lipids, H: hyaline hyphae inside cortical cells with no reaction to the dual staining  
2 method, I: a continuum between melanized and hyaline septate hyphae, J: cell to cell  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

lipids in cortical tissue, L: Intracellular hyphae partially stained with tripan blue); M-O  
Chytridiomycetes sporangia of variable size and morphology in root cortical cells. Bar:  
25µm.









Manuscript



Manuscript







**Table 1.** Sampling sites for *P. frigida*, associated plant species, water salinity (means  $\pm$  SD) and geographical position in three high-altitude Andean wetlands of Northwest Argentina.

Wetland*	Sampling site	Associate plant species	Total dissolved salts (g L <sup>-1</sup> )	Altitude (masl)	Global position							
Brava (5100)	A	--	188 ( $\pm$ 55)	4260	28° 18' 29" S 68° 49' 14" W							
	B	--			28° 16' 27" S 68° 49' 15" W							
	C	<i>Deyeuxia curvula</i> <i>Zameioscirpus atacamensis</i> <i>Ruppia sp.</i> <i>Patosia clandestina</i> <i>Nastanthus caespitosus</i> <i>Ranunculus exilis</i> <i>Calandrinia compacta</i>			188 ( $\pm$ 55)	4260	28° 16' 49" S 68° 50' 23" W					
							Veladero (12)	D	--	191 ( $\pm$ 13)	4327	28° 11' 19" S 69° 00' 25" W
							Mulas Muertas (210)	E	<i>Zameioscirpus atacamensis</i> <i>Deyeuxia curvula</i> <i>Ranunculus exilis</i>	160 ( $\pm$ 39)	4123	28° 17' 24" S 68° 44' 03" W
								F	--			28° 16' 56" S 68° 43' 51" W

\*wetland surface in ha.

**Table 2:** Main soil chemical characteristics at six sampling sites in three high-altitude Andean wetlands of Northwest Argentina

Wetland	Brava			Veladero	Mulas Muertas	
Sampling site	A	B	C	D	E	F
<b>Soil Parameter</b>						
Moisture (%)**	26.1	25.3	35.7	33.1	29.9	25.5
pH**	9.1	9.5	9.0	9.9	8.7	8.0
EC (dS/m)**	141	96	37	112	56	117
OM (%)*	0.43	0.55	2.45	0.32	1,01	0.34
Available P (ppm)*	13.8	15.0	29.5	10.9	19.3	20.8
Na <sup>+</sup> (meq/l) *	459	400	347	433	398	400

\*determined in march

\*\*mean of both sampling seasons

**Table 3:** Mean frequency colonization (F%) of glomeromycetes, septate fungi and chytridiomycetes in *Puccinellia frigida* roots at six sampling sites in three high-altitude Andean wetlands of Northwest Argentina

Fungal type	Glomeromycetes		Septate fungi		Chytridiomycetes	
	November	March	November	March	November	March
Sampling site						
Brava						
A	0	0	48.6±9.0	46.0±11.3	35.0±9.1	35.6±11.0
B	0	0	40.6±21.3	45.0±7.5	32.0±3.6	31.0±20.8
C	3.0±2.6	4.3±2.0	86.3±14.2	100±0.0	89.6±3.5	97.3±4.6
Veladero						
D	0	0	40.0±14.4	44.6±8.0	19.3±4.7	33.3±8.3
Mulas Muertas						
E	5.3±4.0	2.0±1.7	75.3±7.6	98.3±2.8	68.6±7.0	90.3±3.5
F	0	0	25.3±4.5	50.0±14.9	18.3±4.0	25.3±5.0
Mean <sup>1</sup>	1.3±2.7	1.0±1.9	52.7±24.4	64.0±26.7	44.0±27.4	52.1±31.8
Fungal mean	1.2±2.3		58.3±25.9		48.0±29.5	

Values are means ± SD (n = 6-10). <sup>1</sup> Mean for each fungal type at each sampling season

**Table 4:** Two way ANCOVA for each fungal type with Season (March and September) and Sites (A-F) as categorical factors, frequency of colonization as dependent factor, and habitat variables as covariates. P-values are considered significant at the 0.05 level.

	df	Chytrids		Septate fungi		Glomeromycetes	
		F	p	F	p	F	p
Site	5	5.76	<b>0.0023</b>	18.56	<b>&lt;0.0001</b>	12.24	<b>0.0001</b>
Season	1	28.86	<b>&lt;0.0001</b>	11.39	<b>0.0033</b>	0.91	0.35
Site x season	5	14.12	<b>&lt;0.0001</b>	5.25	<b>0.0181</b>	1.83	0.15
OM	1	4.88	<b>0.0403</b>	4.95	<b>0.0345</b>	0.83	0.37
EC	1	4.61	<b>0.0456</b>	4.78	<b>0.0417</b>	0.27	0.61
Moisture	1	2.86	0.11	0.004	0.94	0.3	0.59
pH	1	0.01	0.91	0.01	0.89	1.08	0.31
Available P	1	0.86	0.36	0.78	0.38	0.95	0.34
Na	1	0.04	0.83	0.005	0.94	0.28	0.59
Error	18						

(When the Interaction is significant, means that the season has a different influence at each site on the percent of colonization by each fungal type.)