

Supporting Information

Synthesis and behavior of novel sulfonated water-soluble N-heterocyclic carbene (η^4 -diene) platinum(0) complexes.

Andrea M. Ruiz-Varilla, Edwin A. Baquero, Gustavo F. Silbestri, Camino González-Arellano, Ernesto de Jesús and Juan C. Flores**

Departamento de Química Orgánica y Química Inorgánica, Universidad de Alcalá,
Campus Universitario, 28871 Alcalá de Henares, Madrid (Spain).

camino.gonzalez@uah.es, juanc.flores@uah.es

Contents

NMR and mass spectra of complex 3d	S2
NMR and mass spectra of complex 3e	S4

NMR and mass spectra of compound 3d




Figure S1. ^1H NMR (300 MHz) spectrum of **3d** in D_2O . (Impurities denoted with *)






Figure S2. ^1H NMR (300 MHz) spectrum of **3d** in dms0-d_6 . (Impurities denoted with *)






Figure S5. Calculated and experimental isotopic pattern for the dianion $[3d-2Na]^{2-}$ detected as the most intense signal in the ESI-TOF mass spectra of **3d** in MeOH.

NMR and mass spectra of compound **3e**




Figure S6. 1H NMR (500 MHz) spectrum of **3e** in D_2O . (Impurities denoted with *)




Figure S7. ¹³C NMR (125 MHz) spectrum of **3e** in D₂O. (Impurities denoted with *)




Figure S8. ESI-TOF (negative mode, H₂O) for complex **3e**




Figure S9. Calculated and experimental isotopic pattern for the dianion $[3e-2Na]^{2-}$ detected as the most intense signal in the ESI-TOF mass spectra of **3e** in H_2O .