

# Spectroscopic study of early-type multiple stellar systems

## II. New binary subsystems

M. E. Veramendi and J. F. González

Instituto de Ciencias Astronómicas, de la Tierra y del Espacio (CONICET-UNSJ), Casilla de Correo 49, 5400 San Juan, Argentina  
e-mail: [mveramendi;fgonzalez]@icate-conicet.gob.ar

Received 28 February 2014 / Accepted 5 May 2014

### ABSTRACT

**Context.** This work is part of a long-term spectroscopic study of a sample of 30 multiple stars with early-type components. In this second paper we present the results for six multiple systems in which new stellar components have been detected.

**Aims.** The main aim is to increase the knowledge of stellar properties and dynamical structure of early-type multiple stellar systems.

**Methods.** Using spectroscopic observations taken over a time baseline of more than five years we measured radial velocities by cross-correlations and spectrally separated the double-lined systems. In addition to the discovery of objects with double-lined spectra, the existence of new spectroscopic subsystems were inferred from the radial velocity variations of single-lined components and through the variation of the barycentric velocity of double-lined subsystems. Orbital elements were calculated when possible.

**Results.** Seven new stellar components and two members that we expect to confirm with new observations have been discovered in the six studied multiples. We present orbital parameters for two double-lined binaries and preliminary orbits for three single-lined spectroscopic binaries. Five of the six analysed systems are quadruples, while the remaining has five components distributed on four hierarchical levels. These multiplicity orders are in fact lower limits, since these systems lack high-resolution visual observations and additional hierarchical levels might exist in that separation range.

**Conclusions.** The six analysed systems have a higher degree of multiplicity and a more complex hierarchical structure than previously known, which suggests that high-order multiple systems are significantly more frequent than currently estimated. The long-term spectroscopic monitoring of multiple systems has shown to be useful for the detection of companions in intermediate hierarchical levels.

**Key words.** techniques: radial velocities – binaries: spectroscopic – stars: early-type

## 1. Introduction

The observed properties of multiple stellar systems (three or more components) are empirical evidence that is particularly useful to contrast theoretical models of stellar formation and evolution. However, the observational results published in the literature on systems with early-type components are insufficient, both in quantity and detail, to establish overall conclusions on the frequency and properties of this type of systems (see [Veramendi & González 2014](#), hereafter Paper I, and references therein). For that reason, we carried out a spectroscopic investigation of a sample of 30 systems with catalogued components with types O, B, and A, selected from the Multiple Star Catalogue (MSC; [Tokovinin 1997](#)). In Paper I we explained in detail the selection criteria and presented the full list of the analysed systems together with the information given in the MSC. In that paper we described the analysis and results obtained for six multiples for which we have determined orbital parameters for the first time or have corrected their previously published orbits. In this second paper we present the study and the results obtained for six systems in which we detected new components.

In Sect. 2 we briefly outline the spectroscopic observations and radial velocity (RV) measurements, which were described in detail in Paper I. The analysis and results for the six systems studied in the present paper are presented in Sect. 3. In Sect. 4 we discuss the contribution of this work to the knowledge of the hierarchical structure of the studied systems. We summarize our main conclusions in Sect. 5.

## 2. Observations and data analysis

The spectroscopic observations were carried out using the 2.15 m telescope and the REOSC echelle spectrograph at Complejo Astronómico El Leoncito (CASLEO), San Juan, Argentina. In addition to the ten observing runs mentioned in Paper I, some systems presented in this paper were recently observed during three nights in July 2013. The spectra cover the wavelength range 3700–6300 Å with a resolving power  $R = 13\,300$ , and were reduced by using standard data-reduction procedures within the NOAO/IRAF package.

Spectral types were determined by comparison with standard stars observed at the same spectral resolution. The procedure for measuring RVs has been described in Paper I. Basically, we used cross-correlations and, in the case of double-lined spectroscopic binaries, the spectral separation method by [González & Levato \(2006\)](#).

Based on the spectral morphology and the analysis of measured RVs, we evaluated the multiplicity of the observed stellar components. More specifically, a component with a single-lined spectrum was classified as RV variable if the rms deviation of the measured velocities was at least twice as large as the mean error of the measurements. When it was possible to determine the period of the variation and compute a preliminary orbit, the object was classified as single-lined spectroscopic binary (SB1). The observed components with rms deviation lower than twice the mean measurement error were considered RV constants on the time baseline of our observations. Finally,

**Table 1.** Components of multiple systems.

System	Comp.	HD	ST <sub>pub</sub>	ST	Result
08079–6837	Aa+Ab	68 520 (AB)	B6IV	B6 IV+B8	Orbit SB2 New SB2
	Ba+Bb		0.06	A2 V+A2 V	
08314–3904	Aa	72 436 (AB)	B4V	B4 V	New SB1 with preliminary orbit New RV variable RV constant, probably non-member
	B		–0.13	B6 V	
	C			F0 V	
10209–5603	A	89 890	B3IIIe	B3 III	RV constant (line profile variable) New SB2 (barycentric RV variation?) RV constant
	Ba+Bb		B9IVpSi	A0 IVpSi+A2	
	C			K0 III	
13226–6059	AB	116 087	B3V+...	B3 V	RV constant
	Ca	116 072	B2V	B2 V	New SB1 with preliminary orbit
15185–4753	Ca	135 748	0.13	A2 V	New SB1 with preliminary orbit
20118–6337	Aa+Ab+B	191 056	A0V+...	A1 V+A1 V	New SB2 (Aab) with orbit RV constant
	C		0.09	A1 V	

components whose spectra showed lines belonging to two stars with RVs that are linearly correlated with each other (consistent with orbital motion) were classified as double-lined spectroscopic binaries (SB2).

In new spectroscopic binaries we determined a preliminary orbital period using the *phase dispersion minimization* technique (Stellingwerf 1978) as implemented in the IRAF task *pdm*, which is suitable for irregularly spaced data. We then applied the least-squares method to determine the elements of a Keplerian orbit, including recalculation of the period. Measurement errors were used to weight the data points. The parameter errors were calculated from the rms of the residuals. In general, we fitted the orbital period ( $P$ ), time of primary conjunction ( $T_0$ ), eccentricity ( $e$ ), longitude of periastron ( $\omega$ ), semi-amplitude of RV curve ( $K$ ), and barycentre velocity ( $V_\gamma$ ).

### 3. Results

From the analysis of our spectroscopic observations and measured RVs, we concluded that some of the studied systems are formed by more components than the previously known. In some cases the new components were visible in the spectra, while in others their existence was detected through RV variations. In particular, we present here six multiple systems classified as triples in MSC, in which we detected new components. Each of these is discussed in detail below. Table 1 summarizes the results obtained in each case. The observed components are listed in Col. 2 and are identified by their HD number in Col. 3. For each component, Col. 4 lists the published spectral type or, if not available, the  $B-V$  colour index, and Col. 5 presents the spectral type determined in this work. Finally, a brief description of the results is given in Col. 6.

The measured RVs of the 13 observed components in the six systems are listed in Tables 2 to 5, which correspond to RV constant stars, RV variables, single-lined, and double-lined spectroscopic binaries, respectively.

#### 3.1. WDS 08079–6837

In this system, the components A and B have a separation of 6.1'' and a magnitude difference of  $\Delta V = 2.93$ . Star A (HD 68520) is catalogued as a single-lined spectroscopic binary, for which Sanford (1915) computed an orbit with a period of 14.1683 days.

**Table 2.** Radial velocities of constant stars.

WDS	Subsystem	HJD	RV km s <sup>-1</sup>	Err km s <sup>-1</sup>
08314-3904	C	2 454 545.7377	–13.4	0.6
08314-3904	C	2 454 757.8605	–13.1	0.4
08314-3904	C	2 454 911.6648	–13.7	0.4
10209-5603	C	2 454 522.7974	6.5	0.8
10209-5603	C	2 454 908.7462	5.8	0.5
10209-5603	C	2 454 912.7841	5.5	0.5
10209-5603	C	2 454 960.5717	5.7	0.5
10209-5603	C	2 454 963.6431	7.2	0.7
10209-5603	C	2 455 288.6694	5.6	0.4
13226-6059	AB	2 454 518.7815	–6.0	9.1
13226-6059	AB	2 454 908.7748	–0.4	10.9
13226-6059	AB	2 454 910.8490	–3.4	8.2
13226-6059	AB	2 454 960.6387	–5.3	9.1
13226-6059	AB	2 454 962.6180	–2.7	8.2
13226-6059	AB	2 456 486.4934	–1.3	7.9
13226-6059	AB	2 456 487.4744	4.5	8.1
20118-6337	C	2 454 572.8787	–10.3	5.3
20118-6337	C	2 454 755.5929	–6.2	6.1
20118-6337	C	2 454 961.8300	–2.6	5.7
20118-6337	C	2 454 962.8404	–13.0	5.7
20118-6337	C	2 454 963.8644	–1.9	5.2
20118-6337	C	2 455 084.6463	–14.1	6.2
20118-6337	C	2 456 486.6993	–10.4	5.8
20118-6337	C	2 456 486.8256	–4.6	5.1
20118-6337	C	2 456 487.6327	–7.3	5.8
20118-6337	C	2 456 487.7556	–9.9	4.9
20118-6337	C	2 456 488.6383	–7.8	5.1
20118-6337	C	2 456 488.7647	–9.3	6.0

Medici & Hubrig (2000) detected the secondary component of this subsystem in some observations and determined a mass ratio  $M_1/M_2 = 1.30$ . Recently, Chini et al. (2012) obtained two spectra of this binary, confirming its status as SB2.

The subsystem A was observed frequently, with the aim of detecting spectral features of the secondary and computing orbital parameters for both components. In RV measurements we employed synthetic templates with  $T_{\text{eff}} = 15\,000$  K and  $T_{\text{eff}} = 13\,000$  K. We obtained 26 spectra of this binary. We reconstructed the spectra of the components combining 24 observed

**Table 3.** Radial velocities of variable stars.

WDS	Subsystem	HJD	Phase	RV km s <sup>-1</sup>	Err km s <sup>-1</sup>
08314-3904	B	2 454 520.6609		16.7	1.9
08314-3904	B	2 454 545.6802		10.3	3.3
08314-3904	B	2 454 757.8324		-1.3	1.7
08314-3904	B	2 454 908.6405		23.1	2.4
08314-3904	B	2 454 911.6338		19.9	1.5
08314-3904	B	2 454 960.5078		-2.8	3.4
08314-3904	B	2 455 289.5845		34.5	3.2
08314-3904	B	2 456 336.7409		24.7	4.7
08314-3904	B	2 456 337.5494		12.0	4.3
08314-3904	B	2 456 338.7387		15.3	5.1
10209-5603	A	2 454 519.7156		14.1	3.0
10209-5603	A	2 454 544.7909		13.0	3.3
10209-5603	A	2 454 908.7114		12.6	3.2
10209-5603	A	2 454 911.7532		10.4	3.0
10209-5603	A	2 454 960.5315		12.1	3.1
10209-5603	A	2 454 962.5775		13.4	3.0
10209-5603	A	2 455 288.6449		15.9	3.2
10209-5603	A	2 455 291.7027		11.6	3.3
10209-5603	A	2 456 487.4665		10.6	3.0

spectra, since in the other two the spectral lines of both components were too blended to measure preliminary RVs. However, in a subsequent calculation with the method GL06, we measured RVs for both components even in the two heavily blended spectra. The reconstructed spectra of the components led to a confirmation of the spectral type B6IV published in the MSC for the primary and to an estimation of a spectral type B8 for the secondary.

The period by Sanford (1915) represents our RV data well, therefore we took it as the starting value of the period for the orbital element computation. With all seven parameters free, the orbital fitting gave an eccentricity indistinguishable from zero. Therefore, we adopted a circular orbit and determined final values for the other orbital elements (except for  $\omega$ ). These parameters are listed in Table 6, which also includes the number of spectra ( $N_{sp}$ ) and rms deviation of the velocities ( $\sigma_{RVa}$  and  $\sigma_{RVb}$ ).

We obtained a mass ratio  $q = 0.67$ , lower than that given by Medici & Hubrig ( $q = 0.77$ ) and higher than the value estimated by Sanford ( $q = 0.23$ ). However, the parameters found in this study and those calculated by Sanford for the orbit of the primary component Aa agree well. Figure 1 presents the fitting of the RVs of both components.

It was difficult to observe component B of this multiple system because of the proximity of star A, which is about 15 times brighter. Thus, we only obtained six useful spectra of B, which showed morphological variations indicating two spectral components. We applied cross-correlations with an observed template of spectral type A1V. In only four spectra we measured RVs of both components separately. In the other two we obtained only one broadened peak in the cross-correlation function. There is a linear correlation between the RVs measured for both components, which confirms that they form a double-lined spectroscopic binary. The linear fitting shown in Fig. 2 leads to a mass ratio  $q = 0.93 \pm 0.03$  and a barycentre velocity  $V_\gamma = 2.7 \pm 0.7$  km s<sup>-1</sup>. The few observations do not allow us to derive the orbital period, but given the RV variations detected within the same observing run, it probably is of about a few days.

In conclusion, we found that component B is a binary subsystem and obtained the orbital parameters of Aab through

**Table 4.** Radial velocities of single-lined spectroscopic binaries.

WDS	Subsystem	HJD	Phase	RV km s <sup>-1</sup>	Err km s <sup>-1</sup>
08314-3904	Aa	2 454 519.6730	0.0253	4.5	1.4
08314-3904	Aa	2 454 545.6685	0.3801	4.3	1.4
08314-3904	Aa	2 454 757.8231	0.2761	-17.7	1.4
08314-3904	Aa	2 454 908.6304	0.3346	-2.9	1.4
08314-3904	Aa	2 454 911.6235	0.3755	3.2	1.4
08314-3904	Aa	2 454 960.4976	0.0426	3.4	1.4
08314-3904	Aa	2 455 288.5632	0.5208	32.4	1.4
08314-3904	Aa	2 455 291.6589	0.5631	35.9	1.4
08314-3904	Aa	2 456 336.7289	0.8285	35.5	1.4
08314-3904	Aa	2 456 337.5401	0.8396	33.2	1.4
08314-3904	Aa	2 456 338.7264	0.8558	34.7	1.4
08314-3904	Aa	2 456 696.5787	0.7406	43.52	1.4
13226-6059	Ca	2 454 518.7860	0.1221	-34.2	7.7
13226-6059	Ca	2 454 544.8437	0.8859	-0.3	9.6
13226-6059	Ca	2 454 908.7781	0.3526	-32.8	9.0
13226-6059	Ca	2 454 910.8519	0.8909	14.8	7.4
13226-6059	Ca	2 454 960.6417	0.8148	7.5	7.2
13226-6059	Ca	2 454 962.6213	0.3287	-30.6	8.7
13226-6059	Ca	2 455 288.7134	0.9725	-6.2	8.4
13226-6059	Ca	2 455 289.7776	0.2488	-23.7	7.0
13226-6059	Ca	2 455 291.7775	0.7679	9.9	9.9
13226-6059	Ca	2 456 336.8834	0.0465	-13.4	7.1
13226-6059	Ca	2 456 337.8781	0.3047	-30.9	8.9
13226-6059	Ca	2 456 338.8414	0.5548	-21.4	8.8
13226-6059	Ca	2 456 486.4971	0.8818	-3.6	6.8
13226-6059	Ca	2 456 486.5893	0.9057	1.1	7.3
13226-6059	Ca	2 456 487.4777	0.1363	-28.3	8.6
13226-6059	Ca	2 456 487.5907	0.1657	-29.1	8.9
13226-6059	Ca	2 456 488.5182	0.4064	-36.7	6.8
13226-6059	Ca	2 456 488.5905	0.4252	-31.1	7.2
15185-4753	Ca	2 454 519.8976	0.7531	8.5	1.0
15185-4753	Ca	2 454 520.8523	0.8303	10.3	0.6
15185-4753	Ca	2 454 544.8646	0.7739	9.6	0.7
15185-4753	Ca	2 454 908.8330	0.2341	4.8	0.6
15185-4753	Ca	2 454 911.7887	0.4734	4.7	0.8
15185-4753	Ca	2 454 960.7024	0.4325	5.9	0.6
15185-4753	Ca	2 454 962.6509	0.5902	6.3	0.7
15185-4753	Ca	2 455 083.5063	0.3725	5.3	0.8
15185-4753	Ca	2 455 084.5086	0.4536	5.8	0.7
15185-4753	Ca	2 455 086.4888	0.6139	5.1	0.7
15185-4753	Ca	2 455 087.4896	0.6949	7.6	0.6
15185-4753	Ca	2 456 486.5453	0.9368	7.3	0.6
15185-4753	Ca	2 456 486.6458	0.9449	8.7	0.7
15185-4753	Ca	2 456 487.5409	0.0174	7.6	0.7
15185-4753	Ca	2 456 487.6587	0.0269	6.8	0.8
15185-4753	Ca	2 456 488.5337	0.0977	7.5	0.6
15185-4753	Ca	2 456 488.6623	0.1081	7.5	0.7

fitting the RVs of its two components. Furthermore, the velocities obtained for the barycentres of both subsystems agree well, which confirms that they are physically bound to each other. Therefore, the system WDS 08079–6837, previously classified as a triple system, is in fact a quadruple.

### 3.2. WDS 08314–3904

This system is formed by components A and B separated by 4.47", and a third component C at 30.8" from AB. In the MSC, the physical link between the components has been established on the basis of the criterion of hypothetical dynamical parallax and the similarity between the distances estimated from the spectral type and apparent magnitude.

**Table 5.** Radial velocities of double-lined spectroscopic binaries.




WDS	Subsystem	HJD	Phase	$RV_A$ km s <sup>-1</sup>	$Err_A$ km s <sup>-1</sup>	$RV_B$ km s <sup>-1</sup>	$Err_B$ km s <sup>-1</sup>
08079-6837	Aab	2 454 519.7231	0.3456	-49.8	1.5	88.6	2.0
08079-6837	Aab	2 454 520.6751	0.4128	-28.7	1.5	59.6	2.0
08079-6837	Aab	2 454 544.5800	0.0999	-32.5	1.5	70.8	2.0
08079-6837	Aab	2 454 545.6468	0.1752	-53.3	1.5	97.8	2.0
08079-6837	Aab	2 454 546.6365	0.2450	-61.9	1.5	105.9	2.0
08079-6837	Aab	2 454 572.6090	0.0780	-27.0	1.5	56.9	2.0
08079-6837	Aab	2 454 573.5823	0.1466	-47.8	1.5	83.1	2.0
08079-6837	Aab	2 454 574.5629	0.2158	-58.6	1.5	97.2	2.0
08079-6837	Aab	2 454 576.6181	0.3609	-44.5	1.5	85.5	2.0
08079-6837	Aab	2 454 577.6017	0.4303	-23.1	1.5	54.9	2.0
08079-6837	Aab	2 454 579.5294	0.5663	32.4	1.5	-34.7	2.0
08079-6837	Aab	2 454 908.6100	0.7905	66.7	1.5	-95.5	2.0
08079-6837	Aab	2 454 909.6216	0.8619	56.0	1.5	-72.0	2.0
08079-6837	Aab	2 454 910.6880	0.9371	29.3	1.5	-33.2	2.0
08079-6837	Aab	2 454 911.5975	0.0013	6.5	2.5	5.89	3.0
08079-6837	Aab	2 454 912.6115	0.0729	-27.8	1.5	48.7	2.0
08079-6837	Aab	2 454 960.4905	0.4518	-11.7	1.5	38.8	2.0
08079-6837	Aab	2 454 961.4652	0.5206	16.2	2.5	-5.1	3.0
08079-6837	Aab	2 454 962.5361	0.5962	45.5	1.5	-48.1	2.0
08079-6837	Aab	2 454 963.5350	0.6667	66.6	1.5	-77.8	2.0
08079-6837	Aab	2 455 288.5548	0.6043	48.9	1.5	-53.8	2.0
08079-6837	Aab	2 455 289.5474	0.6743	63.8	1.5	-80.2	2.0
08079-6837	Aab	2 455 291.6380	0.8218	64.0	1.5	-82.0	2.0
08079-6837	Aab	2 455 529.7698	0.6275	55.2	1.5	-67.3	2.0
08079-6837	Aab	2 456 336.5230	0.5623	32.8	1.5	-30.7	2.0
08079-6837	Aab	2 456 338.6493	0.7124	71.3	1.5	-88.7	2.0
08079-6837	Bab	2 454 544.5861		47.7	2.4	-45.3	2.5
08079-6837	Bab	2 454 908.6159		-52.4	1.1	62.2	1.2
08079-6837	Bab	2 454 912.6190		-13.3	2.1	19.9	2.0
08079-6837	Bab	2 454 963.5418		41.9	2.3	-40.3	1.7
10209-5603	Bab	2 454 520.7560		84.9	2.1	-99.8	2.2
10209-5603	Bab	2 454 573.6548		-42.8	2.5	96.4	3.0
10209-5603	Bab	2 454 908.7216		62.1	2.8	-71.7	2.4
10209-5603	Bab	2 454 960.5417		79.0	1.7	-98.1	2.0
10209-5603	Bab	2 455 292.7439		-7.1	1.5	38.0	2.3
10209-5603	Bab	2 456 336.8178		78.9	2.9	-110.7	4.7
10209-5603	Bab	2 456 338.8573		-68.3	2.2	122.7	2.7
20118-6337	Aab	2 454 574.8617	0.8460	56.9	1.9	-58.1	1.1
20118-6337	Aab	2 454 754.5853	0.5966	39.7	1.1	-39.7	1.1
20118-6337	Aab	2 454 961.7888	0.3179	-61.0	1.7	58.5	1.6
20118-6337	Aab	2 454 963.8326	0.1338	-51.9	1.5	49.7	0.9
20118-6337	Aab	2 455 083.6507	0.9685	8.0	2.0	-20.3	2.0
20118-6337	Aab	2 455 084.6143	0.3532	-58.9	1.7	52.7	1.5
20118-6337	Aab	2 455 085.6577	0.7697	65.9	1.5	-69.3	1.4
20118-6337	Aab	2 455 086.6321	0.1587	-60.1	1.7	54.3	2.2
20118-6337	Aab	2 455 087.6014	0.5457	16.6	1.1	-24.3	0.8
20118-6337	Aab	2 455 288.8592	0.8933	41.5	1.4	-43.4	1.0
20118-6337	Aab	2 455 291.8726	0.0963	-40.0	1.0	38.1	1.2
20118-6337	Aab	2 456 486.6665	0.0906	-40.1	1.2	38.8	1.6
20118-6337	Aab	2 456 486.8041	0.1456	-54.9	1.3	56.4	1.4
20118-6337	Aab	2 456 487.6096	0.4671	-18.7	1.6	9.4	1.1
20118-6337	Aab	2 456 487.7811	0.5356	11.9	1.7	-18.9	1.8
20118-6337	Aab	2 456 488.6104	0.8667	50.1	1.5	-53.8	1.0
20118-6337	Aab	2 456 488.7421	0.9193	32.5	1.7	-36.3	1.1

**Table 6.** Definitive orbital parameters of double-lined spectroscopic subsystems.

Subsystem (WDS)	$N_{sp}$	$P$ (days)	$T_0$ (HJD-2 400 000)	$e$	$K_a$ (km s <sup>-1</sup> )	$K_b$ (km s <sup>-1</sup> )	$V_\gamma$ (km s <sup>-1</sup> )	$\sigma_{RVa}$ (km s <sup>-1</sup> )	$\sigma_{RVb}$ (km s <sup>-1</sup> )
08079-6837 A	26	14.1697 ±0.0006	54 514.83 ±0.03	0	66.8 ±0.9	100.0 ±1.2	6.2 ±0.5	1.8	3.0
20118-6337 A	17	2.504836 ±0.000015	54 517.636 ±0.009	0	68.8 ±1.3	69.1 ±1.3	-2.0 ±0.6	1.9	2.2

**Table 7.** Preliminary orbital parameters of single-lined spectroscopic subsystems.

Subsystem (WDS)	$N_{\text{sp}}$	$P$ (days)	$T_0$ (HJD-2400000)	$e$	$\omega$ ( $^\circ$ )	$K_a$ ( $\text{km s}^{-1}$ )	$V_\gamma$ ( $\text{km s}^{-1}$ )	$\sigma_{\text{RVa}}$ ( $\text{km s}^{-1}$ )
08314-3904 A	12	73.25 $\pm 0.04$	55 543.4 $\pm 0.4$	0.155 $\pm 0.016$	190.2 $\pm 9.2$	32.3 $\pm 1.0$	15.5 $\pm 0.5$	1.4
13226-6059 C	18	3.8523 $\pm 0.0010$	55 500.71 $\pm 0.11$	0.24 $\pm 0.11$	346 $\pm 52$	21.2 $\pm 3.2$	-16.5 $\pm 2.4$	5.0
15185-4753 C	17	12.353 $\pm 0.004$	55177.7 $\pm 0.6$	0.4 $\pm 0.1$	324 $\pm 18$	2.4 $\pm 0.4$	6.9 $\pm 0.3$	0.6


**Fig. 1.** Radial velocity curves of WDS 08079-6837 Aab. Filled circles represent our measurements. The continuous line shows the orbital fitting, the dashed line indicates the barycentre velocity.

**Fig. 3.** Radial velocity curve of WDS 08314-3904 Aa. Filled circles represent our measurements. The continuous line shows the orbital fitting, the dashed line indicates the barycentre velocity.

**Fig. 2.** Linear correlation between the RVs of the components of WDS 08079-6837 Bab. The continuous line shows the fitting corresponding to  $\text{RV}_{\text{Ba}} = -0.93 \text{ RV}_{\text{Bb}} + 5.1 \text{ km s}^{-1}$ , which leads to  $q = 0.93 \pm 0.03$  and  $V_\gamma = 2.7 \pm 0.7 \text{ km s}^{-1}$ . Filled circles represent our measurements.


This multiple was observed in eight observing runs, in which we obtained 12 spectra of A (HD 72436). We did not detect any spectral feature that would indicate another spectroscopic component. Radial velocities, measured using a template of spectral type B4V convolved with a rotational profile of  $v \sin i = 150 \text{ km s}^{-1}$ , are shown to be variable.

The best fitting of the RV data was achieved for the eccentric orbit showed in Fig. 3 and Table 7. The fitting of the data is good: the rms deviations are similar to the measurement errors. However, we classified this orbit as “preliminary” since there are only few measurements and additional observations are required to confirm these orbital parameters.

We obtained ten spectra of component B, in which we measured RVs through cross-correlations with an observed template of spectral type B6V and  $v \sin i = 150 \text{ km s}^{-1}$ . The measurements are listed in Table 3. We found an RV variation between different observing runs, although we have not yet identified the corresponding period.

For component C, an object with low rotational velocity, we obtained three spectra with a time baseline of approximately one year. The obtained RVs did not show variations; the mean value is  $-13.4 \pm 0.2 \text{ km s}^{-1}$ .

The similarity between the barycentric velocity of A ( $15.1 \pm 0.5 \text{ km s}^{-1}$ ) and the average of the velocities measured for star B ( $15.2 \text{ km s}^{-1}$ ) supports the hypothesis that there is a physical link



**Fig. 4.** Profile variation of the spectral line Mg II  $\lambda 4481$  in HD 89890. Labels on the right are phases (calculated with  $P = 2.318$  days) and HJD–2 450 000.


between these two binaries. However, the velocity obtained for C differs substantially from these values, suggesting that this object does not belong to WDS 08314–3904. Consequently, the hierarchical structure of this system probably is considerably different from what was previously known: instead of three stars weakly bound to each other, it consists of two close binaries in a wide orbit.

### 3.3. WDS 10209–5603

This system has components A, B, and C catalogued in the MSC. The separation between A and B is  $7.19''$  and their published magnitude difference is  $\Delta V = 3.9$ . Component C is located at  $36.7''$  from AB. In the catalogue, the physical link between the components has been established on the basis of the criterion of a hypothetical dynamical parallax.

Corbally (1984) classified the two companions of the close visual pair and reported star B as a silicon star, giving spectral types B4III and B9.5IV Si for stars A and B, respectively. Pickering et al. (1898) detected an emission in H $\beta$  in a spectrum of component A, but this result has not been confirmed from subsequent observations. Baade (1984) detected variations in the profiles of some spectral lines with a period of about 2.25 days. Since these variations are very similar to those observed in Be stars, HD 89890 is considered to belong to this class of objects, but is currently going through a long B-star phase (Štefl et al. 2002). From RV measurements in 15 spectra, Rivinius et al. (2003) found a period of 2.318 days, which is consistent with the data of Baade. Furthermore, Sterken et al. (1996) measured photometric variations and found several possible periods around 4.6 days, the most prominent being 4.656 days. According to their data, they pointed out that a period of approximately two days is beyond their detection possibilities.

We obtained nine spectra of component A, which exhibits clear line-profile variations. Figure 4 shows the line profile of the Mg II line at  $\lambda 4481$ . The spectra are ordered by phases calculated using a period of 2.318 d and a phase origin set arbitrarily at HJD 2 450 000. The observed variations are clearly consistent with the period published by Rivinius et al. (2003). Because of the asymmetry of line profiles in HD 89890 we measured RVs by



**Fig. 5.** Linear correlation between the RVs of the components of WDS 10209–5603 Bab. Open squares represent measurements made in February 2013. These are fitted by the function  $RV_{Ba} = -0.63 RV_{Bb} + 9.1 \text{ km s}^{-1}$ , which is shown with a long-dashed line. Open triangles represent measurements made in February and April 2008, and filled circles indicate measurements of March and May 2009. Adopting a mass ratio  $q = 0.63$ , these data are fitted by the function  $RV_{Ba} = -0.63 RV_{Bb} + 20.0 \text{ km s}^{-1}$  (short-dashed line), and the function  $RV_{Ba} = -0.63 RV_{Bb} + 17.0 \text{ km s}^{-1}$  (dotted line), respectively. Open circle indicates RVs measured in a spectrum acquired in April 2010.

determining the barycentre of the profile of individual lines. The velocities consigned in Table 3 are the average of nine metallic and He I lines. We include this object in Table 3 considering that an RV variation exists, although in this case, it is not evidence of orbital motion.

Owing to the proximity of component A, which is about 4 mag brighter than B, we were only able to obtain seven spectra of this fainter component on the best observing nights. We detected morphological variations in the observed spectra that suggest two RV variable stars. In the spectral separation process we applied observed templates of spectral types A1V and A9IV for the RV measurement. From the separated spectra we confirm for the primary of this SB2 the classification as a silicon star given by Corbally (1984). The linear correlation between the velocities of both components indicates that they form an SB2 subsystem.

Radial velocities of stars Ba and Bb follow the linear relation expected for the components of a binary system, but a closer inspection of the deviations suggests a long-term variation of the centre-of-mass velocity. We calculated three linear fittings using only RVs measured in a same observing run or in observing runs temporally close to each other. In this way we obtained straight lines with roughly equal slopes but different intercepts, which suggests an RV variation of the barycentre of the spectroscopic binary subsystem. Since the available data are insufficient to determine the period of that variation, we adopted the mass ratio obtained from the fitting of the only two pairs of velocities measured in a same observing run (February 2013). From these data we obtained  $q = 0.63 \pm 0.02$  and  $V_\gamma = 5.6 \pm 1.5 \text{ km s}^{-1}$ . Adopting this value for the mass ratio, we fitted the barycentre velocity for the other observing runs that were temporally close to each other. In Fig. 5 we present the seven pairs of RVs measured and the fittings calculated using six of them, corresponding

to three different observing runs. The centre-of-mass velocity seems to have been decreasing at a rate of  $-3.5 \times 10^{-3} \text{ km s}^{-1} \text{ d}^{-1}$  during the five years covered by our data. The outer period would then be of some thousand days, although probably shorter than  $10^4$  days because with longer periods the RV variations would be smaller than observed. On the other hand, the relatively rapid RV variations suggest an inner orbital period of about 3–4 days. Consequently, the few available data sets suggest an intermediate hierarchical orbit between the visual pair AB and the subsystem Bab. This interesting result encourages us to continue with the observations of this subsystem, despite the difficulties caused by the proximity of HD 89890.

We obtained six spectra of the component C with a time baseline of two years. We did not detect morphological or RV variations between them. From the comparison of these spectra with the available templates we determined a spectral type K0III for this object, which had not been classified previously. The RVs measured by cross-correlations using a synthetic template with  $T_{\text{eff}} = 5500 \text{ K}$  led to a mean value  $6.1 \pm 0.6 \text{ km s}^{-1}$ .


The similarity between the mean velocity of component A, the barycentric velocities obtained for Bab, and the mean velocity found for component C supports the idea that they are gravitationally bound. With the detection of two new hierarchical levels the system WDS 10209–5603 would be, at least, quintuple.

According to theoretical models, a star with spectral type B3III should be significantly younger than a star of spectral type K0III if both of them have followed a normal evolution (early-B type stars evolve into supergiants K1b). Therefore, it is striking that A and C belong to the same physical multiple system. Considering the flux ratio between these objects, estimated from our spectroscopic data and the absolute magnitudes from the Schmidt-Kaler (1982) calibration, we found that the distance moduli of both objects are compatible with each other. This is more evidence, in addition to the similarity of their RVs, that supports the hypothesis that there is a physical link between them. If we assume that both objects have a common origin, then one of them should have had an abnormal evolution. Particularly, component A might have stayed on the main sequence over a longer period of time than a normal star with the same mass, which would mean that it is a blue-straggler star. Moreover, it can be speculated that this blue-straggler might have originated in the merger of the components of a close binary as a result of the influence of a third companion through Kozai cycles and tidal friction (see Sect. 4) (Perets & Fabrycky 2009). The profile variations of its spectral lines limit our detection possibilities of a possible orbital motion only up to periods of a few tens of days. Therefore, we do not rule out an additional companion of A in a hierarchical level below the visual pair AB.

### 3.4. WDS 13226–6059

The components A and B of this multiple system form a visual subsystem with a separation of  $0.17''$ . On the first hierarchical level, the pair AB (HD 116087) and the component C (HD 116072) form a common proper-motion and RV pair, separated by  $60''$ . We observed AB and C during four and seven observing runs, respectively, with a time baseline of 5.4 years.

We took seven spectra of the pair AB, in which a single set of broad spectral lines is visible, without significant morphological variations. In cross-correlations we employed a synthetic template with  $T_{\text{eff}} = 18000 \text{ K}$  and  $v \sin i = 200 \text{ km s}^{-1}$ . We did not measure RV variations greater than the errors given by the task *fxcor*, which were around a mean value of  $8.8 \text{ km s}^{-1}$ . We obtained a mean RV  $-2.1 \pm 3.5 \text{ km s}^{-1}$ .



**Fig. 6.** Radial velocity curves of WDS 13226–6059 Ca. Filled circles represent our measurements. The continuous line shows the orbital fitting, the dashed line indicates the barycentre velocity.


We obtained 18 spectra of component C. In measuring the RVs we employed a synthetic template with  $T_{\text{eff}} = 19000 \text{ K}$  and  $v \sin i = 200 \text{ km s}^{-1}$ . Although we obtained good correlation peaks, the measurement errors were about  $8.0 \text{ km s}^{-1}$  because of its high rotation. The measurements revealed an RV variation, but we did not detect spectral features of a companion. By using *pdm* we obtained tentative periods of 3.85 and 4.537 days. The least-squares fitting of the RV curve led to a slightly lower rms deviation for the former. However, the quality of the orbit is poor because of the low amplitude-to-error ratio of the RV curve, so we do not discard the other possible period. The parameters of the preliminary orbit are listed in Table 7. Parameter uncertainties are significant since RV errors are only  $\sim 3$  times smaller than the semi-amplitude.

The orbital period is relatively short, while the calculated eccentricity could be considered to be significantly different from zero:  $e/\sigma_e \sim 2$ . A statistical test of the null eccentricity hypothesis (as formulated by Lucy & Sweeney 1971) gives a probability of only 5% that the orbit is circular. However, we note that the rms of the residuals ( $\sim 5 \text{ km s}^{-1}$ ) is somewhat smaller than the formal measurement errors ( $\sim 8 \text{ km s}^{-1}$ ). If parameter errors are calculated using this value instead of the rms of the deviations from the fits, the probability of a circular orbit is about 30%. Even if the discrepancy between rms deviations and formal errors is probably due to the overestimation of the latter, we consider that this orbit and particularly its eccentric character should be taken with caution until more observations allow us to obtain a more reliable orbit.

Although the mean velocity of AB and the barycentric velocity obtained for Ca do not agree within the errors, for the moment we do not rule out a physical link between them, at least until the orbital parameters of Ca are confirmed. Therefore WDS 13226–6059 is, at least, a quadruple system.

### 3.5. WDS 15185–4753

This system has components A, B and C catalogued in the MSC. A and B form a visual pair with a separation of  $1.1''$ , without a published orbit. The component C is at  $22.6''$  from AB and



**Fig. 7.** Radial velocity curves of WDS 15185–4753 Ca. Filled circles represent our measurements. The continuous line shows the orbital fitting, the dashed line indicates the barycentre velocity.

forms a common proper-motion system with this pair. Based on infrared observations, Hubrig et al. (2001) reported the detection of an additional component located  $6.15''$  apart from A. From the  $K$  magnitude they estimated that this fourth component is a pre-main sequence star with a mass of approximately  $0.05 M_{\odot}$ . Given the very low mass inferred, Hubrig et al. pointed out that this faint object could be a brown dwarf.

The subsystem AB has often been studied spectroscopically and its published spectral type is B8Ve. Based on observations of high angular resolution ( $\sim 0.1''$ ) with adaptive optics, Oudmajer & Parr (2010) reported the detection of a visual companion of component A, located at  $0.98''$  and position angle of  $305^{\circ}$ . However, this is probably the visual B companion catalogued, which, according with the MSC, is at a position angle of  $311^{\circ}$ .

We carried out observations of the component C (HD 135748), which had no detailed spectroscopic studies. We obtained 17 spectra during six observing runs with a time baseline of 5.4 years. From our spectra, we determine a spectral type A2 V. To measure RVs we employed a template with  $T_{\text{eff}} = 8500$  K and  $v \sin i = 30$  km s $^{-1}$ . The measurement errors were around  $0.7$  km s $^{-1}$  and detected RV variations of a few kilometers per second (Table 4).

The best fitting of our RVs (Fig. 7) was obtained for an eccentric orbit whose parameters are listed in Table 7. However, it should be mentioned that another possible orbit, with a period of about 2.69 days, is also consistent with the observational data. We therefore consider these orbital parameters to be very preliminary. However, the classification of C as a single-lined spectroscopic binary is a robust result: the reduced chi squared is  $\chi^2/(n-1) = 4.85$ .

We took a spectrum of HD 135734 (AB subsystem). We detected two overlapping line sets corresponding to objects of similar spectral type and RV but with very different projected rotational velocity. Levenhagen & Leister (2006) measured a projected RV of  $280 \pm 20$  km s $^{-1}$  for HD 135734, which agrees well with the value previously measured by Chauville et al. (2001) ( $v \sin i = 278 \pm 25$  km s $^{-1}$ ). These results are consistent with the broad-line component detected in our spectra, while the other

has  $v \sin i \approx 50$  km s $^{-1}$ . In cross-correlations we employed a template with spectral type B7V convolved with a rotational profile corresponding to the object with lower rotational velocity. In this way, we measured an RV  $6.8 \pm 1.8$  km s $^{-1}$ . This value agrees with the barycentric velocity of Ca, which confirms the physical link between both subsystems. Therefore, WDS 15185–4753 has, at least, four components, although if the gravitational link between the infrared source detected by Hubrig et al. (2001) and the pair AB is confirmed, this would be a quintuple system.

### 3.6. WDS 20118–6337

This multiple is formed by a visual pair AB (HD 191056) with a separation of  $0.3''$  and a third component C located at  $7.34''$  from AB. In the MSC, the physical link between the components has been established on the basis of the criterion of hypothetical dynamical parallax and the proximity between the distances estimated from the spectral type and apparent magnitude.


We obtained 17 spectra of the subsystem AB, with a time baseline of 5.2 years. At first sight, we detected two components of different brightness showing short-period RV variations. In spectra in which the lines of these two components are well separated we detected the spectral features of a third object, although in all cases these features were blended with the lines of the two brightest components.

We measured the RV of the two brightest components through cross-correlations, using a template of spectral type A1V convolved with a rotational profile corresponding to  $v \sin i = 25$  km s $^{-1}$ . We found that the velocities measured correspond to two stars of equal mass that form a double-lined spectroscopic binary subsystem. Therefore, the difference of intensities between the spectral lines of the components is striking, especially because the component that appears brighter is not the same in all spectra. This is possibly due to a fourth star with a variable RV, whose spectral features are alternately overlapping the lines of one or another component. Initially, this had led to a misleading identification of the components and consequent difficulties in the period estimation of the binary subsystem. After both components of SB2 were correctly identified, we measured their RVs using the separating method by GL06. This technique gave good results, despite the two additional fainter components in the observed spectra. From the separated spectra, we found a spectral type A1V for both components. Figure 8 shows the least-squares fitting of the RVs and Table 6 lists the orbital parameters obtained for the SB2 subsystem. In the same way as for WDS 08079–6837 Aab, we obtained an eccentricity indistinguishable from zero, so we adopted a circular orbit and refitted the remaining orbital elements.

To summarize, based on the analysis of the spectra of the visual pair AB we detected that A is not a single component, but a double-lined spectroscopic binary subsystem (Aab). We did not detect variations in its barycentre velocity over more than five years, so an intermediate hierarchical level between Aab and the visual pair AB is not expected. We estimate that the third component detected between the lines of the SB2 is the visual B, but for the moment we cannot confirm whether this has a spectroscopic companion.

We obtained 12 spectra of component C, which does not have previous RV measurements published in the literature. We determined a spectral type A1V and measured RVs by cross-correlations using a template with  $T_{\text{eff}} = 9750$  K and  $v \sin i = 130$  km s $^{-1}$ . The measurements cover a time baseline of 5.2 years and do not show any significant RV variation. The mean RV obtained for this component is  $-8.1 \pm 3.8$  km s $^{-1}$ .





**Fig. 8.** Radial velocity curves of WDS 20118–6337 Aab. Filled circles represent our measurements. The continuous line shows the orbital fitting, the dashed line indicates the barycentre velocity.

#### 4. Discussion

Figure 9 presents the hierarchical configuration of the analysed systems, considering only the data catalogued in MSC (left) and adding the results of this investigation (right). As can be seen by comparing both panels, this analysis has led to significant changes in the multiplicity order and hierarchical structure of the studied systems. However, it is possible that there are other components inaccessible to our observations and the previous studies. Several of the studied systems present a high period ratio for orbits of consecutive hierarchical levels, which suggests that there might be undiscovered intermediate orbits.

We mark in Fig. 9 separation ranges that remain unexplored and might harbour intermediate hierarchical levels. The longest possible period that could have an intermediate orbit has been determined considering that the period ratio of adjacent orbits should be  $P_{\text{out}}/P_{\text{in}} \gtrsim 10$  to maintain the stability of the system. We employed the published period of the outer orbit. On the other hand, for each subsystem (including spectroscopic singles) we estimated the shortest possible period that could have an outer undetected orbit by evaluating our variability detection limits. Specifically, for a given subsystem, we calculated the expected amplitude of the variations of its barycentric RV caused by a hypothetical body (assumed to have at least  $0.5 M_{\odot}$ ) moving in a circular outer orbit. Assuming random orbital inclinations and random phases, we calculated the probability that during the time span of our observations the RV variations of the observed inner subsystem are larger than the corresponding measurement errors. These calculations were performed as a function of the outer period, so that we finally obtained the longest period for which we have at least a 50% detection probability.

Even though this spectroscopic study has contributed significantly to the knowledge of the hierarchical structure of the studied systems, in all cases it would be interesting to extend the search of additional components to the range of visual separations through high-angular resolution observations. This would contribute not only to the detection of intermediate orbits, but


also to the detection of long-period companions of components classified as single stars. Particularly the components A and C of WDS 10209–5603 and the component C of WDS 20118–6337 could have additional components in intermediate orbits with periods longer than approximately  $10^4$  days that would have gone unnoticed in this study and could only be detected through visual observations (to avoid confusion, we do not present these ranges for single components in Fig. 9). As is generally the case in the study of multiple stars, it is only possible to obtain definitive conclusions about the multiplicity order and the structure of the studied systems by complementing different observation techniques.

In some cases, the companions in intermediate hierarchical levels could affect the dynamical evolution of the inner subsystem through the mechanism known as “Kozai cycles” (Kozai 1962; Lidov 1962; Innanen et al. 1997; Ford et al. 2000). In a hierarchical system with non-coplanar outer and inner orbits, the gravitational perturbation on the inner binary produced by the third body leads to the precession of both orbits. If the relative inclination between the orbital planes is in a given critical range, the eccentricity and inclination of the inner binary oscillate, while the period and semimajor axis are not affected. However, if the period ratio of the outer and inner orbit is too high, this effect is suppressed by general relativity effects or quadrupolar distortion of the stars. In particular, if the outer period exceeds a minimum value given by  $P_{\text{out}}(\text{yr}) \approx [P_{\text{in}}(\text{days})]^{1.4}$  Kozai cycles are not expected to occur (Eggleton & Kisseleva-Eggleton 2006; Makarov & Eggleton 2009). This condition does not depend significantly on the stellar masses and orbital inclination. Since the eccentricity of the inner binary could achieve very high values as a consequence of Kozai cycles (Eggleton & Kisseleva-Eggleton 2006), and considering that the semimajor axis remains constant, the separation of the components at periastron can be significantly reduced. As a result, tidal friction dissipates energy from the orbital motion, which leads to a decrease of the semimajor axis and period. Eventually, the orbit will shrink enough for the tidal friction at periastron to become the dominant effect, circularizing the orbit and suppressing the Kozai cycling.

Taking into account the upper limit for  $P_{\text{out}}$  for the existence of Kozai cycles, for each system we determined the longest period that could have a companion in an intermediate hierarchical level to affect the dynamical evolution of the short-period binary subsystem. Particularly, we analysed whether this range of possible periods has been covered for our study. We conclude that we can rule out a companion that perturbs the inner binary for the subsystems Ca in WDS 15185–4753 and Aab in WDS 20118–6337. For the subsystem Bab of WDS 10209–5603 we considered a tentative period of 3.7 days and obtained that any companion in an intermediate orbit that could be affecting the dynamical evolution of the inner subsystem should be detected in this study. This is probably the case of the orbit detected through the movement of the barycentre of Bab, so it is an interesting question to analyse in the future, when there are enough observations of this close subsystem to determine an orbit.

#### 5. Conclusions

We have presented the results of a spectroscopic analysis of six systems catalogued as triples in the MSC for which we have detected new components. We found that three components classified as single stars are double-lined spectroscopic binaries, and for one of them we calculated a definitive orbit. We obtained RV curves of both components of a catalogued SB2 subsystem



**Fig. 9.** Configuration of the analysed multiple systems. *Left:* considering only the data catalogued in MSC. *Right:* including the results of this study. **a)** WDS 08079–6837; **b)** WDS 08314–3904; **c)** WDS 10209–5603; **d)** WDS 13226–6059; **e)** WDS 15185–4753; and **f)** WDS 20118–6337. Stellar components are represented by circles. Filled circles are stars observed in the present paper. No symbol is plotted for stellar components that are not visible in the spectrum (secondaries of SB1 systems). Dashed lines divide approximately the period axis in spectroscopic (S), visual (V), and common proper-motion (C) ranges. Dotted lines indicate a doubtful physical link between the components. The size of the circles roughly scale with the stellar mass. Green thick lines mark separation intervals that have not been investigated and might harbour additional hierarchical levels.

without previous data of the secondary component, and derived new orbital parameters. On the other hand, we detected that another three components are single-lined spectroscopic binaries, for which we obtained preliminary orbits. Additionally, we found one RV variable. Overall, this spectroscopic study has led to the discovery of seven confirmed new components and two members that we expect to be confirmed with new observations.

This study revealed that five of the analysed systems are quadruples. In one of them, WDS 08314–3904, the membership of one catalogued component was ruled out, but another two additional components were detected. On the other hand, the system WDS 10209–5603 is probably formed by five components, distributed on four hierarchical levels.

These new results reaffirm the conclusion reached in Paper I about the importance of the analysis of spectroscopic observations over a long time-baseline for detecting companions on intermediate hierarchical levels that might influence the dynamical evolution of close binary subsystems. We can also see in several of the systems studied here that these ranges of intermediate periods are essentially inaccessible to visual observations.

This study is a small but interesting evidence that a detailed analysis of the components of multiple systems with early-type components reveals a higher degree of multiplicity and a more complex hierarchical structure than previously known, suggesting that high-order multiple systems are significantly more frequent than it is currently estimated. In particular, the multiplicity order of the analysed systems obtained in this study should be regarded as a lower limit. In fact, in all six analysed systems there exists a separation range around  $P \sim 10^5$  days that remains virtually unexplored. A complete knowledge of these multiples can only be obtained by complementing the spectroscopic results with high-angular resolution visual observations.

*Acknowledgements.* This paper was partially supported by a grant from FONCYT-UNSW PICTO-2009-0125. We are grateful to Natalia Nuñez and Ana Collado, who kindly took some spectra used in this investigation.

## References

- Baade, D. 1984, *A&A*, 134, 105  
 Chauville, J., Zorec, J., Ballereau, D., et al. 2001, *VizieR Online Data Catalog: J/A+A/378/86/*  
 Chini, R., Hoffmeister, V. H., Nasser, A., Stahl, O., & Zinnecker, H. 2012, *MNRAS*, 424, 1925  
 Corbally, C. J. 1984, *ApJS*, 55, 657  
 Eggleton, P. P., & Kisseleva-Eggleton, L. 2006, *Ap&SS*, 304, 75  
 Ford, E. B., Kozinsky, B., & Rasio, F. A. 2000, *ApJ*, 535, 385  
 González, J. F., & Levato, H. 2006, *A&A*, 448, 283  
 Hubrig, S., Le Mignant, D., North, P., & Krautter, J. 2001, *A&A*, 372, 152  
 Innanen, K. A., Zheng, J. Q., Mikkola, S., & Valtonen, M. J. 1997, *AJ*, 113, 1915  
 Kozai, Y. 1962, *AJ*, 67, 591  
 Levenhagen, R. S., & Leister, N. V. 2006, *VizieR Online Data Catalog: J/MNRAS/371/252*  
 Lidov, M. L. 1962, *Planet. Space Sci.*, 9, 719  
 Lucy, L. B., & Sweeney, M. A. 1971, *AJ*, 76, 544  
 Makarov, V. V., & Eggleton, P. P. 2009, *ApJ*, 703, 1760  
 Medici, A., & Hubrig, S. 2000, *Inform. Bull. Variable Stars*, 4827, 1  
 Oudmaijer, R. D., & Parr, A. M. 2010, *MNRAS*, 405, 2439  
 Perets, H. B., & Fabrycky, D. C. 2009, *ApJ*, 697, 1048  
 Pickering, E. C., Fleming, W. P., & Cannon, A. J. 1898, *ApJ*, 8, 116  
 Rivinius, T., Baade, D., & Štefl, S. 2003, *A&A*, 411, 229  
 Sanford, R. F. 1915, *Lick Observatory Bulletin*, 8, 127  
 Schmidt-Kaler, T. 1982, in *Landolt-Bornstein, New Series, Group VI, Vol. 2b*, eds. K. Schaifers, & H. H. Voigt (Berlin: Springer)  
 Stellingwerf, R. F. 1978, *ApJ*, 224, 953  
 Sterken, C., Vogt, N., & Mennickent, R. E. 1996, *A&A*, 311, 579  
 Tokovinin, A. A. 1997, *A&AS*, 124, 75  
 Štefl, S., Rivinius, T., & Baade, D. 2002, in *Radial and Nonradial Pulsations as Probes of Stellar Physics*, eds. C. Aerts, T. R. Bedding, & J. Christensen-Dalsgaard, IAU Colloq., 185, ASP Conf. Ser., 259, 248  
 Veramendi, M. E., & González, J. F. 2014, *A&A*, 563, A138