

Elsevier Editorial System(tm) for Ultrasonics Sonochemistry
Manuscript Draft

Manuscript Number:

Title: Ultrasound promoted Barbier reactions and Csp3 - Csp2 Stille coupling for the synthesis of diarylmethanes and substituted benzophenones

Article Type: Full Length Article

Keywords: benzyltin compounds; Barbier reaction; ultrasound; Stille coupling; diarylmethanes; benzophenones

Corresponding Author: Dr Sandra Delia Mandolesi, Ph. D.

Corresponding Author's Institution: Universidad Nacional del Sur

First Author: Sandra D Mandolesi, Dr

Order of Authors: Sandra D Mandolesi, Dr; Sandra Delia Mandolesi, Ph. D.; Liliana C Koll, Dr

Abstract: Here we present the preparation of a variety of diarylmethanes obtained via ultrasound Stille coupling under palladium catalysis between some substituted aryl compounds and benzyltributyltin compounds generated through sonicated Barbier reaction in a very short time reaction and excellent yield. The study reported below compares different methods to optimize the synthesis of usually unstable benzyltin derivatives and is another contribution to the investigation of Csp3 - Csp2 coupling process involving benzyl-aryl reagents. Substituted carboxylated benzophenones were easily prepared in a very good yield by oxidation of some diarylmethanes.

Highlights

- Sonicated Barbier reaction as the best route for benzyltin compounds.
- Simplicity, mild conditions, rate of reaction and high yields in Barbier reaction.
- Creation of Csp³ benzyl center by sonicated Stille coupling.
- Synthesis of benzophenones by oxidation of diarylmethanes.

Ultrasound promoted Barbier reactions and Csp³ - Csp² Stille coupling for the synthesis of diarylmethanes and substituted benzophenones

Romina A. Ocampo ^{a, b}, Liliana C. Koll ^{a, b*} and Sandra D. Mandolesi ^{a*}

^a Instituto de Química del Sur (INQUISUR), Departamento de Química, Universidad Nacional del Sur, Avenida Alem 1253, 8000 Bahía Blanca, Argentina. ^b CONICET, Argentina.

E-mail: sdmando@criba.edu.ar

Abstract

Here we present the preparation of a variety of diarylmethanes obtained *via* ultrasound Stille coupling under palladium catalysis between some substituted aryl compounds and benzyltributyltin compounds generated through sonicated Barbier reaction in a very short time reaction and excellent yield. The study reported below compares different methods to optimize the synthesis of usually unstable benzyltin derivatives and is another contribution to the investigation of Csp³ - Csp² coupling process involving benzyl-aryl reagents. Substituted carboxylated benzophenones were easily prepared in a very good yield by oxidation of some diarylmethanes.

Keywords: benzyltin compounds, Barbier reaction, ultrasound, Stille coupling, diarylmethanes, benzophenones.

1. Introduction

The extensive use of organotin compounds as reagents or intermediates in organic synthesis [1] has prompted the development of efficient preparation of these type of compounds using rapid and convenient synthetic procedures, such as, for example, the Barbier reaction [2]. As the field of Csp³ - Csp² coupling reactions is always in continuous study from the last thirty years [3], the formation of this type of C-C bond *via* Stille coupling under palladium catalysis is of great interest in organic synthesis and, usually, tributylstannyl compounds are between the most common substrates. Compared with other coupling types, much less have been reported with these relatively more difficult couplings where the alkyl moiety is either on the substrate or on the organometallic partner [4]. The increased focus on the use of sonochemical methods in organic synthesis has been demonstrated by the many reports found in the literature [5]. The presence of ultrasound has shown to enhance the rates of reactions together with reduction of the induction period and less severe conditions. As a result of all previously mentioned, we focussed this work in the synthesis of substituted diarylmethanes taking into account that they play an important role in several biologically active compounds and drugs [6] such as Papaverin (**I**), a muscle relaxing agent, the synthetic antibiotic Trimethoprim (**II**) or the dihydrofolate reductase inhibitor for the potential treatment of cancer, Piritrexim (**III**) (Scheme 1). Besides, diarylmethanes are important precursors of the

corresponding benzophenones, which have demonstrated to have antibacterial activity [7] and, as is already known, the search of new antibiotics is always an important target for medicinal chemistry.

Scheme 1

2. Results and discussion

We based the synthetic strategy for the preparation of the diarylmethanes on the construction of two different aromatic “building blocks”, A and B, that will be coupled through a sonicated Stille reaction. The benzyltin group was located in the A-ring while the B-ring contains a variety of substituted aryl halides. Then, some representative products will be oxidized to the corresponding benzophenones (Figure 1).

Figure 1

To achieve this, benzyl chloride and 3,5-dimethoxybenzyl chloride (**2**) were chosen as starting materials. Compound **2** was obtained (75% yield) from commercially available methyl-3,5-dimethoxybenzoate (**1**) according to Scheme 2.

Scheme 2

The selected substrates providers of the Csp^3 moiety were benzyltri-*n*-butyltin (**3**) and 3,5-dimethoxybenzyltri-*n*-butyltin (**4**). Knowing the ease of decomposition of these type of compounds, we tested four different methods to prepare them: direct synthesis between zinc powder, tri-*n*-butyltinchloride and benzyl halide (I) [8]; Grignard reaction between tri-*n*-butyltin chloride and benzylmagnesium halide (II) [9]; reaction between lithium tri-*n*-butyltin, THF and benzyl halide (III) [10]; and Barbier sonicated reaction in the presence of magnesium turnings, THF, tri-*n*-butyltin chloride and benzyl halide (IV) [2c]. The results are summarized in Table 1. As can be seen in entries 4 and 8, the Barbier reaction was the best procedure for both **3** and **4** compounds, due to its simplicity, mild conditions, rate of reaction and very high yield compared with the other methods [11]. Benzyl chlorides were chosen instead of bromides to prevent the possible reduction to toluene derivatives. Besides, no side products were observed except the Bu_3SnCl formed during the reaction. It is clear that the very short time of reaction prevents the formation of undesired byproducts. The progress of the reaction was followed by thin layer chromatography (TLC) on silica

gel and checked by ^{119}Sn NMR spectra. The use of $(\text{Bu}_3\text{Sn})_2\text{O}$ as electrophile gave very poor yields and $(\text{Bu}_3\text{Sn})_2$ as byproduct. No dimerization of R_3Sn or benzyl species was observed [12].

Table 1

Once **3** and **4** were obtained, (A-ring), we tested the scope of the sonicated Stille coupling catalyzed by palladium complexes. At this point it is important to note that, usually, this coupling is carried out with benzylhalides as the Csp^3 center with a variety of arylstannanes as the Csp^2 source [13]. According to the very good results observed in the synthesis of the benzyltin compounds **3** and **4**, we proposed a reversal of this approach so these substrates will provide the Csp^3 center for the coupling reaction with several substituted aryl halides derivatives (B-ring). Thus, two reaction conditions were monitored: i) $\text{Pd}(\text{PPh}_3)_2\text{Cl}_2$, THF, AsPh_3 , LiCl , r.t., ultrasound and ii) $\text{Pd}_2(\text{dba})_3$, DMF, AsPh_3 , LiCl , r.t., ultrasound. The corresponding diarylmethanes (**5** - **17**) were obtained and the results are shown in Table 2.

Table 2

Except for 1-(3,5-dimethoxybenzyl)-2-methylbenzene, (**13**, entry 9), in all cases the best yields were achieved using the $\text{Pd}_2(\text{dba})_3$ / DMF system (method ii) and no traces of homocoupling byproduct was observed. On the other hand, the Stille reaction performed with the $\text{Pd}(\text{PPh}_3)_2\text{Cl}_2$ / THF system (method i) gave the homocoupling products in almost all cases (entries 1, 2, 3, 5, 6, 8 and 10) and in a high proportion in some of them (entries 2 and 8). It is important to note that the diarylmethanes **8**, **11**, **13**, **15**, **16** and **17** (entries 4, 7, 9, 11, 12 and 13) were the only products observed, no homocoupling byproducts were obtained and the yields were regular to low for most cases. As expected, the increase in the number of substituents on the phenyl groups gave higher time of reaction in both methods suggesting an important steric factor involved. After chromatographic purification with neutral alumina of compounds **5** – **17**, five of them (**5**, **8**, **9**, **13** and **14**) were oxidized [10] as representative cases at the benzylic center in a single reaction step (Scheme 3).

Scheme 3

As is already known [22], the combination of cerium (IV) ammonium nitrate (CAN) in catalytic amounts and potassium bromate can be employed for benzylic oxidation. Thus, compounds **5** and **8** were oxidized, giving the already known benzophenone (**18**) and the expected 3-benzoylbenzoic acid

(19) due to the additional oxidation of the methyl group. Because of this and taking into account that oxygen-bearing substituents like methoxy or nitro were not compatible with the latest reaction conditions [23], oxidation with chromium (VI) oxide (CrO_3) in acetic acid [24] was used in these special cases (9, 13 and 14). The corresponding benzophenones 20-22 were synthesized in very good yields after purification. In the case of substrates 13 and 14, 2-(3,5-dimethoxybenzoyl)benzoic acid (21) and 3-(3,5-dimethoxybenzoyl)benzoic acid (22) were obtained due to the inevitable oxidation of the methyl group.

3. Conclusions

In conclusion, we have demonstrated that the creation of a Csp^3 benzyl center for the synthesis of substituted diarylmethanes structures is possible through trialkylbenzyltin intermediates generated under ultrasound Barbier reaction with excellent yields followed by the corresponding sonicated Stille coupling with substituted aryl halides. The creation of new Csp^3 - Csp^2 bond, usually very difficult to achieve, is well accomplished under mild condition reactions and suitable yields. The use of ultrasound irradiation not only improved the reaction conversion, but made the reaction run smoothly.

4. Experimental

4.1. General methods

All the reactions were performed under nitrogen or argon as indicated. The solvents used were dried and distilled in accordance with standard procedures. ^1H , ^{13}C , and ^{119}Sn NMR spectra were recorded in CDCl_3 on a Bruker ARX 300 Multinuclear instrument (300.1 MHz for ^1H , 75.5 MHz for ^{13}C and 111.9 MHz for ^{119}Sn) at 23° C and calibrated by using signals from solvents referenced to SiMe_4 (^1H , ^{13}C NMR) and with respect to Me_4Sn in the case of ^{119}Sn -NMR spectra. Chemical shifts (δ) are reported in ppm and coupling constants (J) are in Hz. Compounds described in the literature were characterized by comparing their ^1H , ^{13}C and ^{119}Sn NMR spectra to the previously reported data. Unknown compounds were purified and analyzed from a single run and then were repeated to determine an average yield and were characterized by ^1H and ^{13}C NMR spectra, MS and elemental analysis. Mass spectra were obtained with a GC/MS instrument (HP5-MS capillary column, 30 m / 0.25 mm / 0.25 mm) equipped with 5972 mass selective detector operating at 70 eV (EI). High resolution mass spectra (HRMS) were recorded on a Finnigan Mat 900 (HR-EI-MS). IR spectra were recorded on a Nicolet Nexus FT spectrometer instrument. Melting points were determined with a Kofler Hot-Stage apparatus and are uncorrected. The progress of the reaction and the purity of compounds were monitored by TLC analytical silica gel plates (Merck 60 F250). The reactions under ultrasonic conditions were performed in an NDI ULTRASONIC 104X bath operating at 43-47

KHz and 30°C ($\pm 1^\circ\text{C}$). The reaction flask was located in the water bath of the ultrasonic cleaner, and the temperature of the water was controlled by addition or removal of water from the ultrasonic bath. Column chromatography was performed over silica gel 60 70-230 mesh ASTM. Flash chromatography was performed over alumina. All the solvents and reagents were commercially available and analytical grade.

4.2. Synthesis of 3,5-dimethoxybenzyl chloride (**2**) [25]

To a suspension of AlLiH_4 (0.6 g; 15.8 mmol) in dry ethyl ether (15 mL) under nitrogen atmosphere, a solution of commercial methyl-3,5-dimethoxybenzoate (**1**) (2 g; 10.0 mmol) in 10 mL of anhydrous ethyl ether was added dropwise with stirring. After 6 h of refluxing, the mixture was quenched with HCl 10% (20 mL). The organic layer was separated and dried over anhydrous MgSO_4 . The solvent was removed under reduced pressure and the crude product (1.71 g, quantitative yield, 10.2 mmol) was dissolved in CH_2Cl_2 (48 mL) under nitrogen atmosphere and, with magnetic stirring, thionyl chloride freshly distilled (1.5 mL; 20.4 mmol) was added dropwise. The reaction mixture was heated to reflux and then a solution of triethylamine (6 mL) in CH_2Cl_2 (48 mL) was added during 2 h. The reaction mixture was quenched with saturated NaHCO_3 (100 mL) and was extracted with ether (40 mL x 3). The organic layer was separated and dried over anhydrous MgSO_4 and concentrated. The crude residue was purified by flash column chromatography with alumina and 3,5-dimethoxybenzyl chloride (**2**) eluted as a yellowish solid (1.8 g, 9.6 mmol, 94%, m.p.: 45-46°C, lit. [7f] m.p.: 46°C).

4.3 Synthesis of benzyltin derivatives (**3**) and (**4**). Synthesis of benzyltri-*n*-butyltin (**3**)

All the reactions were carried out following the same procedure. One experiment is described in detail in order to illustrate the methods used.

4.3.1 Method I:

Tri-*n*-butyltin chloride (0.54 mL; 2 mmol) was poured to a mixture of zinc powder (0.26 g; 4 mmol) in THF (2 mL) / NH_4Cl saturated aqueous solution (4 mL) contained in a round-bottomed two necked flask equipped with condenser and dropping funnel. With magnetic stirring, benzylbromide (0.48 mL; 4 mmol) was added dropwise at such a rate to maintain a gentle reflux due to the exothermicity of the reaction. During the addition, the disappearance of zinc powder was observed. After stirring 24 h at room temperature, the organic layer was separated and THF removed under reduced pressure. The crude product (heavy liquid) was purified by silica-gel column chromatography doped with 10% KF, eluting (**3**) with hexane as an oil [26] (0.61 g; 1.6 mmol; 80%).

4.3.2 Method II:

Tri- *n*-butyltin chloride (15.3 mL; 13.2 mmol) was dissolved in a mixture of anhydrous benzene (15 mL) and anhydrous ethyl ether (8 mL) in a round bottomed two necked flask equipped with reflux condenser and dropping funnel under nitrogen atmosphere. After cooling to 0°C (in an ice-water bath), benzylmagnesium bromide (15 mL of 1.32 M ethyl ether solution; 19.9 mmol) was added dropwise and then kept 12 h under reflux. The reaction was cooled and quenched by slow addition of 10% HCl solution. The organic layer was dried with anhydrous MgSO₄ and the solvent was removed under reduced pressure. The residue was purified by silica gel column chromatography doped with 10% KF, eluting (**3**) with hexane as an oil (3.6 g; 9.5 mmol; 72%).

4.3.2.1 *3,5-dimethoxybenzyltri-n-butyltin (4)*. Yellowish oil (0.17 g, 0.9 mmol, 46%). ¹H NMR (CDCl₃): δ 7.24-7.20 (s, 1H), 7.16-7.00 (m, 2H), 3.62 (s, 6H); 2.18 (s, 2H, ²J_{Sn-H} = 55.7 CH₂), 1.60-1.22 (m, 6H), 0.99-0.82 (m, 15H), 0.54-0.40 (m, 6H). ¹³C NMR (CDCl₃): δ 159.86, 144.34, 122.90, 110.27, 55.62, 25.06 (³J_{Sn-C} = 20.3 CH₂), 23.09 (²J_{Sn-C} = 53.9 CH₂), 17.29 (¹J_{Sn-C} = 244.9 CH₂), 12.64, 12.29 (¹J_{Sn-C} = 318.8 CH₂). HR-MS (EI): Anal. calcd. for C₂₁H₃₈O₂Sn (442.1894): C, 57.16; H, 8.18%; found: C, 57.15; H, 8.14%.

4.3.3 Method III:

A mixture of tri-*n*-butyltin chloride (1.35 mL; 5 mmol) and lithium chippings (0.38 g; 5 mmol) in anhydrous THF (2.5 mL) was stirred 1 h in a round-bottomed flask equipped with nitrogen atmosphere and reflux condenser. The reaction becomes exothermic and the mixture turned dark green. After 2 h stirring, the mixture was filtered through glass wool to another round bottomed flask under nitrogen atmosphere and cooled to 0°C in an ice-water bath. Benzyl chloride (0.58 mL; 5 mmol) in anhydrous THF (1 mL) was added, followed by 12 h stirring at room temperature. The reaction was quenched with water (5 mL) and the aqueous layer washed with ethyl ether (2 x 2 mL). The combined organic layers were dried over anhydrous MgSO₄ and after removing the solvent, the crude product was purified by 10% KF doped-silica gel column chromatography, eluting (**3**) with hexane as an oil (1.30 g; 3.4 mmol; 68%).

Following the same procedure, compound (**4**) was obtained in 72% yield (0.66 g, 3.6 mmol).

4.3.4 Method IV (Barbier reaction):

A mixture of magnesium turnings (0.32 g; 13 mmol) and tri-*n*-butyltin chloride (3.25 g; 10 mmol) in anhydrous ethyl ether (2 mL) and few drops of pure benzyl chloride was placed in a Schlenk tube under argon atmosphere and submerged into a commercial ultrasonic cleaning bath. Once the reaction starts, the remaining benzyl chloride (1.4 mL, 12 mmoles) was added in ethereal solution (10 mL). When the reaction finished (30 min.), the mixture was washed with a saturated solution of NaCl and extracted with diethyl ether (7 mL). The organic layer was dried over anhydrous MgSO₄

and the solvent was removed under reduced pressure. The product was purified by silica gel column chromatography doped with 10% KF, eluting (**3**) with hexane as an oil (3.42 g; 9 mmol; 90%). Following the same procedure, compound (**4**) was obtained in 87 % yield (2.07 g, 11.3 mmol).

4.4 Synthesis of diarylmethanes (**5**) – (**17**) (ultrasound Stille coupling). Synthesis of 1-benzyl-3-methylbenzene (**8**)

All the reactions were carried out following the same procedure. One experiment is described in detail in order to illustrate the methods used.

4.4.1 Method i: [27]

A solution of benzyltri-*n*-butyltin (**3**) (0.38 g, 1 mmol), Pd(PPh₃)₂Cl₂ (2 mol%), AsPh₃ (30 mg, 0.012 mmol), LiCl (0.13 g, 3 mmol), 3-iodotoluene (0.22 g, 1 mmol) and THF (4 mL) under argon atmosphere was maintained in an ultrasonic bath. After 16 h of reaction no progress in the reaction was seen by TLC analysis. The crude product was filtered through celite to separate the inorganic insolubles salts together with the catalyst. The solvent was distilled off under reduced pressure and product **8** was isolated by column chromatography with alumina doped with 10% of KF to retain tributyltin halides formed during the reaction. **8** eluted with 95:5 (hexane/diethyl ether) as a yellowish oil. (0.12 g, 0.68 mmol, 68%, b.p.: 276.9°C/760 mmHg, lit: ¹⁵b.p.: 279.2/760 mmHg).

4.4.2 Method ii: [28]

A solution of 3,5-dimethoxybenzyltri-*n*-butyltin (**3**) (0.19 g, 0.5 mmol), Pd₂(dba)₃ (7.5 mg, 0.013 mmol), AsPh₃ (7.5 mg, 0.03 mmol), LiCl (31.3 mg, 0.75 mmol), 3-iodotoluene (0.11 g, 0.5 mmol) and DMF (10 mL) under argon atmosphere was maintained in an ultrasonic bath. After 11 h the reaction was complete and no starting product was observed by TLC (SiO₂). The crude product was filtered through celite to separate the inorganic insoluble salts together with the catalyst. The solvent was distilled off under reduced pressure and product **8** was isolated by column chromatography with alumina doped with 10% of KF to retain tributyltin halides formed during the reaction. **8** (0.07 g, 0.39 mmol, 78%) eluted with 95:5 (hexane/diethyl ether).

4.4.2.1 1-(3,5-dimethoxybenzyl)-4-methylbenzene (**12**): yellowish oil, ¹H NMR (CDCl₃): δ 7.19 (d, 2H); 7.05 (d, 2H); 6.55-6.48 (m, 3H); 3.96 (s, 2H); 3.62 (s, 6H); 2.09 (s, 3H). ¹³C NMR (CDCl₃): δ 162.23; 142.05; 139.23; 135.15; 129.02; 128.42; 107.62; 100.90; 57.21; 40.93; 21.14. HR-MS (EI): Anal. calcd. for C₁₆H₁₈O₂ (242.1307): C, 79.31; H, 7.49%, found: C, 79.29; H, 7.53%.

4.4.2.2 1-(3,5-dimethoxybenzyl)-2-methylbenzene (**13**): yellowish oil, ¹H NMR (CDCl₃): δ 7.39-7.30 (m, 2H), 7.21-7.16 (m, 2H), 6.46-6.40 (m, 3H), 3.95 (s, 2H), 3.65 (s, 6H), 2.16 (s, 3H). ¹³C NMR (CDCl₃): δ 161.87, 143.96, 141.89, 135.51, 129.23, 128.60, 127.52, 126.00, 106.94, 99.32, 58.63,

40.12, 20.67. HR-MS (EI): Anal. calcd. for $C_{16}H_{18}O_2$ (242.1307): C, 79.31; H, 7.49%; found: C, 79.34; H, 7.45%.

4.4.2.3 *1-(3,5-dimethoxybenzyl)-3-methylbenzene (14)*: yellowish oil, 1H NMR ($CDCl_3$): δ 7.26-7.00 (m, 4H), 6.34-6.30 (m, 3H), 3.86 (s, 2H), 3.75 (s, 6H), 2.30 (s, 3H). ^{13}C NMR ($CDCl_3$): δ 160.11, 143.61, 140.70, 138.13, 130.40, 130.12, 129.67, 126.05, 107.70, 98.10, 55.70, 42.06, 22.12. HR-MS (EI): Anal. calcd. for $C_{16}H_{18}O_2$ (242.1307): C, 79.31; H, 7.49%; found: C, 79.35; H, 7.50%.

4.5 Synthesis of benzophenones (**18**) – (**22**).

All the reactions were carried out following the same procedure. One experiment is described in detail in order to illustrate the methods used.

4.5.1 *3-benzoylbenzoic acid (19)*: A solution of 1-benzyl-3-methylbenzene (**8**) (0.182 g, 1 mmol), potassium bromate (0.083g, 0.5 mmol) and cerium (IV) ammonium nitrate (CAN) (0.055 g, 0.1 mmol) was added to acetic acid (1.5 mL). The solution was stirred and heated 24 h at 110°C or until no starting product was observed by TLC (SiO_2). The reaction mixture was poured into a cold mixture of water (2 mL) and CH_2Cl_2 (2 mL). The organic layer was separated, washed with water (2 x 3 mL) and dried over anhydrous $MgSO_4$. The solvent was removed under reduced pressure and the product was purified by recrystallization as a white solid from $CHCl_3$ / hexane (0.14 g, 0.71 mmol, 71%, m.p.: 153-154°C, lit.[29] m.p.:157-158°C).

4.5.2 *4-methoxybenzophenone (20)*: To a solution of 1-benzyl-4-methoxybenzene (**9**) (0.16 g, 0.8 mmol) in acetic acid (15 mL), chromium trioxide (0.12 g, 1.2 mmol) was added. The mixture was left at room temperature during 1 h or until no starting product was observed by TLC (SiO_2). The solution was extracted with diethyl ether (3 x 20 mL). The combined organic layers were washed with water (5 x 15 mL) and brine (10 mL). After drying over anhydrous Na_2SO_4 , the solvent was removed off under reduced pressure. The crude product was purified by silica gel column chromatography and **20** eluted as a white solid with hexane/diethyl ether (70/30), (0.15 g, 0.7 mmol, 88%, m.p.: 60-61°C, lit.: [30] m.p.:60-63°C).

4.5.3 *2-(3,5-dimethoxybenzoyl)benzoic acid (21)*: yellowish oil, IR (ν_{max} cm^{-1}): 3448 (O-H), 1683 and 1662 (C=O), 1253 (C-O). 1H NMR ($CDCl_3$): δ 11.80 (s, 1H), 8.17-7.50 (m, 4H), 7.06-6.44 (m, 3H), 3.62 (s, 6H). ^{13}C NMR ($CDCl_3$): δ 193.46, 165.83, 161.55, 137.68, 137.20, 134.75, 131.76, 131.14, 128.30, 127.80, 107.20, 107.81, 58.90. HR-MS (EI): Anal. calcd. for $C_{16}H_{14}O_5$ (242.1307): C, 67.13; H, 4.93%; found: C, 67.10; H, 4.89%.

4.5.4 *3-(3,5-dimethoxybenzoyl)benzoic acid (22)*: yellowish oil, IR (ν_{max} cm^{-1}): 3440 (O-H); 1687 and 1641 (C=O); 1250 (C-O). 1H NMR ($CDCl_3$): δ 10.78 (s, 1H), 8.35-7.46 (m, 4H), 7.23-6.60 (m, 3H),

3.68 (s, 6H); ^{13}C NMR (CDCl_3): δ 192.09, 166.22, 162.40, 142.01, 136.98, 135.44, 132.54, 130.21, 130.06, 129.60, 107.20, 106.71, 59.04. HR-MS (EI): Anal. calcd. for $\text{C}_{16}\text{H}_{14}\text{O}_5$ (242.1307): C, 67.13; H, 4.93%; found: C, 67.18; H, 4.90%.

Acknowledgements

This work was supported by grants from ANPCyT (Capital Federal, Argentina / BID 1728 / OC-AR PICT N° 2467), CONICET (Buenos Aires, Argentina, PIP 112-200801-02272) and Universidad Nacional del Sur (Bahía Blanca, Argentina, PGI 24/Q024). A fellowship from CONICET to RAO is acknowledged. The generous help of Professor M. González Sierra (IQUIR, Rosario, Argentina) in obtaining NMR spectra is gratefully acknowledged.

References


- [1] S. Pratihari, S. Roy, Reactivity and selectivity of organotin reagents in allylation and arylation: nucleophilicity parameter as a guide, *Organometallics* 30 (2011) 3257-3269 and references therein.
- [2] (a) G. Cravotto, P. Cintas, Power ultrasound in organic synthesis: moving cavitation chemistry from academia to innovative and large-scale applications, *Chem. Soc. Rev.* 35 (2006) 180-196 and references therein;
(b) M.E. Wood, Compounds of Group 14 (Ge, Sn and Pb): Product Subclass 8: Tin Halides and Organotin Halides, *Sci. Synth.* 5 (2003) 315-352;
(c) S. Lamandé-Langle, M. Abarbri, J. Thibonnet, A. Duchêne, *J. Org. Chem.* 694 (2009) 2368-2374.
- [3] (a) E. Negishi, L.F. Valente, M. Kobayashi, Palladium-catalyzed cross-coupling reaction of homoallylic or homopropargylic organozincs with alkenyl halides as a new selective route to 1,5-dienes and 1,5-enynes, *J. Am. Chem. Soc.* 102 (1980) 3298-3299;
(b) A.B. Smith III, T.J. Beauchamp, M.J. LaMarche, M.D. Kaufman, Y. Qiu, H. Arimoto, D.R. Jones, K. Kobayashi, Evolution of a Gram-Scale synthesis of (+)-Discodermolide, *J. Am. Chem. Soc.* 122 (2000) 8654-8664;
(c) G. Cahiez, C. Duplais, A. Moyeux, Iron-Catalyzed Alkylation of Alkenyl Grignard Reagents, *Org. Lett.* 9 (2007) 3253-3254;
(d) S. M. Nobre, A.L. Monteiro, Synthesis of diarylmethane derivatives from Pd-catalyzed cross-coupling reactions of benzylic halides with arylboronic acids, *Tetrahedron Lett.* 45 (2004) 8225-8228.
- [4] (a) R. Martin, A. Fürstner, Cross-coupling of alkyl halides with aryl Grignard reagents catalyzed by a low-valent iron complex, *Angew. Chem. Int. Ed.* 43 (2004) 3955-3957;
(b) D.J. Cárdenas, Advances in Functional-Group-Tolerant Metal-Catalyzed Alkyl-Alkyl Cross-Coupling Reactions, *Angew. Chem. Int. Ed.* 42 (2003) 384-387;
(c) T.-Y. Luh, M. Leung, K.-T. Wong, Transition Metal-catalyzed activation of aliphatic C-X bonds in carbon-carbon formation, *Chem. Rev.* 100 (2000) 3187-3204;

- (d) D.J. Cárdenas, Toward efficient and wide-scope metal-catalyzed alkyl-alkyl cross-coupling reactions, *Angew. Chem. Int. Ed.* 38 (1999) 3018-3020 and references therein;
- (e) K.H. Shukla, P. DeShong, Studies on the Mechanism of Allylic Coupling Reactions: A Hammett Analysis of the Coupling of Aryl Silicate Derivatives, *J. Org. Chem.* 73 (2008) 6283-6291;
- (f) A. Herve, A.L. Rodriguez, E. Fouquet, Stille Cross-Coupling of Activated Alkyltin Reagents under "Ligandless" Conditions, *J. Org. Chem.* 70 (2005) 1953-1956.
- [5] (a) T.J. Mason, J.P. Lorimer, *Applied Sonochemistry. The Uses of Power ultrasound in Chemistry and Processing*, Wiley-VCH Verlag GmbH, Weinheim, (2002);
- (b) J.L. Luche, *Synthetic Organic Chemistry*, Plenum Press, New York, (1998);
- (c) R.S. Varma, K.P. Naicker, Solvent-free synthesis of amides from non-enolizable esters and amines using microwave irradiation, *Tetrahedron Lett.* 40 (1999) 6177-6180.
- [6] (a) K.L. McPhail, D.E.A. Rivett, D.E. Lack, M.T. Davies-Coleman, The Structure and Synthesis of Tsitsikammafuran: A New Furanosesquiterpene from a South African *Dysidea* Sponge, *Tetrahedron* 56 (2000) 9391-9396;
- (b) G. Panda, M.K. Parai, S.K. Das, S. Manish Sinha, V. Chaturvedi, A.K. Srivastava, Y. Manju, A. Gaikwad, S. Sinha, Effect of substituents on diarylmethanes for antitubercular activity, *Eur. J. Med. Chem.* 42 (2007) 410-419;
- (d) C.D. Gilmore, K.M. Allan, B.M. Stoltz, Orthogonal synthesis of indulines and isoquinolines via aryne annulations, *J. Am. Chem. Soc.* 130 (2008) 1558-1559;
- (e) D. Subhas, A. Venkat, An efficient benzyltriethylammonium chloride catalysed preparation of electrophilic alkenes: a practical synthesis of trimethoprim, *J. Chem. Res.* 1 (2001) 36-38;
- (f) A. Rosowsky, H. Chen, A Novel Method of Synthesis of 2,4-Diamino-6-arylmethyl-quinazolines Using Palladium(0)-Catalyzed Organozinc Chemistry, *J. Org. Chem.* 66 (2001) 7522-7526.
- [7] (a) G.H. Faulkner, The antibacterial action of some diphenyl derivatives, *Biochem. J.* 38 (1944) 370-372;
- (b) K.H. Hong, G. Sun, Antimicrobial and chemical detoxifying functions of cotton fabrics containing different benzophenone derivatives, *Carbohydr. Polym.* 71 (2008) 598-605;
- (c) H.J. Florestano, M.E. Bahler, Antibacterial activity of a series of diphenylmethanes, *J. Am. Pharmaceut. Assoc.* 42 (2006) 576-578;
- (d) H. T. Balaydın, I. Gülçin, A. Menzek, S. Göksu, E. Şahin, Synthesis and antioxidant properties of diphenylmethane derivative bromophenols including a natural, *J. Enzym. Inhib. Med. Ch.* 25 (2010) 685-695;
- (e) P.G. Wyatt, R.C. Bethell, N. Cammack, D. Charon, N. Dodic, B. Dumaitre, D.N. Evans, D.V.S. Green, P.S. Hopewell, Benzophenone derivatives: a novel series of potent and selective inhibitors of human immunodeficiency virus type I reverse transcriptase, *J. Med. Chem.* 38 (1995) 1657-1665;
- (f) A. Chen, W. Shan, Y. Wu, L. Ren, J. Dong, Z. Ji, Synthesis and Anti-inflammatory Activity of Resveratrol Analogs, *Chem. Pharm. Bull.* 53 (2005) 1587-1590.

- [8] D. Marton, U. Russo, D. Stivanello, G. Tagliavini, Preparation of Benzylstannanes by Zinc-Mediated Coupling of Benzyl Bromides with Organotin Derivatives. *Physicochemical Characterization and Crystal Structures, Organometallics*, 15 (1996) 1645-1650 and references therein.
- [9] (a) B.S Furniss, A. J. Hannaford, P.W.G. Smith, A.R. Tatchell, *Vogel's Textbook of Practical Organic Chemistry*, 5th Edition, Longman Group Ltd. (1989);
(b) S. Boie, M. Gielen, J. Nasielski, J.P. Lieutenant, R. Spielman, *Organometallic Compounds XI. Synthesis and Properties of a Series of Racemic Tetraalkyltins*, *Bull. Soc. Chim. Belg.* 78 (1969) 135-146.
- [10] J. Labadie, J. Stille, Mechanisms of the palladium-catalyzed couplings of acid chlorides with organotin reagents, *J. Am. Chem. Soc.* 105 (1983) 6129-6137.
- [11] (a) C.A. Kraus, R.H. Bullard, Studies relating to alkyl tin compounds. I. Some stannethanes. II. Trimethylbenzyl stannane, *J. Am. Chem. Soc.* 48 (1926) 2131-2136;
(b) F.B. Kipping, CCCXII. An attempt to prepare optically active derivatives of quadrivalent tin, *J. Chem. Soc.* 131 (1928) 2365-2372;
(c) H. Gilman, S.D. Rosemberg, The preparation of triphenyltin-lithium and some of its reactions, *J. Am. Chem. Soc.* 74 (1952) 531-532;
(d) J.B. Peddley, H. Skinner, Thermochemistry of metallic alkyls. Part 9. Heats of bromination of some organotin compounds, *Trans. Faraday Soc.* 55 (1959) 544-547;
(e) M.A. Tius, J. Gomez-Galeno, Aromatic annelation. Synthesis of naphthalenes, *Tetrahedron Lett.* 27 (1986) 2571-2574.
- [12] J.F. Garst, Grignard reagent formation and freely diffusing radical intermediates, *Acc. Chem. Res.* 24 (1991) 95-97.
- [13] (a) H. Sugiyama, F. Yokokawa, T. Shioiri, Total synthesis of mycothiazole, a polyketide heterocycle from marine sponges, *Tetrahedron*, 59 (2003) 6579-6593;
(b) G.A. Holloway, H.M. Hügel, M.A. Rizzacasa, Formal Total Synthesis of Salicylhalamides A and B, *J. Org. Chem.* 68 (2003) 2200-2204;
(c) M.J. Schnermann, D.L. Boger, Total Synthesis of Piericidin A1 and B1, *J. Am. Chem. Soc.* 127 (2005) 15704-15705.
- [14] K. Ofosu-Asante, L. Stock, Selective deuteration of arylalkanes with mixed metal reagents, *J. Org. Chem.* 55 (1990) 3409-3410.
- [15] H. Esaki, F. Aoki, M. Umemura, M. Kato, T. Maegawa, Y. Monguchi, H. Sajiki, Efficient H/D Exchange Reactions of Alkyl-Substituted Benzene Derivatives by Means of the Pd/C–H₂–D₂O System, *Chem. Eur. J.* 13 (2007) 4052-4063.
- [16] R. Kuwano, M. Yokogi, Suzuki–Miyaura Cross-Coupling of Benzylic Carbonates with Arylboronic Acids, *Org. Lett.* 7 (2005) 945-947.
- [17] J.H. Lamneck, P.H. Wise, Dicyclic Hydrocarbons. IX. Synthesis and Physical Properties of the Monomethyldiphenylmethanes and Monomethyldicyclohexylmethanes, *J. Am. Chem. Soc.* 76 (1954) 1104-1106.
- [18] R.J. Taylor, Simple Palladium(II) Precatalyst for Suzuki–Miyaura Couplings: Efficient

- Reactions of Benzylic, Aryl, Heteroaryl, and Vinyl Coupling Partners, *Org. Lett.* 9 (2007) 5397-5400.
- [19] N. Henry, C. Enguehard-Gueiffier, I. Thery, A. Gueiffier, One-Pot Dual Substitutions of Bromobenzyl Chloride, 2-Chloromethyl-6-halogenoimidazo[1,2-*a*]pyridine and -[1,2-*b*]pyridazine by Suzuki–Miyaura Cross-Coupling Reactions, *Eur. J. Org. Chem.* 28 (2008) 4824-4827.
- [20] E. Alacid, C. Najera, First Cross-Coupling Reaction of Potassium Aryltrifluoroborates with Organic Chlorides in Aqueous Media Catalyzed by an Oxime-Derived Palladacycle, *Org. Lett.* 10 (2008) 5011-5014.
- [21] N.F. Janes, F.E. King, J.W.W. Morgan, The chemistry of extractives from hardwoods. Part XXXV. The constitution of Maesopin (2-benzyl-2,4,6,4'-tetrahydrocoumaranone) and its alkali fusion products, *J. Chem. Soc.* (1963) 1356-1363.
- [22] E. Ganin, I. Amer, Selective cerium-catalyzed oxidation of alkyl benzenes to benzyl esters by bromate salts, *J. Molec. Catalysis A: Chem.* 116 (1997) 323-327.
- [23] V. Sridharan, C. Menéndez, Cerium(IV) Ammonium Nitrate as a Catalyst in Organic Synthesis, *Chem. Rev.* 110 (2010) 3805-3849 and references therein.
- [24] S. Yamazaki, Chromium(VI) Oxide-Catalyzed Benzylic Oxidation with Periodic Acid, *Org. Lett.* 1 (1999) 2129-2132.
- [25] (a) S. Sengupta, M.G.B. Drew, R.B. Mukhopadhyay, B. Achari, A. Banerjee, Stereoselective Syntheses of (±)-Komaroviquinone and (±)-Faveline Methyl Ether through Intramolecular Heck Reaction, *J. Org. Chem.* 70 (2005) 7694–7700;
(b) D. Seebach, M. Hayakawa, J. Sakaki, W.B. Schweizer, Derivatives of $\alpha,\alpha,\alpha',\alpha'$ -Tetraaryl-2,2-dimethyl-1,3-dioxolan-4,5-dimethanol (TADDOL) containing nitrogen, sulfur, and phosphorus atoms. New ligands and auxiliaries for enantioselective reactions, *Tetrahedron*, 49 (1993) 1711-1724.
- [26] (a) U. Wannagat, V. Damrath, V. Huck, M. Veith, U. Harder, Sila-Riechstoffe und Riechstoffisostere XII . Geruchsvergleiche homologer Organoelementverbindungen der vierten Hauptgruppe (C, Si, Ge, Sn), *J. Organomet. Chem.* 443 (1993) 153-165;
(b) W.J. Jones, W.C. Davies, S.T. Bowden, C. Edwards, V.E. Davis, L.H. Thomas, 273. Preparation and properties of allyl phosphines, arsines and stannanes, *J. Chem. Soc.* (1947) 1446-1450.
- [27] W.J. Scott, J.K. Stille, Palladium-catalyzed coupling of vinyl triflates with organostannanes. Synthetic and mechanistic studies, *J. Am. Chem. Soc.* 108 (1986) 3033-3040.
- [28] Y. Obora, Y. Tsuji, M. Kobayashi, T. Kawamura, Stille Coupling Reaction Using 4-(Trimethylsilyl)-2-butenylstannanes To Afford Allylic Silanes, *J. Org. Chem.* 60 (1995) 4647-4649.
- [29] W.E. Parham, Y.A. Saded, Synthesis of benzoylbenzoic acids, *J. Org. Chem.* 39 (1974) 2051-2053.
- [30] S. Guo, G. Su, Y. He, Structure of 4-methoxybenzophenone, *Acta Cryst. C*48 (1992) 576-578.

Table 2. Sonicated Stille reaction between benzyltributyltin compounds (**3**) and (**4**) and several substituted aryl halides.



Entry	G	R ₁	X	Reaction conditions ^a	Product	% yield ^b	Time (h)	Ref.
1	H	H	Br	i / ii		39 (10) / 70	12 / 9	[14]
2	H	<i>p</i> -CH ₃	I	i / ii		26 (32) / 67	18 / 12	[15]
3	H	<i>o</i> -CH ₃	I	i / ii		63 (8) / 75	18 / 12	[16]
4	H	<i>m</i> -CH ₃	I	i / ii		68 / 78	16 / 11	[17]
5	H	<i>p</i> -OCH ₃	I	i / ii		45 (40) / 56	18 / 12	[18]
6	H	<i>o</i> -OCH ₃	I	i / ii		67 (20) / 71	18 / 12	[19]
7	H	<i>m</i> -OCH ₃	I	i / ii		45 / 60	20 / 14	[20]
8	-OCH ₃	<i>p</i> -CH ₃	I	i / ii		28 (60) / 59	26 / 20	---
9	-OCH ₃	<i>o</i> -CH ₃	I	i / ii		42 / 39	28 / 19	----
10	-OCH ₃	<i>m</i> -CH ₃	I	i / ii		51 (20) / 65	27 / 20	----
11	-OCH ₃	<i>p</i> -OCH ₃	I	i / ii		45 / 68	28 / 22	[21]
12	-OCH ₃	<i>o</i> -OCH ₃	I	i / ii		55 / 72	30 / 25	[21]
13	-OCH ₃	<i>m</i> -OCH ₃	I	i / ii		52 / 75	36 / 25	[21]

^a Reaction conditions: i) Pd(PPh₃)₂Cl₂, THF, AsPh₃, LiCl, r.t., ultrasound; ii) Pd₂(dba)₃, DMF, AsPh₃, LiCl, r.t., ultrasound. ^b Determined by GC/MS; isolated yields as an average of at least three independent runs and corresponds to reaction conditions i and ii respectively. In parentheses the yield of homocoupling product is shown.

Scheme 1









Figure 1



Scheme 2



Scheme 3



Graphical abstract

