

Send Orders of Reprints at reprints@benthamscience.net

388

Current Neuropharmacology, 2013, 11, 388-413

Natural AChE Inhibitors from Plants and their Contribution to Alzheimer's Disease Therapy

Ana Paula Murray^{a,b,*}, María Belén Faraoni^{a,c}, María Julia Castro^a, Natalia Paola Alza^a and Valeria Cavallaro^a

^aINQUISUR-CONICET, Departamento de Química, Universidad Nacional del Sur, Bahía Blanca, Argentina; ^bResearch Member of CONICET; ^cResearch Member of CIC

Abstract: As acetylcholinesterase (AChE) inhibitors are an important therapeutic strategy in Alzheimer's disease, efforts are being made in search of new molecules with anti-AChE activity. The fact that naturally-occurring compounds from plants are considered to be a potential source of new inhibitors has led to the discovery of an important number of secondary metabolites and plant extracts with the ability of inhibiting the enzyme AChE, which, according to the cholinergic hypothesis, increases the levels of the neurotransmitter acetylcholine in the brain, thus improving cholinergic functions in patients with Alzheimer's disease and alleviating the symptoms of this neurological disorder. This review summarizes a total of 128 studies which correspond to the most relevant research work published during 2006-2012 (1st semester) on plant-derived compounds, plant extracts and essential oils found to elicit AChE inhibition.

Keywords: Alzheimer's Disease, acetylcholinesterase inhibitors, secondary metabolites, plant extracts, essential oils.

INTRODUCTION


Alzheimer's disease (AD) is a progressive neurodegenerative disorder associated with memory impairment and cognitive deficit. It is characterized by low levels of acetylcholine in the brain of AD patients. According to the cholinergic hypothesis, the inhibition of acetylcholinesterase (AChE), an enzyme that catalyzes acetylcholine hydrolysis, increases the levels of acetylcholine in the brain, thus improving cholinergic functions in AD patients. Furthermore, although the general consensus concludes that AChE inhibitors (AChEi) can alleviate AD symptoms, they neither delay nor reverse the disease progress. Most of the drugs currently available for the treatment of AD are AChEi: tacrine (**1**), donepezil (**2**), rivastigmine (**3**) and galanthamine (**4**), all of which have limited effectiveness and some kind of side effect [1]. Tacrine (**1**) and donepezil (**2**), both from synthetic origin, were the first drugs approved for the treatment of cognitive loss in AD patients by US-FDA in 1993 and 1996, respectively. Rivastigmine (**3**) was approved in 2000 (US-FDA) and was designed from the lead compound physostigmine, a natural AChEi alkaloid. Galanthamine (**4**), a natural alkaloid first obtained from *Galanthus* spp. was approved by US-FDA in 2001. Huperzine A (**5**), an alkaloid found in *Huperzia* spp., is an AChEi commercialized as a dietary supplement for memory support and it is used to treat AD symptoms in China. This alkaloid has been thoroughly studied with promising results yielded particularly from the evaluation of cognitive

performance of animals as well as from studies on its efficacy, tolerance and safety.

Taking into account that inhibitors **3**, **4** and **5** are related to natural products and that AChEi are an important therapeutic strategy for the treatment of AD, many research groups have focused their studies on naturally-occurring compounds from plants as potential sources of either new or more effective AChEi. These studies led to the discovery of an important number of secondary metabolites as well as plant extracts, both of which are characterized by their ability to inhibit AChE. On the other hand, the fact that a significantly relevant number of research papers has been recorded in this field during the last decades can be clearly attributed to the development of colorimetric methods which allow a rapid and facile screening of a large number of samples. Ellman's method is the most widely used for the detection of AChEi, even in complex mixtures, and for the quantification of anti-AChE inhibitory activity [2-6].

Several reviews on the newly discovered AChEi obtained from plants, fungus and marine organisms have also been published over the last years [7-10]. The majority of these AChEi belong to the alkaloid group, including indole, isoquinoline, quinolizidine, piperidine and steroidal alkaloids. On the other hand, several non-alkaloidal and potent AChEi have been obtained from natural sources, including terpenoids, flavonoids and other phenolic compounds. Interestingly, although literature demonstrates to be rich in the study on AChEi obtained from plants, this issue keeps on being the center of attention for research as confirmed by the increasing number of studies published every year. Therefore, the purpose of this review is to provide a comprehensive summary of the literature, particularly that published during 2006-2012 (1st semester) on plant-derived compounds, plant

*Address correspondence to this author at the INQUISUR-CONICET, Departamento de Química, Universidad Nacional del Sur, Bahía Blanca, Argentina; Tel: 54 291 4595101 ext. 3538; Fax: 54 291 4595187; E-mail: apmurray@uns.edu.ar




extracts and essential oils which have been reported to inhibit AChE. Readers interested not only in previous findings but also in synthetic/semisynthetic AChEi or natural AChEi of fungal, marine or microbial origin are recommended to see the above-mentioned reviews [i.e. 7-10]. For the sake of brevity and in order to focus our attention on the most relevant findings, only those research papers reporting quantified results (IC_{50} and/or percentage of inhibition at a given concentration) were included. Extracts or essential oils with $IC_{50} > 0.5$ mg/ml were considered weakly active and were therefore not taken into account in the present review. With a few exceptions, only molecules with $IC_{50} < 50$ μ M have been considered. Furthermore, unless otherwise stated, those results on AChE inhibition included in the present review refer to *in vitro* assays carried out with AChE from electric eel.

ALKALOIDS WITH AChE INHIBITORY ACTIVITY


The quinoline alkaloids 3-hydroxy-2,2,6-trimethyl-3,4,5,6-tetrahydro-2H-pyrano[3,2-c] quinoline-5-one (**6**), ribalinine (**7**) and methyl isoplatydesmine (**8**) isolated from the aerial parts of *Skimmia laureola* (Rutaceae) were found to be linear mixed inhibitors of AChE with $K_i = 110.0$, 30.0 and 30.0 μ M, respectively [11]. These alkaloids were also observed to evidence butyrylcholinesterase (BChE) inhibition.




On the other hand, of the several alkaloids that were isolated from the active extracts of *Esenbeckia leiocarpa* (Rutaceae), leptomerine (**9**) and kokusaginine (**10**) with IC_{50} values of 2.5 and 46 μ M, respectively, were observed to elicit AChE inhibitory activity [12]. The isolation of

skimmianine (**11**), a furoquinoline alkaloid with very low AChE inhibitory activity, was also reported by the same authors. This alkaloid was observed in another Rutaceae, *Zanthoxylum nitidum*, exhibiting a moderate AChE inhibitory activity ($IC_{50} = 8.6$ μ g/ml) [13].




Nelumbo nucifera is a well-known medicinal plant belonging to the Nelumbonaceae family which was studied due to its therapeutic potential [14]. N-methylasimilobine (**12**), an aporphine alkaloid with an $IC_{50} = 1.5$ μ g/ml which was found to be a non-competitive inhibitor, was recently isolated from this plant [15]. In a random screening, two extracts of *Beilschmiedia* species were observed to exhibit AChE inhibition and a phytochemical study of *B. alloiophylla* and *B. kunstleri* revealed the presence of several alkaloids with IC_{50} values ranging between 2.0 and 10.0 μ M [16]. The most potent AChEi were found to be 2-hydroxy-9-methoxyaporphine (**13**), laurotetanine (**14**), liriodenine (**15**) and oreobelline (**16**) ($IC_{50} = 2.0$ -5.0 μ M), with anti-AChE activity comparable to huperzine A ($IC_{50} = 1.8$ μ M). A significant AChE inhibitory activity was also observed in





- 12:** R₁=CH₃, R₂=R₅=R₆=H, R₃=OH, R₄=OCH₃
13: R₁=CH₃, R₂=R₄=R₅=H, R₃=OH, R₆=OCH₃
14: R₁=R₂=H, R₃=R₄=R₅=OCH₃, R₆=OH
18: R₁=CH₃, R₂=H, R₃=R₆=OH, R₄=R₅=OCH₃
19: R₁=CH₃, R₂=H, R₃=R₅=OCH₃, R₄=R₆=OH
20: R₁=R₂=R₅=R₆=H, R₃=OH, R₄=OCH₃
21: R₁=H, R₂=R₃=OCH₃, R₄=OH, R₅+R₆=OCH₂O





secoboldine (**17**), boldine (**18**), isoboldine (**19**), asimilobine (**20**) and 3-methoxynordomesticine (**21**) (IC₅₀ = 8.4 - 10.0 μM).


Research on plants from the genus *Corydalis* (Papaveraceae) which are used for the treatment of memory dysfunction in folk medicine reported the presence of benzyloisoquinoline alkaloids with anti-AChE activity [7]. The ethanolic extract obtained from the tuber of *C. turtchaninovii* previously found to elicit AChE inhibition was selected to carry out a chemical study which led to the isolation of the isoquinoline alkaloids stylopine (**22**), epiberberine (**23**), pseudodehydrocorydaline (**24**), pseudocopsitine (**25**) and pseudoberberine (**26**). In the assay with mouse brain cortex as a source of AChE enzyme, the IC₅₀ values obtained for each of these alkaloids were 15.8, 6.5, 8.4, 4.3 and 4.5 μM, respectively [17]. In addition, alkaloids **25** and **26**, the two most active compounds, were found to elicit anti-amnesic activity [17, 18]. Alkaloids with benzyloisoquinoline skeleton from *Corydalis* species having aromatic methylenedioxy groups and a quaternary atom of

nitrogen were observed to show the strongest AChE inhibition [7, 17, 18]. In a more recent work, six protoberberine alkaloids **23**, **27** - **31**, were identified in rhizomes of *Coptis chinensis* which are traditionally used in Chinese medicine for the treatment of various diseases. *Coptis* rhizomes and their alkaloids were reported to have cognitive-enhancing and neuroprotective effects and the analysis of the anti-AChE activity of these alkaloids showed that the IC₅₀ values of berberine (**27**), palmatine (**28**), jateorrhizine (**29**), coptisine (**30**) and groenlandicine (**31**) ranged between 0.44 and 0.80 μM while that of epiberberine (**23**) was slightly higher (IC₅₀ = 1.07 μM) [19]. Of these alkaloids, compounds **27**, **30** and **31** were observed to have an aromatic methylenedioxy group. In this study groenlandicine (**31**) and berberine (**27**) were found to be the most active as BChE inhibitors and epiberberine (**23**) was observed to significantly inhibit β-secretase (BACE1) [19].


The alkaloids (+)-canadoline (**32**) and (+)-canadine (**33**), both isolated from *Corydalis cava* and with an IC₅₀ = 20.1




- 23:** R₁=R₂=OCH₃, R₃=R₄=H, R₅+R₆=OCH₂O
24: R₁=R₂=OCH₃, R₃=CH₃, R₄=R₅=OCH₃, R₆=H
25: R₁+R₂=OCH₂O, R₃=R₆=H, R₄+R₅=OCH₂O
26: R₁+R₂=OCH₂O, R₃=R₆=H, R₄=R₅=OCH₃
27: R₁+R₂=OCH₂O, R₃=R₄=H, R₅=R₆=OCH₃
28: R₁=R₂=R₅=R₆=OCH₃, R₄=R₃=H
29: R₁=OH, R₂=R₅=R₆=OCH₃, R₃=R₄=H
30: R₁+R₂=OCH₂O, R₃=R₄=H, R₅+R₆=OCH₂O
31: R₁=OH, R₂=OCH₃, R₃=R₄=H, R₅+R₆=OCH₂O
34: R₁=R₆=OCH₃, R₂=R₅=OH, R₃=R₄=H




32




33




35: R₁=OCH₃, R₂=R₃=OH, R₄=OCH₃
 36: R₁=R₃=OCH₃, R₂=R₄=OH




37: R₁=R₂=OCH₃
 38: R₁+R₂=OCH₂O



39



40



41: R = H
 42: R = OCH₃
 43: R = OH

and 12.4 μM , respectively, were observed to elicit a moderate inhibitory activity when tested with AChE from human blood [20].

On the other hand, *Stephania venosa* (Menispermaceae), a Thai medicinal plant, was found to show a high AChE inhibitory activity. The ethanolic extract of *S. venosa* was subjected to bioassay-guided fractionation to identify AChEi [21]. The following moderately active quaternary protoberberine alkaloids could be isolated: stepharanine (**34**), cyclanolone (**35**) and *N*-methyl stepholidine (**36**) with IC₅₀ values of 14.10, 9.23 and 31.30 μM , respectively. A similar fractionation approach was followed to identify the compounds responsible for AChE inhibition in *Chelidonium majus* (Papaveraceae) [22]. Three active constituents were identified, namely 8-hydroxydihydrochelerythrine (**37**), 8-hydroxydihydrosanguinarine (**38**) and berberine (**37**). Compounds **37** and **38**, with no previous record as AChEi, were found to elicit significant anti-AChE activity with an IC₅₀ = 0.61 and 1.37 μM , respectively.


Taspine (**39**) was isolated from the alkaloid-enriched extract obtained from *Magnolia x soulangiana* (Magnoliaceae) [23]. This alkaloid was found not only to show a dose-dependent and long-lasting inhibitory effect on AChE (IC₅₀ = 0.33 μM) but also to be more potent than galanthamine (IC₅₀ = 3.2 μM) although its inhibitory activity is comparable to that of tacrine (IC₅₀ = 0.22 μM). Similar observations were obtained when the *in vitro* assay was performed with human AChE (IC₅₀ = 0.54 μM). Compound **39** resulted to be inactive against BChE, acting as a selective AChEi.

Catharanthus roseus (Apocynaceae) is a plant mainly known as a source of vincristine and vinblastine, two alkaloids found in its leaves and appreciated as anticancer compounds. Several other compounds with biological importance can be also found in *C. roseus*. For example, the alkaloid serpentine (**40**), isolated from the roots of this plant,

was reported to be a potent *in vitro* AChEi (IC₅₀ = 0.775 μM) compared with physostigmine (IC₅₀ = 6.45 μM) [24].

A bioassay-guided fractionation from the stems of *Ervatamia hainanensis* (Apocynaceae), a plant used in traditional Chinese medicine, allowed the isolation of several monoterpenoid indole alkaloids, some of them showing a potent AChE inhibitory activity [25]. For example, coronaridine (**41**) and voacangine (**42**), differing from each other only by the methoxy group attached to the aromatic ring, were observed to have an IC₅₀ = 8.6 and 4.4 μM , respectively, these values being similar to that of galanthamine (3.2 μM). On the other hand, 10-hydroxy coronaridine (**43**) was found to evidence a reduced AChE inhibition (IC₅₀ = 29 μM), which was attributed to the introduction of a hydroxyl group to the aromatic ring. The indole alkaloids coronaridine (**41**) and voacangine (**42**), both detected in the stalks of *Tabernaemontana australis* (Apocynaceae), had been formerly identified as AChEi but no inhibition values were reported [26].

The genus *Tabernaemontana* is known for the wide variety of unusual bioactive indole alkaloids it produces. Among them, the bisindole alkaloids isolated from *T. divaricata* roots are an interesting example of new structures with potent AChE inhibitory activity. The crude alkaloid extract obtained from the root of *T. divaricata* was found to yield four bisindole alkaloids **44** - **47** [27]. The analysis of AChE inhibition revealed that 19,20-dihydrotabernamine (**44**) and 19,20-dihydroervahanine A (**45**) strongly inhibit AChE, with an IC₅₀ = 0.227 and 0.071 μM , respectively, thus showing that they are significantly more active than galanthamine (IC₅₀ = 0.594 μM). The fact that inhibition was found to be higher for compound **45** than for compound **44** suggests that the introduction of a carbomethoxy group at C16' increases the enzymatic inhibition. In addition, taking into account that conodurine (**46**) and tabernaegantine (**47**)




were found to show no activity in AChE, it was suggested that the substitution at C11' and C12' is relevant for AChE inhibitory activity [27].

Uncaria rhynchophylla (Rubiaceae) is a Chinese medicine herb used to treat epilepsy. The alkaloid fraction from *U. rhynchophylla* is known for its antiepileptic and neuroprotective effects. Geissoschizine methyl ether (**48**), a strong AChEi, as well as six other weakly active alkaloids were recently isolated from this herb [28]. The active compound **48** was observed to inhibit AChE in a reversible and non-competitive way with an $IC_{50} = 3.7 \mu\text{g/ml}$.


The study of AChE inhibitory activity of Brazilian apocynacea *Himatanthus lancifolius*, commonly known as "agoniada", led to the identification of active extracts in this plant and allowed the isolation of uleine (**49**), an active indole alkaloid, at a high concentration in the alkaloid fraction. The IC_{50} value observed for this alkaloid was $0.45 \mu\text{M}$ [29].

As to the Amaryllidaceae family, phytochemical research conducted in the last decades on this family revealed several




alkaloids with moderate or potent inhibition of AChE [3, 7, 30]. In the search of new natural sources of galanthamine and other Amaryllidaceae alkaloids with anti-AChE activity, bulbs and leaves of *Hippeastrum papilio* collected in the South of Brazil were studied. Galanthamine (**4**), the already known alkaloids narwedine (**50**), haemanthamine (**51**), 11-hydroxyvittatine (**52**), 8-*O*-demethylmaritidine (**53**) and vittatine (**54**) as well as the new alkaloid 11 β -hydroxygalanthamine (**55**) were all isolated and of all of them galanthamine was obtained in significant amounts [31]. Compound **55** was observed to elicit AChE inhibition as other galanthamine-type alkaloids do, with an $IC_{50} = 14.5 \mu\text{M}$. Furthermore, because habranthine, epimer of **55**, was observed to have an anti-AChE activity similar to that of galanthamine, it was concluded that β configuration at C11 is unfavorable for the interaction with AChE [3, 31]. Other potent AChEi, such as *N*-allylnorgalanthamine (**56**) and *N*-(14-methylallyl)norgalanthamine (**57**), were isolated from *Leucojum aestivum*, an amaryllidacea used for the industrial extraction of galanthamine [32]. *N*-alkylated galanthamine derivatives **56** and **57** were isolated together with galanthamine (**4**), epinorgalanthamine (**58**), narwedine (**50**) and lycorine (**59**), from the mother liquors obtained after the industrial production of galanthamine. Alkaloids **56** and **57**, with IC_{50} values of 0.18 and $0.16 \mu\text{M}$, respectively, resulted to be ten times more potent AChEi than galanthamine ($IC_{50} = 1.82 \mu\text{M}$).


The chemical investigation of *Galanthus rizehensis*, a wild-growing species from Turkey, allowed the isolation of two new Amaryllidaceae alkaloid *N*-oxides, incartine *N*-oxide (**60**) and lycorine *N*-oxide (**61**) and seven




- 50: R₁+R₂=O, R₃=R₄=H, R₅=CH₃
 55: R₁=R₃=OH, R₂=R₄=H, R₅=CH₃
 56: R₁=OH, R₂=R₃=R₄=H, R₅=-CH₂CH=CH₂
 57: R₁=OH, R₂=R₃=R₄=H, R₅=-CH₂C(CH₃)=CH₂
 58: R₁=R₃=R₄=R₅=H, R₂=OH




- 51: R₁=OCH₃, R₂=H, R₃=OH, R₄+R₅=CH₂
 52: R₁=R₃=OH, R₂=H, R₄+R₅=CH₂
 53: R₁=OH, R₂=R₃=R₅=H, R₄=CH₃
 54: R₁=OH, R₂=R₃=H, R₄+R₅=CH₂




59




60




61




62




63



64



65



66

known alkaloids namely, 1-acetyl-β-carboline (62), incartine (63), *N*-trans feruloyltyramine (64), lycorine (59), *O*-methylincartine (65), vittatine (54) and 11-hydroxyvittatine (52) [33]. The potential of these alkaloids as AChEi was analyzed but only incartine *N*-oxide (60) was observed to elicit a moderate inhibitory activity (IC₅₀ = 34.50 μM), incartine (63) was observed to be weakly active (IC₅₀ = 106.97 μM) and the other alkaloids were found to be inactive. In a bioassay-guided fractionation of an active extract obtained from bulbs of *Nerine bowdenii*, the Amaryllidaceae alkaloid undulatine (66) was identified as the most active component of the alkaloid fraction, with an IC₅₀ = 37 μM [34].


Although benzylphenethylamine alkaloids were considered to belong exclusively to the Amaryllidaceae, some of them


have been found to belong to other families [35]. A new example of this exception was found through the chemical investigation of *Hosta plantaginea* (Liliaceae) [36]. Seventeen benzylphenethylamine alkaloids, including five new alkaloids, 67-71, along with twelve known compounds [7-deoxy-*trans*-dihydronarcislasine, *O*-methyllycorine, albomaculine, haemanthamine, *O*-demethylhaemanthamine, 8-*O*-demethylmaritadine, haemanthidine, yemenine C, lycorine, pseudolycorine, ungeremine (72) and norsanguinine (73)] were obtained. Some of these alkaloids were analyzed to determine whether they are AChEi or not. Ungeremine (72) (IC₅₀ = 3.85 μM), norsanguinine (73) (IC₅₀ = 1.43 μM) and 8-demethoxy-10-*O*-methylhostasine (69) (IC₅₀ = 2.32 μM) were all found to be potent AChE inhibitors.

After the isolation of the potent AChEi huperzine A (**5**) from *Huperzia serrata* (Lycopodiaceae), several plants belonging to the genus *Lycopodium* have been investigated in an attempt to find alkaloids with unusual skeletons that could have AChE inhibitory activity [7, 8, 37]. Five new *Lycopodium* alkaloids, 11 α -hydroxyfawcettidine (**74**), 2 α ,11 α -dihydroxyfawcettidine (**75**), 8 α ,11 α -dihydroxyfawcettidine (**76**), 2 β -hydroxylycothunine (**77**) and 8 α -hydroxylycothunine (**78**), with the fawcettimine skeleton were isolated from *L. serratum*, along with three known alkaloids, lycothunine (**79**), serratine (**80**) and serratanidine (**81**) [38]. AChE inhibitory activity was analyzed for the alkaloid lycoposerramine-H (**82**) previously isolated from *L. serratum* [39] and for compounds **74**, **75**, **78**. Alkaloids **75** and **82** were observed to inhibit AChE with an IC₅₀ = 27.9 and 16.7 μ M, respectively, while **74** and **78** were observed to show no anti-AChE activity. In another study, three new alkaloids (**83** - **85**) were isolated from *L. carinatum*, a species collected in Malaysia [40]. Carinatamins A (**83**) and B (**84**) were observed to inhibit AChE from bovine erythrocytes with an IC₅₀ = 4.6 and 7.0 μ M, respectively, whereas carinatumin C (**85**) was observed to show no inhibition (IC₅₀ > 100 μ M). Alkaloids **83** and **84** were observed to exhibit an AChE inhibitory activity similar to that of huperzine A and huperzine B (IC₅₀ = 0.8 and 8.0 μ M). Alkaloids from *L. casuarinoides* were also isolated and three new compounds, lycoparins A-C (**86** - **88**), were characterized, of which lycoparin C (**88**) was found to show a moderate AChE inhibitory activity (from bovine erythrocytes) with an IC₅₀ = 25 μ M [41]. Lycoparin A (**86**) and lycoparin B (**87**), both having a carboxylic acid at C-15 and one or two *N*-methyl groups, were found to show no inhibitory activity.

As to *Sarcococca* and *Buxus* species (Buxaceae), they are known to produce steroidal alkaloids, some of which were

observed to evidence strong AChE inhibition [7, 42, 43]. New steroidal alkaloid AChEi from *S. saligna* and *S. hookeriana* were recently found. In the case of *S. saligna*, the study –which was a continuation of previous research [44, 45]– of the bioactive steroidal alkaloids of this species allowed the isolation of five new compounds (**89-93**) and two already known bases (**94** and **95**) [46]. The new alkaloids 5,14-dehydro-*N*_a-demethylsaracodine (**89**), 14-dehydro-*N*_a-demethylsaracodine (**90**), 16-dehydrosarcovagine (**91**), 2,3-dehydrosarsalignone (**92**) and 14,15-dehydrosarcovagine D (**93**), as well as the known compounds sarcovagine C (**94**) and salignarine C (**95**) were analyzed as anti-AChE agents. Only **91**, **92** and **95** were observed to exhibit significant AChE inhibition (IC₅₀ = 12.5, 7.0 and 19.7 μ M, respectively). Compounds **89** - **92**, **94** and **95** were also found to elicit strong and selective BChE inhibition [46]. The bioassay-guided chemical investigation of *S. hookeriana* allowed the isolation of two new pregnane-type steroidal alkaloids, hookerianamide H (**96**) and hookerianamide I (**97**) together with the known alkaloids *N*_a-methylepipachysamine D (**98**), sarcovagine C (**94**) and dictyophlebine (**99**) [47]. Compounds **94**, **96**, **97**, **98** and **99** were tested for their inhibitory properties towards AChE and all of them were observed to elicit significant inhibitory activity (IC₅₀ 2.9 – 34.1 μ M) as well as a potent anti-BChE activity (IC₅₀ 0.3 – 3.6 μ M). Further studies on *S. hookeriana* yielded two new 5 α -pregnane-type steroidal alkaloids, hookerianamides J (**100**) and K (**101**) [48]. Furthermore, eight known steroidal alkaloids, hookerianamide H (**96**) and hookerianamide I (**97**), chonemorphine (**102**), *N*-methylpachysamine A (**103**), epipachysamine-*E*-5-en-4-one (**104**), vagenine A (**105**), 2,3-dehydrosarsalignone (**92**) and sarcovagine C (**94**), were isolated and characterized. Alkaloids **94**, **100**, **101**, **102**, **103** and **104** were analyzed as AChEi. Compounds **100**, **101**, **102**





89: R₁=R₃=R₄=H, R₂=CH₃, R₅=COCH₃, Δ^{5,6}, Δ^{14,15}

90: R₁=R₃=R₄=H, R₂=CH₃, R₅=COCH₃, Δ^{14,15}

91: R₁=R₃=R₄=H, R₂=COCH₃, R₅=CH₃, Δ^{16,17}

92: R₁=R₃=H, R₂=COCCH₃=CHCH₃, R₄=O, R₅=CH₃, Δ^{2,3}, Δ^{5,6}

93: R₁=R₃=H, R₂=COCCH₃=CHCH₃, R₄=O, R₅=CH₃, Δ^{2,3}, Δ^{14,15}

94: R₁=R₃=H, R₂=COCCH₃=CHCH₃, R₄=OCOCH₃, R₅=CH₃

95: R₁=OH, R₂=COCCH₃=CHCH₃, R₃=R₄=H, R₅=CH₃, Δ^{5,6}

96: R₁=R₃=H, R₂=COH, R₄=O, R₅=CH₃, Δ^{2,3}

97: R₁=R₄=R₅=H, R₂=PhCO, R₃=CH₃

98: R₁=R₄=H, R₂=PhCO, R₃=R₅=CH₃

99: R₁=R₃=R₄=H, R₂=R₅=CH₃

100: R₁=R₃=H, R₂=COCH=C(CH₃)₂, R₄=OH, R₅=CH₃, Δ^{16,17}

101: R₁=R₄=H, R₂=R₃=R₅=CH₃, Δ^{4,5}, Δ^{14,15}

102: R₁=R₂=R₃=R₄=H, R₅=CH₃

103: R₁=R₄=H, R₂=R₃=R₅=CH₃

104: R₁=R₃=R₅=H, R₂=COCH=C(CH₃)₂, R₄=O, Δ^{5,6}


105: R₁=R₅=H, R₂=COCH=C(CH₃)₂, R₄=OCOCH₃

and **103** were observed to inhibit AChE moderately (IC₅₀ 22.1 – 48.5 μM) while **104** and **94** were found to be more active inhibitors (IC₅₀ 9.9 and 8.1 μM, respectively).


Phytochemical research on *Buxus hyrcana* allowed the identification of several Buxus alkaloids with cholinesterase inhibitory activity [43, 49]. Three new triterpenoidal alkaloids, namely 17-oxo-3-benzoylbuxadine (**106**), buxhyrcamine (**107**) and 31-demethylcyclobuxoviridine (**108**) along with sixteen known compounds, all tested as AChEi, were isolated and characterized in a recent study on *B. hyrcana* collected from Iran [50]. Weak AChE inhibitory activity was observed for N_b-dimethylcyclobuxoviridine

(**109**), papillozine C (**110**), cyclobuxophylline O (**111**) and arbor-1,9(11)-dien-3-one (**112**) (IC₅₀ = 35.4 - 47.9 μM). In the same *in vitro* assay, 17-oxo-3-benzoylbuxadine (**106**), buxhyrcamine (**107**), homomoenjodaramine (**113**), buxmicrophylline F (**114**), buxrugulosamine (**115**), moenjodaramine (**116**) and N₂₀-formyl-buxaminol E (**117**) were observed to show moderate AChE inhibition (IC₅₀ = 17.6 - 25.5 μM) while spirofor nabuxine (**118**) was found to elicit a strong AChE inhibitory activity (IC₅₀ = 6.3 μM).

The crude methanolic extract of *B. natalensis*, a plant used to improve memory in the elderly by traditional healers in South Africa, was found to elicit AChE inhibition (IC₅₀ =




106




107: R₁=R₂=H

113: R₁=R₂=CH₃


116: R₁=CH₃, R₂=H




108




109



110




111




28 µg/mL). The phytochemical study of this extract yielded seven compounds **119** - **125** which were found to show either moderate or strong AChE inhibition [51]. The alkaloids *O*²-natafuranamine (**119**), *O*¹⁰-natafuranamine (**120**), cyclonatinol (**121**) and 31-demethylbuxaminol (**122**) were isolated and characterized for the first time while buxaminol A (**123**) was isolated for the first time as a natural product. Buxafuranamide (**124**) and buxalongifolamidine (**125**) were already known compounds. Compounds **119**, **120** and **124** were observed to exhibit a significantly higher AChE inhibitory activity compared to the rest, with IC₅₀ values of 3.0, 8.5, and 14.0 µM, respectively. Compounds **121**, **122**, **123** and **125** were observed to be less effective as AChEi (IC₅₀ = 22.9 – 30.2 µM).


The bulbs of *Fritillaria* species (Liliaceae) which are known to be a traditional medicinal herb called “Beimu” in




126




127



128



129




130

China are used as an antitussive, antiasthmatic and expectorant agent. In the past, in a chemical study carried out on alkaloids from *F. imperialis* bulbs new steroidal alkaloids with weak AChE inhibition and great selectivity towards BChE were identified [52]. Thus, taking into account this previous study, the bulbs from five *Fritillaria* species were studied and their alkaloids were identified and evaluated as cholinesterase inhibitors. Eighteen alkaloids were isolated and their effects on human whole blood cholinesterase were assayed. Results showed that *N*-demethyl-puqietinone (**126**) from *F. puziensis*, hupeheninoside (**127**) from *F. hupehensis*, ebeiedinone (**128**) from *F. ebeiensis* var. *purpurea*, yibeinoside A (**129**) from *F. pallidiflora* and chuanbeinone (**130**) from *F. delavayi* showed good AChE inhibition, with IC_{50} values of 6.4, 16.9, 5.7, 6.5 and 7.7 μ M, respectively. However, all of them were weaker AChEi than galanthamine ($IC_{50} = 1.9 \mu$ M). Compounds **127**, **128**, **129** and **130** were found to be stronger inhibitors on plasma BChE than galanthamine, the positive control [53].

In addition, the following steroidal alkaloids: conessine (**131**), isoconessimine (**132**), conessimin (**133**), conarrhimin (**134**) and conimin (**135**) were isolated in a bioassay-guided fractionation from the seeds of *Holarrhena antidysenterica* (Apocynaceae), a common Tibetan drug [54]. Compounds **131**, **133**, **134** and **135** were identified as active constituents against AChE. Conessimin (**133**) was found to be the strongest AChE inhibitor with an $IC_{50} = 4 \mu$ M whereas conessine (**131**), conarrhimin (**134**) and conimin (**135**) were found to be moderate AChE inhibitors ($IC_{50} = 21 - 28 \mu$ M). These findings indicate that the elimination of the *N*-methyl

group of pyrrolidine moiety induces a significant increase of activity while the cleavage of either one or two *N*-methyl groups at C-3 position reduces the inhibitory potency. Compound **133** was selected for a kinetic study through which it was demonstrated that its AChE inhibitory activity is both reversible and non-competitive. Molecular docking simulations of these compounds with AChE helped to understand their interactions with AChE and were consistent with the experimental results obtained [54].



131: $R_1=R_2=R_3=CH_3$

132: $R_1=R_2=CH_3, R_3=H$

133: $R_2=R_3=CH_3, R_1=H$

134: $R_1=R_2=R_3=H$

135: $R_1=R_3=H, R_2=CH_3$


NON-ALKALOIDAL COMPOUNDS WITH AChE INHIBITORY ACTIVITY


In spite of the fact that the majority of the most potent inhibitors known to date are alkaloids, several non-alkaloidal

AChEi from the plant kingdom and with different structural characteristics (terpenoids, sterols, flavonoids and phenolic compounds, etc) have been recognized as promising lead compounds as anti-AD agents [7-10]. Until 2006 only a few diterpenoids demonstrated to inhibit AChE [7]. However, further recent research has reported a larger number of compounds belonging to this group with the ability to exert either moderate or strong AChE inhibitory activity. In addition, a new cassane diterpene named niloticane (**136**) was isolated from the ethyl acetate bark extract of *Acacia nilotica* subsp. *kraussiana* (Fabaceae), a plant used in African traditional medicine [55]. Niloticane (**136**) was found to show an AChE inhibitory activity similar to that of the positive control galanthamine ($IC_{50} = 4$ and $2 \mu M$, respectively). In addition, one new (**137**) and six known (**138** - **143**) labdane-type diterpenoids were identified as AChE inhibitors present in an active extract obtained from *Leonurus heterophyllum* (Lamiaceae) by bioassay-guided fractionation [56]. Anti-AChE activity in **137** - **143** was

analyzed in rat brain cortex as a source of AChE enzyme. Leoheteronin A (**141**) and leopersin G (**143**), both having a 15,16 epoxy group, were observed to be strong inhibitors with IC_{50} values of 11.6 and $12.9 \mu M$, respectively. The new compounds leoheteronin F (**137**) and leoheteronin D (**142**) were found to show moderate inhibition with IC_{50} values of 16.1 and $18.4 \mu M$, respectively. Leoheterin (**138**), hispanone (**139**) and galeopsin (**140**), all having a furan ring at the side chain, were found to be weakly active ($IC_{50} = 38.5 - 42.7 \mu M$).


Asparagus adscendens (Asparagaceae) is a medicinal plant traditionally used as a nerve tonic and remedy for memory impairments in Pakistan. Conyopodiol (**144**), which was isolated from the chloroform fraction of the methanolic extract of *A. adscendens*, was found to elicit AChE and BChE inhibition with an $IC_{50} = 2.17$ and $11.21 \mu M$, respectively [57]. This dual cholinesterase inhibitor was also observed to show potential as a bivalent ligand in molecular docking studies. Four non-competitive AChEi **145**






– **148** were obtained in the chemical investigation of *Ajuga bracteosa* (Lamiaceae), another medicinal plant from Pakistan [58]. The diterpenoid dihydroajugapitin (**148**) was found to be the most active against AChE with an IC₅₀ = 14.0 μM. Compared to compound **148**, lupulin A (**147**), clerodin A (**146**) and dihydroclerodin (**145**) were observed to be less efficient inhibitors (IC₅₀ = 19.2, 26.5 and 35.2, respectively) and diterpenoids **145** - **148** were observed to elicit BChE inhibition. These findings indicate that the presence of a methoxy group at C-15 increases cholinesterase inhibitory potential.

From the methanolic extract of *Haloxylon recurvum* (Chenopodiaceae), a plant used in Pakistan for the treatment of several neuronal disorders, four new C-24 alkylated sterols **149** – **152** and five known sterols **153** – **157** were isolated [59]. Compounds **149** – **157** were analyzed as AChEi and were found to inhibit AChE in a concentration-dependent manner acting as non-competitive inhibitors. Haloxysterol B (**150**) and haloxysterol C (**151**), whose IC₅₀ values were 0.89 and 1.0 μM, respectively, were found to be the most active AChE inhibitors. Their inhibitory activity was observed to be similar to that of galanthamine (IC₅₀ 0.5



162




163

μM). Haloxysterol A (**149**) and 24-ethyl-cholest-6-ene-3,5-diol (**157**) were also observed to show potent AChE inhibition with IC_{50} values of 8.3 and 3.5 μM , respectively. Haloxysterol D (**152**), 5 α ,8 α -epidioxy-(24*S*)-ethyl-cholest-6,9(11),22(*E*)-triene-3 β -ol (**153**), (24*S*)-ethyl-cholest-7,9(11),22(*E*)-triene-3 β -ol (**154**), lawsaritol (**155**) and 24-ethyl-cholest-7-ene-3,5,6-triol (**156**) were found to elicit a moderate anti-AChE activity (IC_{50} = 13.7 - 26.4 μM).

On the other hand, a bioassay-guided fractionation on the bark of *Mesua elegans* (Clusiaceae) allowed the isolation of the anti-AChE components responsible for the activity observed for the extract. Mesuagenin B (**158**) was the most potent inhibitor (IC_{50} = 0.7 μM) and mesuagenin A (**159**), mesuagenin D (**160**) and 5,7-dihydroxy-8-(3-methylbutanoyl)-6-[(*E*)-3,7-dimethylocta-2,6-dienyl]-4-phenyl-2H-chromen-2-one (**161**) were observed to elicit strong AChE inhibition with IC_{50} values of 1.06, 8.73 and 3.06 μM , respectively [60]. This bioassay-guided study is the first report of 4-phenylcoumarins as AChEi.

In the past, some examples of xanthenes with moderate AChE inhibitory activity were reported [7]. Further recent research introduced two new xanthenes, **162** and **163**, to this group of AChEi also with moderate inhibitory activity. Macluraxanthone (**162**) which was obtained from the root of *Maclura pomifera* (Moraceae) was found to elicit non-competitive AChE inhibition (IC_{50} = 8.47 μM) [61]. Furthermore, docking studies yielded results supporting *in vitro* results. Triptexanthoside C (**163**) which was isolated from the methanolic extract of *Gentianella amarella* ssp. *acuta* (Gentianaceae) was observed to elicit AChE inhibition with an IC_{50} = 13.8 μM [62].

The methanolic extract of *Paulownia tormentosa* fruits, with a potent inhibitory activity against AChE, was subjected to bioactivity-guided fractionation which allowed the identification of some geranylated flavonoids, such as



164: $\text{R}_1=\text{R}_2=\text{R}_4=\text{OH}$, $\text{R}_3=\text{OCH}_3$

165: $\text{R}_1=\text{H}$, $\text{R}_2=\text{R}_4=\text{OH}$, $\text{R}_3=\text{OCH}_3$

166: $\text{R}_1=\text{R}_2=\text{H}$, $\text{R}_3=\text{R}_4=\text{OH}$


cholinesterase inhibitors, of which the most active resulted to be 6-geranyl-3,3',5,5',7-pentahydroxy-4'-methoxyflavane (**164**), 6-geranyl-3',5,5',7-tetrahydroxy-4'-methoxyflavanone (**165**) and diplacone (**166**), which were observed to show mixed-type inhibition of human AChE with IC_{50} = 15.6, 22.9 and 7.2 μM , respectively [63]. In addition, the fact that these compounds were also observed to elicit significant BChE inhibition makes them interesting as potential dual inhibitors.

The flavonols present in *Sophora flavescens* (Fabaceae) were studied for several biological activities relevant for AD. Sophoflavescenol (**167**), icaritin (**168**), demethylanhydroicaritin (**169**), 8-C-lavandurylkaempferol (**170**) and kaempferol (**171**) were all found to be good AChE inhibitors, with IC_{50} values of 8.37, 6.47, 6.67, 5.16 and 3.31 μM , respectively [64]. Compounds **167-171** were also found to elicit significant BChE and BACE1 inhibition.

The methanol extract from roots of *Morus lhou* (Moraceae), a polyphenol-rich plant, was found to yield nine flavonoids (**172** - **180**) of which eight showed AChE inhibition [65]. A new flavone, 5'-geranyl-4'-methoxy-5,7,2'-trihydroxyflavone (**172**), was identified as the most potent inhibitor (IC_{50} = 10.95 μM). 5'-geranyl-5,7,2',4'-tetrahydroxyflavone (**173**), kuwanon U (**174**), kuwanon E (**175**), morusin (**176**), cyclomorusin (**178**), neocyclomorusin (**179**) and kuwanon C (**180**) were all observed to be moderate AChE inhibitors (IC_{50} = 16.21 - 36.4 μM) and morusinol (**177**) was observed to be weakly active (IC_{50} = 173.49 μM). C-3 prenylated flavones **176**, **178**, **179** and **180** were found to be noncompetitive inhibitors whereas those unsubstituted at C-3 **172-175** were mixed inhibitors. Flavonoids **172** - **180** were also found to inhibit BChE [65].

On the other hand, three potent AChEi were obtained from *Broussonetia papyrifera*, another plant belonging to the Moraceae family. From the ethanolic extract of the roots of *B. papyrifera* which was found to elicit cholinesterase inhibitory activity, prenylated flavonols **181** - **183** were isolated and characterized [66]. 8-(1,1-dimethylallyl)-5'-(3-methylbut-2-enyl)-3',4',5,7-tetrahydroxyflavonol (**181**), papyriflavonol (**182**) and brousoflavonol (**183**) were observed to inhibit human erythrocyte AChE with IC_{50} values of 0.82, 3.1 and 2.7 μM , respectively. Compound **181**, the most potent, acted as a time-dependent, slow reversible inhibitor.


Isoorientin (**184**) and isovitexin (**185**) were identified as the compounds responsible for the AChE inhibition observed in the extracts from flowers and rhizomes of *Iris pseudopumila* (Iridaceae) from Italy [67]. Compound **184**




167: R₁=CH₃, R₂=H

168: R₁=H, R₂=CH₃


169: R₁=R₂=H




170




171




172




173




174




175




176




177



178



179







180



was observed to be the highest inhibitor with an IC₅₀ = 26.8 μM while **185**, lacking the 3'-hydroxy group in ring B, was observed to show an IC₅₀ value of 36.4 μM. Both compounds were also found to have the ability of significantly inhibiting BChE.

On the other hand, a pterocarpan with moderate AChE inhibition was isolated from the polar extract of *Zygophyllum*



eurypterum (Zygophyllaceae) collected in Pakistan. Atricarpan D [(-)-2,9-dimethoxy-4-(5-oxohexyl)pterocarpan] (**186**) was observed to inhibit AChE with an IC₅₀ = 20.5 μM [68]. Interestingly, three other pterocarpan with similar structure were obtained along with atricarpan D but they were found to be inactive against AChE. Nevertheless, the four pterocarpan were all found to be BChE inhibitors.



A study conducted on AChE and BChE inhibitory activity of coumarins and naphthoquinones obtained from *Mansonia gagei* (Sterculiaceae) proposed a novel class of cholinesterase inhibitor, mansonones or 1,2-naphthoquinones [69]. The level of cholinesterase inhibition observed in this study seemed to correlate to the presence of a fused pyran ring and a substituent at C-6 being present in the molecule. Mansonone E (**187**) was observed to be the most active AChE ($IC_{50} = 23.5 \mu M$) and BChE inhibitor.





In several studies published during the period covered in the present review various phenolic compounds with different structural characteristics were reported as AChEi. Some of them are structurally simple such as gallic acid (**188**, $IC_{50} = 5.85 \mu M$) and ellagic acid (**189**, $IC_{50} = 45.63 \mu M$) [70]. Hopeahainol A (**190**), which was identified as a new compound isolated from *Hopea hainensis*, was observed



to elicit a notable AChE inhibition ($IC_{50} = 4.33 \mu M$) with respect to huperzine A ($IC_{50} = 1.6 \mu M$), as a reversible mixed-type inhibitor [71].

The bioassay-guided fractionation of the extract from *Terminalia chebula* (Combretaceae) fruits allowed the isolation of 1,2,3,4,6-penta-*O*-galloyl- β -D-glucose (**191**) which demonstrated to be a significant AChE inhibitor ($IC_{50} = 29.9 \mu M$) [72]. This gallotanin which has been also isolated from other different sources and which is known by its diverse biological activities, was observed to exert good BChE inhibition and potent antioxidant activity (FRAP assay) in this study.

The bioassay-guided extraction of the stem bark of *Knema laurina* (Myristicaceae) yielded two active fractions (dichloromethane and hexane) which were subjected to chromatographic separation. That latter yielded five alkenyl phenol and salicylic acid derivatives **192** - **196**, of which **192** and **193** were new compounds [73]. Compounds **192**, **195** and **196**, all having salicylic acid moiety, were observed to strongly inhibit AChE with an $IC_{50} = 3.182$, 2.172 and 0.573 μM , respectively. Compounds **193** and **194**, with no carboxyl moiety, were observed to be good AChE inhibitors ($IC_{50} = 17.224$ and 13.114 μM , respectively). These findings suggest that the acidic group is key to good AChE inhibition. It was also observed that anti-AChE activity dramatically decreased when the acidic and the phenolic hydroxy group were methylated. Two catechol alkenyls were isolated from the fruits of *Semecarpus anacardium* (Anacardiaceae), a species used in Ayurvedic medicine for retarding and treatment of memory loss [74]. Compounds **197** and **198** were identified as active components of the dichloromethane extract through a fractionation guided by the detection of AChE inhibition. Microplate assay revealed that these catechol alkenyls are moderate and weak selective AChEi. Compound **197**, with a double bond in the aliphatic chain, was identified as a stronger inhibitor ($IC_{50} = 39.7 \mu M$) with respect to compound **198**, with two double bonds in the aliphatic chain ($IC_{50} = 108 \mu M$).



On the other hand, four structurally diverse AChEi were isolated from the polar extract of *Nelumbo nucifera* (Nelumbonaceae) stamens [75]. Cycloartenol (**199**), *p*-hydroxybenzoic acid (**200**), vanilloloside (**201**) and nuciferoside (**202**) were found to elicit good and noncompetitive inhibition against AChE with an IC₅₀ = 11.89, 20.07, 4.55 and 3.2 μM, respectively. In the same study, compounds **199**, **200** and **202** were observed to exert moderate BChE inhibition and compounds **199** - **202** were found to show no inhibition against BACE1.

PLANT EXTRACTS, FRACTIONS AND ESSENTIAL OILS WITH AChE INHIBITORY ACTIVITY

Table 1 summarizes the studies published from 2006 to 2012 on plant extracts, fractions and essential oils that have been found to be good AChE inhibitors (IC₅₀ < 500 μg/mL). Those plants included in other recent reviews were omitted [76, 77]. Extracts and fractions under further phytochemical studies that led to the discovery of AChE inhibitors were also omitted. Whenever possible, reference is made to the

Table 1. Plant Extracts, Fractions and Essential Oils with AChE Inhibitory Activity

Family and Botanical Name	Type of Extract (Solvent)	Plant's Parts	AChE Inhibition (%)	IC ₅₀	Refs.
Acanthaceae					
<i>Andrographis paniculata</i>	H ₂ O:EtOH	Aerial		222.41 µg/ml	[78]
Amaranthaceae					
<i>Salsola oppositifolia</i>	Alkaloids	Aerial		70.0 µg/ml	[79]
<i>Salsola soda</i>	Alkaloids	Aerial		64.1 µg/ml	[79]
<i>Salsola tragus</i>	Alkaloids	Aerial		30.2 µg/ml	[79]
Amaryllidaceae					
<i>Crinum jagus</i>	MeOH	Leaf	74.25 (42 µg/ml)		[80]
<i>Crinum moorei</i>	50% MeOH PE DCM EtOH	Bulb		21.5 µg/ml 18.9 µg/ml 2.9 µg/ml 22.5 µg/ml	[81]
<i>Nerine undulata</i>	Alkaloids	Bulb		14.3 µg/ml ^a	[82]
<i>Scadoxus multiflorus</i>	Alkaloids	Bulb		313.5 µg/ml ^a	[82]
<i>Sprekelia formosissima</i>	Alkaloids	Bulb		209.7 µg/ml ^a	[82]
<i>Zephyranthes grandiflora</i>	Alkaloids	Bulb		39.2 µg/ml ^a	[83]
Anacardiaceae					
<i>Harpephyllum caffrum</i>	DCM MeOH	Leaf Stem bark Leaf		0.17 mg/ml 0.02 mg/ml 0.12 mg/ml	[84]
<i>Pistacia atlantica</i>	H ₂ O	Leaf		0.87 µg/ml	[85]
<i>Pistacia lentiscos</i>	H ₂ O	Leaf		13.67 µg/ml	[85]
<i>Sclerocarya birrea</i>	DCM MeOH	Young stem Leaf Operculum Young stem		0.15 mg/ml 0.10 mg/ml 0.35 mg/ml 0.47 mg/ml	[84]
<i>Spondias mombin</i>	MeOH	Root bark	64.77 (42 µg/ml)		[80]
Apiaceae					
<i>Centella asiatica</i>	H ₂ O:EtOH	Whole plant		106.55 µg/ml	[78]
Apocynaceae					
<i>Geissospermum vellosii</i>	Alkaloids	Stem bark		2.9 µg/ml	[86]
Araceae					
<i>Colocasia antiquorum</i>	50% MeOH PE DCM	Tuber		7.9 µg/ml 6.4 µg/ml 168.1 µg/ml	[81]
<i>Pinellia ternata</i>	Alkaloids	Tuber		56.2 µg/ml	[87]
Areaceae					
<i>Phoenix dactylifera</i>	Hexane	Seed	52.96 (300 µg/ml)		[88]

Table 1. contd....

Family and Botanical Name	Type of Extract (Solvent)	Plant's Parts	AChE Inhibition (%)	IC ₅₀	Refs.
Asparagaceae					
<i>Leopoldia comosa</i>	Hexane	Bulb		104.9 µg/ml	[89]
Asphodelaceae					
<i>Aloe ferox</i>	50% MeOH PE DCM	Leaf		84.0 µg/ml 37.7 µg/ml 62.6 µg/ml	[81]
Asteraceae					
<i>Achyrocline tomentosa</i>	Organic	Aerial		0.4847 mg/ml	[90]
<i>Arnica chamissonis</i> ssp. <i>foliosa</i>	MeOH Hexane	Flower		43 µg/ml 29 µg/ml	[91]
<i>Chromolaena tequendamensis</i>	MeOH	Whole plant		359.36 mg/l	[92]
<i>Eupatorium viscidum</i>	Organic	Aerial		0.4792 mg/ml	[90]
<i>Pulicaria stephanocarpa</i>	CHCl ₃	Leaf	61.43 (0.2 mg/ml)		[93]
<i>Schistocarpha sinforosi</i>	MeOH	Whole plant		145.31 mg/l	[92]
<i>Trichocline reptans</i>	Organic	Aerial		0.1118 mg/ml	[90]
Berberidaceae					
<i>Berberis darwinii</i>	MeOH	Stem bark		1.23 µg/ml	[94]
Boraginaceae					
<i>Onosma bracteata</i>	MeOH	Leaf	59.73 (250 µg/ml)		[95]
Buddlejaceae					
<i>Buddleja salviifolia</i>	DCM:MeOH (1:1)	Whole plant		0.05 mg/ml	[96]
Burseraceae					
<i>Boswellia socotranao</i>	CHCl ₃	Resin	71.21 (0.2 mg/ml)		[93]
Cistaceae					
<i>Cistus laurifolius</i>	EtOH	Leaf	80.07 (200 µg/ml)		[97]
Combretaceae					
<i>Terminalia bellirica</i>	MeOH	Fruit		14.37 µg/ml	[70]
Convolvulaceae					
<i>Evolvulus alsinoides</i>	H ₂ O:EtOH	Whole plant		141.76 µg/ml	[78]
<i>Ipomoea asarifolia</i>	MeOH	Leaf		0.12 µg/ml	[98]
Crassulaceae					
<i>Kalanchoe brasiliensis</i>	EtOAc	Leaf		0.16 mg/ml	[98]
Cucurbitaceae					
<i>Eureiandra balfourii</i>	MeOH	Tuber	58.61 (0.2 mg/ml)		[93]
Cupressaceae					
<i>Juniperus phoenicea</i>	EtOH	Leaf	53.44 (400 µg/ml)		[99]
<i>Juniperus turbinata</i>	Phenolic	Leaf	83.84 (400 µg/ml)		[99]

Table 1. contd....

Family and Botanical Name	Type of Extract (Solvent)	Plant's Parts	AChE Inhibition (%)	IC ₅₀	Refs.
Ericaceae					
<i>Rhododendron yedoense</i> var. <i>poukhanense</i>	80% MeOH	Bark		169.01 µg/ml	[100]
Eucommiaceae					
<i>Eucommia ulmoides</i>	H ₂ O	Bark		172 µg/ml	[101]
Euphorbiaceae					
<i>Alchornia laxiflora</i>	MeOH	Stem bark	41.12 (42 µg/ml)		[80]
<i>Cephalocroton socotranus</i>	CHCl ₃	Bark	51.1 (0.2 mg/ml)		[93]
<i>Jatropha curcas</i>	MeOH	Leaf		0.25 mg/ml	[98]
<i>Jatropha gossypifolia</i>	MeOH	Leaf		0.05 mg/ml	[98]
Fabaceae					
<i>Acacia nilotica</i>	H ₂ O	Root		0.079 mg/ml ^b	[102]
<i>Acacia raddiana</i>	H ₂ O	Bark		33.91 µg/ml	[85]
<i>Cassia obtusifolia</i>	EtOH	Seed		81.6 µg/ml ^c	[103]
<i>Chamaecrista mimosoides</i>	DCM:MeOH (1:1) H ₂ O	Root		0.03 mg/ml 0.35 mg/ml	[96]
<i>Genista tenera</i>	EtOAc	Aerial	77.0 (70 µg/ml)		[105]
<i>Peltophorum pterocarpum</i>	MeOH	Leaf Stem bark	49.5 (42 µg/ml) 68.85 (42 µg/ml)		[80]
<i>Schotia brachypetala</i>	DCM:MeOH (1:1) H ₂ O	Bark		0.27 mg/ml 0.49 mg/ml	[96]
<i>Senna alata</i>	EtOAc	Leaf		0.08 mg/ml	[98]
<i>Spatholobus suberectus</i>	H ₂ O EtOH	Whole plant		85 µg/ml 9 µg/ml	[104]
<i>Trigonella foenum-graecum</i>	EtOAc Alkaloids	Seed		53.00 µg/ml 9.23 µg/ml	[106]
Gobulariaceae					
<i>Globularia alypum</i>	H ₂ O	Root		16.67 µg/ml	[85]
Guttiferaceae					
<i>Callophyllum inophyllum</i>	MeOH	Root bark	56.52 (42 µg/ml)		[80]
Hypericaceae					
<i>Hypericum perforatum</i>	MeOH	Whole plant		178 µg/ml	[91]
Illiciaceae					
<i>Illicium verum</i>	H ₂ O:EtOH Butanol EtOAc CHCl ₃ Oil	Fruit		58.67 µg/ml 44.94 µg/ml 83.75 µg/ml 103.03 µg/ml 39.89 µg/ml	[107]

Table 1. contd....

Family and Botanical Name	Type of Extract (Solvent)	Plant's Parts	AChE Inhibition (%)	IC ₅₀	Refs.
Lamiaceae					
<i>Cyclotrichium niveum</i>	EtOAc	Whole plant	83.11 (250 µg/ml)		[108]
	DCM		70.82 (250 µg/ml)		
<i>Hyssopus officinalis</i>	Hexane	Whole plant	55.0 (400 µg/ml)		[91]
<i>Lavandula viridis</i>	MeOH	Aerial		244.55 µg/ml	[109]
<i>Marrubium vulgare</i>	Acetone	Aerial	62.70 (25 µg/ml)		[110]
<i>Origanum ehrenbergii</i>	Essential oil	Aerial		0.3 µg/ml	[111]
<i>Origanum majorana</i>	Essential oil	Leaf		36.40 µg/ml	[112]
<i>Origanum syriacum</i>	Essential oil	Aerial		1.7 µg/ml	[111]
<i>Pycnostachys reticulata</i>	50% MeOH	Leaf		28.8 µg/ml	[81]
	EtOH		8.8 µg/ml		
<i>Salvia chionantha</i>	Essential oil	Aerial	56.7 (500 µg/ml)		[113]
<i>Salvia fruticosa</i>	DCM	Whole plant	51.07 (100 µg/ml)		[114]
<i>Salvia lerifolia</i>	Essential oil	Aerial		0.32 µl/ml	[115]
<i>Salvia miltiorrhiza</i>	H ₂ O	Root		50 µg/ml	[104]
	EtOH		5 µg/ml		
<i>Teucrium royleanum</i>	MeOH	Whole plant	52.4 (40 µg/0.2ml)		[116]
Menispermaceae					
<i>Stephania pierrei</i>	EtOH	Tuber		5.68 µg/ml	[117]
<i>Tinospora cordifolia</i>	MeOH	Stem		38.36 µg/ml	[118]
Moraceae					
<i>Dorstenia gigas</i>	CHCl ₃	Leaf	65.12 (0.2 mg/ml)		[93]
<i>Ficus religiosa</i>	MeOH	Stem bark		73.69 µg/ml	[118]
Myristicaceae					
<i>Myristica fragrans</i>	H ₂ O:EtOH	Seed		133.28 µg/ml	[78]
<i>Embelia ribes</i>	MeOH	Root		23.04 µg/ml	[118]
Orchidaceae					
<i>Orchis mascula</i>	MeOH	Root	56.99 (250 µg/ml)		[119]
Paeoniaceae					
<i>Paeonia lactiflora</i>	H ₂ O	Root		20 µg/ml	[104]
	EtOH		8 µg/ml		
<i>Paeonia veitchii</i>	H ₂ O	Root		14 µg/ml	[104]
	EtOH		45 µg/ml		
Papaveraceae					
<i>Corydalis intermedia</i>	MeOH	Whole plant	84 (100 µg/ml)		[120]
	H ₂ O	Tuber	97 (100 µg/ml)		
		Whole plant	57 (100 µg/ml)		
		Tuber	78 (100 µg/ml)		

Table 1. contd....

Family and Botanical Name	Type of Extract (Solvent)	Plant's Parts	AChE Inhibition (%)	IC ₅₀	Refs.
Papaveraceae					
<i>Corydalis solida</i> ssp. <i>laxa</i>	MeOH	Whole plant	89 (100 µg/ml)		[120]
	H ₂ O	Tuber	96 (100 µg/ml)		
		Whole plant	78 (100 µg/ml)		
		Tuber	85 (100 µg/ml)		
<i>Corydalis solida</i> ssp. <i>slivenensis</i>	MeOH	Whole plant	82 (100 µg/ml)		[120]
	H ₂ O	Tuber	97 (100 µg/ml)		
		Whole plant	48 (100 µg/ml)		
		Tuber	87 (100 µg/ml)		
Phyllanthaceae					
<i>Emblica officinalis</i>	MeOH	Fruit		29.26 µg/ml	[70]
Pinaceae					
<i>Pinus halepensis</i>	EtOH	Needle	60.15 (200 µg/ml)		[121]
	Essential oil	Twig	83.91 (200 µg/ml)		
<i>Pinus heldreichii</i> subsp. <i>leucodermis</i>	Essential oil	Needle		51.1 µg/ml	[122]
<i>Pinus nigra</i> subsp. <i>nigra</i>	Essential oil	Needle		94.4 µg/ml	[122]
<i>Pinus nigra</i> subsp. <i>calabrica</i>	Essential oil	Needle		101.5 µg/ml	[122]
<i>Pinus pinaster</i>	Pycnogenol	Bark	63.33 (200 µg/ml)		[121]
Piperaceae					
<i>Piper nigrum</i>	EtOH	Fruit		30.67 µg/ml	[117]
Poaceae					
<i>Cymbopogon jawarancusa</i>	MeOH	Whole plant	72.36 (250 µg/ml)		[95]
<i>Cymbopogon schoenanthus</i>	Essential oil	Fresh leaf (mountain reg./ desert reg.)		0.26 / 0.67 mg/ml	[123]
		Dried leaf (mountain reg./ desert reg.)		0.44 / 0.52 mg/ml	
		Dried root (mountain reg./ desert reg.)		0.27 / 0.32 mg/ml	
<i>Cymbopogon schoenanthus</i>	Hexane	Shoot (mountain reg./ desert reg.)		0.50 / 0.54 mg/ml	[124]
	DCM			0.57 / 0.30 mg/ml	
	EtOAc			0.23 / 0.30 mg/ml	
	MeOH			0.23 / 0.25 mg/ml	
	H ₂ O			0.46 / 0.04 mg/ml	
Polygonaceae					
<i>Fallopia multiflora</i>	H ₂ O	Root		13 µg/ml	[104]
	EtOH			65 µg/ml	
<i>Rheum palmatum</i>	H ₂ O	Root and Rizhome		32 µg/ml	[104]
	EtOH			18 µg/ml	
<i>Ruprechtia apetala</i>	EtOH	Aerial		0.0779 mg/ml	[90]
Portulacaceae					
<i>Portulaca oleracea</i>	Alkaloids	Upper part		29.4 µg/ml	[87]

Table 1. contd....

Family and Botanical Name	Type of Extract (Solvent)	Plant's Parts	AChE Inhibition (%)	IC ₅₀	Refs.
Rhamnaceae					
<i>Rhamnus prinoides</i>	H ₂ O	Root		0.201 mg/ml ^b	[102]
Rosaceae					
<i>Leucosidea sericea</i>	PE	Leaf		0.16 mg/ml	[125]
	DCM	Stem		0.14 mg/ml	
	MeOH			0.24 mg/ml	
	PE			0.26 mg/ml	
Rubiaceae					
<i>Galium odoratum</i>	Hexane	Whole plant	53.1 (400 µg/ml)		[91]
<i>Morinda citrifolia</i>	EtOH	Fruit		138.4 µg/ml	[126]
	CHCl ₃			78.11 µg/ml	
<i>Morinda lucida</i>	MeOH	Leaf	40.15 (42 µg/ml)		[80]
Rutaceae					
<i>Citrus aurantifolia</i>	Essential oil	Leaf		139 µg/ml	[127]
<i>Citrus medica</i>	Essential oil	Peel		171.3 µg/ml	[128]
<i>Ruta graveolens</i>	MeOH	Whole plant	59.1 (100 µg/ml)	34 µg/ml	[91]
	Hexane				
<i>Zanthoxylum coco</i>	Organic	Aerial		0.1579 mg/ml	[90]
Solanaceae					
<i>Solanum leucocarpum</i>	MeOH	Whole plant		204.59 mg/l	[92]
<i>Withania somnifera</i>	MeOH	Root		33.38 µg/ml	[118]
<i>Witheringia coccoloboides</i>	MeOH	Whole plant		220.68 mg/l	[92]
Valerianaceae					
<i>Nardostachys jatamansi</i>	H ₂ O:EtOH	Rhizome		130.11 µg/ml	[78]
	MeOH			47.21 µg/ml	[118]
Zingiberaceae					
<i>Kaempferia parviflora</i>	EtOH	Rhizome		20.64 µg/ml	[117]

DCM: dichloromethane; MeOH: methanol; EtOH: ethanol; PE: petroleum ether; EtOAc: ethyl acetate

^aHuman blood AChE.

^bBovine erythrocyte AChE.

^cMouse brain homogenized.

part of the plant used in each study reported. AChE inhibitory activity is reported in the same way as it was reported by authors and IC₅₀ values were chosen instead of inhibition percentages when both were available.

CONFLICT OF INTEREST

The author(s) confirm that this article content has no conflict of interest.

ACKNOWLEDGEMENTS

CONICET, CIC, UNS and ANPCYT.

REFERENCES

- [1] Chopra, K.; Misra, S.; Kuhad, A. Current perspectives on pharmacotherapy of Alzheimer's. *Expert. Opin. Pharmacother.*, **2011**, *12*, 335-350.
- [2] Ellman, G.L.; Courtney, K.D.; Andres, V.; Featherstone, R.M. A new and rapid colorimetric determination of acetylcholinesterase activity. *Biochem. Pharmacol.*, **1961**, *7*, 88-95.
- [3] Lopez, S.; Bastida, J.; Viladomat, F.; Codina, C. Acetylcholinesterase inhibitory activity of some Amaryllidaceae alkaloids and *Narcissus* extracts. *Life Sci.*, **2002**, *71*, 2521-2529.
- [4] Rhee, I.K.; van de Meent, M.; Ingkaninan, K.; Verpoorte, R. Screening for acetylcholinesterase inhibitors from Amaryllidaceae

- using silica gel thin-layer chromatography in combination with bioactivity staining. *J. Chromatogr. A*, **2001**, *915*, 217-223.
- [5] Marston, A.; Kissling, J.; Hostettmann, K. A rapid TLC bioautographic method for the detection of acetylcholinesterase and butyrylcholinesterase inhibitors in plants. *Phytochem. Anal.*, **2002**, *13*, 51-54.
- [6] Di Giovanni, S.; Borloz, A.; Urbain, A.; Marston, A.; Hostettmann, K.; Carrupt, P.A.; Reist, M. In vitro screening assays to identify natural or synthetic acetylcholinesterase inhibitors: thin layer chromatography versus microplate methods. *Eur. J. Pharm. Sci.*, **2008**, *33*, 109-119.
- [7] Houghton, P.J.; Ren, Y.; Howes, M.J. Acetylcholinesterase inhibitors from plants and fungi. *Nat. Prod. Rep.*, **2006**, *23*, 181-199.
- [8] Williams, P.; Sorribas, A.; Howes, M.J. Natural Products as a source of Alzheimer's drugs leads. *Nat. Prod. Rep.*, **2011**, *28*, 48-77.
- [9] Mukherjee, P.K.; Kumar, V.; Mal, M.; Houghton, P.J. Acetylcholinesterase inhibitors from plants. *Phytochemistry*, **2007**, *14*, 289-300.
- [10] Orhan, G.; Orhan, I.; Subutay-Oztekin, N., Ak, F.; Sener, B. Contemporary anticholinesterase pharmaceuticals of natural origin and their synthetic analogues for the treatment of Alzheimer's disease. *Recent. Pat. CNS Drug. Discov.*, **2009**, *4*, 43-51.
- [11] Rahman, A.U.; Khalid, A.; Sultana, N., Ghayur, M.N.; Mesaik, M.A.; Khan, M.R.; Gilani, A.H.; Choudhary, M.I. New natural cholinesterase inhibiting and calcium channel blocking quinoline alkaloids. *J. Enzyme Inhib. Med. Chem.*, **2006**, *21*, 703-710.
- [12] Cardoso-Lopes, E.M.; Maier, J.A.; da Silva, M.R.; Regasini, L.O.; Simote, S.Y.; Lopes, N.P.; Pirani, J.R.; da Silva Bolzani, V.; Marx Young, M.C. Alkaloids from stems of *Esenbeckia leiocarpa* Engl. (Rutaceae) as potential treatment for Alzheimer Disease. *Molecules*, **2010**, *15*, 9205-9213.
- [13] Yang, Z.; Zhang, D.; Ren, J.; Yang, M. Skimmianine, a furoquinoline alkaloid from *Zanthoxylum nitidum* as a potential acetylcholinesterase inhibitor. *Med. Chem. Res.*, **2012**, *21*, 722-725.
- [14] Mukherjee, P.K.; Mukherjee, D.; Maji, A.K.; Rai, S.; Heinrich, M. The sacred lotus (*Nelumbo nucifera*) - phytochemical and therapeutic profile. *J. Pharm. Pharmacol.*, **2009**, *61*, 407-422.
- [15] Yang, Z.; Zhang, X.; Du, J.; Ma, Z.J.; Guo, F.; Li, S.; Yao, X.J. An aporphine alkaloid from *Nelumbo nucifera* as an acetylcholinesterase inhibitor and the primary investigation for structure-activity correlations. *Nat. Prod. Res.*, **2012**, *26*, 387-392.
- [16] Mollataghi, A.; Coudiere, E.; Hadi, A.H.; Mukhtar, M.R.; Awang, K.; Litaudon, M.; Ata, A. Anti-acetylcholinesterase, anti- α -glucosidase, anti-leishmanial and anti-fungal activities of chemical constituents of *Beilschmiedia* species. *Fitoterapia*, **2012**, *83*, 298-302.
- [17] Hung, T.M.; Na, M.; Dat, N.T.; Ngoc, T.M.; Youn, U.; Kim, H.J.; Min, B.S.; Lee, J.; Bae, K. Cholinesterase inhibitory and anti-amnesic activity of alkaloids from *Corydalis turtschaninovi*. *J. Ethnopharmacol.*, **2008**, *119*, 74-80.
- [18] Hung, T.M.; Ngoc, T.M.; Youn, U.J.; Min, B.S.; Na, M.; Thuong, P.T.; Bae, K. Anti-amnesic activity of pseudocoptisine from *Corydalis tuber*. *Biol. Pharm. Bull.*, **2008**, *31*, 159-162.
- [19] Jung, H.A.; Min, B.S.; Yokozawa, T.; Lee, J.H.; Kim, Y.S.; Choi, J.S. Anti-Alzheimer and antioxidant activities of *Coptidis Rhizoma* alkaloids. *Biol. Pharm. Bull.*, **2009**, *32*, 1433-1438.
- [20] Chlebek, J.; Macáková, K.; Cahlíkovi, L.; Kurfürst, M.; Kunes, J.; Opletal, L. Acetylcholinesterase and butyrylcholinesterase inhibitory compounds from *Corydalis cava* (Fumariaceae). *Nat. Prod. Commun.*, **2011**, *6*, 607-610.
- [21] Ingkaninan, K.; Phengpa, P.; Yuenyongsawad, S.; Khorana, N. Acetylcholinesterase inhibitors from *Stephania venosa* tuber. *J. Pharm. Pharmacol.*, **2006**, *58*, 695-700.
- [22] Cho, K.M.; Yoo, I.D.; Kim, W.G. 8-Hydroxydihydrochelerythrine and 8-Hydroxydihydroanguinarine with a potent acetylcholinesterase inhibitory activity from *Chelidonium majus* L. *Biol. Pharm. Bull.*, **2006**, *29*, 2317-2320.
- [23] Rollinger, J.M.; Schuster, D.; Baier, E.; Ellmerer, E.P.; Langer, T.; Stuppner, H. Taspine: bioactivity-guided isolation and molecular ligand-target insight of a potent acetylcholinesterase inhibitor from *Magnolia x soulangiana*. *J. Nat. Prod.*, **2006**, *69*, 1341-1346.
- [24] Pereira, D.M.; Ferreres, F.; Oliveira, J.M.; Gaspar, L.; Faria, J.; Valentão, P.; Sottomayer, M.; Andrade, P.B. Pharmacological effects of *Catharanthus roseus* root alkaloids in acetylcholinesterase inhibition and cholinergic neurotransmission. *Phytochemistry*, **2010**, *17*, 646-652.
- [25] Zhan, Z.J.; Yu, Q.; Wang, Z.L.; Shan, W.G. Indole alkaloids from *Ervatamia hainanensis* with potent acetylcholinesterase inhibition activities. *Bioorg. Med. Chem. Lett.*, **2010**, *20*, 6185-6187.
- [26] Andrade, M.T.; Lima, J.A.; Pinto, A.C.; Rezende, C.M.; Carvalho, M.P.; Epifanio, R.A. Indole alkaloids from *Tabernaemontana australis* (Muell. Arg) Miens that inhibit acetylcholinesterase enzyme. *Bioorg. Med. Chem.*, **2005**, *13*, 4092-4095.
- [27] Ingkaninan, K.; Changwijit, K.; Suwanborirux, K. Vobasiny-iboga bisindole alkaloids, potent acetylcholinesterase inhibitors from *Tabernaemontana divaricata* root. *J. Pharm. Pharmacol.*, **2006**, *58*, 847-852.
- [28] Yang, Z.; Duan, D.Z.; Du, J.; Yang, M.J.; Li, S.; Yao, X.J. Geissoschizine methyl ether, a corynanthean-type indole alkaloid from *Uncaria rhynchophylla* as a potential acetylcholinesterase inhibitor. *Nat. Prod. Res.*, **2012**, *26*, 22-28.
- [29] Seidl, C.; Correia, B.L.; Stinghen, A.E.; Santos, C.A. Acetylcholinesterase inhibitory activity of uleine from *Himatanthus lancifolius*. *Z. Naturforsch. C.*, **2010**, *65*, 440-444.
- [30] Jin, Z. Amaryllidaceae and Scelletium alkaloids. *Nat. Prod. Rep.*, **2007**, *24*, 886-905.
- [31] de Andrade, J.P.; Berkov, S.; Viladomat, F.; Codina, C.; Zuanazzi, J.A.; Bastida, J. Alkaloids from *Hippeastrum papillo*. *Molecules*, **2011**, *16*, 7097-7104.
- [32] Berkov, S.; Codina, C.; Viladomat, F.; Bastida, J. N-Alkylated galanthamine derivatives: Potent acetylcholinesterase inhibitors from *Leucojum aestivum*. *Bioorg. Med. Chem. Lett.*, **2008**, *18*, 2263-2266.
- [33] Sarikaya, B.B.; Kaya, G.I.; Onur, M.A.; Viladomat, F.; Codina, C.; Bastida, J.; Somer, N.U. Alkaloids from *Galanthus rizehensis*. *Phytochem. Lett.*, **2012**, *5*, 367-370.
- [34] Rijn, R.M.; Rhee, I.K.; Verpoorte, R. Isolation of acetylcholinesterase inhibitory alkaloids from *Nerine bowdenii*. *Nat. Prod. Res.*, **2010**, *24*, 222-225.
- [35] Mulholland, D.A.; Crouch, N.; Decker, B.; Smith, M. T. The isolation of the Amaryllidaceae alkaloid crinamine from *Dioscorea dregeana*. *Biochem. Syst. Ecol.*, **2002**, *30*, 183-185.
- [36] Wang, Y.H.; Zhang, Z.K.; Yang, F.M.; Sun, Q.Y.; He, H.P.; Di, Y.T.; Mu, S.Z.; Lu, Y.; Chang, Y.; Zheng, Q.T.; Ding, M.; Dong, J.H.; Hao, X.J. Benzylphenethylamine Alkaloids from *Hosta plantaginea* with Inhibitory Activity against Tobacco Mosaic Virus and Acetylcholinesterase. *J. Nat. Prod.*, **2007**, *70*, 1458-1461.
- [37] Ma, X.; Gang, D.R. The Lycopodium alkaloids. *Nat. Prod. Rep.*, **2004**, *21*, 752-772.
- [38] Katakawa, K.; Nozoe, A.; Kogure, N.; Kitajima, M.; Hosokawa, M.; Takayama, H. Fawcettimine-related alkaloids from *Lycopodium serratum*. *J. Nat. Prod.*, **2007**, *70*, 1024-1028.
- [39] Takayama, H.; Katakawa, K.; Kitajima, M.; Yamaguchi, K.; Aimi, N. Ten new Lycopodium alkaloids having the lycopodane skeleton isolated from *Lycopodium serratum* Thunb. *Chem. Pharm. Bull. (Tokyo)*, **2003**, *51*, 1163-1169.
- [40] Choo, C.Y.; Hirasawa, Y.; Karimata, C.; Koyama, K.; Sekiguchi, M.; Kobayashi, J.; Morita, H. Carinatamins A-C, new alkaloids from *Lycopodium carinatum* inhibiting acetylcholinesterase. *Bioorg. Med. Chem.*, **2007**, *15*, 1703-1707.
- [41] Hirasawa, Y.; Kato, E.; Kobayashi, J.; Kawahara, N.; Goda, Y.; Shiro, M.; Morita, H. Lycoparins A-C, new alkaloids from *Lycopodium casuarinoides* inhibiting acetylcholinesterase. *Bioorg. Med. Chem.*, **2008**, *16*, 6167-6171.
- [42] Howes, M.-J.R.; Houghton, P.J. Acetylcholinesterase inhibitors of natural origin. *Int. J. Biomed. Pharm. Sci.*, **2009**, *3*, 67-86.
- [43] Devkota, K.P.; Lenta, B.N.; Fokou, P.A.; Sewald, N. Terpenoid alkaloids of the Buxaceae family with potential biological importance. *Nat. Prod. Rep.*, **2008**, *25*, 612-630.
- [44] Rahman, A.U.; Ul-Haq, Z.; Khalid, A.; Anjum, S.; Khan, M.R.; Choudhary, M.I. Pregnane-Type steroidal alkaloids of *Sarcococca saligna*: a new class of cholinesterase inhibitors. *Helv. Chim. Acta*, **2002**, *85*, 678-688.
- [45] Rahman, A.U.; Feroz, F.; Ul-Haq, Z.; Nawaz, S.A.; Khan, M.R.; Choudhary, M.I. New steroidal alkaloids from *Sarcococca saligna*. *Nat. Prod. Res.*, **2003**, *17*, 235-241.
- [46] Rahman, A.U.; Feroz, F.; Naeem, I.; Ul-Haq, Z.; Nawaz, S.A.; Khan, N.; Khan, M.R.; Choudhary, M.I. New pregnane-type

- steroidal alkaloids from *Sarcococca saligna* and their cholinesterase inhibitory activity. *Steroids*, **2004**, *69*, 735-741.
- [47] Devkota, K.P.; Lenta, B.N.; Choudhary, M.I.; Naz, Q.; Fekam, F.B.; Rosenthal, P.J.; Sewald, N. Cholinesterase inhibiting and antiplasmodial steroidal alkaloids from *Sarcococca hookeriana*. *Chem. Pharm. Bull. (Tokyo)*, **2007**, *55*, 1397-1401.
- [48] Devkota, K.P.; Lenta, B.N.; Wansi, J.; Choudhary, M.I.; Kisangau, D.P.; Naz, Q.; Samreen; Sewald, N. Bioactive 5 α -Pregnane-type steroidal alkaloids from *Sarcococca hookeriana*. *J. Nat. Prod.*, **2008**, *71*, 1481-1484.
- [49] Ata, A. In *Studies in Natural Products Chemistry*, 1st Edition, Vol. 38, Rahman, A.U., Ed.; Elsevier B.V.: Amsterdam, **2012**; pp. 225-245.
- [50] Ata, A.; Iverson, C.D.; Kalhari, K.S.; Akhter, S.; Betteridge, J.; Meshkatsadat, M.H.; Orhan, I.; Sener, B. Triterpenoidal alkaloids from *Buxus hyrcana* and their enzyme inhibitory, anti-fungal and anti-leishmanial activities. *Phytochemistry*, **2010**, *71*, 1780-1786.
- [51] Matochko, W.L.; James, A.; Lam, C.W.; Kozera, D.; Ata, A.; Gengan, R. Triterpenoidal alkaloids from *Buxus natalensis* and their acetylcholinesterase inhibitory activity. *J. Nat. Prod.*, **2010**, *73*, 1858-1862.
- [52] Rahman, A.U.; Akhtar, M.N.; Choudhary, M.I.; Tsuda, Y.; Sener, B.; Khalid, A.; Parvez, M. New steroidal alkaloids from *Fritillaria imperialis* and their cholinesterase inhibiting activities. *Chem. Pharm. Bull.*, **2002**, *50*, 1013-1016.
- [53] Lin, B.Q.; Ji, H.; Li, P.; Fang, W.; Jiang, Y. Inhibitors of acetylcholine esterase *in vitro* - Screening of steroidal alkaloids from *Fritillaria* species. *Planta Med.*, **2006**, *72*, 814-818.
- [54] Yang, Z.; Duan, D.Z.; Xue, W.W.; Yao, X.J.; Li, S. Steroidal alkaloids from *Holarrhena antidysenterica* as acetylcholinesterase inhibitors and the investigation for structure-activity relationships. *Life Sci.*, **2012**, *90*, 929-933.
- [55] Eldeen, I.M.; Van Heerden, F.R.; Van Staden, J. *In vitro* biological activities of niloticane, a new bioactive cassane diterpene from the bark of *Acacia nilotica* subsp. *kraussiana*. *J. Ethnopharmacol.*, **2010**, *128*, 555-560.
- [56] Hung, T.M.; Luan, T.C.; Vinh, B.T.; Cuong, T.D.; Min, B.S. Labdane-type diterpenoids from *Leonurus heterophyllus* and their cholinesterase inhibitory activity. *Phytother. Res.*, **2011**, *25*, 611-614.
- [57] Khan, I.; Nisar, M.; Khan, N.; Saeed, M.; Nadeem, S.; Fazal-ur-Rehman; Ali, F.; Karim, N.; Kaleem, W.A.; Qayum, M.; Ahmad, H.; Khan, I.A. Structural insights to investigate Conyopodioidol as a dual cholinesterase inhibitor from *Asparagus adscendens*. *Fitoterapia*, **2010**, *81*, 1020-1025.
- [58] Riaz, N.; Nawaz, S.A.; Mukhtar, N.; Malik, A.; Afza, N.; Ali, S.; Ullah, S.; Muhammad, P.; Choudhary, M.I. Isolation and enzyme-inhibition studies of the chemical constituents from *Ajuga bracteosa*. *Chem. Biodivers.*, **2007**, *4*, 72-83.
- [59] Ahmed, E.; Nawaz, S.A.; Malik, A.; Choudhary, M.I. Isolation and cholinesterase-inhibition studies of sterols from *Haloxylon recurvum*. *Bioorg. Med. Chem. Lett.*, **2006**, *16*, 573-580.
- [60] Awang, K.; Chan, G.; Litaudon, M.; Ismail, N.H.; Martin, M.T.; Gueritte, F. 4-Phenylcoumarins from *Mesua elegans* with acetylcholinesterase inhibitory activity. *Bioorg. Med. Chem.*, **2010**, *18*, 7873-7877.
- [61] Khan, M.T.; Orhan, I.; Senol, F.S.; Kartal, M.; Sener, B.; Dvorská, M.; Smejkal, K.; Slapetová, T. Cholinesterase inhibitory activities of some flavonoid derivatives and chosen xanthone and their molecular docking studies. *Chem. Biol. Interact.*, **2009**, *181*, 383-389.
- [62] Urbain, A.; Marston, A.; Sintro Grilo, L.; Bravo, J.; Purev, O.; Purevsuren, B.; Batsuren, D.; Reist, M.; Carrupt, P.-A.; Hostettmann, K. Xanthenes from *Gentianella amarella* ssp. *acuta* with acetylcholinesterase and monoamine oxidase inhibitory activities. *J. Nat. Prod.*, **2008**, *71*, 895-897.
- [63] Cho, J.K.; Ryu, Y.B.; Curtis-Long, M.J.; Ryu, H.W.; Yuk, H.J.; Kim, D.W.; Kim, H.J.; Lee, W.S.; Park, K.H. Cholinesterase inhibitory effects of geranylated flavonoids from *Paulownia tormentosa* fruits. *Bioorg. Med. Chem.*, **2012**, *25*, 2595-2602.
- [64] Jung, H.A.; Jin, S.E.; Park, J.S.; Choi, J.S. Antidiabetic complications and anti-alzheimer activities of sophoflavescenol, a prenylated flavonol from *Sophora flavescens*, and its structure-activity relationship. *Phytother. Res.*, **2011**, *25*, 709-715.
- [65] Kim, J.Y.; Lee, W.S.; Kim, Y.S.; Curtis-Long, M.J.; Lee, B.W.; Ryu, Y.B.; Park, K.H. Isolation of cholinesterase-inhibiting flavonoids from *Morus thou*. *J. Agric. Food Chem.*, **2011**, *59*, 4589-4596.
- [66] Ryu, H.W.; Curtis-Long, M.J.; Jung, S.; Jeong, I.Y.; Kim, D.S.; Kang, K.Y.; Park, K.H. Anticholinesterase potential of flavonols from paper mulberry (*Broussonetia papyrifera*) and their kinetic studies. *Food Chem.*, **2012**, *132*, 1244-1250.
- [67] Conforti, F.; Rigano, D.; Menichini, F.; Loizzo, M.R.; Senatore, F. Protection against neurodegenerative diseases of *Iris pseudopumila* extracts and their constituents. *Fitoterapia*, **2009**, *80*, 62-67.
- [68] Ahmad, V.U.; Iqbal, S.; Nawaz, S.A.; Choudhary, M.I.; Farooq, U.; Ali, S.T.; Ahmad, A.; Bader, S.; Kousar, F.; Arshad, S.; Tareen, R.B. Isolation of four new pterocarpanes from *Zygophyllum eurypterum* (syn. *Z. atriplicoides*) with enzyme-inhibition properties. *Chem. Biodivers.*, **2006**, *3*, 996-1003.
- [69] Changwong, N.; Sabphon, C.; Ingkaninan, K.; Sawasdee, P. Acetyl- and butyryl-cholinesterase inhibitory activities of Mansorins and Mansonones. *Phytother. Res.*, **2012**, *26*, 392-396.
- [70] Nag, G.; De, B. Acetylcholinesterase inhibitory activity of *Terminalia chebula*, *Terminalia bellerica* and *Embllica officinalis* and some phenolic compounds. *Int. J. Pharm. Pharm. Sci.*, **2011**, *3*, 121-124.
- [71] Ge, H.M.; Zhu, C.H.; Shi, D.H.; Zhang, L.D.; Xie, D.Q.; Yang, J.; Ng, S.W.; Tan, R.X. Hopeahainol A: An acetylcholinesterase inhibitor from *Hopea hainanensis*. *Chemistry*, **2008**, *14*, 376-381.
- [72] Sancheti, S.; Um, B.H.; Seo, S.H. 1,2,3,4,6-penta-O-galloyl- β -D-glucose: A cholinesterase inhibitor from *Terminalia chebula*. *S. Afr. J. Bot.*, **2010**, *76*, 285-288.
- [73] Akhtar, M.N.; Lam, K.W.; Abas, F.; Maulidiani, H.; Ahmad, S.; Shah, S.A.; Rahman, A.U.; Choudhary, M.I.; Lajis, N.H. New class of acetylcholinesterase inhibitors from the stem bark of *Knema laurina* and their structural insights. *Bioorg. Med. Chem. Lett.*, **2011**, *21*, 4097-4103.
- [74] Adhami, H. R.; Linder, T.; Kaehlig, H.; Schuster D.; Zehl, M.; Krenn, L. Cathechol alkenyls from *Semecarpus anacardium*: Acetylcholinesterase inhibition and binding mode predictions. *J. Ethnopharmacol.* **2012**, *139*, 142-148.
- [75] Jung, H.A.; Jung, Y.J.; Hyun, S.K.; Min, B.S.; Kim, D.W.; Jung, J.H.; Choi, J.S. Selective cholinesterase inhibitory activities of a new monoterpene diglycoside and other constituents from *Nelumbo nucifera* stamens. *Biol. Pharm. Bull.*, **2010**, *33*, 267-272.
- [76] Adewusi, E.A.; Moodley, N.; Steenkamp, V. Medicinal plants with cholinesterase inhibitory activity: a review. *Afr. J. Biotechnol.*, **2010**, *9*, 8257-8276.
- [77] Barbosa Filho, J.M.; Medeiros, K.C.P.; Diniz, M.F.; Batista, L.M.; Athayde-Filho, P.F.; Silva, M.S.; da-Cunha, E.V.L.; Silva Almeida, J.R.G.; Quintans-Júnior, L.J. Natural products inhibitors of the enzyme acetylcholinesterase. *Braz. J. Pharmacogn.*, **2006**, *16*, 258-285.
- [78] Mukherjee, P.K.; Kumar, V.; Houghton, P.J. Screening of Indian medicinal plants for acetylcholinesterase inhibitory activity. *Phytother. Res.*, **2007**, *21*, 1142-1145.
- [79] Tundis, R.; Menichini, F.; Conforti, F.; Loizzo, M.R.; Bonesi, M.; Statti, G.; Menichini, F. A potential role of alkaloid extracts from *Salsola* species (Chenopodiaceae) in the treatment of Alzheimer's disease. *J. Enzyme Inhib. Med. Chem.*, **2009**, *24*, 818-824.
- [80] Elufioye, T.O.; Obuoto, E.M.; Sennuga, A.T.; Agbedahunsi, J.M.; Adesanya, S.A. Acetylcholinesterase and butyrylcholinesterase inhibitory activity of some selected Nigerian medicinal plants. *Rev. Bras. Farmacogn.*, **2010**. DOI: 10.1590/S0102-695X2010000400002.
- [81] Fawole, O.A.; Amoo, S.O.; Ndhkala, A.R.; Light, M.E.; Finnie, J.F.; Van Staden, J. Anti-inflammatory, anticholinesterase, antioxidant and phytochemical properties of medicinal plants used for pain-related ailments in South Africa. *J. Ethnopharmacol.*, **2010**, *127*, 235-241.
- [82] Cahlíková, L.; Benešová, N.; Macáková, K.; Urbanová, K.; Opletal, L. GC/MS analysis of three Amaryllidaceae species and their cholinesterase activity. *Nat. Prod. Commun.*, **2011**, *6*, 1255-1258.
- [83] Cahlíková, L.; Valterová, I.; Macáková, K.; Opletal, L. Analysis of Amaryllidaceae alkaloids from *Zephyranthes grandiflora* by GC/MS and their cholinesterase activity. *Rev. Bras. Farmacogn.*, **2011**. DOI: 10.1590/S0102-695X2011005000089.

- [84] Moyo, M.; Ndhkala, A.R.; Finnie, J.F.; Van Staden, J. Phenolic composition, antioxidant and acetylcholinesterase inhibitory activities of *Sclerocarya birrea* and *Harpephyllum caffrum* (Anacardiaceae) extracts. *Food Chem.*, **2010**, *123*, 69-76.
- [85] Benamar, H.; Rached, W.; Derdour, A.; Marouf, A. Screening of Algerian medicinal plants for acetylcholinesterase inhibitory activity. *J. Biol. Sci.*, **2010**, *10*, 1-9.
- [86] Lima, J.A.; Costa, R.S.; Epifânio, R.A.; Castro, N.G.; Rocha, M.S.; Pinto, A.C. *Geissospermum vellosii* stem bark. Anticholinesterase activity and improvement of scopolamine-induced memory deficits. *Pharmacol. Biochem. Be.*, **2009**, *92*, 508-513.
- [87] Yang, Z.; Zhang, D.; Ren, J.; Yang, M.; Li, S. Acetylcholinesterase inhibitory activity of the total alkaloid from traditional Chinese herbal medicine for treating Alzheimer's disease. *Med. Chem. Res.*, **2012**, *21*, 734-738.
- [88] Sekeroglu, N.; Senol, F.S.; Orhan, I.E.; Gulpinar, A.R.; Kartal, M.; Sener, B. *In vitro* prospective effects of various traditional herbal coffees consumed in Anatolia linked to neurodegeneration. *Food Res. Int.*, **2012**, *45*, 197-203.
- [89] Loizzo, M.R.; Tundis, R.; Menichini, F.; Bonesi, M.; Frega, N.G.; Menichini, F. Radical scavenging activity and cholinesterase inhibitory activity of *Leopoldia comosa* (L.) bulbs. *Progr. Nutr.*, **2011**, *13*, 300-303.
- [90] Carpinella, M.C.; Androne, D.G.; Ruiz, G.; Palacios, S.M. Screening for acetylcholinesterase inhibitory activity in plant extracts from Argentina. *Phytother. Res.*, **2010**, *24*, 259-263.
- [91] Wszelaki, N.; Kuciun, A.; Kiss, A.K. Screening of traditional European herbal medicines for acetylcholinesterase and butyrylcholinesterase inhibitory activity. *Acta Pharm.*, **2010**, *60*, 119-128.
- [92] Niño, J.; Hernández, J.A.; Correa, Y.M.; Mosquera, O.M. *In vitro* inhibition of acetylcholinesterase by crude plant extracts from Colombian flora. *Mem. Inst. Oswaldo Cruz.*, **2006**, *101*, 783-785.
- [93] Bakthir, H.; Ali, N.A.A.; Arnold, N.; Teichert, A.; Wessjohann, L. Anticholinesterase activity of endemic plant extracts from Soqotra. *Afr. J. Tradit. Complement. Altern. Med.*, **2011**, *8*, 296-299.
- [94] Habtemariam, S. The therapeutic potential of *Berberis darwinii* stem-bark: Quantification of berberine and *in vitro* evidence for Alzheimer's disease therapy. *Nat. Prod. Commun.*, **2011**, *6*, 1089-1090.
- [95] Ashraf, M.; Ahmad, K.; Ahmad, I.; Ahmad, S.; Arshad, S.; Shah, S.M.; Nasim, F. Acetylcholinesterase and NADH oxidase inhibitory activity of some medicinal plants. *J. Med. Plant. Res.*, **2011**, *5*, 2086-2089.
- [96] Adewusi, E.A.; Moodley, N.; Steenkamp, V. Antioxidant and acetylcholinesterase inhibitory activity of selected southern African medicinal plants. *S. Afr. J. Bot.*, **2011**, *77*, 638-644.
- [97] Akkol, E.K.; Orhan, I.E.; Yeşilada, E. Anticholinesterase and antioxidant effects of the ethanol extract, ethanol fractions and isolated flavonoids from *Cistus laurifolius* L. leaves. *Food Chem.*, **2012**, *131*, 626-631.
- [98] Feitosa, C.M.; Freitas, R.M.; Luz, N.N.N.; Bezerra, M.Z.B.; Trevisan, M.T.S. Acetylcholinesterase inhibition by some promising Brazilian medicinal plants. *Braz. J. Biol.*, **2011**, *71*, 783-789.
- [99] Tavares, L.; McDougall, G.J.; Fortalezas, S.; Stewart, D.; Ferreira, R.B.; Santos, C.N. The neuroprotective potential of phenolic-enriched fractions from four *Juniperus* species found in Portugal. *Food Chem.*, **2012**, *135*, 562-570.
- [100] Lee, S.-H.; Sancheti, S.A.; Bafna, M.R.; Sancheti, S.S.; Seo, S.-Y. Acetylcholinesterase inhibitory and antioxidant properties of *Rhododendron yedoense* var. *poukhanense* bark. *J. Med. Plants Res.*, **2011**, *5*, 248-254.
- [101] Kwon, S.-H.; Lee, H.-K.; Kim, J.A.; Hong, S.I.; Kim, S.-Y.; Jo, T.-H.; Park, Y.-I.; Lee, C.-K.; Kim, Y.-B.; Lee, S.-Y.; Jang, C.-G. Neuroprotective effects of *Eucommia ulmoides* Oliv. bark on amyloid beta₂₅₋₃₅-induced learning and memory impairments in mice. *Neurosci. Lett.*, **2011**, *487*, 123-127.
- [102] Crowch, C.M.; Okello, E.J. Kinetics of acetylcholinesterase inhibitory activities by aqueous extracts of *Acacia nilotica* (L.) and *Rhamnus prinoides* (L'Hér.). *Afr. J. Pharm. Pharmacol.*, **2009**, *3*, 469-475.
- [103] Kim, D.H.; Yoon, B.H.; Kim, Y.W.; Lee, S.; Shin, B.Y.; Jung, J.W.; Kim, H.J.; Lee, Y.S.; Choi, J.S.; Kim, S.Y.; Lee, K.T.; Ryu, J.H. The seed extract of *Cassia obtusifolia* ameliorates learning and memory impairments induced by scopolamine or transient cerebral hypoperfusion in mice. *J. Pharmacol. Sci.*, **2007**, *105*, 82-93.
- [104] Lin, H.Q.; Ho, M.T.; Lau, L.S.; Wong, K.K.; Shaw, P.C.; Wan, D.C. Anti-acetylcholinesterase activities of traditional Chinese medicine for treating Alzheimer's disease. *Chem. Biol. Interact.*, **2008**, *175*, 352-354.
- [105] Rauter, A.P.; Martins, A.; Lopes, R.; Ferreira, J.; Serralheiro, L.M.; Araújo, M.E.; Borges, C.; Justino, J.; Silva, F.V.; Goulart, M.; Thomas-Oates, J.; Rodrigues, J.A.; Edwards, E.; Noronha, J.P.; Pinto, R.; Mota-Filipe, H. Bioactivity studies and chemical profile of the antidiabetic plant *Genista tenera*. *J. Ethnopharmacol.*, **2009**, *122*, 384-393.
- [106] Satheeshkumar, N.; Mukherjee, P.K.; Bhadra, S.; Saha, B.P. Acetylcholinesterase enzyme inhibitory potential of standardized extract of *Trigonella foenum graecum* L and its constituents. *Phytomedicine*, **2010**, *17*, 292-295.
- [107] Bhadra, S.; Mukherjee, P.K.; Kumar, N.S.; Bandyopadhyay, A. Anticholinesterase activity of standardized extract of *Illicium verum* Hook. f. fruits. *Fitoterapia*, **2011**, *82*, 342-346.
- [108] Orhan, I.; Senol, F.S.; Gulpinar, A.R.; Kartal, M.; Sekeroglu, N.; Deveci, M.; Kan, Y.; Sener, B. Acetylcholinesterase inhibitory and antioxidant properties of *Cyclotrichium niveum*, *Thymus praecox* subsp. *caucasicus* var. *caucasicus*, *Echinacea purpurea* and *E. pallida*. *Food Chem. Toxicol.*, **2009**, *47*, 1304-1310.
- [109] Costa, P.; Gonçalves, S.; Andrade, P.B.; Valentão, P.; Romano, A. Inhibitory effect of *Lavandula viridis* on Fe²⁺-induced lipid peroxidation, antioxidant and anti-cholinesterase properties. *Food Chem.*, **2011**, *126*, 1779-1786.
- [110] Orhan, I.E.; Belhatab, R.; Şenol, F.S.; Gulpinar, A.R.; Hoşbaş, S.; Kartal, S. Profiling of cholinesterase inhibitory and antioxidant activities of *Artemisia absinthium*, *A. herba-alba*, *A. fragrans*, *Marrubium vulgare*, *M. astranicum*, *Origanum vulgare* subsp. *glandulosum* and essential oil analysis of two *Artemisia* species. *Ind. Crop. Prod.*, **2010**, *32*, 566-571.
- [111] Loizzo, M.R.; Menichini, F.; Conforti, F.; Tundis, R.; Bonesi, M.; Saab, A.M.; Statti, G.A.; Cindio, B.; Houghton, P.J.; Menichini, F.; Frega, N.G. Chemical analysis, antioxidant, antiinflammatory and anticholinesterase activities of *Origanum ehrenbergii* Boiss and *Origanum syriacum* L. essential oils. *Food Chem.*, **2009**, *117*, 174-180.
- [112] Mossa, A.T.; Nawwar, G.A. Free radical scavenging and antiacetylcholinesterase activities of *Origanum majorana* L. essential oil. *Hum. Exp. Toxicol.*, **2011**, *30*, 1501-1513.
- [113] Tel, G.; Oztürk, M.; Duru, M.E.; Harmandar, M.; Topçu, G. Chemical composition of the essential oil and hexane extract of *Salvia chionantha* and their antioxidant and anticholinesterase activities. *Food Chem. Toxicol.*, **2010**, *48*, 3189-3193.
- [114] Şenol, F.S.; Orhan I.; Celep F.; Kahraman, A.; Doğan, M.; Yilmaz, G.; Şener, B. Survey of 55 Turkish *Salvia* taxa for their acetylcholinesterase inhibitory and antioxidant activities. *Food Chem.*, **2010**, *120*, 34-43.
- [115] Loizzo, M.R.; Menichini, F.; Tundis, R.; Bonesi, M., Conforti, F.; Nadjafi, F.; Statti, G.A.; Frega, N.G.; Menichini, F. *In vitro* biological activity of *Salvia leriifolia* benth essential oil relevant to the treatment of Alzheimer's disease. *J. Oleo Sci.*, **2009**, *58*, 443-446.
- [116] Ahmad, B.; Mukarram Shah, S.M.; Khan, H.; Hassan Shah, S.M. Enzyme inhibition activities of *Teucrium royleanum*. *J. Enzym. Inhib. Med. Ch.*, **2007**, *22*, 730-732.
- [117] Tappayuthpijarn, P.; Itharat, A.; Makchuchit, S. Acetylcholinesterase inhibitory activity of Thai traditional nootropic remedy and its herbal ingredients. *J. Med. Assoc. Thai.*, **2011**, *94*, S183-S189.
- [118] Vinutha, B.; Prashanth, D.; Salma, K.; Sreeja, S.L.; Pratiti, D.; Padmaja, R.; Radhika, S.; Amit, A.; Venkateshwarlu, K.; Deepak, M. Screening of selected Indian medicinal plants for acetylcholinesterase inhibitory activity. *J. Ethnopharmacol.*, **2007**, *109*, 359-363.
- [119] Ashraf, M.; Ahmad, K.; Ahmad, I.; Ahmad, S.; Arshad, S.; Shah, S.M.; Nasim, F. Acetylcholinesterase and NADH oxidase inhibitory activity of some medicinal plants. *J. Med. Plant. Res.*, **2011**, *5*, 2086-2089.
- [120] Adersen, A.; Gauquin, B.; Gudiksen, L.; Jäger, A.K. Screening of plants used in Danish folk medicine to treat memory dysfunction

- for acetylcholinesterase inhibitory activity. *J. Ethnopharmacol.*, **2006**, *104*, 418-422.
- [121] Ustun, O.; Senol, F.; Kurkcuoglu, M.; Orhan, I.; Kartal, M.; Baser, K. Investigation on chemical composition, anticholinesterase and antioxidant activities of extracts and essential oils of Turkish *Pinus* species and pycnogenol. *Ind. Crop. Prod.*, **2012**, *38*, 115-123.
- [122] Bonesi, M.; Menichini, F.; Tundis, R.; Loizzo, M.R.; Conforti, F.; Passalacqua, N.G.; Statti, G.A.; Menichini, F. Acetylcholinesterase and butyrylcholinesterase inhibitory activity of *Pinus* species essential oils and their constituents. *J. Enzym. Inhib. Med. Ch.*, **2010**, *25*, 622-628.
- [123] Khadri, A.; Serralheiro, M.L.M.; Nogueira, J.M.F.; Neffati, M.; Smiti, S.; Araújo, M.E.M. Antioxidant and antiacetylcholinesterase activities of essential oils from *Cymbopogon schoenanthus* L. Spreng. Determination of chemical composition by GC-mass spectrometry and ¹³C NMR. *Food Chem.*, **2008**, *109*, 630-637.
- [124] Khadri, A.; Neffati, M.; Smiti, S.; Falé, P.; Lino, A.R.L.; Serralheiro, A.M.; Araújo, M.E.M. Antioxidant, antiacetylcholinesterase and antimicrobial activities of *Cymbopogon schoenanthus* L. Spreng (lemon grass) from Tunisia. *LWT-Food Sci. Technol.*, **2010**, *43*, 331-336.
- [125] Aremu, A.O.; Amoo, S.O.; Ndhkala, A.R.; Finnie, J.F.; Stadenm J.V. Antioxidant activity, acetylcholinesterase inhibition, iridoid content and mutagenic evaluation of *Leucosidea sericea*. *Food Chem. Toxicol.*, **2011**, *49*, 1122-1128.
- [126] Pachauri, S.D.; Tota, S.; Khandelwal, K.; Verma, P.R.; Nath, C.; Hanif, K.; Shukla, R.; Saxena, J.K.; Dwivedi, A.K. Protective effect of fruits of *Morinda citrifolia* L. on scopolamine induced memory impairment in mice: A behavioral, biochemical and cerebral blood flow study. *J. Ethnopharmacol.*, **2012**, *139*, 34-41.
- [127] Chaiyana, W.; Okonogi, S. Inhibition of cholinesterase by essential oil from food plant. *Phytomedicine*, **2012**, *19*, 836-839.
- [128] Menichini, F.; Tundis, R.; Bonesi, M.; de Cindio, B.; Loizzo, M.R.; Conforti, F.; Statti, G.A.; Menabeni, R.; Bettini, R.; Menichini, F. Chemical composition and bioactivity of *Citrus medica* L. cv. Diamante essential oil obtained by hydrodistillation, cold-pressing and supercritical carbon dioxide extraction. *Nat. Prod. Res.*, **2011**, *25*, 789-799.

Received: October 02, 2012

Revised: February 25, 2013

Accepted: February 25, 2013